

***Healthy Heart, Healthy Family:* A NHLBI/HRSA Collaborative Employing Community Health Workers to Improve Heart Health**

Ritabelle Fernandes, MD, MPH
Kathryn L. Braun, DrPH
Jovonni R. Spinner, MPH, CHES
Cynthia Sturdevant, MD, MPH
Sharonne J. Ancheta, BSN
Sheryl R. Yoshimura, RD
Merlita Compton, MPH
Jo-Hsi Wang, BS
Carolyn J. Lee, MBA

Abstract: Kokua Kalihi Valley, a federally qualified health center in Hawaii, collaborated with the National Heart, Lung, and Blood Institute to test the efficacy of community health workers (CHWs) to deliver the *Healthy Heart, Healthy Family* curriculum to low-income Filipinos with cardiovascular disease (CVD) risk factors. At 12 months, significant improvements were seen in health behaviors, knowledge, and self-efficacy in managing chronic diseases. We also observed decreases in total cholesterol from 186.25 mg/dl to 170.88 mg/dl ($p=.001$), low-density lipoprotein from 114.43 mg/dl to 103.04 mg/dl ($p=.013$), and fasting blood glucose from 117.95 mg/dl to 109.07 mg/dl ($p=.034$). Although these changes were statistically significant, they are small and not clinically meaningful in reducing CVD risk. The high-density lipoprotein was 3.3 mg/dl lower (worse) at 12 months ($p=.003$), mean values for blood pressure, BMI, and waist circumference increased. Community health workers can be trained to deliver evidence-based curricula that improve health behaviors and increase self-efficacy in managing chronic diseases.

Key words: Asian Americans, cardiovascular disease, cardiovascular risk factors, community health workers, cultural competency, Filipino Americans, health education.

RITABELLE FERNANDES, CYNTHIA STURDEVANT, SHARONNE J. ANCHETA, SHERYL R. YOSHIMURA, MERLITA COMPTON, JO-HSI WANG, and CAROLYN J. LEE are affiliated with the Kokua Kalihi Valley Comprehensive Family Services in Honolulu; **KATHRYN L. BRAUN** is with the Office of Public Health Studies at the John A. Burns School of Medicine, University of Hawaii; and **JOVONNI R. SPINNER** is with the Division of Application of Research Discoveries, National Heart, Lung, and Blood Institute, National Institutes of Health in Bethesda, Maryland. Please address correspondence to Ritabelle Fernandes, MD, MPH; Kokua Kalihi Valley; 2239 N. School St.; Honolulu, HI 96819; (808) 791-9416; rfernandes@kkv.net.

Cardiovascular disease (CVD) is the leading cause of death among Filipino Americans, and CVD mortality rates are higher among Filipino Americans than among other ethnic groups.¹ In addition to high prevalence of CVD, Filipino Americans are more likely to have hypertension than other Asian American groups.² For Filipino Americans who are also facing poverty, limited English language and cultural barriers make preventative care and treatment for CVD especially challenging.

Federally qualified health centers (FQHC) provide preventive and primary care to the uninsured and underserved, with the mission to reduce health disparities and improve outcomes.³ Kokua Kalihi Valley Comprehensive Family Services (KKV) is a FQHC in Kalihi Valley, Honolulu, a health professional shortage area.⁴ This neighborhood has attracted large populations of immigrants from the Philippines who come to the United States (U.S.) for family reunification, better health care, work, and educational opportunities. More than 30% of KKV's patients are low-income Filipinos. Many arrive with chronic conditions that have progressed to advanced stages, and arrive uninsured and unable to navigate the U.S. health care system.

Community health workers (CHW) are important in FQHCs, providing assistance to people with low health literacy and limited English proficiency. They are typically from the community they serve, may be bilingual, and have some health or social service training.⁵ Their educational backgrounds vary, from high school graduates to college degrees. At KKV, CHWs are salaried employees occupying a wide range of positions, including outreach worker, case manager, medical assistant, and educator. Community health workers have various functions that include improving access to care, health education, patient navigation, case management, interpretation, social support, community advocacy, referrals, and linking families to community resources. The use of CHWs as extenders of the chronic care disease management team is essential in helping reduce cultural barriers and improving care coordination for low-income, immigrant clients.⁶

Healthy Heart, Healthy Family is a curriculum to prevent and control CVD specifically created for Filipino Americans by the National Heart, Lung, and Blood Institute (NHLBI) at the National Institutes of Health,⁷ adapted from existing NHLBI community group education curricula. The curriculum is designed to be taught by CHWs to small groups of 8–12 people. Eleven educational lessons (Box 1) aim to help participants build skills to make practical, lasting changes to help fight CVD and improve health. Trainers use a facilitative teaching style to keep participants motivated and to promote adult learning. Features include heart-healthy bingo and other interactive activities, cultural adaptations, Tagalog translations, use of picture cards, and six easy-to-read bilingual booklets on heart-healthy living.

The objective of this 12-month study was to evaluate the impact of the CHW-delivered *Healthy Heart, Healthy Family* curriculum on low-income Filipinos with CVD risk factors in Hawaii. Evaluation measures included 1) clinical outcomes; 2) knowledge of healthy habits; 3) satisfaction with curriculum; 4) quality of life; and 5) self-efficacy.

Box 1.**HEALTHY HEART, HEALTHY FAMILY
EDUCATIONAL LESSONS**

1. Knowledge is power: Know your risk for heart disease
2. Act in time to heart attack signs
3. Get energized! Say YES to physical activity
4. Help your heart: Control your high blood pressure
5. Be heart smart: Keep your cholesterol in check
6. Keep your heart in mind: Aim for a healthy weight
7. Protect your heart: Prevent and control diabetes
8. Welcome heart healthy eating into your home
9. Eat in a heart healthy way—Even when time or money is tight
10. Take control of your health: Enjoy living smoke free
11. Review and graduation

Methods

Study design. A one-group pre-post-test, design was used. Institutional review board approval was obtained from the University of Hawaii. A total of 99 participants provided written consent for participation and review of their medical records.

Participant recruitment. Recruitment strategies included informational flyers and posters, direct invitation by KKV physicians, and community outreach. From June 2008 to July 2009, 99 Filipinos adults with CVD risk factors (such as hypertension, diabetes, high cholesterol, obesity, family history, smoking, and sedentary lifestyle) were recruited.

CHW recruitment and training. Three CHWs from KKV attended a training sponsored by NHLBI and the Health Resources and Services Administration to learn how to lead the program, and later trained eight other CHWs. A total of 11 CHWs participated in this study. All were fluent in English, Tagalog, and/or Ilokano, the two Filipino languages spoken in Hawaii. Five occupied social service positions, four were medical assistants, and two were educators. As part of their job description at KKV, CHWs perform home visits and outreach to a panel of 20–25 clients, so have a good understanding of the context within which behavior change needs to take place.

Activities. The curriculum was delivered to six small groups of participants, each with two CHW facilitators. Sessions, each two hours in duration, were held weekly for 11 consecutive weeks. Incentives (water bottles and pedometers) and healthy snacks were provided. Mean attendance was seven sessions, with 65% of participants attending nine or more sessions. To encourage participants to adopt recommendations for exercise, physical activity (e.g., twice weekly Chairobics) and community gardening programs were offered and led by CHWs. After the 11 weeks, CHWs organized monthly activities promoting heart-healthy behaviors around themes such as Heart Day, Family Day, and Mother's Day.

Measures. Most measures were provided with the curriculum. Demographic variables included age, gender, education, marital status, primary language, time in U.S., poverty level, insurance, and housing. Participation in sessions, exercise, and monthly activities was tracked. Data to assess effectiveness were collected at baseline, 6 months, and 12 months. Disease and medication histories were taken, and clinical measures included weight, height, body mass index (BMI), waist circumference, blood pressure, fasting blood glucose, fasting lipid profile, and glycosylated hemoglobin. A pre-post health habits survey was used. The 25 food consumption questions (scored on a 4-point scale from 1=never to 4=most of the time) asked about sodium consumption, fat consumption, and weight management. Nine items on physical activity, three on smoking, and four on alcohol use were scored yes/no or ordinally. Thirteen knowledge items were scored *yes/no/don't know*. The final item was a story about a family whose members were at different stages of readiness to adopt a healthy lifestyle, based on the Transtheoretical Model of Behavior Change.⁸

We supplemented the NHLBI forms with two other validated tools. The six-item Stanford Self-efficacy Scale for chronic diseases asked about the level of confidence a participant had in managing his/her disease and keeping symptoms from interfering with daily activities. Response options were on a Likert scale from 1=*not at all confident* to 10=*totally confident*.⁹ The 16-item Flanagan Quality of Life Scale (QOLS) asked participants to rate their life in relation to five conceptual domains: material and physical well-being; relationships; social, community, and civic activities; personal development; and fulfillment and recreation.¹⁰ It is scored on a Likert scale from 1=*terrible* to 7=*delighted*. The QOLS, created originally by American psychologist John Flanagan in the 1970s, has been adapted for use in chronic illness groups. It is a valid instrument for measuring quality of life across patient groups and cultures and is conceptually distinct from health status or other causal indicators of quality of life.¹¹

Analysis. Data were analyzed using SPSS.¹² To assess changes in knowledge, behavioral, and clinical measures, we compared baseline and 12-month values using paired t-tests for continuous variables and chi-squared tests for categorical variables. Clinical data were adjusted for age and gender based on the 2000 Census.

Results

The mean age of the 99 participants enrolled in the study was 68.5 years, with a median of 69 years and a range of 25–85 years; 16.2% were male and 83.8% were female (Table 1). Seventy percent were at or below 100% of federal poverty level. Most participants lived with extended family, and only 3% lived in public housing. Nearly half had limited education, of eighth-grade or less. The sample had a mean time in the U.S. of 15.9 years, and 77% were from the Northern Philippines. Tagalog was the first language for about 8%, and Ilokano was the first language of most others. Participants had many risk factors for CVD; for example, 81.5% had high cholesterol, 81.5% had hypertension, 34.8% had diabetes, 32.6% had family history of CVD, 40.2% were overweight, and 15.2% were obese.

Between baseline and the 12-month follow up, seven participants were lost to follow up. Thus, analysis of over-time changes was limited to 92 individuals for whom we had

Table 1.
BASELINE CHARACTERISTICS OF INITIAL SAMPLE, DROPOUTS, AND FINAL SAMPLE

Characteristic		Initial sample N=99	Dropouts N=7	Final sample N=92
Gender	Male	16 (16.2%)	1 (14.3%)	15 (16.3%)
	Female	83 (83.8%)	6 (85.7%)	77 (83.7%)
Public housing resident	No	96 (97.0%)	6 (85.7%)	90 (97.8%)
Poverty level (federal)	<100%	70 (70.7%)	5 (71.4%)	65 (70.7%)
	100–199%	17 (17.2%)	1 (14.3%)	16 (16.2%)
	200 or higher	10 (10.1%)	1 (14.3%)	9 (9.1%)
Education	8th grade or less	48 (48.5%)	1 (14.3%)	47 (51.1%)
	High school graduate	24 (24.3%)	1 (14.3%)	21 (22.8%)
	1–3 years of college	14 (14.1%)	3 (42.9%)	11 (12.0%)
	College graduate	13 (13.1%)	0	13 (14.1%)
Primary language	Tagalog	8 (8.1%)	1 (14.3%)	7 (7.6%)
	Ilokano	81 (81.8%)	6 (85.7%)	75 (81.5%)
	Both	6 (6.1%)	0	6 (6.5%)
	Other	4 (4.0%)	0	4 (4.3%)
Cardio Vascular Disease risk factors	High cholesterol	81 (81.8%)	6 (85.7%)	75 (81.5%)
	Hypertension	80 (80.8%)	6 (85.7%)	74 (81.5%)
	Diabetes	34 (34.3%)	2 (28.6%)	32 (34.8%)
	Family history	32 (32.3%)	2 (28.6%)	30 (32.6%)
	Overweight	39 (39.4%)	2 (28.6%)	37 (40.2%)
	Obese	15 (15.3%)	1 (14.3%)	14 (15.2%)
	Sedentary	14 (14.1%)	1 (14.3%)	13 (14.1%)
	Smoking	5 (5.0%)	0	5 (5.4%)

complete data. There were no significant differences in demographic characteristics of the full (n=99) and follow-up (n=92) groups. The seven participants who dropped out shared central characteristics (six were female and the majority spoke Ilokano, and had many risk factors for CVD (see Table 2).

Participants showed small improvements on three of the clinical measures, which were adjusted for age and gender based on the 2000 U.S. Census (Table 2 and Figure 1). Mean total cholesterol decreased from 186.25 mg/dl at baseline to 170.88 mg/dl at 12 months (p=.001); mean low-density lipoprotein decreased from 114.43 mg/dl to 103.04 mg/dl (p=.013); and mean fasting blood glucose decreased from 117.95 mg/dl to 109.07 mg/dl, (p=.034). However, the high-density lipoprotein (HDL) was 3.3 mg/dl lower (worse) at 12 months (p=.003). Although these changes were statistically significant, they are small and not clinically meaningful in reduction of CVD risk.

Table 2.**AGE-ADJUSTED CLINICAL MEASURES
AT BASELINE AND 12 MONTHS (N=92)**

Variable	Baseline	12 Months	p-value	Standard error
Total Cholesterol (mg/dl)	186.25	170.88	.001	4.399
Low density lipoprotein (mg/dl)	114.43	103.04	.013	4.401
High density lipoprotein (mg/dl)	44.29	40.99	.003	1.069
Triglycerides (mg/dl)	139.86	136.51	.610	6.541
Systolic BP (mmHg)	122.60	124.87	.376	2.471
Diastolic BP (mmHg)	74.24	74.29	.922	1.208
Hemoglobin A1C (%)	6.65	6.61	.317	0.045
Body Mass Index	28.56	30.73	.457	2.898
Waist circumference (cms)	36.23	36.63	.182	0.298
Fasting blood glucose (mg/dl)	117.95	109.07	.034	4.080

Additionally, the mean value for hemoglobin A1C (%) did not change. At 12 months, mean values for BMI and waist circumference had increased, but increases were not statistically significant.

Blood pressure reductions were seen at six months, but were not sustained at 12 months. The mean systolic blood pressure decreased from 122.60 mmHg at baseline to 116.61 mmHg at six months ($p=.001$), and mean diastolic blood pressure decreased from 74.24 mmHg at baseline to 71.35 mmHg at six months ($p=.03$). Between baseline and six months, there was also a significant decrease ($p=.043$) in the proportion of participants who were categorized as either being normotensive, pre-hypertensive, or as Stage 1 and 2 hypertensive (Figure 2). These improvements were not sustained, as mean blood pressure readings at 12 months were very similar to those at baseline (124.87 mmHg and 74.29 mmHg, respectively).

Changes in self-efficacy scores are shown in Table 3. Statistically significant improvements were seen in participants' confidence in managing fatigue ($p=.005$), emotional distress ($p=.01$), and keeping CVD from affecting everyday life ($p=.013$). Improvements were seen in confidence to manage physical discomfort and reduce doctor visits, but these were not statistically significant. There were no significant changes in the quality of life of the participants except for in material comforts.

Findings from the health habits questionnaire revealed significant improvements on nine of the 25 food consumption items. For example, compared with baseline, significantly greater proportions of participants were reading nutrition labels, eating fruits for dessert, snacking on vegetables and fruits, replacing salt with herbs and spices, using monounsaturated and polyunsaturated cooking oils, and baking instead of frying at 12 months (Table 4). Additionally, the number of participants performing physical activity in the form of aerobic exercise increased from 16.7% at baseline to




Figure 1. Clinical measures at baseline and 12 months (n= 92).

- TC = Total cholesterol
- LDL = Low-density lipoprotein
- HDL = High-density lipoprotein
- SBP = Systolic blood pressure
- DBP = Diastolic blood pressure
- HbA1c = Glycosylated hemoglobin
- BMI = Body mass index
- WC = Waist circumference
- FBG = Fasting blood glucose

43.2% at 12 months ($p<.001$). Only five participants reported smoking at baseline, and this number did not change. Eight reported drinking alcohol at baseline, but only three reported drinking at the 12-month follow up.

Compared with baseline, significantly greater proportions of participants got correct answers on 12 of the 13 CVD knowledge questions at 12 months and felt more confident in their abilities to cook heart-healthy foods (Table 5). On the stages-of-change story questions, significantly more participants reported identifying with family members in the preparation, action, and maintenance stages of readiness ($p<.001$), rather than the pre-contemplation and contemplation stages (not shown in table). Of the 92 participants, all but three reported being satisfied or very satisfied with the curriculum, and 91% reported sharing curriculum information with their families (not shown in table).

We compared knowledge and health habit outcomes across the six intervention groups because groups were conducted by different CHWs. Group 2 participants, who had the poorest health habits at baseline, showed the most gains. Other groups showed more modest gains. Overall, however, this analysis suggested that CHWs did an equitable job of improving knowledge and health behaviors.




Figure 2. Stages of hypertension at baseline, 6 months, and 12 months (n=92).

Discussion

Our study produced several important findings. First, the *Healthy Heart, Healthy Family* curriculum designed for Filipino Americans was attractive to and appreciated by low-income Filipino immigrants in Hawaii. In addition to CVD risk factors, our participants presented with many challenges, including limited English proficiency, low health literacy, poverty, and unfamiliarity with CVD risk and the U.S. health care system. The low attrition rate and the active and sustained engagement of the participants in this program suggest that the curriculum resonated well with this group.

Table 3.

SELF-EFFICACY FOR MANAGING CHRONIC DISEASE (N=92)

Variable	Baseline	12 months	p-value	Standard error
Fatigue	7.69	8.38	.005	.239
Physical discomfort	7.76	8.25	.062	.256
Emotional distress	7.56	8.77	.01	1.029
Other health problems	7.73	8.13	.13	.261
Reduce doctor visits	8.19	8.56	.068	.198
Keep illness from affecting everyday life	8.10	8.63	.013	.212

Table 4.
FREQUENCY AND PROPORTION PERFORMING HEALTH BEHAVIORS ALL OR MOST OF THE TIME (N=92)

Behavior	Baseline n (%)	12 months n (%)	p-value
Use vegetable oil spray	40 (43.5%)	56 (60.9%)	.002
Read nutrition labels	52 (56.5%)	65 (70.7%)	.02
Remove the skin off the chicken	46 (50%)	70 (76.1%)	<.001
Use monounsaturated and polyunsaturated cooking oils	64 (69.6%)	74 (80.4%)	.03
Aerobic exercise	15 (16.7%)	36 (39.1%)	<.001
Replace salt with herbs and spices	5 (5.5%)	15 (16.3%)	.02
Snack on fruits and vegetables	56 (60.9%)	73 (79.4%)	.001
Bake instead of fry	31 (33.7%)	55 (59.8%)	<.001
Eat fruits for dessert	55 (59.8)	74 (80.4%)	.001

Second, the study shows that CHWs were able to learn the curriculum and train others in the curriculum. The *Healthy Heart Healthy Family* manual guided their delivery and served as a valuable training resource to promote behavior modification and help participants build skills to make practical and lasting changes to fight heart disease.

Employing CHWs in program delivery also conferred other benefits. For example, we attribute the low attrition and high participation rates to the linguistic and cultural competence of the CHWs, including one CHW who has been working at KKV for nearly 30 years and is well connected within the Filipino community. Because our CHWs knew the participants’ culture and their CHW roles, they were able to reinforce the curriculum’s messages through regular outreach and support. They also were successful in linking participants to services at the FQHC, including clinical care and health promotion activities. Finally, the CHWs were familiar with Filipino cultural values of *pakikisama* (togetherness), *bayanihan* (community spirit), and *utang-na-loob* (obligation and reciprocity). They overtly recognized and encouraged these values, which helped foster cohesiveness between the CHWs and the participants. Education and group wellbeing both are highly valued in Filipino culture. Thus, participants developed a sense of responsibility to the group and the CHW instructors, which facilitated their retention and active participation in the program.

Third, our findings confirmed that CHW interventions can improve participant knowledge and behavior.^{6,13} Many immigrants to the U.S. are unfamiliar with CVD risk factors, but are eager to learn about ways to prevent and control disease. Many are also unfamiliar with the U.S. health care system, and benefit greatly from the ability of the CHWs to understand them, help increase their health care knowledge and skills, and improve their access to care.¹³

Fourth, we demonstrated that CHW interventions can lead to improvements in

Table 5.**FREQUENCY AND PROPORTION WITH CORRECT ANSWERS ON CVD KNOWLEDGE (N=92)**

Question	Baseline n (%)	12 months n (%)	p-value
Can high waist measure increase risk of Cardio Vascular Disease?	42 (45.7%)	56 (60.9%)	.03
Can Body Mass Index tell you if you are overweight?	41 (44.6%)	69 (75.0%)	<.001
Does your liver make all the cholesterol your bodyneeds to keep you healthy?	26 (28.3%)	56 (60.1%)	<.001
Can eating foods high in sodium increase your risk of Hypertension?	67 (72.3%)	88 (95.6%)	<.001
Does lard have a low amount of saturated fat?	27 (29.3%)	59 (64.1%)	<.001
Can eating too much saturated fat and trans fat raise your cholesterol?	63 (68.4%)	82 (89.1%)	<.001
Is blood pressure of 140/90 considered high?	71 (77.2%)	84 (91.3%)	.003
Do people know their cholesterol is high because they have gained weight?	72 (78.3%)	82 (89.1%)	.056
Is being physically active a way to reduce your risk for health disease?	79 (85.9)	87 (94.6%)	.036
Do you think only people with high blood cholesterol should follow a health healthy diet?	53 (57.6%)	69 (75.0%)	.009
Can nonsmokers die from secondhand smoke?	70 (76.1%)	85 (82.4%)	<.001
Is having a fasting blood sugar of 126 mg/dl or higher considered diabetes?	44 (47.8%)	71 (77.2%)	<.001
Is having a waist measurement greater than 35 inches healthy for a woman?	33 (35.9%)	53 (57.6%)	.002
How confident are you in your ability to cook heart healthy foods? ^a	59 (64.1%)	77 (83.7%)	.003

^aShown are those who were confident or very confident.

healthy behaviors. People of all ethnicities and cultures suffer from CVD. In a 52-country study, Yusef et al. confirmed that abnormal lipid values, smoking, hypertension, diabetes, abdominal obesity, psychosocial factors, low fruit and vegetable consumption, and lack of regular physical activity account for most of the CVD risk in all ages and both sexes worldwide.¹⁴ Unfortunately, we did not realize the meaningful clinical outcomes we were hoping for, possibly because of the advanced age of the participants and the short follow-up (12 months).

Limitations of this study were its small sample size and the fact that data on health habits were collected from participants and may be subject to recall bias. Additionally, participants may have over-reported their adherence to program recommendations in

an effort to please the CHWs (social desirability bias). Further, there was no attempt at randomization and no control group due to limited funds for this study. Because the *Health Heart, Healthy family* promoted a low-salt and low-cholesterol diet, it is possible that some participants restricted fats dramatically, which could have contributed to the increase in HDL (as some low fat diets decrease HDL apolipoprotein transport rates).¹⁵

The study's strengths include its unique setting and population whose CVD needs have not been studied previously. Although the population of Asian Americans in the U.S. has increased dramatically over the last few decades, limited data are available on Filipinos and their heart health.¹⁶ We present a successful model of education and services delivered by CHWs that positively improved knowledge of cardiovascular health and behaviors to reduce CVD risk among low-income, immigrant Filipinos.

Poor health of people worldwide is largely associated with poverty, overcrowding, poor hygiene, and environmental contamination, aggravated by inadequate health promotion and disease prevention. As these groups move from traditional to modern lifestyles, they acquire diseases such as obesity, cardiovascular disease, and diabetes.¹⁷ Common traits of successful CVD programs include dedicated focus on the population, widespread community involvement (often through the employment of indigenous CHWs), and regularly scheduled contact between the program and participants.¹⁸ Community health workers are increasingly important as health educators and as bridges between the health care system and underserved communities.

The cultural and bilingual tools of CHWs enable them to build trust, serve as health care navigators, and connect clients to clinical and health promotion services. We demonstrated that CHWs can be trained to deliver an evidence-based, CVD risk-reducing curriculum to Filipino Americans. Although participants did not realize improvements in clinical measures, they gained knowledge about CVD risk and improved CVD-risk-related behaviors.

Acknowledgments

The research reported in this study was supported by grant H80CS00776-06-03, National Heart, Lung, and Blood Institute (NHLBI) and Health Resources and Services Administration (HRSA)/Bureau of Primary Health Care (BPHC) collaboration.

Notes

1. National Institutes of Health. Asian American and Pacific Islander workshops summary report on cardiovascular health (NIH Pub no. 00-3793). Bethesda, MD: U.S. Department of Health and Human Services, 2000.
2. Ye J, Rust G, Baltrus P, et al. Cardiovascular risk factors among Asian Americans: results from a national health survey. *Ann Epidemiol.* 2009 Oct;19(10):718–23.
3. O'Malley AS, Forrest CB, Politzer RM, et al. Health care trends, 1994–2001: what do they portend for the federal growth initiative? *Health Aff.* 2005 Mar–Apr;24(2):465–72.
4. Health Resources and Services Administration. Shortage. Washington, DC: Health Resources and Services Administration, 2011.
5. Love MB, Gardner K, Legion V. Community health workers: who they are and what they do. *Health Educ Behav.* 1997 Aug;24(4):510–22.

6. Zuvekas A, Nolan L, Tumaylle C, et al. Impact of community health workers on access, use of services, and patient knowledge and behavior. *J Ambulatory Care Manage.* 1999 Oct;22(4):33–44.
7. U.S. Department of Health and Human Services. *Healthy heart, healthy family: a community health worker's manual for the Filipino community.* Bethesda, MD: U.S. Department of Health and Human Services, 2010.
8. McLeroy KR, Bibeau D, Steckler A, et al. An ecological perspective on health promotion program. *Health Educ Q.* 1988 Winter;15(4):351–77.
9. Lorig KR, Sorbel DS, Ritter PL, et al. Effect of a self-management program for patients with chronic disease. *Eff Clin Pract.* 2001 Nov–Dec;4(6):256–62.
10. Flannagan JC. Measurement of quality of life: current state of the art. *Arch Phys Med Rehabil.* 1982 Feb;63(2):56–59.
11. Burckhardt CS, Anderson KL. QOLS: reliability, validity and utilization. *Health Qual Life Outcomes.* 2003 Oct 23;1:60.
12. IBM Corporation. *IBM SPSS Statistics version 20.* Armonk, NY: IBM Corporation, 2011.
13. Agency for Health Care Research and Quality. Outcomes of community health worker interventions. *Evid Rep Technol Assess.* 2009 Jun;(181):1–144, A1–2, B1–14 passim.
14. Yusuf S, Hawken S, Ounpuu S, et al. Effect of potentially modifiable risk factors associated with myocardial infarction in 52 countries (the INTERHEART study): case control study. *Lancet.* 2004 Sep 11;364(9438):937–52.
15. Brinton EA, Eisenberg S, Breslow JL. A low-fat diet decreases high density lipoprotein (HDL) cholesterol levels by decreasing HDL apolipoprotein transport rates. *J Clin Invest.* 1990 Jan;85(1):144–51.
16. National Heart, Lung, and Blood Institute. Report on a National Heart, Lung, and Blood Institute Workshop: heterogeneity in cardiometabolic risk in Asian American in the U.S. *J Am Coll Cardiol.* 2010 Mar 9;55(10):966–73.
17. Gracey M, King M. Indigenous health part 1. Determinants and disease patterns. *Lancet.* 2009 Jul 4;374(9683):65–75.
18. Huffman MD, Galloway JM. Cardiovascular health in indigenous communities: successful programs. *Heart Lung Circ.* 2010 May–Jun;19(5–6):351–60.