

PROOF COVER SHEET

Author: Laurence J.M. Coiffard

Article title: Silymarin, a molecule of interest for topical photoprotection

Article no: GNPL-637219

Enclosures: 1) Query sheet
2) Article proofs

Dear Author,

1. Please check these proofs carefully. It is the responsibility of the corresponding author to check these and approve or amend them. A second proof is not normally provided. Taylor & Francis cannot be held responsible for uncorrected errors, even if introduced during the production process. Once your corrections have been added to the article, it will be considered ready for publication.

For detailed guidance on how to check your proofs, please see <http://journalauthors.tandf.co.uk/production/checkingproofs.asp>.

2. Please review the table of contributors below and confirm that the first and last names are structured correctly and that the authors are listed in the correct order of contribution. This check is to ensure that your name will appear correctly online and when the article is indexed.

Sequence	Prefix	Given name(s)	Surname	Suffix
1		Celine	Couteau	
2		Clotilde	Cheignon	
3		Eva	Paparis	
4		Laurence J.M.	Coiffard	

Queries are marked in the margins of the proofs. Unless advised otherwise, submit all corrections and answers to the queries using the CATS online correction form, and then press the “Submit All Corrections” button.

AUTHOR QUERIES

General query: You have warranted that you have secured the necessary written permission from the appropriate copyright owner for the reproduction of any text, illustration, or other material in your article. (Please see <http://journalauthors.tandf.co.uk/preparation/permission.asp>.) Please check that any required acknowledgements have been included to reflect this.

<AQ1/> Please check “complimentary” has been changed to “complementary” in sentence “There are different ways ...complimentary”.

<AQ2/> Please provide “Conclusion” section.

<AQ3/> Please check reference “Diffey & Robson. (1989)” is listed, but not cited in the text.

How to make corrections to your proofs using Adobe Acrobat


Taylor & Francis now offer you a choice of options to help you make corrections to your proofs. Your PDF proof file has been enabled so that you can edit the proof directly. This is the simplest and best way for you to ensure that your corrections will be incorporated. To do this, please follow these instructions:

1. Check which version of Adobe Acrobat you have on your computer. You can do this by clicking on the “Help” tab, and then “About”.

If Adobe Reader is not installed, you can get the latest version free from <http://get.adobe.com/reader/>.

- If you have Adobe Reader 8 (or a later version), go to “Tools”/ “Comments & Markup”/ “Show Comments & Markup”.
- If you have Acrobat Professional 7, go to “Tools”/ “Commenting”/ “Show Commenting Toolbar”.

2. Click “Text Edits”. You can then select any text and delete it, replace it, or insert new text as you need to. It is also possible to highlight text and add a note or comment.



3. Make sure that you save the file when you close the document before uploading it to CATS. A full list of the comments and edits you have made can be viewed by clicking on the “Comments” tab in the bottom left-hand corner of the PDF.

If you prefer, you can make your corrections using the CATS online correction form.

Silymarin, a molecule of interest for topical photoprotection

Celine Couteau, Clotilde Cheignon, Eva Papisaris and Laurence J.M. Coiffard*

Faculty of Pharmacy, Université de Nantes, Nantes Atlantique Universités, LPiC, MMS, EA2160,
1 rue G. Veil – BP 53508, Nantes, F-44000 France

5 (Received 6 July 2011; final version received 29 August 2011)

10 Some UV-filters have side effects. For example, oestrogenic effect was demon-
strated for 4-methylbenzylidene camphor. Given that secondary metabolites are
known for their UVB photoprotective properties in plants that contain them, we
15 chose to study silymarin as an agent which could potentially be used in sun
products. This determination is based on the physical determination of the
reduction of the energy in the UV range, through a film of product which has
previously been spread on an adequate substrate. About 15 mg of O/W emulsion
containing silymarin at various concentrations was applied on roughened
20 PMMA plates and the transmission measurements are carried out using a
spectrophotometer equipped with integrating sphere. Incorporated in O/W
creams, at a concentration of 10% (w/w), silymarin gives a Sun Protection Factor
similar to that of octylmethoxycinnamate, which is why it is predominantly used
in Europe. Overall, these results demonstrate that silymarin is a promising new
sunscreen agent.

20 **Keywords:** silymarin; SPF; PF-UVA; *in vitro* method

1. Introduction

1 There are different ways of protecting the skin against the negative effects of the sun and
they are complementary. These complementary ways are the photoprotection coming from
clothes and the topical formulations (Ghazi, Couteau, & Coiffard, 2010, 2011).

25 Topical products are formulated with organic and/or inorganic sun filters. Their
efficacy is known, but they can lead to unwanted effects on humans and/or on the
environment (Danovaro et al., 2008; Knobler et al., 1989; Schlumpf, Jarry, Wuttke, Ma, &
Lichtensteiger, 2004; Soeborg, Ganderup, Kristensen, Bjerregaad, & Pedersen, 2006).
30 In this context, finding alternatives would be relevant. Plants exposed to sunlight produce
secondary metabolites to protect their genome. Some already appear as interesting sources
of new photoprotective molecules for topical application. For example, boldine, which is
an alkaloid coming from the boldo tree, is well known for its antioxidant properties
(Hidalgo, Farah, Carrasco, & Fernández, 2005; O'Brien, Carrasco-Pozo, & Speisky, 2006).
35 Aromatic compounds from certain lichens (1'-chloropannarine, epiphorelic acids I and II,
calicine) demonstrate, *in vitro*, a comparable efficacy as the octylmethoxycinnamate, the
UVB filter which is mostly used within Europe (Couteau, Faure, Fortin, Papisaris, &
Coiffard, 2007a; Couteau, Pommier, Papisaris, & Coiffard, 2007b). The flavonoids, natural
colourants of plants, are also an interesting family, in particular chlorogenic acid, baicalin,

*Corresponding author. Email: laurence.coiffard@univ-nantes.fr

luteolin, apigenin and puerarin (Choquet, Couteau, Papis, & Coiffard, 2009). From the marine environment, mycosporins such as amino acids would provide a photoprotection in the UVA range (Conde, Churio, & Previtali, 2000; Vaid & Katiyar, 2010; Zhang & Wu, 2007). We took particular interest in the silymarin, coming from the *Silybum marianum*, known for its antioxidant properties (Gazak, Svobodova, & Psotova, 2004; Katiyar, 2005).

2. Results and discussion

The silymarin spectrum presents a peak of absorption at 286 nm. The influence of different concentrations of silymarin on the efficacy in the UVB and UVA ranges was studied. The results obtained in terms of efficiency are extremely promising, as the use level of 10% (w/w) of silymarin allows a Sun Protection Factor (SPF) close to 9 (Figure 1). Thanks to this value measured, the silymarin can be compared to a UV filter such as the octylmethoxycinnamate (Couteau et al., 2007a), which is a very efficient sun filter and the reason why it is predominantly used in Europe. The silymarin positions itself as a UVB filter. It is less efficient in the UVA range (Figure 1), as confirmed by the spectrum measured with the aqueous solution. The combinations with the zinc oxide and the titanium dioxide give interesting results. The measured SPF values of the preparations, formulated with filters according to their respective regulations, are higher than 10 (Table 1). After 2 h of irradiation at 650 W m⁻², the silymarin proved to be photostable. Indeed, in the case of a silymarin-based sun product, more than 90% efficacy is kept. Therefore, the silymarin shows a photostability higher than most of the organic UV filters

b/w in print
colour online




Figure 1. Efficacy of silymarin in UVB (1) and UVA (2) ranges.

Table 1. Data concerning the combination between silymarin and inorganic sun filters, before and after irradiation during 2 h at 650 W/min.

Product associated with silymarin	SPF ₀ ± SD	PF-UVA ₀ ± SD	λ _{c0} (nm)	SPF _{2h} ± SD	PF-UVA _{2h} ± SD	λ _{c2h} (nm)
10% (w/w)						
Z-Cote max (10% w/w)	12.37 ± 4.39	7.32 ± 2.34	378	11.84 ± 3.84	7.18 ± 2.06	378
Eusolex T-Oléo (10% w/w)	16.30 ± 2.98	7.61 ± 0.97	377	16.49 ± 2.95	17.00 ± 1.39	378

60 sold on the market at the moment (Couteau et al., 2007b). Other studies have already
demonstrated that the silymarin would be interesting in the prevention of photocarcino-
65 genesis phenomena (Jiang, Wang, Onodera, & Ikejima, 2009; Nichols & Katiyar, 2010;
Rancan et al., 2002; Zudaire & Roy, 2001). By its efficacy, its photostability and the other
properties already displayed, the silymarin is an ingredient to consider and to emphasise in
the topical photoprotection domain.

3. Experimental

3.1. Materials

Silymarin, with a purity of 80% as total flavonolignans determined by HPLC, was
obtained from Bioserae (Bram, France). Cetiol® HE, stearic acid, glycerin, parabens and
70 triethanolamine (TEA) were purchased from Cooper (Melun, France). Xanthan gum
(Keltrol® BT) was obtained from Kelco (Lille Skensved, Denmark). Zinc oxide (Z-Cote
Max®) was obtained from BASF (Ludwigshafen, Germany). Titanium dioxide (Eusolex®
T-aqua) was obtained from Merck (Damstadt, Germany). Polymethylmethacrylate
(PMMA) plates were purchased from Europlast (Aubervilliers, France).

3.2. Methods

75 Aqueous solutions were scanned at wavelengths between 200 and 400 nm using a
spectrophotometer double-beam (Hitachi UV-visible, model U-2000). The spectra were
measured against a pure water sample in quartz cells with a 1-cm optical path length.

An O/W emulsion was prepared in the laboratory by adding known concentrations of
80 silymarin into the formulation components. A detailed description of the protocol is
described in a previously published paper (Couteau et al., 2007b). The plates were
irradiated for 2 h with a solar simulator (Suntest CPS+; Atlas, Moussy le Neuf, France)
device equipped with a xenon arc lamp (1500 W) and special glass filters restricting the
transmission of light below 290 nm. The light source emission was maintained at
85 650 W m^{-2} in accordance to the global solar spectral irradiance (Couteau et al., 2007a).
Before and after irradiation, the SPF and the PF-UVA of the creams were measured
2 *in vitro*.

References

- 90 Choquenot, B., Couteau, C., Papis, E., & Coiffard, L.J.M. (2009). Flavonoids and polyphenols, molecular
families with sunscreen potential: Determining effectiveness with an *in vitro* method. *Natural Products
Communications*, 4, 227–230.
- Conde, F.R., Churio, M.S., & Previtali, C.M. (2000). The photoprotector mechanism of mycosporine-like amino
acids. Excited-state properties and photostability of porphyra-334 in aqueous solution. *Journal of
Photochemistry and Photobiology B: Biology*, 56, 139–144.
- 95 Couteau, C., Faure, A., Fortin, J., Papis, E., & Coiffard, L.J.M. (2007a). The photoprotector mechanism of
mycosporine-like amino acids. Excited-state properties and photostability of porphyra-334 in aqueous
solution. *Journal of Pharmaceutical and Biomedical Analysis*, 44, 270.
- Couteau, C., Pommier, M., Papis, E., & Coiffard, L.J.M. (2007b). Study of the efficacy of 18 sun filters
authorized in European Union tested *in vitro*. *Pharmazie*, 62, 449–452.
- 100 Danovaro, R., Bongiorno, L., Corinaldesi, C., Giovannelli, D., Damiani, E., Astolfi, P., . . . , Pusceddu, A. (2008).
Sunscreens cause coral bleaching by promoting viral infections. *Environmental Health Perspective*, 116,
441–447.
- Diffey, B.L., & Robson, J. (1989). A new substrate to measure sunscreen protection factors throughout the
ultraviolet spectrum. *Journal of Society of Cosmetic Chemists*, 40, 127–133.
- 105 Gazak, R., Svobodova, A., & Psotova, A. (2004). Oxidised derivatives of silybin and their antiradical and
antioxidant activity. *Bioorganic and Medicinal Chemistry*, 12, 5677–5687.

4 C. Couteau et al.

- Ghazi, S., Couteau, C., & Coiffard, L.J.M. (2010). What level of protection can be obtained using sun protective clothing? Determining effectiveness using an *in vitro* method. *International Journal of Pharmacy*, 397, 144–147.
- 110 Ghazi, S., Couteau, C., & Coiffard, L.J.M. (2011). How to guarantee adequate sun protection for a young sports person. *Journal der Deutschen Dermatologischen*, 9, 470–474.
- Hidalgo, M.E., Farah, M., Carrasco, L., & Fernández, E. (2005). Photostability and photoprotection factor of boldine and glaucine. *Journal of Photochemistry and Photobiology B: Biology*, 80, 65–69.
- 115 Jiang, Y.Y., Wang, H.J., Onodera, S., & Ikejima, T. (2009). The protective effect of silibinin against mitomycin C-induced intrinsic apoptosis in human melanoma A375-S2 cells. *Journal of Pharmacologic Sciences*, 11, 137–146.
- Katiyar, S. (2005). Silymarin and skin cancer prevention anti-inflammatory antioxidant and immunomodulatory effects review. *International Journal of Oncology*, 26, 169–176.
- 120 Knobler, E., Almeida, L., Ruzkowski, A.M., Held, J., Harber, L., & DeLeo, V. (1989). Photoallergy to benzophenone. *Archives of Dermatology*, 125, 801–804.
- Nichols, J.A., & Katiyar, S.K. (2010). Skin photoprotection by natural polyphenols: Anti-inflammatory, antioxidant and DNA repair mechanisms. *Archives of Dermatology Research*, 302, 71–83.
- O'Brien, P., Carrasco-Pozo, C., & Speisky, H. (2006). Boldine and its antioxidant or health-promoting properties. *Chemico-Biological Interactions*, 159, 1–17.
- 125 Rancan, F., Rosan, S., Boehm, K., Fernández, E., Hidalgo, M.E., Quihot, W., & Oltmanns, U. (2002). Protection against UVB irradiation by natural filters extracted from lichens. *Journal of Photochemistry and Photobiology B: Biology*, 68, 133–139.
- Schlumpf, M., Jarry, H., Wuttke, W., Ma, R., & Lichtensteiger, W. (2004). Estrogenic activity and estrogen receptor beta binding of the UV filter 3-benzylidene camphor. *Comparison with 4-methylbenzylidene camphor*. *Toxicology*, 199, 109–120.
- 130 Soeborg, T., Ganderup, N.C., Kristensen, J.K., Bjerregaard, P., & Pedersen, K.L. (2006). Distribution of the UV filter 3-benzylidene camphor in rat following topical application. *Journal of Chromatography B*, 834, 117–121.
- Vaid, M., & Katiyar, S.K. (2010). Molecular mechanisms of inhibition of photocarcinogenesis by silymarin, a phytochemical from milk thistle (*Silybum marianum* L. Gaertn.) (Review). *International Journal of Oncology*, 36, 1053–1060.
- 135 Zhang, L., Li, L., & Wu, Q. (2007). Protective effects of mycosporine-like amino acids of *Synechocystis* sp. PCC 6803 and their partial characterization. *Journal of Photochemistry and Photobiology B: Biology*, 86, 240–245.
- 140 Zudaire, L., & Roy, S. (2001). Photoprotection and long-term acclimation to UV radiation in the marine diatom *Thalassiosira weissflogii*. *Journal of Photochemistry and Photobiology B: Biology*, 62, 26–34.