

Zeitschrift für

Gerontologie + Geriatric

mit **European Journal of Geriatrics**

**Elektronischer Sonderdruck für
M. Noll-Hussong**

Ein Service von Springer Medizin

Z Gerontol Geriat 2012 · 45:404–410 · DOI 10.1007/s00391-012-0303-9

© Springer-Verlag 2012

zur nichtkommerziellen Nutzung auf der
privaten Homepage und Institutssite des Autors



**M. Noll-Hussong · H. Glaesmer · S. Herberger · K. Bernardy · C. Schönfeldt-
Lecuona · A. Lukas · H. Guendel · T. Nikolaus**

Die Früchte des Krieges

Somatoforme Schmerzstörungen und Vorgeschichten von
früher Kriegstraumatisierung bei Älteren



BUNDESVERBAND
GERIATRIE



Z Gerontol Geriat 2012 · 45:404–410
 DOI 10.1007/s00391-012-0303-9
 Received: 18 August 2011
 Revised: 9 September 2011
 Accepted: 26 Januar 2012
 Published online: 9 July 2012
 © Springer-Verlag 2012

M. Noll-Hussong¹ · H. Glaesmer² · S. Herberger³ · K. Bernardy^{4,7} ·
 C. Schönfeldt-Lecuona⁵ · A. Lukas⁶ · H. Guendel¹ · T. Nikolaus⁶

¹ Klinik und Poliklinik fuer Psychosomatische Medizin und Psychotherapie, University of Ulm

² Department of Medical Psychology and Medical Sociology, University of Leipzig

³ Klinik fuer Psychosomatische Medizin und Psychotherapie, Klinikum Muenchen-Harlaching

⁴ Department of Anaesthesiology, Intensive Care and Pain Therapy,

Saarland University Hospital, Homburg/Saar

⁵ Klinik fuer Psychiatrie III, University of Ulm

⁶ Agaplesion Bethesda Klinik Ulm, University of Ulm

⁷ Department of Psychosomatic Medicine, MediClin Bliestal Clinics, Bliestal

The grapes of war

Somatoform pain disorder and history of early war traumatization in older people

According to the US National Institute on Aging, approximately 500 million people worldwide were at least 65 years old by 2006. This number represents 12% of the world's population. By 2030, the figure is expected to climb to 20%. As health conditions also improve in developing countries, life expectancies will generally rise, and the number of people with mental problems due to age-related processes will accordingly increase.

Since the concept of “gerontopsychosomatics” was created in 1990, there has been increasing interest in the bio-psychosocial implications of mind–body interactions during the aging process in the second half of adulthood. However, integrated treatment strategies that involve psychotherapy adjusted for elderly patients with “psychosomatic diseases” such as somatoform disorders are lacking [48].

Irrespective of age, good health does not imply an absence of symptoms. Experiencing symptoms is part of daily life and an endemic human experience. Of the general population, 80–90% report ≥1 symptom per week, and a typical adult has 1 medically unexplained symptom every 5–7 days. Of college students, 80% report ≥1 symptom within 3 days and about 70% of adults complain of back pain during their lifetime. However, only a small minority require medical consultation [8].

Various labels have been given to medically unexplained symptoms, including “functional somatic symptoms,” “somatization,” and “somatoform disorders” [52]. It seems as though each medical specialty has at least one “functional syndrome,” and it has been suggested that the existence of specific somatic syndromes is largely an artifact of medical specialization. For example, for gastroenterologists, abdominal pain with altered intestinal habit is labeled irritable bowel syndrome (IBS), and for rheumatologists, chronic widespread pain, allodynia, and fatigue is considered fibromyalgia [52].

In this context and reflecting the old-fashioned dualistic dichotomy between “body” and “soul,” the official classification systems distinguish between somatic (“functional”) and mental (“somatoform”) symptoms and disorders, respectively (in terms of a double classification of human diseases) [33]. The main feature of somatoform disorders is the repeated presentation of physical symptoms along with persistent requests for medical investigations, in spite of repeated negative findings and reassurances by physicians that the symptoms have no physical basis [33]. If physical disorders are present, these disorders do not explain the nature and extent of the symptoms or the patient's distress [13] and preoccupation. In

this context, severe and disabling symptoms of pain that remain medically unexplained after in-depth examination (symptoms that also cannot be explained fully by the direct effect of a substance or attributable to another mental disorder (i.e., panic disorder)) are referred to as somatoform pain disorder [32]. These patients frequently consult primary care providers. In specialized clinics, 30–40% of patients in the gastroenterology and neurology departments and up to 66% of those in the gynecology department suffer from somatoform pain disorder [52]. The German national survey on psychiatric disorders, conducted as part of the national health survey of 1999, indicated that somatoform disorders are among the leading mental disorders (after anxiety and affective disorders), with a 4-week cross-sectional prevalence of 7.5% [53]. Internationally, the prevalence of the disorder ranges between 9% and 20% [38]. In a representative study of 630 persons older than 60 years who were asked to complete the SOMS 2 (*Screening for Somatoform Symptoms*) questionnaire, 71.8% reported at least one symptom, 50.5% experienced at least four symptoms, and 23.4% suffered from at least eight symptoms. Patients with medically unexplained physical symptoms are so-called high utilizers of the health care system, incurring costs

Hier steht eine Anzeige.






Fig. 1 ▲ Vulnerability factors for the development of somatoform pain disorder in older people are female gender, low education level, trauma, poor coping strategies, negative affect, and alexithymia. *Alexithymia* refers to a state of deficiency in understanding, processing, or describing emotions. Adapted from [48]

that are up to 10 times higher than those of average patients. Adjusting the findings for the presence of mental and somatic comorbidity had relatively little effect on this association [4].

Considering multimorbidity and aging, there is a strong relationship between chronic somatic diseases and mental disorders such as somatoform disorders. Psychiatric morbidity is generally high, and many patients have chronic psychosocial problems. In a geriatric acute hospital, a minimum of 20% of the patients aged 60 or older fulfill the case criteria for a psychogenic illness. As intelligence is negatively associated with the number of somatoform symptoms, reported symptoms increase with decreasing cognitive performance, as indicated by the Mini-Mental State Examination (MMSE). Moreover, somatoform disorder has a high comorbidity with major depression and anxiety disorders [34, 39], and there seems to be an association between somatoform and substance use disorders [22]. Research has suggested that the frequency of somatoform symptoms is much higher in people over 60 years than in persons who are younger than 60 [48]. Multimorbidity and polypharmacy are common problems in older people and are significantly correlated with higher mortality, increased disability, and functional decline. In a study

identifying multimorbidity patterns using exploratory tetrachoric factor analysis based on claims data of 63,104 males and 86,176 females in the age group 65+ years, the following three multimorbidity patterns were identified:

- cardiovascular/metabolic disorders (prevalence: female 30%, male 39%),
- anxiety/depression/somatoform disorders and pain (34%, 22%), and
- neuropsychiatric disorders (6%, 0.8%) [47].

Somatoform complaints and disorders occur more frequently in elderly women than in elderly men [48].

It has been estimated that 50% of older people who live independently and 75% of those who live in nursing homes suffer from chronic pain. However, the etiology of this persistent pain is often not entirely clear. Among the somatoform symptoms, pain complaints such as back pain, joint pain, pain in the limbs, and headache or facial pain are the most frequent [24]. In the German general population, the lifetime prevalence of pain-predominant somatoform disorders is 33.7%, with 12.3% of these individuals being significantly impaired. The prevalence of pain was found to be greater with increasing age, lower education, lower income, rural residency, and residency in Eastern Germa-

ny. Although the prevalence of somatoform pain is high, the majority of patients do not receive adequate psychotherapeutic care but are inadequately treated by somatic treatments, e.g., reflected by so-called doctor (s)hopping [24]. Altogether, somatoform pain disorder is both a frequent and neglected condition in older people [19].

Characteristics of somatoform pain disorder in older people

Reports indicate that the characteristics of older adults with chronic pain may differ from those of younger persons [37]. For example, pain in older people is more often localized in several body regions [37], and symptoms often do not become clinically prevalent until middle age [44]. Moreover, elderly patients tend to report more somatic complaints and avoid expressing the emotional dimension of distress. When examining the differences between “young olds” and “old olds,” somatoform pain increases with increasing age [25], e.g., in chronic widespread pain [2]. A study of 115 geriatric in-patients that aimed to identify links between regions of experienced pain, subjective pain intensity, and psychological variables revealed that 61.7% of the study sample reported intense pain in at least one body region. These patients differed significantly from those with less pain in terms of their level of functional capacity and need for assistance in daily activities. They also displayed a significantly more negative attitude towards aging than patients with minimal pain. Furthermore, emotional distress (like pain itself), but also comorbid depression and anxiety play important roles in memory complaints in patients with chronic pain, as well as rumination, actually the intrusive component of catastrophizing. Finally, earlier pain experiences (e.g., in the context of a physical disorder or traumatic events during war) may “pave the way” for somatoform pain syndromes [10].

Burden of traumatization of early war experiences

Somatoform symptoms are more prevalent in traumatized patients than nontrau-

matized patients [43]. From a lifespan perspective, sexual or physical abuse in early life is associated with psychopathology in later adulthood [29], especially the severity of somatization [41] and chronic pain [18]. Against the background of World War II, the extent and long-term effects of war-related traumatic experiences (e.g., experiences during military war service, bombing, or wartime rape) in the elderly generation are of special interest [36]. An investigation of the long-term effects of the British evacuation of children during World War II on the mental health of the now older people found significant associations between early experiences and lifetime mental health [46]. A recent study examining the occurrence of traumatic experiences and the prevalence rates of post-traumatic stress disorder (PTSD) and partial PTSD, defined according to the DSM-IV, in a randomly selected sample of the German general population aged 60 years and over showed that PTSD was apparent in 3.4% of the sample. When partial PTSDs were included, the total rose to 7.2% of the elderly population. It is important to note that the most frequently mentioned traumatic incidents of the generation examined in this study were war-related traumas experienced as a child or early adolescent during World War II. As a person's age increased, the prevalence of war-related traumatic experiences also increased from 19.2% in those aged 60–64 up to 59.7%(!) in those who were 75 and older [17]. Much lower prevalence rates were found in the Swiss older people who largely escaped war-related trauma [35]. An examination looking at a more recent conflict found that Persian Gulf War veterans' reports of traumatic events [14] were clearly associated with somatoform diagnoses. Veterans who handled dead bodies had a 3-fold higher risk of receiving a diagnosis of somatoform disorder [31]. A higher proportion of veterans of the Persian Gulf War of 1991 reported symptoms of pain than did military comparison groups [50]. In fact, reports of pain symptoms could be easily continued with regard to refugees [27], child soldiers [30], prisoners of war [11], the "unspoken secret" of sexual violence in World War II [28], and veterans of former and recent conflicts [1, 26].

Z Gerontol Geriat 2012 · 45:404–410 DOI 10.1007/s00391-012-0303-9
© Springer-Verlag 2012

M. Noll-Hussong · H. Glaesmer · S. Herberger · K. Bernardy · C. Schönfeldt-Lecuona · A. Lukas · H. Guendel · T. Nikolaus

The grapes of war. Somatoform pain disorder and history of early war traumatization in older people

Abstract

Persistent pain is not a normal part of aging. Nevertheless, many older patients have long-lasting, more or less medically unexplained pain symptoms and, consequently, are often severely disabled, incur high health care costs, and have high comorbidity rates. Moreover, the effects of early traumatization, especially due to wars, and even below the level of posttraumatic stress disorder (PTSD) are apparent. However, the developmental and neurobiological underpinnings of somatoform pain disorder, especially in pain-prone elderly patients, and its correlations with a history of war traumatization even decades

after the incident remain unclear. Furthermore, a management strategy for this disorder tailored to older people and their special needs is lacking. Adequate therapeutic regimens such as adjusted psychotherapeutic procedures for elderly patients can only be promoted through a better understanding of the neurobiological and biographical underpinnings of this still controversial disorder.

Keywords

Somatoform disorders · Chronic pain · Posttraumatic stress disorders · Aged · Neurobiology

Die Früchte des Krieges. Somatoforme Schmerzstörungen und Vorgeschichten von früher Kriegstraumatisierung bei Älteren

Zusammenfassung

Anhaltende Schmerzen sind kein normaler Bestandteil des Alterns. Dennoch haben viele ältere Patienten dauerhaft eine mehr oder weniger medizinisch nicht erklär-bare Schmerzsymptomatik. Sie sind damit oft schwer beeinträchtigt, verursachen hohe Kosten im Gesundheitswesen und zeigen eine hohe Komorbiditätsrate. Auch sind die Folgen früher Traumatisierungen, insbesondere durch Kriege und sogar unterhalb der Ausprägung einer posttraumatischen Belastungsstörung (PTSD), offensichtlich. Jedoch bleiben die Entstehungs- und neurobiologischen Grundlagen der somatoformen Schmerzstörung, vor allem in schmerzgeneigten, älteren Patienten, und ihre Zusammenhänge mit einer Vorgeschichte von Kriegstraumatisierung – sogar Jah-

rzehnte nach dem Ereignis – unklar. Darüber hinaus fehlt für diese Erkrankung eine Behandlungsstrategie, die auf ältere Menschen und ihre besonderen Bedürfnisse zugeschnitten wäre. Adäquates therapeutisches Vorgehen, wie angepasste psychotherapeutische Prozeduren für ältere Patienten, können nur durch ein besseres Verständnis der neurobiologischen und biographischen Grundlagen dieser nach wie vor kontroversen Störung gefördert werden.

Schlüsselwörter

Somatoforme Störungen · Chronische Schmerzen · Posttraumatische Belastungsstörungen · Alte Menschen · Neurobiologie

In addition to developmental and psychological considerations, the (neuro)biological links between somatoform pain disorder and histories of early traumatization have been intensely researched [7, 42]. For example, dysfunctions of the hypothalamic–pituitary–adrenal stress axis are described. However, the mechanisms still remain largely unclear (■ Fig. 1).

Neurobiology of somatoform pain disorder and a history of early traumatization

Reaching a greater neurobiological understanding of aging and disease prevention strategies seems to be crucial for maintaining brain health. However, the neural mechanisms of non-nociceptive pain have rarely been studied through positron emission tomography (PET), SPECT, or functional magnetic resonance imaging




Fig. 2 ◀ Ongoing conflicts worldwide 2010. Adapted from [54]

(fMRI). In this context, a dysfunction of neural pain processing circuits has been suggested as one underlying pathophysiological factor in somatoform pain disorder [6]. A fMRI study that investigated the cerebral processing of noxious heat stimuli as an objective marker of pain sensation in 12 middle-aged women with somatoform pain disorder showed a pain-related hypoactive state of the ventromedial prefrontal/orbitofrontal cortex and a hyperactive state of the parahippocampal gyrus, amygdala, and anterior insula in the patient group compared to the control group. These findings of altered cerebral processing of experimentally induced pain in patients with somatoform pain disorder support the hypothesis of dysfunction in the pain matrix, especially in affect-regulating regions [21]. Remarkably, altered pain processing has also been found in veterans with PTSD [16]. Independent of dementia processes, middle-aged patients with somatoform pain disorder show gray-matter loss in pain-processing structures [51]. To date, only two neuroimaging studies in the “grey area” between somatoform pain and early traumatization (and, furthermore, only in the middle age) have been performed. Comparing the neural substrates of empathy-induced pain in multisomatoform pain patients “with vs. without” a history of sexual abuse during childhood revealed higher activations in left lateral and medial superior frontal gyrus as well as a nonsignificant activation of the right supplementary motor area in abused patients. The nonabused partici-

pants showed a higher activation of the left hippocampus [42]. Ringel et al. [45] found that pain ratings during rectal distension in patients with IBS are associated with the activation of dorsal cingulate regions implicated in homeostatic afferent processing, and prior abuse enhances this activation. Patients with IBS and a history of abuse report more pain, greater activation in the posterior and middle dorsal cingulate subregions, and reduced activity in the left supragenual anterior cingulate cortex, a region implicated in pain inhibition and arousal. Again, these findings suggest a possible explanation for the clinical observation of greater pain reporting and poorer outcomes in IBS patients with a history of abuse [45]. It should be noted, as an important point, that also the epigenetic transmission of the impact of early stress (like traumatization) across generations has recently been discussed [15].

Treatment options

To date there are no evidence-based recommendations for the treatment of somatoform disorders in general for older people, nor in primary and secondary care, or for psychotherapy [48]. Against this background, there is a similar lack of treatment options for somatoform pain disorder. In general, there is little available evidence on the use of psychosocial interventions by general practitioners beyond depression and with patients over 65 years of age. Remarkably, the Cochrane review “Effectiveness and cost effectiveness

of counseling in primary care” excluded “psychosomatic problems such as pain” [5]. Treatment seems to be more effective in patients in secondary care than in primary care. This may be because secondary care patients have more severe disease, receive a different treatment regimen, or have a more closely supervised intervention. The few controlled studies in middle-aged patients [40] seem to favor cognitive-behavioral therapy (CBT) in group settings. Of note, Arnold et al. [3] showed that the acceptance of CBT is significantly higher in older people. Furthermore, special psychotherapeutic treatments for traumatization such as EMDR (eye movement desensitization and reprocessing) may be also efficacious in the treatment of chronic pain [20, 49], but further research is urgently necessary. In general, non-pharmacological treatments that involve the active participation of patients, such as exercise and psychotherapy, seem to be more effective than those that involve passive physical measures such as massage, injections, and operations. Pharmacological agents with CNS action are more consistently effective than drugs aimed at the restoration of peripheral dysfunction. A careful balance between biomedical, organ-oriented, and cognitive interpersonal approaches might be the most appropriate intervention. Especially for chronic low back pain, it has been shown that an expressly “intensive multidisciplinary bio-psycho-social rehabilitation with a functional restoration approach improves pain and function”. Finally, the iatrogenic component in

the maintenance of somatoform pain disorder should also be considered, for which reason the dynamics of the patient–doctor relationship should not be overlooked [23].

Need for further research

With Eric Kandel and having psychotherapy in mind, we would claim:

“The need for a biological perspective is self-evident. [...] We would argue that research is ... needed into the mechanisms of psychotherapeutic action on the biological, cognitive, and behavioural levels, yet psychotherapy research lags far behind that on pharmacotherapy. [...] This is in part due to the cost, inconvenience and difficulty of conducting and evaluating a complete course of psychotherapy under controlled conditions” [12].

Regrettably, 70 years after World War II, there remains insufficient evidence-based and scientific knowledge about how to treat these older patients adequately and effectively [48]. Moreover, considering the 80 ongoing wars and (permanent) conflicts worldwide (e.g., in Lebanon, Afghanistan, Iraq, and Sudan (■ Fig. 2)), the prevention of somatoform disorders is a challenge. The investigation of the neurobiological link between somatoform pain disorder in older people and a history of early traumatization using both biological (such as fMRI and genetics) and psychological (such as questionnaires and the study of behavior) instruments is ambitious, but it could lead to a better scientific understanding of this mental disorder and improved clinical decision-making. The development of both tailored and optimized, differential intense psychotherapeutic procedures could ease a great deal of pain in a growing group of patients who have been neglected thus far.

Conclusion

The burden of medically unexplained pain, especially in older people, urgently requires further research. As histories of early traumatization (e.g., war experiences) already below the level of PTSD play an important role in the etiology of chronic pain disorder, both the “forgot-

ten” elderly who suffer from historical war experiences and younger individuals who take part in conflicts [9], genocides (e.g., in Rwanda), and migration in a globalizing and still militant world should be considered

Corresponding address

M. Noll-Hussong

Klinik und Poliklinik fuer Psychosomatische Medizin und Psychotherapie, University of Ulm Am Hochstr. 8, 89081 Ulm Germany
minohu@gmx.net

Conflict of interest. On behalf of all authors, the corresponding author states that there are no conflicts of interest.

References

1. Afari N, Harder LH, Madra NJ et al (2009) PTSD, combat injury, and headache in Veterans returning from Iraq/Afghanistan. *Headache* 49:1267–1276
2. Aggarwal VR, Mcbeth J, Zakrzewska JM et al (2006) The epidemiology of chronic syndromes that are frequently unexplained: do they have common associated factors? *Int J Epidemiol* 35:468–476
3. Arnold IA, Speckens AE, Van Hemert AM (2004) Medically unexplained physical symptoms: the feasibility of group cognitive-behavioural therapy in primary care. *J Psychosom Res* 57:517–520
4. Barsky AJ, Orav EJ, Bates DW (2005) Somatization increases medical utilization and costs independent of psychiatric and medical comorbidity. *Arch Gen Psychiatry* 62:903–910
5. Bower P, Rowland N (2006) Effectiveness and cost effectiveness of counselling in primary care. *Cochrane Database Syst Rev* 3:CD001025
6. Browning M, Fletcher P, Sharpe M (2011) Can neuroimaging help us to understand and classify somatoform disorders? A systematic and critical review. *Psychosom Med* 73:173–184
7. Buffington CA (2009) Developmental influences on medically unexplained symptoms. *Psychother Psychosom* 78:139–144
8. Creed F, Barsky A (2004) A systematic review of the epidemiology of somatisation disorder and hypochondriasis. *J Psychosom Res* 56:391–408
9. Difede J, Barchas JD (2010) Psychiatric and neurologic aspects of war: an overview and perspective. *Ann NY Acad Sci* 1208:1–9
10. Elklit A, Christiansen DM (2009) Predictive factors for somatization in a trauma sample. *Clin Pract Epidemiol Ment Health* 5:1
11. Engdahl B, Dikel TN, Eberly R et al (1998) Comorbidity and course of psychiatric disorders in a community sample of former prisoners of war. *Am J Psychiatry* 155:1740–1745
12. Etkin A, Pittenger C, Polan HJ et al (2005) Toward a neurobiology of psychotherapy: basic science and clinical applications. *J Neuropsychiatry Clin Neurosci* 17:145–158
13. Fink P, Schroder A (2010) One single diagnosis, bodily distress syndrome, succeeded to capture 10 diagnostic categories of functional somatic syndromes and somatoform disorders. *J Psychosom Res* 68:415–426
14. Ford JD, Campbell KA, Storzbach D et al (2001) Posttraumatic stress symptomatology is associated with unexplained illness attributed to Persian Gulf War military service. *Psychosom Med* 63:842–849
15. Franklin TB, Ruggie H, Weiss IC et al (2010) Epigenetic transmission of the impact of early stress across generations. *Biol Psychiatry* 68:408–415
16. Geuze E, Westenberg HG, Jochims A et al (2007) Altered pain processing in veterans with posttraumatic stress disorder. *Arch Gen Psychiatry* 64:76–85
17. Glaesmer H, Gunzelmann T, Braehler E et al (2010) Traumatic experiences and post-traumatic stress disorder among elderly Germans: results of a representative population-based survey. *Int Psychogeriatr* 22:661–670
18. Goldberg RT, Pachas WN, Keith D (1999) Relationship between traumatic events in childhood and chronic pain. *Disabil Rehabil* 21:23–30
19. Grabe HJ, Meyer C, Hapke U et al (2003) Somatoform pain disorder in the general population. *Psychother Psychosom* 72:88–94
20. Grant M, Threlfo C (2002) EMDR in the treatment of chronic pain. *J Clin Psychol* 58:1505–1520
21. Gundel H, Valet M, Sorg C et al (2008) Altered cerebral response to noxious heat stimulation in patients with somatoform pain disorder. *Pain* 137:413–421
22. Hasin D, Katz H (2007) Somatoform and substance use disorders. *Psychosom Med* 69:870–875
23. Henningsen P, Zipfel S, Herzog W (2007) Management of functional somatic syndromes. *Lancet* 369:946–955
24. Hessel A, Beutel M, Geyer M et al (2005) Prevalence of somatoform pain complaints in the German population. *Psychosoc Med* 2:Doc03
25. Hessel A, Geyer M, Gunzelmann T et al (2003) Somatoform complaints in elderly of Germany. *Z Gerontol Geriatr* 36:287–296
26. Kerns RD, Dobscha SK (2009) Pain among veterans returning from deployment in Iraq and Afghanistan: update on the Veterans Health Administration Pain Research Program. *Pain Med* 10:1161–1164
27. Kuwert P, Braehler E, Glaesmer H et al (2009) Impact of forced displacement during World War II on the present-day mental health of the elderly: a population-based study. *Int Psychogeriatr* 21:748–753
28. Kuwert P, Freyberger HJ (2007) The unspoken secret: sexual violence in World War II. *Int Psychogeriatr* 19:782–784
29. Kuwert P, Klauer T, Eichhorn S et al (2010) Trauma and current posttraumatic stress symptoms in elderly German women who experienced wartime rapes in 1945. *J Nerv Ment Dis* 198:450–451
30. Kuwert P, Spitzer C, Trader A et al (2007) Sixty years later: post-traumatic stress symptoms and current psychopathology in former German children of World War II. *Int Psychogeriatr* 19:955–961
31. Labbate LA, Cardena E, Dimitreva J et al (1998) Psychiatric syndromes in Persian Gulf War veterans: an association of handling dead bodies with somatoform disorders. *Psychother Psychosom* 67:275–279
32. Lahmann C, Henningsen P, Noll-Hussong M (2010) Somatoform pain disorder – overview. *Psychiatr Danub* 22:453–458
33. Lahmann C, Henningsen P, Noll-Hussong M et al (2010) Somatoform disorders. *Psychother Psychosom Med Psychol* 60:227–233 quiz 234
34. Leiknes KA, Finset A, Moum T et al (2007) Current somatoform disorders in Norway: prevalence, risk factors and comorbidity with anxiety, depression and musculoskeletal disorders. *Soc Psychiatry Psychiatr Epidemiol* 42:698–710
35. Maercker A, Forstmeier S, Enzler A et al (2008) Adjustment disorders, posttraumatic stress disorder, and depressive disorders in old age: findings from a community survey. *Compr Psychiatry* 49:113–120

-
36. Maercker A, Forstmeier S, Wagner B et al (2008) Post-traumatic stress disorder in Germany. Results of a nationwide epidemiological study. *Nervenarzt* 79:577–586
 37. Mailis-Gagnon A, Nicholson K, Yegneswaran B et al (2008) Pain characteristics of adults 65 years of age and older referred to a tertiary care pain clinic. *Pain Res Manag* 13:389–394
 38. Meyer C, Rumpf HJ, Hapke U et al (2000) Lifetime prevalence of mental disorders in general adult population. Results of TACOS study. *Nervenarzt* 71:535–542
 39. Muller JE, Wentzel I, Nel DG et al (2008) Depression and anxiety in multisomatoform disorder: prevalence and clinical predictors in primary care. *S Afr Med J* 98:473–476
 40. Nickel M, Cangoez B, Bachler E et al (2006) Bioenergetic exercises in inpatient treatment of Turkish immigrants with chronic somatoform disorders: a randomized, controlled study. *J Psychosom Res* 61:507–513
 41. Nickel R, Egle UT (2006) Psychological defense styles, childhood adversities and psychopathology in adulthood. *Child Abuse Negl* 30:157–170
 42. Noll-Hussong M, Otti A, Laeer L et al (2010) Aftermath of sexual abuse history on adult patients suffering from chronic functional pain syndromes: an fMRI pilot study. *J Psychosom Res* 68:483–487
 43. Paras ML, Murad MH, Chen LP et al (2009) Sexual abuse and lifetime diagnosis of somatic disorders: a systematic review and meta-analysis. *JAMA* 302:550–561
 44. Raphael KG, Widom CS (2011) Post-traumatic stress disorder moderates the relation between documented childhood victimization and pain 30 years later. *Pain* 152:163–169
 45. Ringel Y, Drossman DA, Leserman JL et al (2008) Effect of abuse history on pain reports and brain responses to aversive visceral stimulation: an FMRI study. *Gastroenterology* 134:396–404
 46. Rusby JS, Tasker F (2009) Long-term effects of the British evacuation of children during World War 2 on their adult mental health. *Aging Ment Health* 13:391–404
 47. Schafer I, Von Leitner EC, Schon G et al (2010) Multimorbidity patterns in the elderly: a new approach of disease clustering identifies complex interrelations between chronic conditions. *PLoS One* 5:e15941
 48. Schneider G, Heuft G (2011) Medically unexplained and somatoform complaints and disorders in the elderly: a systematic review of the literature. *Z Psychosom Med Psychother* 57:115–140
 49. Silver SM, Rogers S, Russell M (2008) Eye movement desensitization and reprocessing (EMDR) in the treatment of war veterans. *J Clin Psychol* 64:947–957
 50. Thomas HV, Stimpson NJ, Weightman A et al (2006) Pain in veterans of the Gulf War of 1991: a systematic review. *BMC Musculoskelet Disord* 7:74
 51. Valet M, Gundel H, Sprenger T et al (2009) Patients with pain disorder show gray-matter loss in pain-processing structures: a voxel-based morphometric study. *Psychosom Med* 71:49–56
 52. Wessely S, Nimnuan C, Sharpe M (1999) Functional somatic syndromes: one or many? *Lancet* 354:936–939
 53. Wittchen HU, Muller N, Pfister H et al (1999) Affective, somatoform and anxiety disorders in Germany—initial results of an additional federal survey of “psychiatric disorders”. *Gesundheitswesen* 61(Spec No):216–222
 54. Wikipedia; http://en.wikipedia.org/wiki/File:Ongoing_Conflicts.png, accessed 17 February 2012