

Debate

Open Access

Functional illness in primary care: dysfunction versus disease

Nefyn Williams*¹, Clare Wilkinson¹, Nigel Stott² and David B Menkes³

Address: ¹Department of Primary Care and Public Health, Cardiff University, North Wales Clinical School, Wrecsam, UK, ²Department of Primary Care and Public Health, Cardiff University, UK and ³Department of Psychological Medicine, Waikato Clinical School, University of Auckland, New Zealand

Email: Nefyn Williams* - williamsnh@cf.ac.uk; Clare Wilkinson - wilkinsonc@cf.ac.uk; Nigel Stott - stottoxwich@aol.com; David B Menkes - menkesd@waikatodhb.govt.nz

* Corresponding author

Published: 15 May 2008

Received: 4 October 2007

BMC Family Practice 2008, 9:30 doi:10.1186/1471-2296-9-30

Accepted: 15 May 2008

This article is available from: <http://www.biomedcentral.com/1471-2296/9/30>

© 2008 Williams et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: The Biopsychosocial Model aims to integrate the biological, psychological and social components of illness, but integration is difficult in practice, particularly when patients consult with medically unexplained physical symptoms or functional illness.

Discussion: This Biopsychosocial Model was developed from General Systems Theory, which describes nature as a dynamic order of interacting parts and processes, from molecular to societal. Despite such conceptual progress, the biological, psychological, social and spiritual components of illness are seldom managed as an integrated whole in conventional medical practice. This is because the biomedical model can be easier to use, clinicians often have difficulty relinquishing a disease-centred approach to diagnosis, and either dismiss illness when pathology has been excluded, or explain all undifferentiated illness in terms of psychosocial factors. By contrast, traditional and complementary treatment systems describe reversible functional disturbances, and appear better at integrating the different components of illness. Conventional medicine retains the advantage of scientific method and an expanding evidence base, but needs to more effectively integrate psychosocial factors into assessment and management, notably of 'functional' illness. As an aid to integration, pathology characterised by structural change in tissues and organs is contrasted with dysfunction arising from disordered physiology or psychology that may occur independent of pathological change.

Summary: We propose a classification of illness that includes orthogonal dimensions of pathology and dysfunction to support a broadly based clinical approach to patients; adoption of which may lead to fewer inappropriate investigations and secondary care referrals and greater use of cognitive behavioural techniques, particularly when managing functional illness.

Background

Our conceptual models of illness shape our consulting behaviour [1]. This paper aims to stimulate debate by developing the biopsychosocial model in primary health care, particularly with regard to undifferentiated illness.

Starfield's definition of primary care [2] as '...first contact, continuous, comprehensive and co-ordinated care provided to populations undifferentiated by gender, disease system or organ system' has achieved international acceptance. Such a broadly defined discipline requires a broad working definition of health or illness, yet such breadth is

not always evident in daily practice. A particular challenge to conventional practice is posed by medically unexplained physical symptoms (MUPS), estimated to constitute one in five primary care consultations [3], and a persistent cause of presentation in 2.5% [4]. Despite notional acceptance of the biopsychosocial model, contemporary practice tends to frame such presentations as either psychological or physical, which is compounded by current classification systems [5]. Newer approaches include: assessing, organising and treating these biological, psychological and social components within a four-dimensional grid [6]; addressing unmet psychiatric needs delivered either by psychiatrists in a collaborative care model [7], or by training primary care clinicians in a re-attribution programme [8]. In these interventions the emphasis is shifted from physical to the psychological, but true integration remains elusive. In order to make the biopsychosocial model easier to use in everyday practice, we suggest that illness in primary care be considered in two interacting dimensions:

1. Pathology; characterised by demonstrable structural change in tissues and organs
2. Dysfunction; arising from disordered physiology or psychology, which may occur independent of pathological change

Appropriate care requires that both dimensions are appraised by a skilled clinician, mindful that both may manifest, and interact, in the psychosocial context of the individual patient. Before discussing the implications of this approach for clinical practice, medical education and research, we first consider the development of the biopsychosocial model from general systems theory.

Discussion

General Systems Theory

General systems theory offers an intuitively appealing model for understanding illness presenting in primary care. In contrast to determinism, which states that every event has an antecedent cause, nature is seen as a dynamic order of interacting parts and processes [9]. In this hierarchy of systems (e.g., biochemicals to organelle, cell, tissue, organ, organism, family, community, society), there is both vertical and horizontal interaction. For example, the endocrine system interacts with both the nervous and immune systems. Each level has properties not present in its individual parts, but arising from the relationship between the parts. Thus people are more than the sum of their organs and tissues, and society more than the sum of its individuals. If equilibrium is disturbed, reactive forces reverberate in the system until a new equilibrium is established. This model can explain how each level in the hierarchy interacts with increasing complexity, and forms a

continuum from the sub-cellular level to the roles and relationships constituting society [10]. It even works to promote concepts of global connectedness [11]. Additional explanatory detail is needed at the level of the individual human being, arguably the level of greatest complexity gain due to sentience and self-concept [12,13].

The biopsychosocial model

General systems theory provides a structure for systematic thinking about a complex world, and has been used in health care to describe how the physical, psychological and social components of illness can be considered together in primary care [10,14-16], and other branches of medicine [17]. The resulting biopsychosocial model attempts to integrate these three components of illness, and can be distinguished from the biomedical model, where psychosocial aspects are considered separately, if at all [18,19]. A biopsychosocial approach requires broad definitions of health [20,21], provides a framework to address functional illness consisting of MUPS and functional somatic syndromes, as well as the importance of life events and spiritual or existential aspects of care.

Medically Unexplained Physical Symptoms (MUPS)

GPs are often presented with illness which defies biomedical understanding and thus differentiation because it is self-limiting; distorted by early treatment or changed circumstance; intermediate between existing labels; a mild or atypical variant of known pathology; not yet understood; or some combination of these [22-24]. Diary studies reveal that individuals tend to develop a new physical symptom on average every five to seven days; the vast majority of which do not result in a consultation. Those consulting doctors are more likely to have had a recent stressful life event [22]. Similarly, individuals with high levels of subjective distress are more likely to notice and complain about internal bodily sensations [25,26]. Psychological distress is linked to both the number and severity of unexplained symptoms [27] and health care use [28]. MUPS are common and are often associated with psychological morbidity, but most patients presenting with them do not have definite psychiatric illness [29,30].

Functional somatic syndromes

Clusters of unexplained symptoms are often grouped into syndromes and given 'diagnostic' labels such as irritable bowel syndrome, fibromyalgia or chronic fatigue syndrome. They have been classified according to the secondary care division of specialties and sub-specialties. Thus gastroenterologists label irritable bowel syndrome; cardiologists non-cardiac chest pain; urologists urethral syndrome etc. Many syndrome labels correspond to and can be seen to justify specialists' favourite treatments. Thus non-specific back pain may be described as 'facet syndrome' by anaesthetists who inject joints, 'somatic dys-

function' by osteopaths who treat spinal areas by manipulation, 'instability' by spinal surgeons who fuse spinal vertebrae, 'somatisation' by psychiatrists and 'illness behaviour' by psychologists. Many of these syndromes have common core clinical features [22,31], and often co-exist [32,33], so it is uncertain how much they are distinct clinical entities, or variations on a theme of general body distress [33-36]. As with MUPS, these syndromes tend to have a strong psychological component. A stepped care approach has been advocated using: non-pharmacological treatments such as exercise and cognitive behavioural therapy, which are more effective than passive treatments such as injections and operations, and pharmacological treatments targeted on the central nervous system such as tricyclic antidepressants [37].

Psychosocial factors

Psychosocial factors are also important in disease of known pathology. Social isolation, unemployment and other stressful life events are independently associated with higher mortality from all causes [38]. Depression predicts reduced survival following myocardial infarction [39,40], possibly due to an association with fatigue [41] or non-compliance [42], although other studies have failed to replicate this finding [43]. The well-established link between poverty and ill health cannot be fully explained by health behaviours or specific aetiological factors [44,45]. In a more positive sense, social capital has a pervasive beneficial effect on quality of life, measures of physical health, and resilience to stress [46]. Psychological factors, notably stress, are known to affect the neuroendocrine and immune systems [47] and have been linked to cancer and HIV progression [48,49]. Similarly, psychological disorder appears to enhance the development of inflammation and infection in atherosclerotic plaques [50] and activate autoimmune disease such as rheumatoid arthritis [51]. Neuroendocrine and immune effects of extreme stress or depression may be better understood as normal-range responses to the human condition, which in turn have deleterious effects on physical health.

Social context

One of the most important functions of medicine through the ages has been as a source of refuge for the sick. This includes hospital admission for healing, relief or protection; official sanction of the sick role in terms of certification or sick pay; refuge from distressing symptoms and the fear of serious disease by the provision of reassurance, symptom relief, and sometimes cure [52].

Spiritual dimension

The biopsychosocial model also has an interpretative function, in that it can be used as a source of meaning to individuals' experience of illness [53]. The need to find deeper meaning to life, illness and death is an ancient

characteristic of *Homo sapiens*, and the abstract thought involved in this process appears unique to our species. Closely aligned to the phenomenon is an instinct for spiritual practice and religious worship. Spiritual or existential discomfort is rarely formally considered by doctors, except in palliative care [54], or dynamic psychotherapy. Spiritual needs have been defined by one author [55] as "the needs and expectations that all humans have to find meaning, purpose, and value in life". These belief systems may or may not be part of religious faith, but 'spiritual care' is about helping people whose sense of meaning and purpose is challenged by their experience of illness [55].

Difficulties using the biosychosocial model

The biomedical model can be easier to use

Although integrating biological, psychological and social components of illness can accommodate the complexity of MUPS and functional syndromes, it is frequently misunderstood or inadequately applied in clinical practice. It is often viewed as abstract and impractical, perhaps because there is misleading simplicity in the key components of the model.

In routine clinical care, the biological, psychological and social components of illness may be interpreted and managed separately rather than in an integrated manner. Significant symptoms of anxiety or depression are present in 25-52% of primary care patients [56,57], but many go unrecognised because over half present with physical rather than psychological symptoms [58-60], and genuine integration of treatments aimed at both soma and psyche is the exception rather than the rule. This is partly due to time constraints [61], and because clinicians are creatures of habit, using rules of thumb or heuristics to guide their assessments or management decisions, which in turn are heavily influenced by their medical training, methods of classification [62] and clinical experience [63]. Unsurprisingly, many doctors find dualistic practice easier and less stressful. The biomedical model, in which "disease can be viewed independently from the person who is suffering from it, and from his or her social context" [10] remains deeply entrenched in contemporary medical practice and teaching.

Difficulty letting go of the disease-centred approach to diagnosis

There are many examples of unexplained symptoms given spurious pathological labels. This can be on clinical grounds, such as 'vertebrobasilar insufficiency' for dizziness on neck extension, or 'sciatica' for any pain radiating down the leg. It can arise from investigations revealing minor degrees of abnormality, such as mild gastritis on endoscopy accounting for epigastric pain, marginally abnormal biochemistry explaining lethargy, or mild hypertension as a cause of headache. Moreover, investigations can reveal supposedly pathological changes preva-

lent in large numbers of the asymptomatic general population, such as cervical and lumbar spondylosis on plain radiographs [64], or intervertebral disc abnormalities on magnetic resonance imaging (MRI) [65].

Over-enthusiastic pathological diagnoses can be damaging to patients in a number of ways, especially when patients collude with or even drive the process. Patients may be over-investigated, referred for unnecessary opinions and procedures, which may have serious adverse consequences [66] and waste resources [67]. Pathological labelling can increase psychological morbidity and the likelihood of somatic fixation [68]. Abnormal ideas about the aetiology of symptoms can lead to unnecessary surgery [69] and inappropriate self-management. For example, as a consequence of being told from radiographic findings that musculoskeletal symptoms are due to 'wear and tear', patients commonly avoid beneficial physical activity. Similarly, anxiety symptoms attributed to a stressful stimulus often lead to avoidance, the exact opposite of what behavioural therapy has demonstrated to be helpful.

Defining illness too narrowly in pharmacological terms

There may be a temptation in practice to explain illness entirely in pharmacological and biochemical terms. For example, sumatriptan is useful in acute migraine by interacting with serotonin receptors, so discussion of migraine aetiology and management may be restricted to the serotonin system. A similarly restricted view on the aetiology of depression and schizophrenia results from focusing on the mode of action of antidepressants and antipsychotics on monoamine neurotransmitters [70].

Dismissing illness when pathology has been excluded

Non-pathological conditions can be just as disabling as pathologies when generic outcome measures are compared across different conditions [71,72]. Despite this, there is a tendency for medical practitioners, after excluding important pathological diagnoses, to discount patients' symptoms and lose interest in their care. Qualitative studies report that doctors may resort to 'victim blaming' after pathology is excluded [73]. These reactions not only damage the doctor-patient relationship, but encourage abnormal illness behaviour, notably including the quest for 'legitimate' symptoms and investigations, thereby worsening outcomes [74,75].

Explaining all undifferentiated illness in terms of psychosocial factors

When no pathological cause can be found for unexplained symptoms, it is tempting to conclude that 'it is all in the mind', especially when there is a strong link with psychological distress. The success of cognitive behavioural therapy, psychiatric consultation within primary care [7] and re-attribution [8] to treat unexplained symptoms has led many to presume that psychological and

social factors are paramount, but psychological therapies may be rehabilitative in some patients irrespective of the primary cause of their symptoms. There is also evidence of disturbed physiology in many cases of unexplained symptoms such as fatigue [76]. Moreover, the possibility of undifferentiated illness reflecting a missed physical diagnosis should always be considered [77]. Evidence that psychological therapies may affect brain function in a manner similar to that caused by drug treatments suggests common pathways of symptom genesis and relief [78], and reinforces hypotheses of functional integration of physiology and psychology.

Classification systems for functional illness

Mental disorders are very prevalent in primary care and commonly occur together. Functional illness often exists without other mental illness [30]. Traditional classification systems developed in specialised secondary care settings concentrate on its extreme and chronic manifestations. They tend to either classify it with psychiatric disorders or as separate syndromes based on pre-defined checklists, prolonged symptom duration and after organ pathology has been excluded [63]. Newer classification systems are more useful for primary care in that they are split from psychiatry and milder and shorter duration of MUPS are included in a separate category [5]. However even this new classification does not aid integration and presumes that functional illness is a separate entity.

Homo Sapiens is body, mind and spirit

As discussed earlier, a narrowly defined clinical role can be easier and less stressful for doctors. Integrating the biological, psychological and social components of healing is difficult enough without the added time and complexity of considering a relevant spiritual or existential dimension. What insights are available from other medical systems to guide us?

Insights from traditional and complementary medical systems

Reversible functional disturbance

Nineteenth century physicians Beard and Dubois [79] described an integrated view of unexplained symptoms and syndromes, and saw them as manifestations of reversible functional disturbance. A non-judgemental, supportive, empathic approach was advocated, which used the strength of the doctor-patient relationship to provide reassurance that symptoms were likely to improve, and to encourage patient autonomy. There has been a long tradition of such a patient-centred approach in central Europe [80], advocated more recently in English-speaking countries [79]. The extended re-attribution and management model is an example of a patient-centred integrative approach used in Danish primary care [8].

Eastern medical traditions

Traditional Chinese medicine uses a highly integrated model of illness. Instead of diagnosing disease, traditional Chinese medical practitioners describe a 'pattern of disharmony' or an imbalance in the patient's body. A symptom is not traced back to its source, but is interpreted in terms of the patient's entire bodily pattern. A person who is well or 'in harmony' has no distressing symptoms and is in physical, mental and spiritual balance; in illness, the symptom is only one part of a complete bodily imbalance that can be seen in other aspects of the individual's life and behaviour [81]. Similar principles of maintaining an energetic equilibrium are involved in Ayurvedic medicine [82].

Somatic dysfunction

Osteopathy also has an integrated model of illness, based on the concept that symptoms can arise from abnormal functioning of the musculoskeletal system, not dependent on structural pathological processes [83]. Pathology can be defined by its location, structural change, and how this disturbs physiological processes; dysfunction, by contrast, is the result of the interplay of different structures in various locations. The diagnostic task in pathological diagnosis is to localise the lesion exactly, and to determine its nature; in dysfunction to determine the chain of abnormal physiology and psychology, and to assess the importance of the individual links which may be useful in explanation or management. Pathological disease alters the anatomy, sometimes microscopic, of affected tissue; whereas dysfunction in the absence of pathology is a disorder of physiology or psychology. By analogy with computers, pathology is a problem with hardware; dysfunction with software. Modern technology enables clinicians to diagnose pathological conditions more effectively, and with more objectivity. In dysfunction technology is of limited use, integrative clinical skills and effective communication are decisive [84].

Contrasting pathology with dysfunction

Dysfunction in any body system

We propose that the concept of dysfunction can be expanded to include functional syndromes in all of the body's systems, including the brain. Most psychological disturbance is a variant, or an extreme version of commonly experienced cognitions, emotions or behaviours. There is a continuum with normality. In primary mood and anxiety disorders, no structural pathological changes in the brain have been found, although abnormalities of neurotransmitter function have been postulated [85] and are consistent with the observed effects of some treatments [86]. In common mental disorders, including mood, substance use and personality disorders, it is impossible to localise the problem to any single brain structure; the disorders, or at least their manifestations,

are often reversible. In contrast, pathological disease brings about irreversible structural change in the brain's morphology, such as in chronic organic brain syndromes (including dementia caused by stroke, demyelination or Alzheimer-type degeneration) and probably also schizophrenia [87]. This distinction between pathology and dysfunction is not completely clear cut, because in some cases dysfunction appears to progress to structural pathology. There is evidence, for example, that untreated depression or seizures may lead to irreversible brain changes [88,89].

Pathology and dysfunction as separate dimensions

Evidence for links between body and mind continues to accumulate, vitiating the previously predominant Cartesian dualistic model [22]. One consequence is the biopsychosocial model described above. We propose a further extension for the classification of unexplained symptoms and functional syndromes. Although it seems intuitive that illnesses may arise predominantly from the psyche or the soma, functional illness initiation and progression typically involves both domains. Our proposed classification thus includes orthogonal dimensions of pathology and dysfunction (Figure 1). For this purpose pathology is narrowly defined as pathological processes that cause gross or microscopic structural change in any of the body's tissues, including the brain, resulting in abnormality of function. Dysfunction, by contrast, is abnormal functioning of the body, caused by, or manifested as disturbed physiological or psychological processes independent of known structural pathology.

Implications for clinical practice, audit, and research

We conclude that routinely contrasting dysfunction with disease facilitates the routine use of the biopsychosocial model in everyday clinical assessment and management, and has implications for teaching and research. Adoption in primary care should be marked by changes in consulting behaviour, some of which would be readily discernible by clinical audit, such as reductions in pathological labelling, fewer investigations and secondary care referrals. Adoption might lead to more explanation of physiological and psychological mechanisms [90,91], and greater use of non-pharmacological interventions such as cognitive behavioural therapy, graded exercise and counselling [37]. Some of these treatments may need to be provided outside conventional medical practice, but patient confidence in the therapeutic plan will be an essential ingredient, particularly if clinicians follow the recently published National Institute for Clinical Excellence (NICE) guideline on chronic fatigue syndrome, where unprecedented emphasis has been placed on the importance of management plans that are understood and approved of by the patient and their carers [92].




Figure 1
Hypothetical scatter plot of dysfunction versus pathology in primary care consultations. Dysfunction. abnormal functioning, either physiological or psychological, which is reversible and not dependent on pathological processes. **Pathology:** abnormal functioning caused by structural pathological change, either gross or microscopic.

Summary

The biological, psychological and social components of illness are seldom managed as an integrated whole in conventional medical practice. Traditional and complementary medical systems appear better at integrating these three components. As an aid to integration, pathology characterised by structural change in tissues and organs is contrasted with dysfunction arising from disordered physiology or psychology.

Abbreviations

MUPS: Medically Unexplained Physical Symptoms; HIV: Human Immunodeficiency Virus; MRI: Magnetic Resonance Imaging; NICE: National Institute of Clinical Excellence.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

This debate paper arose from a series of discussions between NW and CW concerning the nature of dysfunction in complementary medicine and its relevance to pri-

mary care, particularly in relation to medically unexplained symptoms and psychological disturbance. NW wrote an initial draft which was amended by CW and then passed on to NS and DM who made substantial contributions and changes.

Acknowledgements

We would like to acknowledge colleagues from the Department of Primary Care and Public Health, Cardiff University for their comments and suggestions on various drafts of this paper.

References

1. Kleinman A, Eisenberg L, Good B: **Culture, illness and care. Clinical lessons from anthropological and cross-cultural research.** *Ann Int Med* 1978, **88**:251-258.
2. Starfield B: **Is primary care essential?** *Lancet* 1994, **344**:1129-1133.
3. Peveler R, Kilkenny L, Kinmouth A: **Medically unexplained physical symptoms in primary care: a comparison of self-report screening questionnaires and clinical opinion.** *J Psychosom Res* 1997, **42**:245-252.
4. Verhaak PFM, Meijer SA, Visser AP, Walters G: **Persistent presentation of medically unexplained symptoms in general practice.** *Fam Prac* 2006, **23**:414-420.
5. Rosendal M, Fink P, Falkoe E, Hansen HS, Olesen F: **Improving the classification of medically unexplained symptoms in primary care.** *Eur J Psychiat* 2007, **21**:25-36.

6. de Jonge P, Huysse FJ, Slaets JJP, Sollner W, Stiefel FC: **Operationalization of biopsychosocial case complexity in general health care: the INTERMED project.** *Aust NZ J Psychiatry* 2005, **39**:795-799.
7. Feltz-Cornelis CM van der, van Oppen P, Ader HJ, van Dyck R: **Randomised controlled trial of a collaborative care model with psychiatric consultation for persistent medically unexplained symptoms in general practice.** *Psychother psychosom* 2006, **75**:282-289.
8. Fink P: **Assessment and treatment of functional disorders in general practice: the extended reattribution and management model – an advanced educational program for nonpsychiatric doctors.** *Psychosomatics* 2002, **43**:93-131.
9. von Bertalanffy L: *General systems theory* New York: George Braziller; 1968.
10. McWhinney IR: *A textbook of family medicine* 2nd edition. New York: Oxford University Press; 1997.
11. Wroe M, Doney M: *The rough guide to a better world* London: Rough Guides Ltd; 2004.
12. Galin D: **The structure of subjective experience: Sharpen the concepts and terminology.** In *Toward a science of consciousness: The first Tucson discussions and debates. Complex adaptive systems* Edited by: Hameroff SRK, Alfred W, Scott AC. Cambridge MA US: MIT Press; 1996:121-140.
13. Chambers T: **Theory and the organic bioethicist.** *Theoretical Med Bioethics* 2001, **22**:123-134.
14. Stott NCH, Davis RH: **The exceptional potential in each primary care consultation.** *J Roy Coll Gen Pract* 1979, **29**:201-205.
15. *Report 27 – The nature of general medical practice* London: RCGP; 1996.
16. Olesen F, Dickinson J, Hjortdahl P: **General practice – time for a new definition.** *BMJ* 2000, **320**:354-357.
17. Waddell G: **A new clinical model for the treatment of low back pain.** *Spine* 1987, **12**:632-644.
18. Engel GL: **The need for a new medical model: a challenge for biomedicine.** *Science* 1977, **196**:129-135.
19. Engel GL: **The clinical application of the biopsychosocial model.** *Am J Psychiatry* 1980, **137**:535-543.
20. World Health Organisation: *Constitution* Geneva: WHO; 1946.
21. Seedhouse D: *Liberating medicine* Chichester: John Wiley & Sons; 1991.
22. Hungin P: **The therapeutic illusion: self-limiting illness.** In *Oxford Textbook of Primary Medical Care* 1st edition. Edited by: Jones R, Britten N, Culpepper L, Gass DA, Grol R, Mant D, Silagy C. Oxford: Oxford University Press; 2004:210-214.
23. Kroenke K, Mangelsdorff D: **Common symptoms in ambulatory care: incidence, evaluation, therapy and outcome.** *Am J Med* 1989, **86**:262-266.
24. Crimlisk HL, Bhatia K, Cope H, David A, Marsden CD, Ron MA: **Slater revisited: 6 year follow-up study of patients with medically unexplained motor symptoms.** *BMJ* 1998, **316**:582-586.
25. Pennebaker JW: *The psychology of physical symptoms* New York: Springer-Verlag; 1983.
26. Barsky AJMD, Borus JFMD: **Functional Somatic Syndromes.** *Ann Intern Med* 1999, **130**:910-921.
27. Katon WW, Sullivan M, Walker E: **Medical symptoms without identified pathology: Relationship to psychiatric disorders, childhood and adult trauma, and personality traits.** *Ann Intern Med* 2001, **134**:917-925.
28. Frostholm L, Fink P, Christensen KS, Toft T, Oernboel E, Olesen F, Weinman J: **The patients' illness perceptions and the use of primary health care.** *Psychosom Med* 2005, **67**:997-1005.
29. Burton C: **Beyond somatisation: a review of the understanding and treatment of medically unexplained physical symptoms (MUPS).** *Br J Gen Pract* 2003, **53**:231-239.
30. Toft T, Fink P, Oernboel E, Christensen K, Frostholm L, Olesen F: **Mental disorders in primary care: prevalence and co-morbidity among disorders. Results from the functional illness in primary care (FIP) study.** *Psychol Med* 2005, **35**:1175-1184.
31. Aaron LA, Buchwald D: **A review of the evidence for overlap among unexplained clinical conditions.** *Ann Intern Med* 2001, **134**:868-881.
32. Jones R, Lydeard S: **Irritable bowel syndrome in the general population.** *BMJ* 1992, **304**:87-90.
33. Wessely S, Nimnuan C, Sharpe M: **Functional somatic syndromes; one or many?** *Lancet* 1999, **354**:936-939.
34. Fink P, Toft T, Hansen MS, Ornbol E, Olesen F: **Symptoms and syndromes of bodily distress: an exploratory study of 978 internal medical, neurological and primary care patients.** *Psychosomatic Med* 2007, **69**:30-39.
35. Nimnuan C, Rabe-Hesketh S, Wessely S, Hotopf M: **How many functional somatic syndromes?** *J Psychosom Res* 2001, **51**:549-557.
36. Kanaan RAA, Lepine JP, Wessely SC: **The association or otherwise of the functional somatic syndromes.** *Psychosom Med* 2007, **69**:855-859.
37. Henningsen P, Zipfel S, Herzog W: **Management of functional somatic syndromes.** *Lancet* 2007, **369**:946-955.
38. Fiscella K: **Is lower income associated with greater biopsychosocial morbidity? Implications for physicians working with underserved patients.** *J Fam Pract* 1999, **48**:372-377.
39. Frasure-Smith N, LesPerance F, Telajac M: **Depression following MI.** *JAMA* 1993, **270**:1819-1825.
40. Guck TP, Kavan MG, Elsasser GN, Barone EJ: **Assessment and treatment of depression following myocardial infarction.** *Am Family Physician* 2001, **64**:641-648.
41. Irvine J, Basinski A, Baker B, Jandciu S, Paquette M, Cairns J, Connolly S, Roberts R, Gent M, Dorian P: **Depression and risk of sudden cardiac death after acute myocardial infarction: testing for the confounding effects of fatigue.** *Psychosom Med* 1999, **61**:729-737.
42. Ziegelstein RC, Fauerbach JA, Stevens SS, Romanelli J, Richter DP, Bush DE: **Patients with depression are less likely to follow recommendations to reduce cardiac risk during recovery from a myocardial infarction.** *Arch Int Med* 2000, **160**:1818-1823.
43. Lane D, Carroll D, Ring C, Beevers DG, Lip GY: **Mortality and quality of life 12 months after myocardial infarction: effects of depression and anxiety.** *Psychosom Med* 2001, **63**:221-30.
44. Sloggett A, Joshi H: **Higher mortality in deprived areas: community or personal disadvantage?** *BMJ* 1994, **309**:1470-1474.
45. Smith GD, Morris J: **Income inequality and mortality: why are they related?** *BMJ* 1996, **312**:987-988.
46. Kawachi I: **Social capital and community effects on population and individual health.** In *Socioeconomic status and health in industrial nations: Social, psychological and biological pathways* Edited by: Adler NE, Marmot M, McEwen BS, Stewart J. New York: New York Academy of Sciences; 1999:120-130.
47. Haleem S, Kaplan H, Pollack M: **Moderating effects of gender and vulnerability on the relationships between financial hardship, low education and the immune response.** *Stress Med* 2000, **16**:167-177.
48. Garsson B, Goodkin K: **On the role of immunological factors as mediators between psychosocial factors and cancer progression.** *Psychiatry Research* 1999, **85**:51-61.
49. Leserman J: **The Effects of Stressful Life Events, Coping, and Cortisol on HIV Infection.** *CNS Spectrums* 2003, **8**:25-30.
50. Sher L: **The role of the immune system and infection in the effects of psychological factors on the cardiovascular system.** *Can J Psychiatry* 1998, **43**:954-955.
51. Zautra AJ, Hoffman JM, Matt KS, Yocum D, Potter PT, Castro WL, Roth S: **An examination of individual differences in the relationship between interpersonal stress and disease activity among women with rheumatoid arthritis.** *Arthritis Care Res* 1998, **11**:271-279.
52. Stott NCH: *Primary health care; bridging the gap between theory and practice* Berlin: Springer-Verlag; 1983.
53. Toon PD: *Towards a philosophy of general practice: a study of the virtuous practitioner* Occasional Paper 78: RCGP; 1999.
54. Zhukovsky DS: **A model of palliative care: the palliative medicine program of the Cleveland Clinic Foundation.** *World Health Organisation Demonstrations Project* 2000, **8**:268-77.
55. Murray SA, Kendall M, Boyd K, Worth A, Benton TF: **General practitioners and their possible role in providing spiritual care: a qualitative study.** *Brit J Gen Pract* 2003, **53**:957-959.
56. Spitzer RL, Williams JB, Kroenke KE, Linzer M, deGruy FV, Hahn SR, et al.: **Utility of a new procedure for diagnosing mental disorders in primary care. The PRIME-MD 1000 study.** *JAMA* 1994, **272**:1749-1756.
57. Kessler D: **Cross sectional study of symptom attribution and recognition of depression and anxiety in primary care.** *BMJ* 1999, **318**:436-439.

58. Goldberg DP, Bridges K: **Somatic presentations of psychiatric illness in primary care setting.** *J Psychosom Res* 1988, **32**:137-144.
59. Bridges K, Goldberg D, Evans B, Sharpe T: **Determinants of somatization in primary care.** *Psychol Med* 1991, **21**:473-483.
60. Zimmerman C, Tansella M: **Psychosocial factors and physical illness in primary care: promoting the biopsychosocial model in medical practice.** *J Psychosom Res* 1996, **40**:351-358.
61. Zantinge EM, Verhaak PFM, Kerstens JJ, Bensing JM: **The workload of GPs: consultations of patients with psychological and somatic problems compared.** *Br J Gen Pract* 2005, **55**:604-614.
62. Kahneman D, Tversky A: **On the reality of cognitive illusions.** *Psych Review* 1996, **103**:582-591.
63. Fink P, Rosendal M, Olesen F: **The classification of somatisation and functional somatic syndromes in primary care.** *Aust NZ J Psychiatry* 2005, **39**:772-781.
64. Van Tulder MW, Assendelft WJJ, Koes BW, Bouter LM: **Spinal radiographic findings and non-specific low back pain.** *Spine* 1997, **22**:427-434.
65. Jarvik JG, Hollingworth W, Heagerty P, Haynor DR, Deyo RA: **The longitudinal assessment of imaging and disability of the back (LAIDBack) study.** *Spine* 2001, **26**:1158-1166.
66. Mayou R, Sharpe M: **Treating medically unexplained physical symptoms.** *BMJ* 1997, **315**:561-562.
67. Law GT, Wong CY, Chan YC, Shum TT, Lok PS, Wong KY: **Cost-effectiveness of CT thorax and bronchoscopy in haemoptysis with normal CXR for exclusion of lung cancer.** *Australasian Radiology* 2002, **46**:381-383.
68. Kouyanou K, Pither CE, Wessely S: **Iatrogenic factors and chronic pain.** *Psychosom Med* 1997, **59**:597-604.
69. Fink P: **Surgery and medical treatment in persistent somatizing patients.** *J Psychosom Res* 1992, **36**:439-447.
70. Healy D: *Let them eat Prozac* Toronto: James Lorimer; 2003.
71. Creed F, Ratcliffe J, Fernandez L, Tomenson B, Palmer S, Rigby C, Guthrie E, Read N, Thompson D: **Health-related quality of life and health care costs in severe, refractory irritable bowel syndrome.** *Ann Intern Med* 2001, **134**:860-868.
72. Fanuele JC, Birkmeyer NJO, Abdu WA, Toteson TD, Weinstein J: **The impact of spinal problems on the health status of patients: Have we underestimated the effect?** *Spine* 2000, **25**:1509-1514.
73. Pill R, Stott N: **Concepts of illness causation and responsibility: some preliminary data from a sample of working class mothers.** *Soc Sci Med* 1982, **16**:43-52.
74. Smith RC: **A clinical approach to the somatizing patient.** *J Family Pract* 1985, **21**:294-301.
75. Pridmore S, Skerritt P, Ahmadi J: **Why do doctors dislike treating people with somatoform disorder?** *Australasian Psychiatry* 2004, **12**:134-138.
76. Chaudhuri A, Behan PO: **Fatigue in neurological disorders.** *Lancet* 2004, **364**:978-988.
77. de Torrente de la Jara G, Pecoud A, Favrat B: **Musculoskeletal pain in female asylum seekers and hypovitaminosis D3.** *BMJ* 2004, **329**:156-157.
78. Beutel ME, Stern E, Silbersweig DA: **The emerging dialogue between psychoanalysis and neuroscience: neuroimaging perspectives.** *J Amer Psychoanalytic Assoc* 2003, **51**:773-801.
79. Sharpe M, Carson A: **"Unexplained" somatic symptoms, functional syndromes, and somatization: do we need a paradigm shift?** *Ann Intern Med* 2001, **134**:926-930.
80. Sollner W, Stix P, Stein B, Franz M, Lampe A, Herzog T: **Quality criteria for psychosomatic consultation-liaison service.** *Wiener Medizinische Wochenschrift* 2002, **152**:528-534.
81. Kaptchuk TJ: *Chinese medicine: the web that has no weaver* London: Rider; 1983.
82. Chopra A, Doiphode W: **Ayurvedic medicine. Core concept, therapeutic principles, and current relevance.** *Med Clin North Am* 2002, **86**:75-89.
83. Greenman PE: *Principles of manual medicine* 2nd edition. Baltimore MA: Williams and Wilkins; 1996.
84. Lewit K: *Manipulative therapy in rehabilitation of the locomotor system* 2nd edition. Oxford: Butterworth-Heinemann; 1991.
85. Diehl DJ, Gershon S: **The role of dopamine in mood disorders.** *Comprehensive Psychiatry* 1992, **33**:115-120.
86. Yatham LN, Liddle PF, Shiah IS, Lam RVV, Ngan E, Scarrow G, Imperial M, Stoessl J, Sossi V, Ruth TJ: **PET study of [¹⁸F] 6-fluoro-L-dopa uptake in neuroleptic and mood-stabilizer-naïve first-episode nonpsychotic mania: Effects of treatment with divalproex sodium.** *Amer J Psychiatry* 2002, **159**:768-774.
87. Haroutunian V, Davis KL: **Introduction to the Special Section: myelin and oligodendrocyte abnormalities in schizophrenia.** *Int J Neuropsychopharm* 2007, **10**:499-502.
88. Palsson S, Aevansson O, Skoog I: **Depression, cerebral atrophy, cognitive performance and incidence of dementia. Population study of 85-year-olds.** *Brit J Psychiatry* 1999, **174**:249-253.
89. Lloyd AJ, Grace JB, Jaros E, Perry RH, Fairbairn AF, Swann AG, O'Brien JT, McKeith IG: **Depression in late life, cognitive decline and white matter pathology in two clinico-pathologically investigated cases.** *Int J Geriatric Psychiatry* 2001, **16**:281-287.
90. Thomas KB: **General practice consultations: is there any point in being positive?** *BMJ* 1987, **294**:1200-1202.
91. Dowrick C, Katona C, Peveler R, Lloyd H: **Somatic symptoms and depression: diagnostic confusion and clinical neglect.** *Br J Gen Pract* 2005, **55**:829-830.
92. Turnbull N, Shaw EJ, Baker R, Dunsdon S, Costin N, Britton G, Kuntze S, Norma R: *Chronic fatigue syndrome/myalgic encephalomyelitis (or encephalopathy): diagnosis and management of chronic fatigue syndrome/myalgic encephalomyelitis (or encephalopathy) in adults and children* London: Royal College of General Practitioners; 2007.

Pre-publication history

The pre-publication history for this paper can be accessed here:

<http://www.biomedcentral.com/1471-2296/9/30/prepub>

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

