

Water Deficit Stress Tolerance in Some of Barley Genotypes and Landraces under Field Conditions

Alireza POUR ABOUGHADAREH^{1*}, Mohammad Reza NAGHAVI², Marouf KHALILI³

¹Payame Noor University of Ilam Center, Department of Agronomy, Ilam, Iran; a.paraboghadareh@yahoo.com (*corresponding author)

²Payame Noor University of Ghare Aghaj Center, Department of Agronomy, East Azerbaijan, Iran

³Payame Noor University of Mahabad Center, Department of Agronomy, West Azerbaijan, Iran

Abstract

In order to investigate the effects of water deficit stress on some agro-morphological and physiological traits of seven genotypes and five landraces of barley, a split plot experiment was conducted a completely randomized block design with four replications in 2011-2012 cropping season at experimental field of College of Agriculture, Payame Noor University of Center, West Azerbaijan, Iran. Analysis of variance revealed significant differences among genotypes and landraces for all of the studied traits, and showed highly significant effects of water deficit stress on all the studied traits. Based on correlation analysis, all the characters included in the study except leaf temperature and number of grain per spike showed significant positive correlation with grain yield under both conditions. In terms of physiological characters high significant correlation coefficient was found between relative water content and grain yield under stress condition; however, high significant correlation coefficient observed between fluorescence chlorophyll and grain yield under non-stress condition. 1000-grain weight had negative correlation with all characters except leaf temperature under non-stress condition. However, this character had significant and positive correlation with fluorescence chlorophyll, relative water content, SPAD value, spike length, number of spike per plant under stress condition. Considering the grain yield potential 'Naghadeh', 'Piranshahr' and 'Mahabad' landraces together with 'CW3117-77' genotype were ranked as the superior group of drought tolerant barley. On the other hand, 'Naghadeh' landrace was identified as a barley landrace that could be grown under both normal and water deficit stress conditions with high grain yield.

Keywords: agro-morphological traits, barley, correlation, drought tolerance, physiological characteristics

Introduction

Barley (*Hordeum vulgare* L.) is a major crop ranked fourth in the world-wide production of cereals. It is considered a primary staple food or feed crop in the semi-arid tropics of Asia, Africa, and South America. The grain is normally used as food and animal fodder, but recently it has been used as raw material for the production of beer. Barley is typically cultivated in the arid and semi-arid regions of Iran generally in areas with low precipitation that are not suitable for wheat (Baik and Ullrich, 2008). Drought is a significant limiting factor for agricultural productivity and generally inhibits plant growth through reduced water absorption and nutrient uptake. For improving the drought tolerance of crop varieties by plant breeding, it is necessary to identify genotypes with tolerance to drought stress during all growth stages. Landraces are still cultivated in traditional crop-growing areas. There is renewed interest in landraces and primitive cultivars as important sources of genetic variation (Brush, 1995) mainly because of the trend toward greater uniformity that has narrowed the genetic base of modern cultivars, thus increasing their vulnerability to biotic and abiotic stress (Moghaddam *et al.*, 1997).

Decreased water availability generally results in reduced growth and final yield in crop plants. Plant drought tolerance is a highly complex trait that involves multiple genetic, physiological and biochemical mechanisms (Baik and Ullrich, 2008; Erdei *et al.*, 2002). Drought affects morphological, physiological, biochemical and molecular processes in plants resulting in growth inhibition. The extent of these changes is dependent on the time, stage and severity of environmental stress (Cao *et al.*, 2011). Measurements of different physiological processes of plants responses to drought are important information on the various strategies of the plant intended to remove or to reduce the harmful effects of water deficit in soil or plant tissues. Water deficit conditions cause water losses within the plant and result in relative water content (RWC) reduction. Therefore, RWC is widely used as one of the most reliable indicators for defining both the sensitivity and the tolerance of plants to water deficit (Rampino *et al.*, 2006; Sanchez-Rodriguez *et al.*, 2010). Rong-Hua *et al.* (2006) concluded that chlorophyll content could be considered as a reliable indicator in screening barley genotypes for drought tolerance. Experiments with a host of plants and different photosynthetic metabolism processes, which can be induced by varieties of plants and many biotic and abi-

otic factors, can directly or indirectly produce modification to fluorescence induction kinetics (Crudace, 2000; Percival and Baker, 1991). In addition, Slapakauskas and Ruzgas (2005) reported that measuring of chlorophyll provides information on quantitative and quantitative changes in photosynthesis.

The search for traits related to drought tolerance is an important step in cereals breeding and production. Field experiments investigating the yields of different cultivars under water deficit conditions are the most reliable way to assess their drought tolerance. Dencic *et al.* (2000) reported that many morphological and physiological characteristics were affected by drought stress. Also, they reported agronomic traits such as grain yield and its components are the major selection criteria for evaluating drought tolerance under field condition. Gupta *et al.* (2001) studied physiological and yield attributes of two wheat genotypes with stress at boot and anthesis. They indicated that number of grains, grain yield, biological yield, and harvest index decreased to a greater extent when water stress was imposed at anthesis stage. Therefore, physiological, agromorphological and biochemical approaches have a great importance in order to understand the complex responses of plants to water deficiency and develop rapidly new varieties. A physiological approach would be the most attractive way to develop new varieties (Araus *et al.*, 2008) but breeding for specific, sub-optimal environments involves a deeper understanding of yield-determining process.

Generally, different strategies have been proposed for the selection of relative drought tolerance and resistance, so, some researchers have proposed selection under non-stress conditions (Betran *et al.*, 2003; Rajaram and Van Ginkle, 2001; Richards, 1996), others have suggested selection in the target stress conditions (Ceccarelli and Grando, 1991; Rathjen, 1994) while, several of them have chosen the midway and believe in selection under both non-stress and stress conditions (Byrne *et al.*, 1995; Clarke *et al.*, 1992; Fernandez, 1992; Fischer and Maurer, 1978). In addition to, indices such as SSI (Fischer and Maurer, 1978), TOL (Rosielle and Hamblin, 1981), STI and GMP (Fernandez, 1992), DRI (Bidinger *et al.*, 1978), MSTI (Farshadfar and Sutka, 2002) have been reported for selection of drought tolerance genotypes.

In the present study, five landraces together seven genotypes of barley were used to assess drought tolerance. The drought tolerance has been evaluated by physiological

indices, i.e. leaf RWC, RCC, chlorophyll fluorescence and agro-morphological characteristics related to grain yield, under water deficit stress conditions.

Materials and methods

Plant material and growth conditions

To investigate the effects of water deficit stress, a research was carried out at experimental field of College of Agriculture, Payame Noor University of Mahabad Center, West Azerbaijan, Iran (latitude 36.46°N, longitude 45.43°E, Altitude 1385 m above sea level) during growing season of 2011-2012. The climate is characterized by mean annual precipitation of 330 mm, mean annual temperature of 12°C. Plant materials consisted of twelve genotypes and landraces of barley provided from Agricultural and Natural Resources Research Center of West Azerbaijan, Iran. The names of used in this investigation are shown in Tab. 1.

Experiment was conducted in split plot within a randomized complete block design with four replications. The experimental treatments consisted of irrigation levels as the main plot at second levels: irrigation after 70 mm evaporation from class A pan (without stress), irrigation after 150 mm evaporation from class A pan (water deficit stress) and twelve genotypes and landraces of barley as the sub plot were considered. Each plot contained 4 rows with 25 cm apart and 1m in length. All plots were irrigated after sowing and subsequent irrigations in beginning in tillering. Weeds were controlled by hand during crop growth and development.

Crop sampling and calculation

Agronomic characteristics and physiological criteria including: spike length (cm), number of spike per plant, number of grain per spike, 1000-grains weight (gr), grain yield (gr per plant), leaf temperature, fluorescence chlorophyll, relative chlorophyll content (SPAD value) and relative water content (RWC) were measured after of physiology maturity by selected 10 plants of each experimental plot randomly. For measuring physiological criteria was used the flag leaf. The chlorophyll content in the flag leaf was determined using a chlorophyll meter (SPAD-502, Japan). Five flag leaves of each cultivar grown in stress and non-stress conditions were measured after tillering stage. Three measurements in the middle of the flag leaf were

Tab. 1. Names of barley genotypes/landraces studied in this experiment

Code	Name	Code	Name
1	'Badia Kavirr' (genotype)	7	'CB' (genotype)
2	'168.4 Lign131/Abrabi Abiad' (genotype)	8	'Sahand' (landrace)
3	'Mahabad' (landrace)	9	'Makoobi' (landrace)
4	'Rihaner' (genotype)	10	'Piranshahr' (landrace)
5	'Naghadeh' (landrace)	11	'CW3117-77-5-9-5' (genotype)
6	'Atlas46/Kavir' (genotype)	12	'Rihane-05' (genotype)

made randomly for each plant, and the average sample was used for analysis. Relative water content was determined according to Turner (1986), where fresh leaves were taken from each cultivar and each replication after tillering stage and weighed immediately to record fresh weight (FW). Then they were placed in distilled water for 4 h and weighed again to record turgid weight (TW). After that subjected to oven drying at 70°C for 24h to record dry weight (DW). The RWC was calculated using the following equation:

$$RWC = ((FW - DW) / (TW - DW)) \times 100$$

The chlorophyll fluorescence was measured by a chlorophyll fluorometer (Opti Science, OS-30MSA). Also, leaf temperature measurement were made using the infrared thermometer.

Statistical analysis

Simple analysis variance was performed for data using SPSS software. Mean comparisons were conducted using Duncan's multiple rang test. Percentage of reduction in characteristics due to drought stress was calculated as following:

$$C = \frac{\bar{X}ns - \bar{X}ds}{\bar{X}ns} \times 100$$

Where $\bar{X}ns$ the mean of characteristic in given genotype/landrace under non-stress condition and $\bar{X}ds$ is the mean of characteristic in given genotype/landrace under water deficit stress condition. Analysis of correlation coefficient between grain yield and other characteristics was used to determine the principle components influencing final grain yield (Fayaz and Arzani, 2011).

Stress susceptibility index (SSI) was calculated for each cultivar following Fischer and Maurer (1978):

$$SSI = (1 - Yds/Yns) / (1 - (\bar{Y}ns/\bar{Y}ns))$$

Where Yds and Yns are the grain yield under stress and non-stress conditions, $\bar{Y}ns$ and $\bar{Y}ns$ are the average grain yield of all genotype/landrace under stress and non-stress conditions.

Results and discussion

Analysis of variance revealed significant ($p \leq 0.01$) differences among genotypes and landraces for all of the studied traits except of 1000-grains weight, and showed highly significant effects of water deficit stress on all the studied traits (Tab. 2). Water deficit stress \times genotypes interactions effect was also highly significant ($p \leq 0.01$) for all traits, suggesting different response of genotypes/landraces to each conditions. The maximum and minimum value of coefficient of variation belonged to number of spike per plant and plant height, respectively.

Means of agro-morphological and physiological characteristics under stress and non-stress conditions as well as reduction percent of reduction in the concerned characteristics due to water deficit stress are shown in Tab. 3. Leaf temperature ranged from 22.70 for 'Atlas46/Kavir' genotype and 'Sahand' landrace to 25.62 for 'CB' genotype under non stress condition; however, leaf temperature were highest under stress condition and ranged from 27.50 for 'Naghadeh' to 30.50 for 'Rihane-05'. Fluorescence chlorophyll ranged from 0.74 for 'CB' genotype to 0.80 for 'Atlas46/Kavir' genotype and 'Sahand' landrace under stress conditions, however, this characteristics were lowest under stress condition and ranged from 0.69 for 'Rihane-05' to 0.74 for Naghadeh landrace. Also, SPAD value ranged from 39.95 to 43.07 for as46/Kavir under non-stress condition and ranged from 33.150 for 'Rihane-05' to 40.77 for 'Naghadeh' under stress condition. The highest reduction (21.90%) in SPAD value due to stress was observed in 'Rihane-05' genotype. Under non-stress condition RWC ranged from 73.32 for 'CB' genotype to 81.46 for 'Sahand'; however, under stress condition this characteristic ranged from 69.08 for 'Rihane-05' to 78.93 for 'Naghadeh' landrace. The decline in RWC in 'Rihane-05' and 'Atlas46/Kavir' genotypes due to the stress was significantly less than all other genotypes and landraces. Decline of RWC reported in many researches (Farshadfar, 2012; Farshadfar et al., 2012). Ahmadi et al. (2012) showed that, RWC decline due to stress conditions. Plant height ranged from

Tab. 2. Analysis of variance for agro-morphological traits in genotypes/landraces of barley grown under water deficit stress and normal conditions

Source of variation	df	MS									
		SL	NGS	NS	GW	Y	Ph	Tem	Flu	RCC	RWC
Replication	3	0.19	21.25	0.84	18.43	1.61	0.56	1.28	≈0†††	0.62	0.47
Stress	1	76.50**	135.37	140.16**	4501.82**	439.59**	1230.51**	666.76**	0.09**	553.44**	979.24**
Error 1	3	0.16	21.51	0.64	17.49	1.36	0.73	1.031	≈0††	0.716	0.46
Genotype/Landrace	11	1.30**	25.80**	1.61**	23.77	2.12*	271.63**	2.176**	0.001**	7.13**	13.50**
Interaction effect	11	2.89**	26.78**	3.92**	25.99*	5.44**	27.28**	4.221**	0.001**	13.08**	23.55**
Error 2	66	0.42	8.93	0.58	12.93	0.91	0.23	0.59	≈0†	1.55	3.11
CV (%)		7.84	8.49	21.96	10.45	21.86	0.88	2.9	1.72	3.16	2.31

*and **: Significant at the 0.05 and 0.01 probability level, respectively; SL, NGS, NS, GW, Y and Ph indicate; spike length, number of grain per spike, 1000-grains weight, grain yield per plant and plant height, respectively
 †, †† and ††† indicate; 1.6E-4, 7.3E-5 and 1.05E-4; Tem, Flu, RCC and RWC indicate; leaf temperature, fluorescence chlorophyll, relative chlorophyll content and relative water content, respectively

50.55 for 'Rihane-05' to 67.48 for 'Badia Kavirr' genotype under non-stress condition and ranged from 40.85 for '168.4 Lign131' genotype to 59.48 for 'Mahabad' landrace under stress conditions. The highest reduction (29.29%) in plant height due to stress was observed in '168.4 Lign131' genotype. The decrease in plant height under drought stress condition could be due to decrease in relative turgidity and dehydration of protoplasm which is associated with loss of turgor and reduced cell division and cell expansion (Bayoumi *et al.*, 2008). Number of grains per spike ranged from 31.75 for 'Rihane' genotype to 35.50 for 'Badia Kavirr' under non-stress condition, and ranged from 31.50 for 'Naghadeh' landrace to 43.50 for

'Rihane-05' under stress condition. The highest reduction (2.87%) in grains per spike due to stress was observed in 'Piranshahr' landrace. In the study conducted by Pour Aboughadareh (2012), in the wild wheat number of grains per spike showed lower reduction under stress conditions than other component yield. 'Rihane' and 'Atlas46/Kavir' barley genotype produced the highest 1000-grain weight and spike length under non-stress condition, while under stress condition the highest 1000-grain weight and spike length related to 'Naghadeh' landrace and 'CB' genotype, respectively. The highest reduction in spike length and 1000-grain weight due to stress condition was observed in 'Rihane-05' genotype and 'Naghadeh' landrace, respec-

Tab. 3. Mean of agro-morphological and physiological characteristics in genotypes and landraces of barley under normal (N) and water deficit stress (S) conditions and percent of reduction (%R)

Code		Tem	Flu	RCC	RWC	SL	NS	NGS	GW	Ph	Y
1	N	23.12ab	0.79ab	42.62ab	80.76ab	9.65ab	5.25ab	35.50a	38.82ab	67.48a	7.17ab
	S	30.12cde	0.70cd	0.70cde	35.15cd	6.63cde	1.25cd	38.00bc	26.30bcd	64.20a	1.24cd
	%R	-30.27	11.11	11.11	17.54	31.35	76.19	-7.04	32.26	4.85	82.79
2	N	25.12de	0.75cd	40.67ef	77.48ef	8.58cd	3.50c	33.50a	42.35ab	57.78f	4.94d
	S	29.50bcde	0.71bc	0.71bcd	36.30bcd	7.13bcd	2.00bcd	37.75bc	26.37bcd	40.85k	1.98cd
	%R	-17.41	5.45	5.46	10.76	16.91	42.86	-12.69	37.72	29.29	59.89
3	N	24.37bcde	0.76bcd	41.20cde	78.63cde	8.80bcd	4.25bc	35.00a	38.80ab	64.28b	5.77bcd
	S	29.12bc	0.71bc	0.71bc	37.35bc	7.60bc	2.25bc	35.50bcd	28.07abc	59.48b	2.22bc
	%R	-19.48	6.47	6.48	9.34	13.64	47.06	-1.43	27.64	7.47	61.53
4	N	24.62cde	0.76cd	41.07def	78.16def	8.93bcd	4.00bc	31.75a	45.57a	59.65e	5.79bcd
	S	28.50b	0.72b	0.72ab	38.40b	8.00ab	3.00b	32.25d	31.15a	50.83f	2.98b
	%R	-15.73	4.60	4.61	6.51	10.36	25.00	-1.57	31.65	14.79	48.47
5	N	23.37abc	0.78ab	42.27abcd	80.38abcd	9.43abc	4.75abc	35.75a	38.07b	60.70d	6.45abcd
	S	27.50a	0.74a	0.74a	40.77a	9.03a	4.25a	31.50d	31.47a	55.60d	4.21a
	%R	-17.64	5.90	5.91	3.55	4.24	10.53	11.89	17.33	8.40	34.65
6	N	22.75a	0.80a	43.07a	81.50a	10.10a	5.75a	32.00a	45.17ab	55.48h	8.25a
	S	30.25de	0.70cd	0.70de	34.70cd	6.48de	1.25cd	40.75ab	24.35cd	49.13g	1.24cd
	%R	-32.96	12.48	12.48	19.44	35.89	78.26	-27.34	46.10	11.45	84.95
7	N	25.62e	0.74d	39.95f	77.32e	8.35d	3.50c	33.50a	42.85ab	60.45d	4.97cd
	S	28.50b	0.72b	0.72ab	38.60b	8.15ab	3.00b	34.50cd	28.60abc	51.45f	2.95b
	%R	-11.21	2.58	2.58	3.38	2.40	14.29	-2.99	33.26	14.89	40.67
8	N	22.75a	0.80a	42.77ab	81.46ab	10.00a	5.75a	33.25a	42.37ab	57.28g	8.05a
	S	29.25bcd	0.71bc	0.71bcd	36.52bcd	7.20bcd	2.00bcd	36.75bcd	27.32abc	53.45e	1.96cd
	%R	-28.57	11.01	11.01	14.61	28.00	65.22	-10.53	35.52	6.68	75.65
9	N	24.12abcd	0.77bc	41.62bcde	79.40bcde	8.95bcd	4.75abc	34.00a	41.07ab	54.40i	6.58abcd
	S	29.12bc	0.72bc	0.72bc	37.27bc	7.48bcd	2.25bc	36.25bcd	27.35abc	46.55h	2.21bc
	%R	-20.72	7.28	7.29	10.45	16.48	52.63	-6.62	33.41	14.43	66.36
10	N	23.37abc	0.79ab	42.22abcd	80.53abcd	9.28abcd	4.75abc	34.75a	39.52ab	61.50c	6.48abcd
	S	28.62b	0.72b	0.72b	38.20bc	8.03ab	2.50b	33.75cd	29.55ab	57.50c	2.45b
	%R	-22.45	8.37	8.37	9.53	13.48	47.37	2.88	25.24	6.50	62.16
11	N	23.75abcd	0.78ab	42.17abcd	80.33abcd	9.30abcd	5.00ab	34.25a	40.85ab	52.18j	6.92abc
	S	28.62b	0.72b	0.72b	38.07b	7.95ab	2.50b	36.25bcd	27.40abc	44.03i	2.47b
	%R	-20.52	7.94	7.94	9.72	14.52	50.00	-5.84	32.93	15.62	64.35
12	N	23.37abc	0.79ab	42.45abc	80.67abc	9.55abc	5.00ab	35.00a	39.60ab	50.55k	6.89abc
	S	30.50d	0.69d	0.69e	33.15d	5.83e	1.00d	43.50a	22.77d	42.73j	0.99d
	%R	-30.48	11.85	11.86	21.91	39.01	80.00	-24.29	42.49	15.48	85.63

Different letter in each column indicate significant difference at $p \leq 0.05$.

Tem, Flu, RCC, RWC, SL, NGS, NS, GW, Y and Ph indicate; leaf temperature, fluorescence chlorophyll, relative chlorophyll content (SPAD), relative water content, spike length, number of grain per spike, 1000-grains weight, grain yield per plant and plant height, respectively.

tively. Riaz and Choudhy (2003) reported that genotypes with high 1000-grain weight under irrigated conditions may not be superior for this trait under moisture stress conditions. This is possible due to the limitation of moisture which forces plant to complete its grain filling in relatively shorter duration (Fayaz and Arzani, 2011). Number of grain per spike and 1000-grain weight decreased under water deficit stress condition. Probably water deficit caused male sterility which may in turn reflected in abortion of terminal and basal florets; hence, reduction in number of grain per spike and flowed by grain weight (Saini and Aspinall, 1981). In durum wheat Garcia del moral *et al.* (2005) observed 18.5% of reduction in number of grain per spike due to the negative effect of moisture stress. Also, 'Atlas46/Kavir' 'Rihane-05' genotypes produced the highest and lowest grain yield than all genotypes and landraces under non-stress and stress, respectively. The highest reduction (85%) related to 'Rihane-05' genotype. Kirby and Jones (1997) and Giunta *et al.* (1993) also reported that mean decreased in grain yield under drought stress. In addition, decline of grain yield and components yield reported in

many researches (Gooding *et al.*, 2003; Ozturk and Aydin, 2004; Shah and Paulsen, 2003; Wardlaw, 2002).

Correlation studies are useful in measuring the strength and the direction of these relationships among the different characters and grain yield (Gashaw *et al.*, 2007). So, to study the relationship, simple correlation between each pair of the characteristics for both conditions was calculated (Tab. 4). Based on correlation analysis, all the characters included in the study except leaf temperature and number of grain per spike showed significant ($p \leq 0.01$) positive correlation with grain yield under both conditions. Garcia del Moral *et al.* (2003) reported that the number of spikes per squares meter in durum wheat was positively related to grain yield under rain fed conditions. In terms of physiological characters high significant ($p \leq 0.01$) correlation coefficient was found between relative water content and grain yield under stress condition; however, high significant ($p \leq 0.01$) correlation coefficient was found between fluorescence chlorophyll and grain yield under non-stress condition. Significant ($p \leq 0.01$) and negative correlation coefficient was found between grain

Tab. 4. Correlation coefficient between physiological traits, grain yield and its components under normal (above diameter) and water deficit stress conditions (below diameter)

	Tem	Flu	RCC	RWC	SL	NS	NGS	GW	Y	Ph
Tem	1	-0.96**	-0.97**	-0.95**	-0.91**	-0.75**	-0.12 ^{ns}	0.12 ^{ns}	-0.73**	0.07 ^{ns}
Flu	-0.97**	1	0.97**	0.98**	0.92**	0.85**	0.14 ^{ns}	-0.15 ^{ns}	0.82**	-0.14 ^{ns}
RCC	-0.98**	0.98**	1	0.95**	0.92**	0.79**	0.19 ^{ns}	-0.18 ^{ns}	0.77**	-0.13 ^{ns}
RWC	-0.96**	0.98**	0.97**	1	0.91**	0.84**	0.15 ^{ns}	-0.16 ^{ns}	0.82**	-0.15 ^{ns}
SL	-0.97**	0.97**	0.99**	0.96**	1	0.84**	0.07 ^{ns}	-0.05 ^{ns}	0.83**	-0.11 ^{ns}
NS	-0.91**	0.92**	0.90**	0.92**	0.89**	1	0.06 ^{ns}	-0.06 ^{ns}	0.98**	-0.14 ^{ns}
NGS	0.76**	-0.79**	-0.78**	-0.77**	-0.77**	-0.72**	1	-0.98**	-0.07 ^{ns}	0.13 ^{ns}
GW	-0.72**	0.76**	0.74**	0.74**	0.72**	0.70**	-0.98**	1	0.086	-0.15 ^{ns}
Y	-0.91**	0.92**	0.90**	0.92**	0.88**	1.00**	-0.71**	0.70**	1	-0.17 ^{ns}
Ph	-0.15 ^{ns}	0.15 ^{ns}	0.20 ^{ns}	0.17 ^{ns}	0.20 ^{ns}	0.11 ^{ns}	-0.28*	0.27 ^{ns}	0.11 ^{ns}	1

** and ^{ns}: Significant at the 0.01 probability level and Non-significant, respectively.

Tem, Flu, RCC, RWC, SL, NGS, NS, GW, Y and Ph indicate: leaf temperature, fluorescence chlorophyll, relative chlorophyll content (SPAD), relative water content, spike length, number of grain per spike, 1000-grains weight, grain yield per plant and plant height, respectively.

Tab. 5. Grain yield and stress susceptibility index (SSI) of genotypes and landraces of barley under normal (Yn) and water deficit stress (Ys) conditions

Genotype/landrace	Yp (gr per plant)	Ys (gr per plant)	SSI
'Badia Kavirr'	7.17	1.24	1.26
'168.4 Lign131/Abrabi Abiad'	4.94	1.98	0.91
'Mahabad'	5.77	2.22	0.94
'Rihaner'	5.79	2.98	0.74
'Naghadeh'	6.45	4.21	0.53
'Atlas46/Kavir'	8.25	1.24	1.29
'CB'	4.97	2.95	0.62
'Sahand'	8.05	1.96	1.15
'Makoobi'	6.58	2.21	1.01
'Piranshahr'	6.48	2.45	0.95
'CW3117-77-5-9-5'	6.92	2.47	0.98
'Rihane-05'	6.89	0.99	1.30

yield per plant and number of grain per spike under stress condition. 1000-grain weight also had negative correlation with all characters except leaf temperature under non-stress condition. However, this character had significant ($p \leq 0.01$) and positive correlation with fluorescence chlorophyll, relative water content, SPAD value, spike length, number of spike per plant under stress condition (Tab. 4). These results are in agreement with the previous reported results on landraces of wheat under stress condition (Farshadfar *et al.*, 2012). Moghaddam *et al.* (1997) reported negative correlation between number of grain per spike and 1000-grain weight. In the studies conducted by Sinha and Sharma (1979) and Belay *et al.* (1993), grain yield was positively correlated with each of the three primary grain yield components, with either positive or negative correlation between grain yield and plant height.

Stress susceptibility index (SSI) was used as a selection criterion of drought tolerant in terms of minimization of yield reduction caused by drought stress as compared with non-stress conditions. Calculated SSI varied in from 0.62 to 1.30 for genotypes, and ranged from 0.53 to 1.15 for landraces (Tab. 5). 'Naghadeh' and 'Sahand' landraces that had the lowest and highest SSI values were found to be the most tolerant and the most susceptible landraces, respectively. Also, 'CB' and 'Rihane-05' genotypes that had the lowest and highest value were found to be the most tolerant and susceptible genotypes, respectively. This was in agreement with conclusions made based on agro-morphological and physiological characteristics. Fayaz and Arzani (2011) reported cultivars with low SSI values are moisture resistant because they have lesser reduction in grain yield under stress compared with non-stress condition. Nevertheless, this index per se appears to have serious limitations for the quantification of genotype reaction to moisture conditions, because it is based on minimizing yield reduction in stress compared with non-stress conditions. Therefore, selection for low SSI would tend to reduce yield in non-stress conditions (Dencic *et al.*, 2000). However, barley genotypes and landraces identified as the stress tolerant by SSI may have tolerant mechanism, and can be used as sources of drought stress resistance in barley breeding programs for development of secondary germplasm with high grain yield potential. Considering the grain yield potential 'Naghadeh', 'Mahabad', 'Piranshahr' landraces and CW3117-77 genotypes were ranked as the superior group of drought tolerant barley. On the other hand, 'Naghadeh' landrace was identified as a barley landrace that could be grown under both normal and water deficit stress conditions with high grain yield.

Conclusions

This paper proposes a strategy to select the traits to be used in breeding programs. The result obtained from this study could be useful for barley breeders and seed producer in order to increase grain yield in water deficit con-

ditions. The changes of characters relationship in barley seed under different irrigation condition should be considered for variety selection and every plant breeding program of the plant. Screening drought tolerant genotypes and landraces using compare mean and SSI discriminated 'Naghadeh' landrace and 'CW3117-77' genotype as the most drought tolerant. Therefore they are recommend to be used as parents for improvement of drought tolerance in other cultivars.

References

- Ahmadi M, Farshadfar E, Veisi S (2012). Evaluation of genetic diversity in bread landrace of bread wheat under irrigated and rainfed conditions. *Inter J Agri Crop Sci* 4(21):1627-1636.
- Araus JL, Salfer M P, Royo C, Serett M D (2008). Breeding for yield potential and stress adaptation in cereals. *Critical Rev in Plant Sci* 27:377-412.
- Baik BK, Ullrich SE (2008). Barley for food: Characteristics, improvement, and renewed interest. *J Cereal Sci* 48(2):233-242.
- Bayoumi TY, Eid MH, Metwali MM (2008). Application of physiological and biochemical indices as a screening technique for drought tolerance in wheat genotypes. *Afr J Biotech* 7:2341-2352.
- Belay G, Tesemma T, Becker HC, Merker A (1993). Variation and interrelationships of agronomic traits in Ethiopian tetraploid wheat landraces. *Euphytica* 71:181-188.
- Betran FJ, Beck D, Banziger M, Edmeades GO (2003). Genetic analysis of inbred and hybrid grain yield under stress and non-stress environments in tropical maize. *Crop Sci* 43:807-817.
- Brush SB (1995). *In situ* conservation of landraces in centers of crop diversity. *Crop Sci* 35:346-354.
- Byrne PF, Bolanos J, Edmeades GO, Eaton DL (1995). Grains from selection under drought versus multi location testing in related tropical maize populations. *Crop Sci* 35:63-69.
- Cao HX, Sun CX, Shao HB, Lei XT (2011). Effects of low temperature and drought on the physiological and growth changes in oil palm seedlings. *African Journal of Biotechnology* 10:2630-2637.
- Ceccarelli S, Grando S (2000). The environment selection and environmental sensitivity in barley. *Euphytica* 57:57-167.
- Clarke J M, DePauw RM, Townley-Smith TF (1992). Methods evaluation of quantification for drought tolerance in wheat. *Crop Sci* 32:723-728.
- Crudace AJ (2000). The investigation of the in vivo behavior of maize herbicide-Isoxaflutole. PhD thesis. University of Essex, Colchester, UK.
- Dencic S, Kastori R, Kobiljski B, Duggan B (2000). Evaluation of grain yield and its components in wheat cultivars and landraces under near optimal and drought conditions. *Euphytica* 113:43-52.

- Erdei L, Tari I, Csiszar J, Pecsvaradi A, Horvath F, Szabo M, Ordog M, Cseuz L, Zhiponova M, Szilak L, Gyorgyey L (2002). Osmotic stress responses of wheat species and cultivars differing in drought tolerance: some interesting genes (advices for gene hunting). *Acta Biol Szeged* 46:63-65.
- Farshadfar E (2012). Application of integrated selection index and rank sum for screening drought tolerant genotypes in bread wheat. *Inter J Agri Crop Sci* 4(6):325-332.
- Farshadfar E, Farshadfar M, Dabiri S (2012). Comparison between effective selection criteria of drought tolerance in bread wheat landraces of Iran. *Annals of Biological Research* 3 (7):3381-3389.
- Fayaz N, Arzani A (2011). Moisture stress tolerance in reproductive growth stages in triticale (*X Triticosecale* Wittmack) cultivars under field conditions. *Crop Breeding Journal* 1(1):1-12.
- Fernandez GCJ (1992). Effective selection criteria for assessing plant stress tolerance, 257-270 p. In: Proceedings of the international symposium on adaptation of vegetables and other food crops in temperature and water stress, Taiwan 13-16 August 1992.
- Fischer RA, Maurer R (1978). Drought resistance in spring wheat cultivars; I. Grain yields responses. *Aust J Agric Res* 29:897-912.
- Garci A, Del Moral LF, Rharrabti Y, Villegas D, Royo C (2003) Evaluation of grain yield and its components in durum wheat under mediterranean conditions: An ontogenic approach. *Agron J* 95:266-274.
- Gashaw A, Mohammed H, Singh H (2007). Selection criterion for improved grain yields in Ethiopian durum wheat genotypes. *Afr Crop Sci J* 15(1):25-31.
- Giunta F, Motzo R, Deidda M (1993). Effect of drought on yield and yield components of durum wheat and triticale in Mediterranean environment. *Field Crops Res* 33(4):399-409.
- Gooding MJ, Ellis RH, Shewry, PR, Schofield JD (2003). Effects of restricted water availability and increased temperature on the grain filling, drying and quality of winter wheat. *J Cereal Sci* 37:295-309.
- Gupta NK, Gupta S, Kumar A (2001). The effect of water stress on physiological attributes and its relationship with growth and yield of wheat cultivars at different stages. *Crop Sci* 186:55-62.
- Kirby EJM, Jones HG (1997). The relations between the main shoot and tillers in barley. *J Agri Sci* 88:381-389.
- Moghaddam M, Ehdaie B, Waines JG (1997). Genetic variation and interrelationships of agronomic characters in landraces of bread wheat from southeastern Iran. *Euphytica* 95:361-369.
- Ozturk A, Aydin F (2004). Effect of water stress at various stages on some quality characteristics of winter wheat. *J Agron Crop Sci* 190:93-99.
- Percival MP, Baker NR (1991). Herbicides and photosynthesis. In: Baker NR, Percival MP (Eds.). *Herbicides*. Elsevier, Amsterdam.
- Pour Aboughadareh AR (2012). Evaluation of *Triticum boeoticum* and *Triticum urartu* populations based on morphological characteristics under normal and water stress conditions. M.Sc Thesis. University of Tabriz, Tabriz, Iran.
- Rajaram S, Van Ginkle M (2001). Mexico, 50 years of international wheat breeding. In: Bonjean AP, Angus WJ (Eds.). *The World Wheat Book: A History of Wheat Breeding*. Lavoisier Publishing, Paris, France.
- Rampino P, Pataleo S, Gerardi C, Mita G, Perrotta C (2006). Drought stress response in wheat: physiological and molecular analysis of resistant and sensitive genotypes. *Plant Cell Environ* 29:2143-2152.
- Rathjen AJ (1994). The biological basis of genotype \times environment interaction: its definition and management. Proceedings of the Seventh Assembly of the Wheat Breeding Society of Australia, Adelaide, Australia.
- Riaz R, Choudhary MA (2003). Genetic analysis of some economic traits of wheat under drought condition. *Asian J Plant Sci* 2:790-796.
- Richards RA (1996). Defining selection criteria to improve yield under drought. *Plant Growth Regul* 20:157-166.
- Rong-Hua L, Peiguo G, Baum M, Grando S, Ceccarelli S (2006). Evaluation of chlorophyll content and fluorescence parameters as indicators of drought tolerance in barley. *Agric Sci Chin* 5:751-757.
- Saini HS, Aspinall D (1981). Effect of water deficit on sporogenesis in wheat (*Triticum aestivum* L.). *Ann Bot* 43:623-633.
- Sanchez-Rodriguez E, Rubio-Wilhelmi M, Cervilla LM, Blasco B, Rios JJ, Rosales MA, Romero L, Ruiz JM (2010). Genotypic differences in some physiological parameters symptomatic for oxidative stress under moderate drought in tomato plants. *Plant Sci* 178:30-40.
- Shah NH, Paulsen GM (2003). Interaction of drought and high temperature on photosynthesis and grain-filling of wheat. *Plant Soil* 257:219-226.
- Sinha GCP, Sharma NN (1979). Correlation regression and path analysis studies in wheat varieties. *Indian J Agron* 25:225-229.
- Slapakauskas V, Ruzgas V (2005). Chlorophyll fluorescence characteristics of different winter wheat varieties (*Triticum aestivum* L.). *Agronomy Research* 3(2):203-209.
- Turner NC (1986). Crop water deficit: A decade of progress. *Adv Agron* 39:1-51.
- Wardlaw IF (2002). Interaction between drought and chronic high temperature during kernel filling in wheat in a controlled environment. *Ann of Bot* 80:469-476.