



Adipocyte size as a determinant of metabolic disease and adipose tissue dysfunction

Sofia Laforest, Jennifer Labrecque, Andréanne Michaud, Katherine Cianflone & André Tchernof

To cite this article: Sofia Laforest, Jennifer Labrecque, Andréanne Michaud, Katherine Cianflone & André Tchernof (2015) Adipocyte size as a determinant of metabolic disease and adipose tissue dysfunction, *Critical Reviews in Clinical Laboratory Sciences*, 52:6, 301-313, DOI: 10.3109/10408363.2015.1041582

To link to this article: <http://dx.doi.org/10.3109/10408363.2015.1041582>



Published online: 17 Aug 2015.



Submit your article to this journal [↗](#)



Article views: 138



View related articles [↗](#)



View Crossmark data [↗](#)

REVIEW ARTICLE

Adipocyte size as a determinant of metabolic disease and adipose tissue dysfunction

Sofia Laforest^{1,2,3}, Jennifer Labrecque^{1,2,3}, Andréanne Michaud^{1,2,3}, Katherine Cianflone³, and André Tchernof^{1,2,3}

¹CHU de Québec Research Center, Quebec, Canada, ²School of Nutrition, Laval University, Quebec, Canada, and ³Quebec Heart and Lung Institute, Quebec, Canada

Abstract

Obesity is a heterogeneous disease and is associated with comorbidities such as type 2 diabetes mellitus, cardiovascular disease and cancer. Several studies have examined the role of dysfunctional adipose tissue in the pathogenesis of obesity, highlighting the contrasting properties and impact of distinct fat compartments, sometimes with contradictory results. Dysfunctional adipose tissue involves enlargement, or hypertrophy, of pre-existing fat cells, which is thought to confer increases in cardiometabolic risk, independent of the level of obesity *per se*. In this article, we critically analyze available literature that examined the ability of adipocyte cell size to predict metabolic disease and adipose tissue dysfunction in humans. Many studies demonstrate that increased fat cell size is a significant predictor of altered blood lipid profiles and glucose–insulin homeostasis independent of adiposity indices. The contribution of visceral adiposity to these associations appears to be of particular importance. However, available studies are not unanimous and many fat depot-specific aspects of the relationship between increased fat cell size and cardiometabolic risk or parameters of adipose tissue dysfunction are still unresolved. Methodological factors such as the approach used to express the data may represent significant confounders in these studies. Additional studies should consider the fact that the relationship between fat cell size and common adiposity indices is non-linear, particularly when reaching the obese range. In conclusion, our analysis demonstrates that fat cell size is a significant predictor of the cardiometabolic alterations related to obesity. We propose that adipocyte hypertrophy, especially in the visceral fat compartment, may represent a strong marker of limited hyperplastic capacity in subcutaneous adipose tissues, which in turn is associated with the presence of numerous cardiometabolic alterations.

Abbreviations: **3T3-L1:** Mouse embryonic pre-adipocyte cell line; **ApoB:** apolipoprotein B; **ATGL:** adipose triglyceride lipase; **BFM:** body fat mass; **BMI:** body mass index; **CD11b:** macrophage and neutrophil cell marker; **CD11c:** dendritic cell and macrophage cell marker; **CD31:** endothelial cell marker; **CD68:** macrophage cell marker; **DEXA:** dual-energy X-ray absorptiometry; **FCS:** fat cell size; **GLUT4:** Glucose transporter type 4; **HOMA-IR:** homeostatic model assessment of insulin resistance; **HIF- α :** hypoxia-inducible factor 1; **HSL:** hormone-sensitive lipase; **IGF-1:** insulin-like growth factor 1; **IL-6:** interleukin 6; **IRS-1:** insulin receptor substrate 1; **MeSH:** Medical Subject Heading; **NAFLD:** non-alcoholic fatty liver disease; **NF- κ B:** nuclear factor kappa B; **NGT:** normal glucose tolerant; **OM:** omental; **PCOS:** polycystic ovary syndrome; **PLIN:** perilipin (lipid droplet-associated protein); **SC:** subcutaneous; **SVF:** stroma-vascular fraction; **T2DM:** type 2 diabetes mellitus; **TNF- α :** tumor necrosis factor alpha; **VEGFA:** vascular endothelial growth factor A; **VEGF-R2:** vascular endothelial growth factor receptor 2; **vWF:** von Willebrand Factor; **WHR:** waist-to-hip ratio

Keywords

Adipocyte hypertrophy, adipokines, metabolic syndrome, obesity, subcutaneous fat, visceral fat

History

Received 9 January 2015
Revised 17 March 2015
Accepted 04 April 2015
Published online 17 August 2015

Introduction

Over one-third of the world population is now struggling with overweight or obesity and the disease burden frequently associated with these conditions¹. Obesity is a well-known risk factor of metabolic diseases, such as type 2 diabetes mellitus (T2DM), dyslipidemia, coronary heart disease, hypertension, non-alcoholic fatty liver disease (NAFLD)

Referees: Dr. Susan K. Fried, Faculty of Medicine, Boston University, Boston, MA, USA; Dr. Susanne Votruba, The National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK), Bethesda, MD 20892-2560, USA

Address for correspondence: André Tchernof, PhD, CHU de Québec Research Center, 2705 Laurier Blvd, R-4779, Québec G1V 4G2, Canada. Tel: +1 418 654-2296. E-mail: andre.tchernof@criucpq.ulaval.ca

and stroke². It has also been linked with dementia, obstructive sleep apnea and numerous types of cancer². However, a significant proportion of obese individuals, 15–30% depending on the populations examined and the definition of metabolic health, do not develop alterations associated with obesity, at least not in the short term^{3,4}. These subsets of apparently healthy obese subjects are named metabolically healthy obese.

Differences in body fat distribution likely account for many discrepancies between metabolically healthy and unhealthy individuals. Jean Vague proposed as early as 1947 that patients with upper-body obesity had an increased risk for metabolic diseases, while individuals with lower-body (femoral-gluteal) obesity had a lower risk of such alterations⁵. Since then, body fat distribution indices have been recognized as stronger predictors of metabolic alterations compared to overall obesity, emphasizing their impact on global metabolic health². Specifically, studies have shown that excess fat accumulation on visceral anatomical structures such as the greater omentum or mesentery is a strong predictor of a detrimental metabolic status; whereas accumulation of gluteal and femoral fat is viewed as protective when total adiposity is accounted for². In fact, despite a high body fat mass, low visceral fat accumulation is a major feature of the metabolically healthy profile observed in some obese individuals^{6,7}.

Adipose tissue expansion in a given fat compartment occurs through adipocyte hyperplasia, hypertrophy or a combination of both. Recruitment of new cells through differentiation of pre-adipocytes (hyperplasia) is now generally considered as a mechanism that protects against metabolic alterations. Indeed, adipocyte hyperplasia is associated with preferential accumulation of subcutaneous (SC) adipose tissue and the gynoid phenotype^{8–12}. On the other hand, visceral adipocytes and the android fat distribution phenotype are more commonly associated with adipocyte hypertrophy, that is, enlargement of existing fat cells^{8,9,11,12}.

Increased fat cell size (FCS) has been demonstrated to be associated with metabolic impairment in patients as early as the 1970s^{13,14}. In 1979¹⁵, adipocyte hypertrophy was proposed as a valuable parameter for characterization of metabolic disturbances. Since then, FCS has been linked to numerous variables used for assessing adipose tissue biology as well as to metabolic alterations independent of total adiposity^{9,11,16–25}. To the best of our knowledge, studies on the potential role of adipocyte hypertrophy as a biological marker of metabolic disease and adipose tissue dysfunction have never been reviewed in detail. This article proposes an extensive analysis of the studies on this topic and demonstrates that adipocyte size may, indeed, represent an important determinant of adipose tissue dysfunction and a potential marker of pathologies that may or may not be linked with total adiposity.

Literature search

To perform this critical analysis, the first step was to determine all possible Medical Subject Heading (MeSH) terms to be used in searches on the PubMed database. The term “adipocyte” was combined with the following keywords: size, hypertrophy, white, area, diameter, morphology, cellularity, dysregulation and volume. The same approach was then repeated but this

time using the term “fat cell” instead of “adipocyte”. The articles were selected depending on the following specific criteria: (1) emphasis was placed on human studies directly relevant to the research topic; and (2) only articles on white adipose tissue were selected. Relevant articles from the reference list of identified papers were added. In total, 172 studies were retained, which were published between 1970 and March 2015. A record of the articles was kept and they were listed according to the research criteria used to find them. An exhaustive analysis of the population, the methods and major findings of each study was performed. We further selected the studies that had analyzed the ability of FCS to predict obesity-related metabolic complications or parameters of adipose tissue function. The latter studies represent the main body of literature used for this article. Analyses of cell size variations among sexes, and obesity degrees as assessed by body mass index (BMI) and cell sizing techniques, were performed using 76 publications in which the variables of interest were available. FCS measurements in these studies were converted to the same unit. Units of volume (μL , pL and μm^3) and units of surface (μm^2) were converted to a diameter in micrometers (μm) assuming a spherical or circular cell shape. Units of mass (μg) were first converted to volume using the density of triolein (density = 0.915 g/mL) and then to cell diameter in μm .

Methods used to measure fat cell size

This section provides a brief overview of the main techniques used to assess FCS. We focus mainly on the most commonly used techniques: collagenase digestion, osmium tetroxide fixation and histological analysis. We also provide analyses of FCS variation according to sex, obesity level and measurement technique.

Collagenase digestion

Collagenase digestion was developed by Rodbell as a means of separating mature adipocytes from the stroma-vascular fraction (SVF)²⁶. It is now used as the first step in many experiments, including cell cultures and cell sizing. Briefly, adipose tissue is digested by collagenase and mature adipocytes are separated from the SVF by floatation in an aqueous solution. Pictures of mature adipocytes can be taken with a phase contrast microscope to assess adipocyte size. This method has been the most frequently used. However, some drawbacks have prevented it from becoming a gold standard in determining FCS. Small adipocytes do not float as easily as the average-sized cells, due to their low lipid content²⁷. In addition, adipocytes tend to break in unfixed tissue because of their fragility²⁷; however, the introduction of adenosine in the solution minimized this bias²⁸. Centrifugation during the floatation step can bias the recovery of small adipocytes, therefore this step can be omitted²⁹. Finally, blue methylene staining may be required to assess viability of the cells and facilitate identification of lipid droplets versus undamaged fat cells²⁶.

Osmium tetroxide fixation and Multisizer Counter analysis

This technique was developed in 1968 by Hirsh and Gallian³⁰ based on fact that osmium tetroxide fixes intracellular lipids

and allows staining of very fragile cell types. In brief, adipose tissue can be digested by collagenase and subsequently fixed by osmium tetroxide, or these steps can be simultaneous if a collidine-HCl solution is used to separate the SVF from mature adipocytes instead of collagenase. Adipocytes are then analyzed with a counter that measures cell size by fluctuation of electrical resistance. This technique allows the study of large distributions of adipocyte sizes as well as analysis of cell subpopulations such as very small fat cells^{31,32}. However, even though this technique allows for fixation of very small adipocytes (15–25 μm), a threshold value (most often 25 μm) is used to discriminate between mature adipocytes and artifacts. Multilobular fat cells also tend to rupture during the fixation process which can lead to an underestimation of mean FCS, especially in obese individuals³¹. This technique is also time-consuming and comes at a relatively high price, as it requires proper handling of osmium tetroxide, a hazardous chemical^{27,30,31}.

Histological analysis

Adipocyte cell size analysis can also be performed on histological slides. Type and concentration of fixatives, paraffin-embedding, time of fixation, slice thickness and type of coloration may vary among laboratories. Of importance, this is the only technique that allows examination of global tissue architecture. However, many potential biases are inherent to this approach and many assumptions have to be made in order to assess cell size. Cell distribution has to be considered uniform, which may not be entirely true in obese and young individuals²⁷. Furthermore, fixation agents are known to induce significant shrinkage of the cells^{33,34}. Likewise, it must be assumed that cells are perfect spheres showing their largest diameter²⁷. This technique has been extensively used recently due to the possibility of simultaneously performing immunostaining for various cellular markers.

Other techniques

Other methods have been used in various contexts to appreciate variation in FCS. Flow cytometry as well as scanning electron microscopy have been used by a few teams^{35–39}. These specialized methods are not in common use to measure FCS in clinical research.

Cell size analysis software

Promising semi-automatic methods have arisen with the purpose of limiting user-dependent biases and time of use^{40–42}. Information on a few completely automated programs has also been published^{43,44}. Obstacles such as the time needed to obtain high-quality images may slightly complicate their use, along with the presence of the SVF and other tissue artifacts in some samples. Moreover, user-dependent variation still needs to be addressed.

Overall, among these approaches to assess FCS, none are without drawbacks. Characterization of very large and very small adipocytes remains a challenge for all methods available. Moreover, the number of adipocytes examined in

each study varied from 50 to 20 000, which at times limited study comparisons. As a result, no single method has arisen in our literature search as the gold standard for adipocyte cell sizing.

Population and methodological variation

Our survey of the literature demonstrates that adipocyte size varies between men and women as well as between fat depots (visceral versus SC). Moreover, as a function of increasing obesity level, mean adipocyte size increases, reaching a plateau at a certain level of adiposity. Arner and colleagues have already shown that the relationship between SC FCS and body fat mass (BFM) is curvilinear in men and women⁴⁵. Moreover, adipocyte hypertrophy is negatively correlated with adipocyte hyperplasia, when adjusted for the predicted adipocyte volume at a given BFM, and subjects with higher fat cell size than predicted by the curve have lower rates of adipogenesis⁴⁵. This suggests an important impact of adipose morphology, independent of BFM. However, a complete review of FCS variation and obesity indices such as BMI has not been published yet.

Using all the studies in which mean FCS and mean BMI were available, we summarize sex-, depot- and obesity-related variation in adipocyte size in humans and address potential differences related to the method used for analysis. As shown in Figure 1(A), omental (OM) and abdominal SC FCS increase in both sexes with increasing obesity level as assessed by BMI and reach a plateau around BMI values of $\sim 30 \text{ kg/m}^2$. When plotting data from men (Figure 1B) and women (Figure 1C) separately, we find that the plateau is reached at lower BMI values in men ($\sim 25 \text{ kg/m}^2$) compared to women ($\sim 35 \text{ kg/m}^2$). Moreover, OM FCS appears to be $\sim 20\%$ lower than SC FCS in women, whereas this depot difference is not apparent in men. These findings suggest a clear difference among sexes regarding the expansion of adipose tissue. Men are more prone to hypertrophic obesity, as the plateau is seen at lower BMI values. On the other hand, women show signs of both hypertrophic and hyperplastic expansion as obesity level increases. Similar observations can be made with both the SC and OM adipocyte size curves.

When examining FCS as a function of the measurement technique, we found that the use of histological slides generally leads to lower mean FCS by approximately 15% across all BMI values (Figure 1D). Collagenase digestion and osmium fixation appear to generate similar mean FCS (Figure 1D). These patterns can be observed in both depots as well. All techniques showed a general pattern of BMI-related increase, with a plateau reached at higher obesity levels.

In this analysis, population differences noted in FCS variation closely reflect those observed in many individual studies as reviewed here. Adipocyte size is clearly influenced by sex, anatomical localization and obesity levels. As discussed in the sections below, these variations need to be considered critically when examining the relationship between fat cell hypertrophy in a given fat compartment and metabolic diseases or parameters of adipose tissue function.




Figure 1. Mean FCS of the abdominal subcutaneous (SC) and omental (OM) depot in 176 subgroups of individuals in 78 published studies. Sizes of the symbols reflect study samples size. Published studies on FCS were screened and inclusion criteria were availability of BMI and FCS for the same subgroup. A total of 6355 individuals were included in the analysis. Non-linear regressions were determined taking study sample size into consideration. (A) OM and SC FCS in both in males and females. (B) OM and SC FCS as a function of BMI in males. (C) OM and SC FCS as a function of BMI in females. (D) FCS variation as a function of BMI and the technique used to assess FCS. Each curve combines OM and SC FCS. Collagenase digestion, $n = 4778$; Osmium tetroxide fixation, $n = 1052$; Histological analysis, $n = 2533$.

Adipocyte size as a potential biomarker of cardiometabolic alterations or disease endpoints independent of common adiposity indices

Common adiposity indices have been widely used to assess the presence of cardiometabolic risk factors in overweight or obese individuals. BMI is certainly the most widely used; however, it has its limitations. BMI does not take into account body fat distribution and only provides a crude assessment of body composition. Other anthropometric indices have been examined along with more invasive measurements of body fat distribution and/or body composition². An important question that has been addressed in many original publications is whether adipocyte size also predicts the metabolic complications frequently associated with obesity or abdominal obesity. Considering that FCS is strongly related to body composition and fat distribution, whether it predicts metabolic alterations independent of overall or regional adiposity has also been tested in a number of studies. The next sections provide a review of the studies that have addressed the link between FCS and cardiometabolic risk factors or disease endpoints.

Blood lipid profile

Imbeault et al.⁴⁶ reported that when matched for visceral adipose tissue area, abdominal but not femoral SC FCS

predicted an altered lipid profile in men but not in women. Specifically, men with large adipocytes had hypertriglyceridemia and higher LDL-apolipoprotein B (ApoB) levels. Another study reported that in patients with first-degree relatives of T2DM individuals, elevated SC FCS correlated with lower HDL-cholesterol concentration, but not with high total cholesterol or LDL-cholesterol concentrations when adjusted for BMI²⁰. In homozygous twins discordant for obesity level, after adjustment for total BFM, SC FCS correlated positively with LDL-cholesterol concentrations⁴⁷. Interestingly, in the obese twin characterized by adipocyte hypertrophy, higher LDL-cholesterol and lower HDL-cholesterol levels were found compared to the lean co-twin, but this difference was not found in twins characterized by hyperplasia. However, in both cases, the obese co-twin had 61% larger adipocytes than the lean twin⁴⁷. In obese women, Ledoux et al.²¹ did not find a correlation between FCS in either the SC or visceral depot and blood lipid concentrations when adjusting for BMI or waist-to-hip ratio (WHR).

An elegant study by Hoffsted et al.¹⁶ showed that in obese women, OM, but not SC FCS, was associated with plasma ApoB, total cholesterol, LDL-cholesterol and triglyceride concentrations independent of BMI, BFM and body fat distribution indices obtained by dual-energy X-ray absorptiometry (DEXA). In another study, visceral and SC fat cell

volumes were positively correlated with triglyceride levels and negatively with HDL-cholesterol concentrations in bariatric patients after control for total BFM⁹. In a sample of women ranging in adiposity from lean to moderately obese, after control for BMI, total BFM and visceral adipose tissue area measured by computed tomography, we found that OM FCS predicted triglyceride concentration whereas SC FCS did not¹⁷. Yet, in female and male Indians, another group reported no significant correlation between total cholesterol and triglyceride concentrations and SC or OM FCS after adjustment for BMI, BFM, waist circumference, total abdominal adipose tissue area and SC adipose tissue area⁴⁸. However, in that particular study, associations between visceral adipose tissue accumulation and metabolic parameters were scarce as visceral adipose tissue only correlated with HDL-cholesterol and triglyceride concentrations in men. In sum, a number of studies have shown associations between high FCS either in the SC or visceral fat compartment and an altered lipid profile. Some of these associations appeared to be independent of total or abdominal adiposity in some cases. Further studies in both genders that take ethnicity as well as the OM and SC fat compartment into account are needed to generalize these findings.

Glucose homeostasis, insulin resistance and T2DM

The relationship between SC FCS and insulin resistance is well documented. A number of studies have shown a clear association between abdominal SC FCS and markers of insulin resistance, in both women and men^{11,15,24,45,49–54}. In overweight South African women with normal glucose tolerance (NGT), no link was found between abdominal or gluteofemoral SC FCS and insulin resistance markers such as basal plasma insulin, insulin area and glucose area after a 100 g oral glucose load⁵⁵. Nevertheless, another study demonstrated that SC FCS was increased in obese subjects with T2DM when compared with obese control subjects matched for BMI, age, sex and ethnicity⁵⁶. Also, when controlling for BMI, SC FCS was positively correlated with homeostatic model assessment of insulin resistance (HOMA-IR) and negatively with post-glucose insulin sensitivity⁵⁷.

This was challenged by McLaughlin et al.⁵⁸ who reported that mean diameter of the larger SC cells assessed with osmium tetroxide was not different between insulin resistant versus insulin sensitive patients (119 versus 115 μm , respectively). However, insulin resistant patients were characterized by a larger population of small cells (a high ratio of small-to-large cells), a feature that can only be assessed with the osmium technique and which could possibly reflect arrested development of the very small adipocytes toward fully mature cells⁵⁸. Another study found no link between SC FCS and HOMA-IR²¹. On the other hand, in a sample of T2DM patients matched to NGT patients for BMI, Pasarica et al. found larger SC fat cells measured with osmium tetroxide in T2DM patients (124 μm diameter compared to 115 μm , respectively)⁵⁹. In that study, there was an increase of the small cell population in the diabetic subgroup and mean FCS was positively correlated with HOMA-IR. Similar results were obtained by Yang et al.⁶⁰. Mean fat cell volume was inversely correlated with insulin sensitivity measured by euglycemic, hyperinsulinemic

clamp after adjustment for BMI in a sample of first-degree relatives of T2DM patients⁶⁰. In non-obese first-degree relatives of T2DM patients, SC FCS was negatively correlated with GLUT4 protein expression and positively with circulating insulin levels, whereas BMI was not⁶¹. We found that SC GLUT4 mRNA abundance was lower in women with hypertrophic obesity⁶². Moreover, another group reported that after adjustment for BMI and percent body fat, SC FCS was associated with insulin resistance, but not with fasting glucose concentration in first-degree relatives of T2DM patients²⁰. Arner and colleagues found that in women characterized by SC adipocyte hypertrophy, adipocyte volume, adjusted for BFM, was an important and independent correlate of HOMA-index and fasting insulin levels⁴⁵.

Depending on how anthropometric adjustments are performed, SC FCS is not always an independent predictor of alterations in indices of glucose–insulin homeostasis. Our group reported that SC FCS was not a significant predictor of fasting glucose or fasting insulin in non-obese women when adjusted for adiposity and visceral adipose tissue area measured by computed tomography¹⁷. The adjustment for variation in visceral adipose tissue area was critical in the latter analysis, raising the possibility that lack of control for this variable in other studies may have led to the finding of significant associations. Similarly, Azuma et al.¹⁸ found that in T2DM patients, increased FCS was associated with insulin resistance independent of total BFM, but when they adjusted for body fat distribution measurements (leg fat mass or trunk fat mass), the relationship became non-significant. These results are not unanimous. Glycemia, insulinemia and glucose disposal rate were associated with SC FCS independent of age, BMI, BFM or body fat distribution indices obtained by DEXA in a sample of obese women¹⁶. Similar results were found in obese men and women, after adjustment for total body fat content⁹. SC adipocyte hypertrophy also explained the difference in glucose disposal rate between South Asians and Caucasians, after control for total body fat content, intraperitoneal and SC fat¹⁹. In another study, SC FCS was associated with markers of insulin resistance in NGT patients but not in T2DM patients, indicating that this relation may become non-linear when a certain degree of disease severity is reached²⁴. Such phenomena could contribute to the discrepancies noted among some studies.

The relationship between visceral FCS (OM adipocytes in most studies) and insulin resistance has also been of interest. Our group was the first to report a relationship between fasting insulin, HOMA-IR and OM FCS in non-obese women¹⁷. However, once adjusted for adiposity and body fat distribution, there was no significant association. The close correlation between OM FCS and visceral adipose tissue area measured by computed tomography explained the latter finding. Similar results were obtained in female and male Indians when adjusting for BMI, BFM, waist circumference, total and SC adipose tissue area⁴⁸. Another study found an association between fasting glycemia, HOMA-IR⁶³ and OM FCS; however, statistical adjustment for anthropometric values was not performed in that study⁵².

In obese patients, the association between visceral FCS and markers of insulin resistance remains uncertain. OM FCS seems to predict fasting glycemia, fasting insulin and

HOMA-IR independent of BMI or WHR in a few studies^{21–23}. Hoffstedt et al. found that visceral adipocyte hypertrophy was not associated with plasma insulin, fasting glycemia and glucose disposal after control for age, BMI, BFM and DEXA-measured body fat distribution indices¹⁶. The same group found that visceral fat cell volume (to a lesser extent than SC FCS) was correlated with insulin levels, insulin-induced glucose disposal and insulin sensitivity after control for total BFM, in both sexes⁹. To add to these results, no significant relationship was found between OM FCS and HOMA-IR in non-diabetic obese adults, yet OM FCS was associated with these variables independently of BMI and WHR *in vitro* when glucose uptake in freshly isolated adipocytes stimulated by insulin was measured²⁴. In morbidly obese patients with insulin resistance, a greater proportion of adipocytes exceeding 100 μm diameter was also observed along with a lower proportion of small adipocytes compared to insulin sensitive patients^{7,52,64}. This relation was challenged recently by van Beek and colleagues⁶⁵, who reported no difference in adipocyte size between subjects with T2DM compared to those with NGT.

Overall, many studies have reported significant associations between SC FCS and various markers of insulin resistance. Some reported that this association was independent of concomitant elevations in total BFM, pointing toward a specific role of adipocyte hypertrophy and perhaps adipose tissue dysfunction, as a marker of insulin resistance. On the other hand, most of the studies that adjusted their analysis for well-measured body fat distribution markers, such as visceral adipose tissue accumulation, have shown that the association between SC FCS and indices of insulin resistance was no longer significant. We propose that excess visceral adipose tissue accumulation and SC fat cell hypertrophy may represent markers of a common phenomenon: limited hyperplastic capacity of adipose tissues. Yet, the relationship with visceral adipocyte hypertrophy remains equivocal in most studies. The non-linear nature of the relationship between FCS and obesity level or insulin resistance may help to explain discrepancies among the various studies that examined this relationship.

Metabolic syndrome features

Three studies examined the association between FCS and the number of metabolic syndrome features. In bariatric surgery patients, no difference was noted in FCS of patients with or without the metabolic syndrome^{25,66}. The first study assessed the metabolic syndrome on the basis of the Harmonized criteria⁶⁷. Three features or more were required to qualify as metabolically unhealthy. The second study considered metabolic health status based on the following criteria: (1) fasting glucose > 100 mg/dL; (2) total insulin > 19 $\mu\text{U}/\text{mL}$; (3) triglycerides > 150 mg/dL; (4) HDL-cholesterol < 39 mg/dL; (5) systolic blood pressure > 140 mmHg; and (6) diastolic blood pressure > 90 mmHg. Patients were considered as metabolically unhealthy when at least one feature was present. Visceral adipocyte hypertrophy was observed in patients with low HDL-cholesterol levels and high fasting glucose concentrations in one of these studies²⁵. Also, the number of metabolic syndrome features increased with visceral FCS²⁵, consistent

with another study²² (adapted version of Ref.⁶⁸) that associated adipocyte hypertrophy with a detrimental metabolic state in bariatric patients. The fact that only severely obese patients were studied may have led to low-level associations between FCS and the number of metabolic syndrome features, considering that FCS values reach a plateau in the obese range. Another study reported that high mesenteric FCS was associated with a 1.79-fold increase in the risk of having metabolic syndrome (ATP III criteria) in overweight men⁶³. More studies are needed to assess whether FCS is associated with the onset of metabolic syndrome independent of adiposity.

NAFLD and liver fat accumulation

Six studies found an association between liver fat accumulation and SC adipocyte size^{22,47,69–72}. Two studies reported the opposite^{73,74}. One of these studies included a homogenous population of T2DM patients⁷⁴ and the other examined males before and after weight gain⁷³. When patients were matched for BMI, age, sex and/or BFM, SC adipocyte hypertrophy was an independent marker of liver fat^{47,69–72}. Moreover, OM adipocyte hypertrophy could independently predict stages of NAFLD^{22,75}. In one study, the extent of fatty liver disease was not predicted by FCS, but this finding was consistent with the lack of association between FCS and chronic intermittent hypoxia, the best and only predictor of liver disease stages in that sample⁷⁶.

Polycystic ovary syndrome

There are a few studies on adipocyte hypertrophy and polycystic ovary syndrome (PCOS) in the literature. Two teams noted a 25% increase of SC FCS in PCOS women compared to controls when matched for age and BMI^{77,78}. Another group found similar results for visceral adipocyte size, even if women with PCOS were younger than controls in this particular study⁷⁹. Moreover, in PCOS women, lipolysis responsiveness was reduced in SC and increased in visceral adipocytes when compared to controls⁷⁹. Additional studies are required to assess whether adipocyte hypertrophy in PCOS could represent the missing link between adipose tissue dysfunction and the metabolic alterations observed in this condition.

Cardiovascular endpoints

To the best of our knowledge, there is no data available on FCS and cardiovascular endpoints. One study related visceral adipocyte volume positively to arterial stiffness, a known cardiovascular disease risk factor⁸⁰. Mean SC gluteal FCS has also been positively correlated with mean arterial blood pressure⁸¹. Ledoux et al.²¹ found in patients with hypertension that mean OM and SC abdominal cell size were increased. Other studies did not report such an association.

FCS as a predictor of metabolic alterations in weight loss studies

A large number of studies have been performed on weight loss through various modalities (bariatric surgery, diet and/or exercise) and have documented the impact of such

interventions on adipocyte size reductions. However, only a few studies have attempted to link the favorable metabolic effect of weight loss to reduced FCS. Some studies have shown that weight loss-induced decreases in FCS were associated with changes in plasma levels of leptin, adiponectin, glucose, insulin, triglycerides, cholesterol, LDL-cholesterol as well as with changes in HOMA-IR, glucose disposal rate and systolic/diastolic blood pressure independent of total BFM^{70,82–85}.

Cancer

The link between obesity and the risk of mortality from cancer is well known^{86,87}. However, the biological processes underlying this association are still being investigated. Breast adipose tissue has been of interest recently due to its relative proximity with cancer cells. Hypertrophic adipocytes can induce a permanent pro-tumorous microenvironment as a source of: (1) growth factors such as estrogen, IGF-1 and leptin; (2) constant energy supply; and (3) pro-inflammatory cytokines such as IL-6 and TNF- α . In fact, *in vitro* studies have shown that mature adipocytes are able to sustain and promote tumor growth⁸⁸. Therefore, hypertrophic mammary adipocytes could reflect dysfunctional adipose tissue contributing to a microenvironment that favors cancer growth. Only one team has examined FCS in mammary fat tissue in women with breast cancer and reported a positive correlation with BMI and other markers of inflammation, such as increased NF- κ B binding and number of macrophages⁸⁹. More studies are necessary to understand the link among obesity, FCS and breast cancer.

Adipose tissue dysfunction and hypertrophic adipocytes

An increasingly large body of evidence points toward dysfunctional adipose tissue as a major determinant of metabolic impairments in individuals with abdominal obesity. Hypertrophy of the adipose cells may represent a marker, and perhaps to some extent, a driver of adipose tissue dysfunction. After providing an overview of how FCS relates to metabolic abnormalities, we now review the basis of adipose tissue dysfunction and its relationship with adipocyte hypertrophy.

Lipid metabolism

The traditional belief has been that lipid metabolism is stimulated in large adipocytes, both from the standpoint of fatty acid uptake and fatty acid release through lipolytic pathways. For example, elegant studies comparing populations of small (35 μ m) versus large (50 μ m) diameter adipocytes from the same anatomical location have shown that fatty acid synthase and lipoprotein lipase activities were increased in large compared to small adipocytes when expressed per number of cells⁹⁰. In that study, the lipolytic response to the β -adrenergic agonist isoproterenol was also increased in large versus small adipocytes. Interestingly, the beta(1)-integrin/ERKs signaling pathway was also activated in large adipocytes and had been proposed as a putative mechanism of the adaptation of adipose tissue function to cell size^{90,91}. Consistent with these results, we found that adipocyte

isoproterenol-stimulated lipolysis was higher in women with hypertrophic adipocytes, independent of overall adiposity and body fat distribution⁶². Furthermore, Laucikienė et al. demonstrated that hormone-induced lipolysis rates were increased in large (100 μ m) compared to small (82 μ m) diameter SC cells. They also observed that the protein levels of hormone-sensitive lipase (HSL), perilipin (lipid droplet-associated protein, PLIN) and adipose triglyceride lipase (ATGL) were increased in large adipocytes⁹². In contrast, opposite results were obtained in lean and obese children, in whom a significant negative correlation was found between basal lipolysis of isolated adipocytes (expressed per number of cells) and abdominal SC FCS⁵⁴. The latter association was lost upon statistical control for BMI, and lipolytic responsiveness to isoproterenol was not associated with FCS⁵⁴. A study using an original fluorescence-based technique to assess lipid uptake in individual cells has reached opposite conclusions with SC adipose tissue explants from monkeys under insulin-stimulated conditions⁹³. Specifically, small cells of the explants responded to insulin by increasing lipid uptake, whereas adipocytes with cell diameters >80–100 μ m were insulin resistant. Data were expressed per cell area in that study. It was proposed that such a mechanism could protect adipocytes from lipid overload and restrict further expansion of adipose tissue⁹³. Additional studies in 3T3-L1 cells have shown highly dynamic lipid trafficking among cellular compartments and between lipid droplets⁹⁴. Interestingly, rates of exchange were lower in cells with larger lipid droplets compared to those with smaller lipid droplets, suggesting that cells with large lipid droplets are less efficient in transporting and possibly metabolizing fatty acids than those with small lipid droplets⁹⁴. An *in vivo* study was performed by the group of Jensen and collaborators to assess rates of lipid uptake in small, medium or large adipocytes with radioactive and stable-isotope-labeled fatty acids⁹⁵. Interestingly, when expressed per lipid weight, no difference in lipid uptake was found among small (83 μ m), medium (103 μ m) or large (117 μ m) diameter adipocytes. On the other hand, when expressed per cell number, larger adipocytes had higher rates of lipid uptake⁹⁵. These experiments did not include insulin-stimulated conditions.

Considering the above-cited studies, it is difficult to reach firm conclusions regarding the impact of cell size on lipid metabolism in adipose tissue. Much confusion may arise from the method chosen to express the data. For example, in a study of diacylglycerol acyltransferase activity in isolated microsomal fractions from SC and OM adipose tissues, we have shown no relationship between maximal activity of this enzyme and adipocyte size when data were expressed per cell number⁹⁶. Yet, when expressed per mg of tissue, activity of the enzyme in OM tissue was lower in patients with excess visceral adipose tissue accumulation and large OM adipocytes⁹⁶. Consistent results were obtained in other *in vivo* studies by Votruba and collaborators, showing that net lipid uptake from a meal expressed per adipose tissue weight, but not per cell, was decreased as a function of adipocyte size in both upper and lower body SC fat⁹⁷. These results suggest that a given amount of tissue seems to take up fewer fatty acids and synthesize triglyceride less efficiently when adipocytes are larger, despite lack of a difference or increased

metabolism on a per cell basis. The study by Serra and collaborators⁹⁸ also reported a positive relationship between abdominal SC FCS and adipose tissue heparin releasable lipoprotein lipase activity (expressed per number of cells) in the same depot, but only in obese women with a low relative accumulation of visceral fat. The authors concluded that high visceral fat accumulation relative to total abdominal fat reflected lower triglyceride storage capacity, at least in the SC fat compartment⁹⁴. Inconsistencies among studies may also be explained by a loss of heterogeneity in insulin-stimulated lipid uptake of individual cells with increasing FCS and levels of obesity. This has been demonstrated by Varlamov and collaborators in white adipose tissue explants of rhesus macaques⁹⁹. Additional factors such as an obesity-related decrease in the ability to increase adipose tissue blood flow in response to a meal may also interfere with lipid storage in a given fat compartment¹⁰⁰, which could help to explain some of the discrepancies among *in vitro* and *in vivo* studies.

Glucose metabolism

Some studies have shown that there is no difference in small versus large fat cells from the same patient for protein levels of IRS-1 and GLUT4^{101,102}, whereas another study reported increased GLUT4 protein in large versus small adipocytes from the same compartment⁹⁰. Insulin stimulation significantly increased GLUT4 translocation in small (81 μm) but not in large (114 μm) diameter cells from the same individual, indicating an insulin resistant state consistent with the reduction of insulin-stimulated lipid uptake described in the previous section on lipid metabolism¹⁰¹. Consistent with these results, we reported a decrease in GLUT4 mRNA transcripts in SC adipose tissue of women with hypertrophic OM adipocytes compared to those with adipocyte hyperplasia⁶².

Adiponectin and leptin

Some studies have shown that expression and secretion of adipose-derived cytokines or adipokines may vary as a function of adipocyte size and location^{103,104}. A full review of all cytokines is beyond the scope of this review. This section focuses on leptin and adiponectin, two adipokines secreted almost exclusively by adipocytes. Inflammatory cytokines will be discussed in the subsection on inflammation.

Many reports have shown a negative association between SC FCS and adiponectin release^{19,61,69,103,105,106}, even after adjustment for SC adipose tissue area¹⁹. One study reported no association between OM FCS or SC FCS and adiponectin levels²⁴. No association was found between SC FCS and serum adiponectin in another study in children⁵⁴. Another group reported that this association was non-significant when expressed as a function of cell surface¹⁰³. In South Asians, adiponectin decreased at lower obesity levels compared to Caucasians, likely due to ethnicity-related differences in FCS or in body fat distribution⁶⁹. Of interest, our group found that adiponectin release by isolated OM mature adipocytes (expressed as a function of lipid weight of the cell suspension) was reduced with increasing BFM and OM FCS¹⁰⁴. On the other hand, adiponectin release by SC fat cells remained unaffected by differences in total BFM or SC FCS¹⁰⁴.

Adipocytes appear to secrete more leptin as their size increases, at least for those derived from the SC fat compartment^{19,24,103,107–109}. This is consistent with other studies showing that SC FCS is an independent predictor of plasma leptin concentration²⁴ and that SC FCS is the second most important predictor of leptin concentration after total BFM¹⁰⁷. SC FCS predicted 16.2% of the variance in leptin levels¹⁰⁹. SC FCS seems to be related to leptin levels, even when expressed per cell surface units¹⁰³ an association that is fully explained by differences in SC adipose tissue areas¹⁹. However, in OM adipocytes, the association is equivocal. OM fat cells express lower levels of leptin mRNA¹¹⁰ and it was thought that FCS and/or differences in tissue innervation contributed to this phenomenon. Recent studies showed no correlation²⁴ or even a negative relationship¹¹¹ between serum leptin levels and OM FCS.

Inflammation

Hypertrophic adipocytes overexpress NF- κ B¹¹² and TNF- α ^{106,113,114} independent of BMI and total BFM¹¹⁵. Other studies demonstrated that levels of the inflammation marker C-reactive protein were correlated with the presence of larger adipocytes in obese individuals^{51,106,114}. However, this association was not found in lean individuals when they were analyzed separately from the obese subgroup¹¹⁴.


The presence and number of macrophages in adipose tissues relate to their dysfunction and systemic inflammation. Macrophages in crown-like structures are often observed in adipose tissues of mice on a high fat diet, but their occurrence in human fat is far less common¹¹⁶. We observed only a few crown-like structures in obese and more specifically in SC adipose tissue¹¹⁶. No correlation was found between the number of crown-like structures and adipocyte hypertrophy in humans^{65,74,76,117}.

We examined 40 women for whom SC and visceral adipose tissue was obtained and assessed the number of macrophages in the tissue using the CD68 marker and found positive correlations between percentage of macrophages in SC and OM adipose tissue, as well as SC and OM FCS, respectively¹¹⁶. We also examined the CD11c and CD11b markers to distinguish subtypes of macrophages. We found that visceral and SC adipose tissue area was strongly correlated with CD11b and CD11c expression in SC tissue, but not in visceral fat. Consistent with this result, OM and SC FCS were associated with expression of both CD11b and CD11c, but in SC tissue only. Increased SC FCS was associated positively with CD68 + macrophages in children. When FCS was categorized into tertiles, there was a 3-fold increase in the number of macrophages between the first (<116 μm) and last (>130 μm) FCS tertile⁵⁴. Further characterization of adipose tissue immune cells is needed to better understand how cell size relates to the low-grade, chronic inflammation state of obesity.

Hypoxia/angiogenesis

Hypoxia and mitochondrial dysfunction could explain some of the mechanisms underlying adipose tissue dysfunction. Reactive oxygen species production could trigger an inflammatory response and cellular death. Hallgren et al.¹¹⁸ found

Figure 2. Obesity is a multifactorial disease characterized by expansion of adipose tissue occurring through adipocyte hypertrophy (enlargement of pre-existing cells) or adipocyte hyperplasia (generation of new cells through adipogenesis). Limited expandability of adipose tissue through hyperplasia leads to increases in FCS (adipocyte hypertrophy), which represents a critical marker of central adiposity, adipose tissue dysfunction and concomitant metabolic disease risk.



that large diameter cells (88 μm) from both lean and obese individuals consumed more O_2 than small diameter cells (66 μm) regardless of the phenotype. This finding was not supported by recent work from Yin et al. who also showed that for lean patients, FCS increases consumption rate of O_2 , but this relation was not present in obese patients¹¹⁹. Moreover, obese patients had a lower O_2 consumption rate than their lean counterpart, regardless of FCS¹¹⁹. This finding is consistent with a few other studies that found no link between FCS and angiogenic capacity as measured by expression of von Willebrand Factor (vWF), hypoxia-inducible factor 1 (HIF- α) or vascular endothelial growth factor A (VEGFA)^{21,61,120}. One study reported a positive association between FCS and VEGF receptor 2 (VEGF-R2) expression and with the number of vessels per 10 adipocytes¹²¹. We also examined women characterized by OM adipocyte hypertrophy and found higher expression of vWF and CD31 in both fat compartments compared to women with hyperplasia⁶². Another study found a negative association between FCS (OM/SC) and capillary density¹²². The relationship between dysfunctional adipocytes and angiogenesis necessitates further investigation since it appears in the obese phenotype, but its association with FCS remains unclear.

Adipogenesis

A complete review of adipogenesis is beyond the scope of this article and has already been done by our group¹²³. Adipogenic capacity is depot- and obesity-specific¹²³ and is apparently reduced in hypertrophic obesity¹²⁴.

Elegant studies by Arner and colleagues have shown that adipocyte hypertrophy is negatively correlated with adipocyte hyperplasia, when adjusted for the predicted adipocyte volume at a given BFM⁴⁵. Subjects with higher FCS than predicted by the regression curve had a lower rate of adipogenesis. These individuals also had a more deleterious metabolic profile. These results support the hypothesis that

each individual has a different plateau of maximum FCS and when it is reached, metabolic alterations arise.

In our recent analysis of published data, we confirmed the notion that adipogenesis is reduced in overweight and obese women with adipocyte hypertrophy¹²³. Moreover, in another study¹²⁴, we initiated *in vitro* primary pre-adipocyte cultures of cells obtained from the SC and OM adipose tissues of 35 women. We assessed adipogenic capacity by measuring lipid accumulation (oil red staining) and glyceraldehyde 3-phosphate dehydrogenase (G3PDH) activity as a terminal differentiation marker. We found that lower SC pre-adipocyte differentiation capacity was related to increased OM FCS, excess visceral adipose tissue accumulation, high VLDL lipid content and slightly elevated fasting glycemia¹²⁴. These results are consistent with the notion that limited expandability of SC adipose tissue is related to adipocyte hypertrophy, particularly in the visceral fat compartment, and the concomitant presence of metabolic alterations.

Conclusions and perspectives

In conclusion, our analysis demonstrates that FCS may represent a significant biomarker of the cardiometabolic alterations related to obesity. In particular, many studies demonstrated that FCS predicts alterations in the blood lipid profile and glucose-insulin homeostasis independent of adiposity indices. The contribution of visceral adiposity to these associations seems to be of particular significance. We propose that excess visceral adipose tissue accumulation and adipocyte hypertrophy, either in the SC or visceral fat compartment depending on the studies, may represent strong markers of a common phenomenon: limited hyperplastic capacity of adipose tissues, which in turn associates with the presence of numerous cardiometabolic alterations (Figure 2).

However, we also emphasize that available studies are far from unanimous, in particular when addressing the relationship between FCS and parameters of adipose tissue function.

A number of methodological factors such as the approach used to express the data may have confounded these analyses. Additional factors to be considered for future studies on this topic include analyses of cell subpopulations and the plateauing of cell size with increasing obesity levels. These issues are addressed below.

Most investigators have used average FCS as the primary variable in their analyses, which has become the default standard. However, this measure has been revealed to be incomplete when global tissue physiology is studied. Other approaches to better characterize the dynamics of cell populations have already been discussed, including osmium fixation and Multisizer Counting. This approach could represent a very useful method to characterize cell populations in more detail. Interestingly, the fraction of very small cells has become of interest in patients with severe obesity when it was found to be increased in the presence of the metabolic syndrome. Adipose tissue expansion could lead to changes in the various cell populations that may be missed when examining mean FCS. Analysis of cell size distribution patterns, of the ratio of small-to-large cells, as well as minimal and maximal cell sizes are potential alternatives.

Another important factor to consider in analyses based on adipocyte size is the patient population under study. As shown in Figure 1, mean FCS increases with adiposity level and reaches a plateau at higher obesity levels. Accordingly, studies in obese and severely obese individuals have shown that mean FCS only slightly differs among these subgroups of patients. In these cases, it is expected that the predictive ability of FCS will be reduced. The relationship between FCS and adiposity indices becomes non-linear, which may limit the effectiveness of regression studies to control for the effect of adiposity. More importantly, the association between FCS and metabolic parameters may also become non-linear in the obese or very obese range. This phenomenon could indicate that the “cardiometabolic pathology” has reached another stage, where adipose tissue storage capacity is very limited and other features such as ectopic fat accumulation will become better predictors of metabolic status than FCS *per se*. Further studies are needed to better understand the impact of adipocyte hypertrophy on metabolic health and the ways to manage it.

Declaration of interest

The authors have nothing to disclose. Work cited in this study was supported by operating grants from the Canadian Institutes of Health Research (CIHR) to AT (MOP-64182) and KC (MOP-63336). SL is the recipient of studentships from the Centre de recherche en endocrinologie moléculaire et génomique humaine (CREMOGH) and from the Institut universitaire de cardiologie et de pneumologie de Québec (IUCPQ). AM is funded by a scholarship from Fonds de la recherche du Québec en santé (FRQS).

References

- Ng M, Fleming T, Robinson M, et al. Global, regional, and national prevalence of overweight and obesity in children and adults during 1980–2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet* 2014;384:766–81.
- Tchernof A, Despres JP. Pathophysiology of human visceral obesity: an update. *Physiol Rev* 2013;93:359–404.
- Blüher M. The distinction of metabolically ‘healthy’ from ‘unhealthy’ obese individuals. *Curr Opin Lipidol* 2010;21:38–43.
- Kramer CK, Zinman B, Retnakaran R. Are metabolically healthy overweight and obesity benign conditions?: a systematic review and meta-analysis. *Ann Intern Med* 2013;159:758–69.
- Vague J. Sexual differentiation, a factor affecting the forms of obesity. *Presse Méd* 1947;55:339–40.
- Brochu M, Tchernof A, Dionne IJ, et al. What are the physical characteristics associated with a normal metabolic profile despite a high level of obesity in postmenopausal women? *J Clin Endocrinol Metab* 2001;86:1020–5.
- Kloting N, Fasshauer M, Dietrich A, et al. Insulin-sensitive obesity. *Am J Physiol Endocrinol Metab* 2010;299:E506–15.
- Drolet R, Richard C, Sniderman AD, et al. Hypertrophy and hyperplasia of abdominal adipose tissues in women. *Int J Obes* 2008;32:283–91.
- Ryden M, Andersson DP, Bergstrom IB, Arner P. Adipose tissue and metabolic alterations—Regional differences in fat cell size and number matter, but differently – a cross sectional study. *J Clin Endocrinol Metab* 2014;99:E1870–6.
- Tchoukalova YD, Votruba SB, Tchkonina T, et al. Regional differences in cellular mechanisms of adipose tissue gain with overfeeding. *Proc Natl Acad Sci USA* 2010;107:18226–31.
- Kissebah AH, Vydellingum N, Murray R, et al. Relation of body fat distribution to metabolic complications of obesity. *J Clin Endocrinol Metab* 1982;54:254–60.
- Joe AW, Yi L, Even Y, et al. Depot-specific differences in adipogenic progenitor abundance and proliferative response to high-fat diet. *Stem Cells* 2009;27:2563–70.
- Bernstein RS, Grant N, Kipnis DM. Hyperinsulinemia and enlarged adipocytes in patients with endogenous hyperlipoproteinemia without obesity or diabetes mellitus. *Diabetes* 1975;24:207–13.
- Bjorntorp P, Grimby G, Sanne H, et al. Adipose tissue fat cell size in relation to metabolism in weight-stable, physically active men. *Horm Metab Res* 1972;4:182–6.
- Haller H, Leonhardt W, Hanefeld M, Julius U. Relationship between adipocyte hypertrophy and metabolic disturbances. *Endokrinologie* 1979;74:63–72.
- Hoffstedt J, Arner E, Wahrenberg H, et al. Regional impact of adipose tissue morphology on the metabolic profile in morbid obesity. *Diabetologia* 2010;53:2496–503.
- Veilleux A, Caron-Jobin M, Noel S, et al. Visceral adipocyte hypertrophy is associated with dyslipidemia independent of body composition and fat distribution in women. *Diabetes* 2011;60:1504–11.
- Azuma K, Heilbronn LK, Albu JB, et al. Adipose tissue distribution in relation to insulin resistance in type 2 diabetes mellitus. *Am J Physiol Endocrinol Metab* 2007;293:E435–42.
- Chandalia M, Lin P, Seenivasan T, et al. Insulin resistance and body fat distribution in South Asian men compared to Caucasian men. *PLoS One* 2007;2:e812.
- Henninger AM, Eliasson B, Jenndahl LE, Hammarstedt A. Adipocyte hypertrophy, inflammation and fibrosis characterize subcutaneous adipose tissue of healthy, non-obese subjects predisposed to type 2 diabetes. *PLoS One* 2014;9:e105262.
- Ledoux S, Coupaye M, Essig M, et al. Traditional anthropometric parameters still predict metabolic disorders in women with severe obesity. *Obesity* 2010;18:1026–32.
- O’Connell J, Lynch L, Cawood TJ, et al. The relationship of omental and subcutaneous adipocyte size to metabolic disease in severe obesity. *PLoS One* 2010;5:e9997.
- Hardy OT, Perugini RA, Nicoloso SM, et al. Body mass index-independent inflammation in omental adipose tissue associated with insulin resistance in morbid obesity. *Surg Obes Relat Dis* 2011;7:60–7.
- Lundgren M, Svensson M, Lindmark S, et al. Fat cell enlargement is an independent marker of insulin resistance and ‘hyperleptinaemia’. *Diabetologia* 2007;50:625–33.
- Faria G, Pestana D, Aral M, et al. Metabolic score: insights on the development and prediction of remission of metabolic syndrome after gastric bypass. *Ann Surg* 2014;260:279–86.

26. Rodbell M. Metabolism of isolated fat cells. I. Effects of hormones on glucose metabolism. *J Biol Chem* 1964;239:375–80.
27. Ashwell M, Priest P, Bondoux M, et al. Human fat cell sizing – a quick, simple method. *J Lipid Res* 1976;17:190–2.
28. Honnor RC, Dhillon GS, Londos C. cAMP-dependent protein kinase and lipolysis in rat adipocytes. I. Cell preparation, manipulation, and predictability in behavior. *J Biol Chem* 1985; 260:15122–9.
29. Church CD, Berry R, Rodeheffer MS. Isolation and study of adipocyte precursors. *Methods Enzymol* 2014;537:31–46.
30. Hirsch J, Gallian E. Methods for the determination of adipose cell size in man and animals. *J Lipid Res* 1968;9:110–19.
31. DeMartinis FD. Very small fat cell populations determined by a modified osmium tetroxide-urea. *Am J Physiol* 1985;249:C89–96.
32. Etherton TD, Thompson EH, Allen CE. Improved techniques for studies of adipocyte cellularity and metabolism. *J Lipid Res* 1977; 18:552–7.
33. Bjurulf P. Atherosclerosis and body-build with special reference to size and number of subcutaneous fat cells. *Acta Med Scand Suppl* 1959;349:1–99.
34. Schremmer CN. [Weight changes of various tissues following formaline fixation]. *Frankf Z Pathol* 1967;77:299–304.
35. Majka SM, Miller HL, Helm KM, et al. Analysis and isolation of adipocytes by flow cytometry. *Methods Enzymol* 2014;537: 281–96.
36. Lee YH, Chen SY, Wiesner RJ, Huang YF. Simple flow cytometric method used to assess lipid accumulation in fat cells. *J Lipid Res* 2004;45:1162–7.
37. Bernstein RL, Hyun WC, Davis JH, et al. Flow cytometric analysis of mature adipocytes. *Cytometry* 1989;10:469–74.
38. Julien P, Despres JP, Angel A. Scanning electron microscopy of very small fat cells and mature fat cells in human obesity. *J Lipid Res* 1989;30:293–9.
39. Guo W, Bigornia S, Leizerman I, et al. New scanning electron microscopic method for determination of adipocyte size in humans and mice. *Obesity* 2007;15:1657–65.
40. Bjornheden T, Jakubowicz B, Levin M, et al. Computerized determination of adipocyte size. *Obes Res* 2004;12:95–105.
41. Galarraga M, Campion J, Munoz-Barrutia A, et al. Adiposoft: automated software for the analysis of white adipose tissue cellularity in histological sections. *J Lipid Res* 2012;53:2791–6.
42. Tchoukalova YD, Harteneck DA, Karwoski RA, et al. A quick, reliable, and automated method for fat cell sizing. *J Lipid Res* 2003;44:1795–801.
43. Chen HC, Farese Jr RV. Determination of adipocyte size by computer image analysis. *J Lipid Res* 2002;43:986–9.
44. Osman OS, Selway JL, Kepczynska MA, et al. A novel automated image analysis method for accurate adipocyte quantification. *Adipocyte* 2013;2:160–4.
45. Arner E, Westermark PO, Spalding KL, et al. Adipocyte turnover: relevance to human adipose tissue morphology. *Diabetes* 2010;59: 105–9.
46. Imbeault P, Lemieux S, Prud'homme D, et al. Relationship of visceral adipose tissue to metabolic risk factors for coronary heart disease: is there a contribution of subcutaneous fat cell hypertrophy? *Metabolism* 1999;48:355–62.
47. Heinonen S, Saarinen L, Naukkarinen J, et al. Adipocyte morphology and implications for metabolic derangements in acquired obesity. *Int J Obes* 2014;38:1423–31.
48. Meena VP, Seenu V, Sharma MC, et al. Relationship of adipocyte size with adiposity and metabolic risk factors in Asian Indians. *PLoS One* 2014;9:e108421.
49. Bjorntorp P, Bengtsson C, Blohme G, et al. Adipose tissue fat cell size and number in relation to metabolism in randomly selected middle-aged men and women. *Metabolism* 1971;20:927–35.
50. Cotillard A, Poitou C, Torcivia A, et al. Adipocyte size threshold matters: link with risk of type 2 diabetes and improved insulin-resistance after gastric bypass. *J Clin Endocrinol Metab* 2014;99: E1466–70.
51. Despres JP, Nadeau A, Tremblay A, et al. Role of deep abdominal fat in the association between regional adipose tissue distribution and glucose tolerance in obese women. *Diabetes* 1989;38: 304–9.
52. Monickaraj F, Gokulakrishnan K, Prabu P, et al. Convergence of adipocyte hypertrophy, telomere shortening and hypoadiponectinemia in obese subjects and in patients with type 2 diabetes. *Clin Biochem* 2012;45:1432–8.
53. Weyer C, Foley JE, Bogardus C, et al. Enlarged subcutaneous abdominal adipocyte size, but not obesity itself, predicts type II diabetes independent of insulin resistance. *Diabetologia* 2000;43: 1498–506.
54. Landgraf K, Rockstroh D, Wagner IV, et al. Evidence of early alterations in adipose tissue biology and function and its association with obesity-related inflammation and insulin resistance in children. *Diabetes* 2015;64:1249–61.
55. Joffe BI, Goldberg RB, Feinstein J, et al. Adipose cell size in obese Africans: evidence against the existence of insulin resistance in some patients. *J Clin Pathol* 1979;32:471–4.
56. Dubois SG, Heilbronn LK, Smith SR, et al. Decreased expression of adipogenic genes in obese subjects with type 2 diabetes. *Obesity* 2006;14:1543–52.
57. Roberts R, Hodson L, Dennis AL, et al. Markers of de novo lipogenesis in adipose tissue: associations with small adipocytes and insulin sensitivity in humans. *Diabetologia* 2009;52:882–90.
58. McLaughlin T, Sherman A, Tsao P, et al. Enhanced proportion of small adipose cells in insulin-resistant vs insulin-sensitive obese individuals implicates impaired adipogenesis. *Diabetologia* 2007; 50:1707–15.
59. Pasarica M, Xie H, Hymel D, et al. Lower total adipocyte number but no evidence for small adipocyte depletion in patients with type 2 diabetes. *Diabetes Care* 2009;32:900–2.
60. Yang J, Eliasson B, Smith U, et al. The size of large adipose cells is a predictor of insulin resistance in first-degree relatives of type 2 diabetic patients. *Obesity* 2012;20:932–8.
61. Hammarstedt A, Graham TE, Kahn BB. Adipose tissue dysregulation and reduced insulin sensitivity in non-obese individuals with enlarged abdominal adipose cells. *Diabetol Metab Syndr* 2012;4:42.
62. Michaud A, Boulet MM, Veilleux A, et al. Abdominal subcutaneous and omental adipocyte morphology and its relation to gene expression, lipolysis and adipocytokine levels in women. *Metabolism* 2014;63:372–81.
63. Kranendonk ME, van Herwaarden JA, Stupkova T, et al. Inflammatory characteristics of distinct abdominal adipose tissue depots relate differently to metabolic risk factors for cardiovascular disease: distinct fat depots and vascular risk factors. *Atherosclerosis* 2015;239:419–27.
64. Srdic B, Stokic E, Korac A, et al. Morphological characteristics of abdominal adipose tissue in normal-weight and obese women of different metabolic profiles. *Exp Clin Endocrinol Diabetes* 2010; 118:713–18.
65. van Beek L, Lips MA, Visser A, et al. Increased systemic and adipose tissue inflammation differentiates obese women with T2DM from obese women with normal glucose tolerance. *Metabolism* 2014;63:492–501.
66. Lackey DE, Burk DH, Ali MR, et al. Contributions of adipose tissue architectural and tensile properties toward defining healthy and unhealthy obesity. *Am J Physiol Endocrinol Metab* 2014;306: E233–46.
67. Alberti KG, Eckel RH, Grundy SM, et al. Harmonizing the metabolic syndrome: a joint interim statement of the International Diabetes Federation Task Force on Epidemiology and Prevention; National Heart, Lung, and Blood Institute; American Heart Association; World Heart Federation; International Atherosclerosis Society; and International Association for the Study of Obesity. *Circulation* 2009;120:1640–5.
68. Alberti KG, Zimmet P, Shaw J. Metabolic syndrome—a new worldwide definition: a Consensus Statement from the International Diabetes Federation. *Diabet Med* 2006;23:469–80.
69. Anand SS, Tarnopolsky MA, Rashid S, et al. Adipocyte hypertrophy, fatty liver and metabolic risk factors in South Asians: the Molecular Study of Health and Risk in Ethnic Groups (mol-SHARE). *PLoS One* 2011;6:e22112.
70. Larson-Meyer DE, Heilbronn LK, Redman LM, et al. Effect of calorie restriction with or without exercise on insulin sensitivity, beta-cell function, fat cell size, and ectopic lipid in overweight subjects. *Diabetes Care* 2006;29:1337–44.
71. Petaja EM, Sevastianova K, Hakkarainen A, et al. Adipocyte size is associated with NAFLD independent of obesity, fat distribution, and PNPLA3 genotype. *Obesity* 2013;21:1174–9.

72. Koska J, Stefan N, Votruba SB, et al. Distribution of subcutaneous fat predicts insulin action in obesity in sex-specific manner. *Obesity* 2008;16:2003–9.
73. Johannsen DL, Tchoukalova Y, Tam CS, et al. Effect of eight weeks of overfeeding on ectopic fat deposition and insulin sensitivity: testing the “adipose tissue expandability” hypothesis. *Diabetes Care* 2014;37:2789–97.
74. Jansen HJ, Vervoort GM, van der Graaf M, et al. Liver fat content is linked to inflammatory changes in subcutaneous adipose tissue in type 2 diabetes patients. *Clin Endocrinol* 2013;79:661–6.
75. Wree A, Schlattjan M, Bechmann LP, et al. Adipocyte cell size, free fatty acids and apolipoproteins are associated with non-alcoholic liver injury progression in severely obese patients. *Metabolism* 2014;63:1542–52.
76. Aron-Wisnewsky J, Minville C, Tordjman J, et al. Chronic intermittent hypoxia is a major trigger for non-alcoholic fatty liver disease in morbid obese. *J Hepatol* 2012;56:225–33.
77. Manneras-Holm L, Leonhardt H, Kullberg J, et al. Adipose tissue has aberrant morphology and function in PCOS: enlarged adipocytes and low serum adiponectin, but not circulating sex steroids, are strongly associated with insulin resistance. *J Clin Endocrinol Metab* 2011;96:E304–11.
78. Faulds G, Ryden M, Ek I, et al. Mechanisms behind lipolytic catecholamine resistance of subcutaneous fat cells in the polycystic ovarian syndrome. *J Clin Endocrinol Metab* 2003;88:2269–73.
79. Ek I, Arner P, Ryden M, et al. A unique defect in the regulation of visceral fat cell lipolysis in the polycystic ovary syndrome as an early link to insulin resistance. *Diabetes* 2002;51:484–92.
80. Arner P, Backdahl J, Hemmingsson P, et al. Regional variations in the relationship between arterial stiffness and adipocyte volume or number in obese subjects. *Int J Obes* 2015;39:222–7.
81. Achimastos A, Raison J, Levenson J, Safar M. Adipose tissue cellularity and hemodynamic indexes in obese patients with hypertension. *Arch Intern Med* 1984;144:265–8.
82. Andersson DP, Eriksson Hogling D, Thorell A, et al. Changes in subcutaneous fat cell volume and insulin sensitivity after weight loss. *Diabetes Care* 2014;37:1831–6.
83. Lofgren P, Andersson I, Adolfsson B, et al. Long-term prospective and controlled studies demonstrate adipose tissue hypercellularity and relative leptin deficiency in the postobese state. *J Clin Endocrinol Metab* 2005;90:6207–13.
84. Rizkalla SW, Prifti E, Cotillard A, et al. Differential effects of macronutrient content in 2 energy-restricted diets on cardiovascular risk factors and adipose tissue cell size in moderately obese individuals: a randomized controlled trial. *Am J Clin Nutr*. 2012; 95:49–63.
85. Varady KA, Tussing L, Bhutani S, Braunschweig CL. Degree of weight loss required to improve adipokine concentrations and decrease fat cell size in severely obese women. *Metabolism* 2009; 58:1096–101.
86. Calle EE, Kaaks R. Overweight, obesity and cancer: epidemiological evidence and proposed mechanisms. *Nat Rev Cancer* 2004; 4:579–91.
87. Renehan AG, Tyson M, Egger M, et al. Body-mass index and incidence of cancer: a systematic review and meta-analysis of prospective observational studies. *Lancet* 2008;371:569–78.
88. D’Esposito V, Passaretti F, Hammarstedt A, et al. Adipocyte-released insulin-like growth factor-1 is regulated by glucose and fatty acids and controls breast cancer cell growth in vitro. *Diabetologia* 2012;55:2811–22.
89. Morris PG, Hudis CA, Giri D, et al. Inflammation and increased aromatase expression occur in the breast tissue of obese women with breast cancer. *Cancer Prev Res* 2011;4:1021–9.
90. Farnier C, Krief S, Blache M, et al. Adipocyte functions are modulated by cell size change: potential involvement of an integrin/ERK signalling pathway. *Int J Obes Relat Metab Disord* 2003;27:1178–86.
91. Farnier C, Krief S, Blache M, et al. The signaling pathway for beta1-integrin/ERKs is involved in the adaptation of adipocyte functions to cell size. *Ann N Y Acad Sci* 2002;973: 594–7.
92. Laurencikiene J, Skurk T, Kulyte A, et al. Regulation of lipolysis in small and large fat cells of the same subject. *J Clin Endocrinol Metab* 2011;96:E2045–9.
93. Varlamov O, Somwar R, Cornea A, et al. Single-cell analysis of insulin-regulated fatty acid uptake in adipocytes. *Am J Physiol Endocrinol Metab* 2010;299:E486–96.
94. Somwar R, Roberts Jr CT, Varlamov O. Live-cell imaging demonstrates rapid cargo exchange between lipid droplets in adipocytes. *FEBS Lett* 2011;585:1946–50.
95. Rajjo TI, Harteneck DA, Jensen MD. Direct free fatty acid storage in different sized adipocytes from the same depot. *Obesity* 2014; 22:1275–9.
96. Cote JA, Nadeau M, Leboeuf M, et al. Adipose tissue diacylglycerol acyltransferase activity and blood lipoprotein triglyceride enrichment in women with abdominal obesity. *Atherosclerosis* 2014;233:172–7.
97. Votruba SB, Mattison RS, Dumesic DA, et al. Meal fatty acid uptake in visceral fat in women. *Diabetes* 2007;56:2589–97.
98. Serra MC, Ryan AS, Sorkin JD, et al. High adipose LPL activity and adipocyte hypertrophy reduce visceral fat and metabolic risk in obese, older women. *Obesity (Silver Spring)* 2015;23:602–7.
99. Varlamov O, Chu M, Cornea A, et al. Cell-autonomous heterogeneity of nutrient uptake in white adipose tissue of rhesus macaques. *Endocrinology* 2015;156:80–9.
100. McQuaid SE, Hodson L, Neville MJ, et al. Downregulation of adipose tissue fatty acid trafficking in obesity: a driver for ectopic fat deposition? *Diabetes* 2011;60:47–55.
101. Franck N, Stenkula KG, Ost A, et al. Insulin-induced GLUT4 translocation to the plasma membrane is blunted in large compared with small primary fat cells isolated from the same individual. *Diabetologia* 2007;50:1716–22.
102. Jernas M, Palming J, Sjöholm K, et al. Separation of human adipocytes by size: hypertrophic fat cells display distinct gene expression. *FASEB J* 2006;20:1540–2.
103. Skurk T, Alberti-Huber C, Herder C, Hauner H. Relationship between adipocyte size and adipokine expression and secretion. *J Clin Endocrinol Metab* 2007;92:1023–33.
104. Drolet R, Belanger C, Fortier M, et al. Fat depot-specific impact of visceral obesity on adipocyte adiponectin release in women. *Obesity* 2009;17:424–30.
105. Vargas G, Chandalia M, Jiang Y, et al. Heterogeneity in subcutaneous adipose tissue morphology and metabolic complications in overweight and obese women. *Metab Syndr Relat Disord* 2013;11:276–82.
106. Bahceci M, Gokalp D, Bahceci S, et al. The correlation between adiposity and adiponectin, tumor necrosis factor alpha, interleukin-6 and high sensitivity C-reactive protein levels. Is adipocyte size associated with inflammation in adults? *J Endocrinol Invest* 2007;30:210–14.
107. Wahlen K, Sjölin E, Lofgren P. Role of fat cell size for plasma leptin in a large population based sample. *Exp Clin Endocrinol Diabetes* 2011;119:291–4.
108. Van Harmelen V, Reynisdottir S, Eriksson P, et al. Leptin secretion from subcutaneous and visceral adipose tissue in women. *Diabetes* 1998;47:913–17.
109. Couillard C, Mauriege P, Imbeault P, et al. Hyperleptinemia is more closely associated with adipose cell hypertrophy than with adipose tissue hyperplasia. *Int J Obes Relat Metab Disord* 2000; 24:782–8.
110. Hube F, Lietz U, Igel M, et al. Difference in leptin mRNA levels between omental and subcutaneous abdominal adipose tissue from obese humans. *Horm Metab Res* 1996;28:690–3.
111. Arner P, Andersson DP, Thorne A, et al. Variations in the size of the major omentum are primarily determined by fat cell number. *J Clin Endocrinol Metab* 2013;98:E897–901.
112. Maury E, Noel L, Detry R, Brichard SM. In vitro hyperresponsiveness to tumor necrosis factor-alpha contributes to adipokine dysregulation in omental adipocytes of obese subjects. *J Clin Endocrinol Metab* 2009;94:1393–400.
113. Hube F, Birgel M, Lee YM, Hauner H. Expression pattern of tumour necrosis factor receptors in subcutaneous and omental human adipose tissue: role of obesity and non-insulin-dependent diabetes mellitus. *Eur J Clin Invest* 1999;29:672–8.
114. Winkler G, Kiss S, Keszthelyi L, et al. Expression of tumor necrosis factor (TNF)-alpha protein in the subcutaneous and visceral adipose tissue in correlation with adipocyte cell volume, serum TNF-alpha, soluble serum TNF-receptor-2 concentrations and C-peptide level. *Eur J Endocrinol* 2003;149:129–35.

115. Arner E, Ryden M, Arner P. Tumor necrosis factor alpha and regulation of adipose tissue. *N Engl J Med* 2010;362:1151–3.
116. Michaud A, Drolet R, Noel S, et al. Visceral fat accumulation is an indicator of adipose tissue macrophage infiltration in women. *Metabolism* 2012;61:689–98.
117. El Bouazzaoui F, Henneman P, Thijssen P, et al. Adipocyte telomere length associates negatively with adipocyte size, whereas adipose tissue telomere length associates negatively with the extent of fibrosis in severely obese women. *Int J Obes* 2014;38:746–9.
118. Hallgren P, Sjostrom L, Hedlund H, et al. Influence of age, fat cell weight, and obesity on O₂ consumption of human adipose tissue. *Am J Physiol* 1989;256:E467–74.
119. Yin X, Lanza IR, Swain JM, et al. Adipocyte mitochondrial function is reduced in human obesity independent of fat cell size. *J Clin Endocrinol Metab* 2014;99:E209–16.
120. Villaret A, Galitzky J, Decaunes P, et al. Adipose tissue endothelial cells from obese human subjects: differences among depots in angiogenic, metabolic, and inflammatory gene expression and cellular senescence. *Diabetes* 2010;59:2755–63.
121. Lemoine AY, Ledoux S, Queguiner I, et al. Link between adipose tissue angiogenesis and fat accumulation in severely obese subjects. *J Clin Endocrinol Metab* 2012;97:E775–80.
122. Faber DR, Moll FL, Vink A, et al. Adipose tissue quantity and composition contribute to adipokine concentrations in the subclavian vein and the inferior mesenteric vein. *Int J Obes* 2012;36:1078–85.
123. Lessard J, Tchernof A. Depot-and obesity-related differences in adipogenesis. *Clin Lipidol* 2012;7:587–96.
124. Lessard J, Laforest S, Pelletier M, et al. Low abdominal subcutaneous preadipocyte adipogenesis is associated with visceral obesity, visceral adipocyte hypertrophy, and a dysmetabolic state. *Adipocyte* 2014;3:197–205.