



Characterization of the Structure of *tert*-butyl[1-hydroxy-2-methyl-3-(1*H*-1,2,4-triazol-1-yl)]propan-2-ylcarbamate Using 2D Heteronuclear NMR Experiments

Younas Aouine¹, Hassane Faraj¹, Anouar Alami^{1*}, Abdelilah El-Hallaoui¹ and Mohamed Akhazzane²

¹Laboratoire de Chimie Organique Fès, Faculté des Sciences Dhar El Mahraz, Université Sidi Mohamed Ben Abdellah, Morocco

²Cité de l'innovation, Université Sidi Mohamed Ben Abdellah, Fès, Morocco

Abstract

The alkylation of 1*H*-1,2,4-triazole with an *O*-tosyloxazoline derivative, followed by an oxazoline ring-opening reaction and protection of amine function, leads normally to obtain two regioisomers of β -aminoalcohols. After purification by column chromatography of the crude reaction product, only a single product is obtained. Hence, there is need of its identification by spectroscopic study.

Keywords

Oxazoline, 1*H*-1,2,4-triazole, β -aminoalcohol, 2D heteronuclear NMR experiments

Introduction


Triazoles constitute an important class of biologically active heterocyclic compounds that have received a great deal of attention since their discovery. Diverse compounds derived from 1,2,4-triazoles have a wide spectrum activities, including antimicrobial [1,2] and antibacterial properties [3,4], human antifungal agents [5], anticancer agents [6], antiviral [7], antitumor activity [8] and in agricultural science as potent fungicides, herbicides and insecticides [9,10]. Amino acids containing the 1,2,4-triazole moiety and their derivatives represent a well-known group of organic compounds also presenting biological activity. Thus β -(1,2,4-triazol-1-yl)-L-alanine is known as an important metabolite in plants of the fungicide myclobutanil [11-13] and β -(3-amino-1,2,4-triazol-1-yl)-L-alanine is a metabolite of the weed killer 3-amino-1,2,4-triazole [14].

Materials and Methods

NMR spectra (¹H, ¹³C and ¹⁵N) were recorded on a Bruker AM 300 (operating at 300.13 MHz for ¹H, at 75.47 MHz for ¹³C and at 30.41 MHz for ¹⁵N) spectrometer (Centre Universitaire Régional d'Interface, Fez-Morocco). NMR data are listed in ppm and are reported relative to tetra-methylsilane (¹H, ¹³C).

Discussion

It is reported that afforded the corresponding 1- and 4-alkylated isomers, with prevalence of the N₁-isomer [15-17]. Reaction of 1*H*-1,2,4-triazole with **1** and K₂CO₃, was carried out in the presence of a catalytic amount of tetrabutylammonium bromide in *N,N*-dimethylformamide at 120°C for 12 hours. The previous reaction stage is followed by an oxazoline ring-opening reaction carried out in acidic medium. The aminoalcohol derivative **2**, which is the aim of this paper, is obtained after addition of *tert*-butoxycarbonyl anhydride Boc to the intermediate product in a mixture of water/dioxane (1/2) at (0 < T < 5°C) in the presence of triethylamine [18]. Its structure was established on the basis of NMR spectroscopy (¹H, ¹³C and ¹⁵N), in addition to MS data and elemental analysis.



The definite assignment the chemical shifts of protons, carbons and nitrogens of compound **2** are shown in table 1 and table 2 (Figure 1,

*Corresponding author: Anouar Alami, Laboratoire de Chimie Organique Fès, Faculté des Sciences Dhar El Mahraz, Université Sidi Mohamed Ben Abdellah, Morocco, Tel: +212-661 796 480, Fax: +212-535 733 171, E-Mail: alamianouar@yahoo.fr


Received: January 12, 2016; Accepted: March 12, 2016; Published: March 16, 2016

Copyright: © 2016 Aouine Y, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Citation: Aouine Y, Faraj H, Alami A, El-Hallaoui A, Akhazzane M (2016) Characterization of the Structure of *tert*-butyl[1-hydroxy-2-methyl-3-(1*H*-1,2,4-triazol-1-yl)]propan-2-ylcarbamate Using 2D Heteronuclear NMR Experiments. Int J Exp Spectroscopic Tech 1:004

Table 1: ^1H (300 MHz) and ^{13}C (75.47 MHz) NMR spectral data for **2** in DMSO-*d*₆, including results obtained by homonuclear 2D shift-correlated ^1H COSY and heteronuclear 2D shift-correlated HMQC ($^1J_{\text{CH}}$)^a. Chemical shifts (δ , ppm) and coupling constants (*J*, Hz, in parenthesis)^b.

Position	δ_{H}	δ_{C}	Correlation	
			^1H - ^1H	^1H - ^{13}C
(NH) 1	6.29 (s)	-	1-H	-
2	-	56.66	-	-
3	3.35, 3.45 (AB (2dd), 10.8; 5.6)	64.85	H ¹ -3, H ² -3, O-H	H ¹ -3, H ² -3, C-3
(OH) 4	4.92 (t, 5.6)	-	O-H, H ¹ -3, H ² -3	-
5	1.18 (s)	20.43	H ¹ -5, H ² -5, H ³ -5	H ^{1,2,3} -5; C-5
6	4.35, 4.51 (AB, 14)	51.74	H ¹ -6, H ² -6	H ¹ -6, H ² -6, C-6
9	7.95 (s)	151.61	H ³ triazole -9	H ³ triazole -9, C-9
11	8.23 (s)	145.24	H ⁵ triazole -11	H ⁵ triazole -11, C-11
12	-	154.97	-	-
14	-	78.43	-	-
15; 17; 18	1.39 (s)	28.68	H ^{1,2,3} -15	H ^{1,2,3} -15; C-15
			H ^{1,2,3} -17	H ^{1,2,3} -17; C-17
			H ^{1,2,3} -18	H ^{1,2,3} -18; C-18

^aCorrelation from C to the indicated hydrogens.^bChemical shifts and coupling constants (*J*) obtained of 1D ^1H -NMR spectrum.**Figure 1:** Homonuclear ^1H - ^1H 2D spectrum for compound **2** between 3 and 6.5 ppm.**Table 2:** Listing of ^{15}N (400 MHz) NMR spectral data for **2** in DMSO-*d*₆, including results obtained by heteronuclear single quantum coherence shift-correlated (HSQC) and heteronuclear multiple bond coherence shift-correlated (HMBC).

Position	δ_{H}	δ_{N}	Correlation
(NH) 1	7.02 (s)	93.29	H-1, N-1
5	1.24 (s)	93.29	H ^{1,2,3} -5, N-1
6	4.39, 4.83 (AB, 14)	93.29	H ^{1,2} -6, N-1
		214.54	H ^{1,2} -6, N-7
		299.12	H ^{1,2} -6, N-8
9	7.96 (s)	214.54	H ³ triazole-9, N-7
		251.97	H ³ triazole-9, N-10
		299.12	H ³ triazole-9, N-8
11	8.20 (s)	214.54	H ⁵ triazole-11, N-7
		251.97	H ⁵ triazole-11, N-10

Chemical shifts (δ , ppm) and coupling constants (*J*, Hz, in parenthesis) obtained of 1D ^1H -NMR spectrum. ^{15}N NMR spectrum of the compound **2** shows four signals corresponding to four nitrogen atoms and assigning each signal to its corresponding nitrogen atom is carried out based mainly on the spectral data ^1H - ^{15}N HMBC (Figure 3).

Figure 2 and Figure 3).

This interaction ^1H - ^{15}N has allowed us to make the following observations:




- A signal at 93.29 ppm attributed to the carbamate nitrogen N-1: correlation between CH_3 , CH_2 -triazole and the amidic nitrogen.

- A signal at 214.54 ppm attributed to the N-7 of the 1,2,4-triazole ring: correlation between the two triazole protons, CH_2 -triazole and N-7 nitrogen at position 1 of the 1,2,4-triazole ring.

- A signal at 251.97 ppm attributed to the N-10 of the 1,2,4-triazole ring: interaction between the two triazole protons and the N-10 located in the position 4 of the 1,2,4-triazole ring.

- A signal at 299.12 ppm attributed to N-8 of the 1,2,4-triazole ring: interaction between CH_2 -triazole, triazole proton H-9 and N-8 at position 2 of the 1,2,4-triazole ring.

Further, the analysis of ^1H - ^{15}N HMBC spectrum of compound **2** confirms that the nucleophilic substitution reaction of 1,2,4-triazole



Citation: Aouine Y, Faraj H, Alami A, El-Hallaoui A, Akhazzane M (2016) Characterization of the Structure of *tert*-butyl[1-hydroxy-2-methyl-3-(1*H*-1,2,4-triazol-1-yl)]propan-2-ylcarbamate Using 2D Heteronuclear NMR Experiments. Int J Exp Spectroscopic Tech 1:004

and *O*-tosyloxazoline derivative **1** is carried out on the nitrogen in position 1 of the 1,2,4-triazole ring and its structure has been clearly identified (Figure 4).

References

1. Xu W, Song B, Bhadury P, Song Y, Hu D (2010) Synthesis and crystal structure of novel sulfone derivatives containing 1,2,4-triazole moieties. *Molecules* 15: 766-779.
2. Karabasanagouda T, Adhikari AV, Shetty NS (2007) Synthesis and antimicrobial activities of some novel 1,2,4-triazolo[3,4-b]-1,3,4-thiadiazoles and 1,2,4-triazolo[3,4-b]-1,3,4-thiadiazines carrying thioalkyl and sulphonyl phenoxy moieties. *Eur J Med Chem* 42: 521-529.
3. Tehranchian S, Akbarzadeh T, Fazeli MR, Jamalifar H, Shafiee A (2005) Synthesis and antibacterial activity of 1-[1,2,4-triazol-3-yl] and 1-[1,3,4-thiadiazol-2-yl]-3-methylthio-6,7-dihydrobenzo[c]thiophen-4(5H) ones. *Bioorg Med Chem Lett* 15: 1023-1025.
4. Foroumadi A, Soltani F, Moshafi MH, Ashraf-Askari R (2003) Synthesis and in vitro antibacterial activity of some N-(5-aryl-1,3,4-thiadiazole-2-yl)piperazinyl quinolone derivatives. *Farmaco* 58: 1023-1028.
5. Chen Q, Zhu XL, Jiang LL, Liu ZM, Yang GF (2008) Synthesis, antifungal activity and CoMFA analysis of novel 1,2,4-triazolo[1,5-a]pyrimidine derivatives. *Eur J Med Chem* 43: 595-603.
6. Holla BS, Poojary KN, Rao BS, Shivananda MK (2002) New bis-aminomercaptotriazoles and bis-triazolothiadiazoles as possible anticancer agents. *Eur J Med Chem* 37: 511-517.
7. Al-Soud YA, Al-Dweri MN, Al-Masoudi NA (2004) Synthesis, antitumor and antiviral properties of some 1,2,4-triazole derivatives. *Farmaco* 59: 775-783.
8. DemirbaÅY N, Ugurluoglu R, DemirbaÅY A (2002) Synthesis of 3-alkyl(aryl)-4-alkylidenamino-4,5-dihydro-1H-1,2,4-triazol-5-ones and 3-alkyl-4-alkylamino-4,5-dihydro-1H-1,2,4-triazol-5-ones as antitumor agents. *Bioorg Med Chem* 10: 3717-3723.
9. Crispino GA, Goudar JS (2001) Process for the preparation of a triazolinone herbicide. US Pat 6268506 B1.
10. Song BA, Hu DY, Zheng S, Huang RM, Yang S, et al. (2001) Synthesis and bioactivity of phosphorodithioate compounds containing 1,2,4-triazole. *Chin J Org Chem* 21: 524-529.
11. Arnold M, Evrard D, Leander J, Lodge D, Schoep D, et al. (1993) Tetrazole amino acids as competitive NMDA antagonists. *Bioorg Med Chem Lett* 3: 43-48.
12. Ebert B, Lenz S, Brehm L, Bregnedal P, Hansen JJ, et al. (1994) Resolution, absolute stereochemistry, and pharmacology of the S-(+)- and R-(-)-isomers of the apparent partial AMPA receptor agonist (R,S)-2-amino-3-(3-hydroxy-5-phenylisoxazol-4-yl)propionic acid [(R,S)-APPA]. *J Med Chem* 37: 878-884.
13. Ikegami F, Murakoshi I (1994) Enzymic synthesis of non-protein b-substituted alanines and some higher homologues in plants. *Phytochemistry* 35: 1089-1104.
14. Dunnill P, Fowden L (1965) The amino acids of seeds of the Cucurbitaceae. *Phytochemistry* 4: 933-944.
15. Bulger PG, Cotterell IF, Cowden CJ, Davies AJ, Dolling UH (2000) An investigation into the alkylation of 1,2,4-triazole. *Tetrahedron Lett* 41: 1297-1301.
16. Boukallaba K, Elachqar A, El Hallaoui A, Alami A, El Hajji S, et al. (2006) Synthesis of new alpha-heterocyclic alpha-aminophosphonates. *Phosphor Sulfur Silicon* 108: 819-823.
17. Bentama A, El Hadrami EM, El Hallaoui A, Elachqar A, Lavergne JP, et al. (2003) Synthesis of new alpha-heterocyclic alpha-aminoesters. *Amino Acids* 24: 423-426.
18. Aouine Y, Faraj H, Alami A, El-Hallaoui A, Elachqar A, et al. (2011) Simple and efficient synthesis of racemic 2-(tert-butoxycarbon-ylamino)-2-methyl-3-(1H-1,2,4-triazol-1-yl)propanoic acid, a new derivative of Î²-(1,2,4-triazol-1-yl) alanine. *Molecules* 16: 3380-3390.