

Effects of Dietary Supplemental Ascorbic Acid and Cholecalciferol on Bone Characteristics of Hens at the Late Laying Stage

Ogunwole O. A.^{1,*}, Adedeji B. S.¹, Olumide M. D.², Mosuro A. O.¹, I. E. Adewemimo¹

¹Agricultural Biochemistry and Nutrition Unit, Department of Animal Science, University of Ibadan, Nigeria

²Department of Animal Science, School of Agriculture and Industrial Technology, Babcock University, Ilishan Remo, Ogun State, Nigeria

Abstract Effect of dietary inclusion of vitamin C and D₃ supplementation on bone characteristics of hens aged 70 weeks at the late laying stage. Lohman Brown (n=180) were randomly allotted to 12 dietary treatments in triplicates of 5 birds each. The formulated basal diet was supplemented with three levels of vitamin C (0, 300 and 600 mg/kg) and four levels of vitamin D₃ (0, 1000, 2000 and 3000 IU/kg). The experiment was a 3x4 factorial arrangement in a completely randomised design. Two hens per replicate were selected, sacrificed and bone physico-chemical attributes assessed. Tibia length of hens on 2000 IU D₃ (115.60) was significantly higher (P < 0.05) than those on 0 (114.76), 1000 (114.47) and 3000 IU D₃ (113.66) supplementation. Bone robusticity index of hens on supplemental vitamin C were higher (P < 0.05) at 0 mg (30.36) than at 300 (29.10) and 600 mg (28.60) while inclusion of D₃ at 1000 (29.20), 2000 (29.32) and 3000 IU (28.86) were similar (P > 0.05) but lower than when supplemented at 0 IU (30.04). Bone calcium, phosphorus, magnesium, potassium and ash differed significantly (P < 0.05) at various inclusion levels of vitamin C and D₃ while the effect of interaction of supplemental vitamin C and D₃ on tibia bones of the hens was only significant (P < 0.05) for tibia calcium. Tibia calcium ranged between 12.69 to 16.83% in hens on combined dietary supplement of 0 IU D₃+ 600 mg vitamin C and 3000 IU D₃+600 mg vitamin C, respectively. In conclusion, dietary supplements of vitamin C and D₃ improved the physical and chemical characteristics of spent laying hens' bone.

Keywords Robusticity index, Tibia bone composition, Supplemental vitamin, Lohmann Brown hens, Laying hens bone attributes

1. Introduction

Deficiency and loss in bone mineral structure is a function of skeletal problem (osteoporosis) in modern highly productive laying chickens [1]. Bone fracture is one of the most serious skeletal problems in modern laying hens [2]. Randall and Duff [3] reported osteoporosis as the major reason for bone loss and subsequent fractures in laying hens. The role of nutrition in the maintenance of bone integrity cannot be undermined. Birds that have laid consistently over a long time are usually susceptible to bone brittleness and fracture. Adequate supplemental vitamin C or D₃ may help reduce the incidence of skeletal abnormality often observed in spent birds. Diets of hens well fortified with the required nutrients for bone health will impact the skeletal integrity of

hens at the late laying stage.

Available reports on the roles of vitamins on skeletal integrity in laying hens were inconsistent. Earlier report by Newman and Leeson [4] revealed that dietary vitamin C had little effects on bone breaking strength of laying leghorn hens. Improved bone parameters were reported in broiler chickens [5] while [6] did not observe any influence of dietary vitamin C on tibia bone physical parameters. The influence of dietary vitamin D on tibia and femur bones of chickens has been similarly reported [7-10]. Increased bone strength was only achieved in broiler chickens when dietary vitamin C supplementation was concomitant with vitamin D [11]. Dietary ascorbic acid affects bone resorption marker enzymes [11] and thus its concentration may affect the integrity of bone matrix. Fritts and Waldroup [10] reported that vitamin C influences the conversion of vitamin D into its metabolic active form calcitriol which is essential for calcium and phosphorous regulation and calcification processes.

However, information on the combined effect of dietary vitamin C and D₃ on bone health is scanty in literature.

* Corresponding author:

droaogunwole@gmail.com (Ogunwole O. A.)

Published online at <http://journal.sapub.org/food>

Copyright © 2018 The Author(s). Published by Scientific & Academic Publishing

This work is licensed under the Creative Commons Attribution International

License (CC BY). <http://creativecommons.org/licenses/by/4.0/>

Therefore, this study was aimed at assessing the effects of supplemental vitamin C and D₃ on bone characteristics of spent laying hens.

2. Materials and Methods

2.1. Experimental Site

The experiment was carried out at the Poultry Unit, Teaching and Research Farm, University of Ibadan, Ibadan, Nigeria. The university is located in Ibadan, in the tropical rain forest zone of Nigeria within latitude 7° 26 N and longitude 3° 54 E, with a mean altitude of 277 meters above sea level.

2.2. Experimental Animal and Diets

A total of 180 Bovan brown hens aged 12 weeks were randomly allocated into 12 treatments of three replicates of five hens per replicate in a 3x4 factorial arrangement of a completely randomized design. The basal diet for the study was supplemented with three levels of vitamin C (0, 300 and 600 mg/kg) and four levels of vitamin D (0, 1000, 2000 and 3000 IU/kg) and their combination to produce 12 treatments. The basal experimental diet composition is shown in Table 1. The experiment lasted 60 weeks.

Table 1. Gross composition (%) of basal diet fed to the laying hens

Ingredient	% Inclusion (Kg)
Maize	50.00
Soya bean meal	22.00
Wheat offal	11.08
Palm kernel cake	11.24
Oyster shell	3.03
Di-calcium phosphate	1.70
Vitamin-mineral premix	0.25
Table Salt	0.30
Biotronics	0.30
DL-methionine	0.15
L-lysine	0.15
Total	100.00
Calculated nutrients	
ME (Kcal/kg)	2702.90
Crude protein (%)	17.39
Crude fibre (%)	5.25
Methionine (%)	0.40
Lysine (%)	0.98
Calcium (%)	1.80
Available phosphorus (%)	0.52

Vitamin-mineral premix*: Vitamin A–10,000IU, Vitamin D₃–2000IU, Vitamin E–40mg, Vitamin K–1.43 mg, Vitamin B1–0.7mg, Vitamin B2–4mg, Vitamin B6–2.5mg, Vitamin B12–0.2mg, Niacin–10mg, Panthothenic–10,000mg, Folic acid –0.25mg, Biotin–100mg, Choline Chloride–300mg, Manganese–80mg, Zinc–60mg, Iron–40mg, Copper–80mg, Iodine–0.8mg, Selenium–0.2mg, Cobalt–0.3mg, Antioxidant–100mg

2.3. Data Collection

At week 70, two hens per replicate were selected and sacrificed. They were carefully dissected into primal cuts. The left and right tibia of each bird was removed with the drumsticks and flesh intact. The drumsticks were labeled and immersed in boiling water (100°C) for 10 minutes. After cooling to room temperature, the drumsticks were defleshed manually and then air dried for 24 hours at room temperature. Bone physical parameters measured were: bone weight, bone length and robusticity index while bone chemical composition measured included calcium, phosphorus, magnesium and potassium. The bone weight/length index was obtained by dividing the tibia weight by its length. The tibiotarsal and robusticity indexes were determined according to Adebisi *et al.* [12]. The percentage ash was determined relative to dry weight of tibia while calcium, phosphorus, magnesium and potassium composition were analysed according to AOAC [13].

2.4. Statistical Analysis

Data were subjected to polynomial regression, descriptive statistics and analysis of variance (SAS, 2003) while means were separated using Tukey's HSD option of the same package at $\alpha_{0.05}$.

3. Results

The main effect of dietary supplementation of vitamin C on bone physical characteristics of laying chickens is shown in Table 2. Tibia length, tibia weight, tibia length/weight index and robusticity index differed significantly ($P<0.05$) across the treatments. Tibia length of hens fed supplemental vitamin C at 0 (116.70) and 300 mg (114.35) were similar ($P>0.05$). They were however higher ($P<0.05$) than 113.11 cm in hens fed 600 mg vitamin C. Tibia weight increased significantly ($P<0.05$) with supplemental vitamin C from 6.73g in hens fed 0 mg vitamin C to 7.04 and 7.11g in hens fed 300 and 600 mg vitamin C, respectively. Tibia length/weight index in hens on 0 mg dietary vitamin C (57.52) was significantly lower ($P<0.05$) than those on 300 (61.49) and 600 mg (63.00) supplemental vitamin C. Robusticity index was significantly higher ($P<0.05$) in hens on 0 mg dietary vitamin C (30.36) compared with 29.10 and 28.60 in hens fed 300 and 600 mg vitamin C, respectively. Robusticity index of bone in hens fed 300 mg (29.10) and 600 mg (28.60) dietary vitamin C were similar ($P<0.05$) but were lower ($P<0.05$) than those on 0 mg dietary vitamin C (30.36).

The main effect of inclusion of dietary vitamin D₃ supplementation on bone physical characteristics of laying chickens is shown in Table 3. Tibia length, tibia weight, tibia length/tibia weight, and robusticity index differed significantly ($P<0.05$) with dietary supplement of vitamin D₃. Tibia length of hens on 0 IU (114.76) was not significantly different ($P>0.05$) from those on 1000 (114.47) and 3000 IU

(113.66) vitamin D but were lower ($P < 0.05$) than those on 2000 IU D₃ (115.60). Tibia length of hens on 1000 and 2000 IU however were not significantly different ($P > 0.05$). Tibia weight of hens on 0 IU (6.46) was similar ($P > 0.05$) to those of hens on 1000 IU (6.96) but lower than 7.12 and 7.30 in hens on 2000 and 3000 IU, respectively.

Tibia length/weight index of hens on 0 IU (56.26) was significantly lower compared with hens fed 1000 (60.90), 2000 (62.83) and 3000 IU D₃ (62.69). Significantly higher ($P < 0.05$) bone robusticity index was observed in hens fed 0 IU D₃ (30.04) compared with 29.20, 29.32 and 28.86 in hens on 1000, 2000 and 3000 IU D₃, respectively.

The relationship between increased supplemental vitamin C in the diet of hens and the bone robusticity index was negatively correlated and is shown in Figure 1 and further explained by equation 1.

$$Y = -0.001x + 30.49 \quad (R^2 = 0.799) \quad (1)$$

The relationship between increased supplementation of dietary vitamin D₃ and bone robusticity index is shown in Figure 2 and explained by equation 2. Supplemental vitamin D₃ was negatively and significantly ($P < 0.05$) correlated with the bone robusticity index of laying hens.

$$Y = -5E-0.7x^2 + 0.001x + 30.37 \quad (R^2 = 0.998) \quad (2)$$

Table 2. Main effect of supplemental vitamin C on laying hens' bone physical characteristics

Parameters	Inclusion levels (mg)			SEM	P values
	0	300	600		
Tibia length (cm)	116.70 ^a	114.35 ^{ab}	113.11 ^b	1.37	0.04
Tibia weight (g)	6.73 ^b	7.04 ^a	7.11 ^a	0.53	0.03
Tibia length/weight index	57.52 ^b	61.49 ^a	63.00 ^a	4.41	0.04
Robusticity index	30.36 ^a	29.10 ^b	28.60 ^b	0.71	0.03

^{a,b,c} means on the same row with different superscripts are significantly different ($P < 0.05$)

Table 3. Main effect of supplemental vitamin D on laying hens' bone physical characteristics

Parameters	Inclusion levels (IU)				SEM	P Values
	0	1000	2000	3000		
Tibia length (cm)	114.76 ^b	114.47 ^{ab}	115.60 ^a	113.66 ^b	1.58	0.04
Tibia weight (g)	6.46 ^b	6.96 ^{ab}	7.12 ^a	7.30 ^a	0.60	0.04
Tibia length/weight index	56.26 ^b	60.90 ^a	62.83 ^a	62.69 ^a	5.10	0.04
Robusticity index	30.04 ^a	29.20 ^b	29.32 ^b	28.86 ^b	0.81	0.03

^{a,b,c} means on the same row with different superscripts are significantly different ($P < 0.05$); SEM: Standard Error of Means




Figure 1. Relationship between supplemental vitamin C and bone robusticity index of laying hens




Figure 2. Relationship between supplemental vitamin D₃ and bone robusticity index of laying hens

The main effect of dietary inclusion of vitamin C supplementation on bone chemical characteristics of laying chicken is shown in Table 4. Calcium, phosphorus, magnesium, potassium and ash differed significantly ($P < 0.05$) with the increased dietary inclusion of vitamin C. Tibia calcium increased with dietary supplementation of vitamin C. Hens on 0 mg supplemental vitamin C had tibia calcium of 15.16 that was similar ($P > 0.05$) to 14.80 in those on 300 mg but significantly higher ($P < 0.05$) than 13.65 in hens on 0 mg vitamin C. Tibia phosphorus in hens fed 600 mg vitamin C (15.33) was higher than those at 0 (14.28) and 300 mg (14.37) supplementation. Significantly higher ($P < 0.05$) bone magnesium was obtained at 300 mg (0.47) supplemental level of vitamin C compared with 0.45 and 0.43 in hens on 0 and 600 mg vitamin C supplementation, respectively. Tibia potassium of hens fed 0 mg (0.40) vitamin C supplementation was significantly higher ($P < 0.05$) than in those fed 300 (0.35) and 600 mg (0.37) supplemental vitamin C. Tibia ash of hens on 600 mg supplemental vitamin C was significantly higher ($P < 0.05$) than 52.88 and 53.18 in hens fed 0 and 300 mg vitamin C, respectively.

Table 4. Main effect of supplemental vitamin C on bone chemical composition of laying hens

Parameters (%)	Inclusion levels (mg)			SEM	P value
	0	300	600		
Calcium	13.65 ^b	14.80 ^{ab}	15.16 ^a	0.51	0.01
Phosphorus	14.28 ^b	14.37 ^b	15.33 ^a	0.58	0.02
Magnesium	0.45 ^b	0.47 ^a	0.43 ^b	0.03	0.04
Potassium	0.40 ^a	0.35 ^b	0.37 ^b	0.02	0.01
Ash	52.88 ^b	53.18 ^b	55.40 ^a	2.09	0.04

^{a,b,c} means on the same row with different superscripts are significantly different ($P < 0.05$); SEM: Standard Error of Means

The main effect of dietary inclusion of vitamin D₃ supplementation on bone chemical characteristics of spent

laying chickens is shown in Table 5. Tibia calcium, phosphorus, magnesium, potassium and ash differed significantly ($P < 0.05$) with varying inclusion of supplemental vitamin D₃. Tibia calcium was significantly higher ($P < 0.05$) at 3000 IU (15.21) than 13.52 at 0 IU supplemental level but similar to those on 1000 (14.98) and 2000 IU D₃ (14.44). Tibia phosphorus at 2000 IU (15.34) was higher than 13.76, 14.70 and 14.84 in hens on 0, 1000 and 3000 IU supplemental D₃, respectively. Tibia magnesium of hens was higher at 1000 IU (0.48) compared with those on 1000 IU and other treatments. Potassium in the tibia of laying hens (0.39, 0.38 and 0.37) on 0, 2000 and 3000 IU D₃ were similar ($P > 0.05$) and higher than 0.33 in hens on 1000 IU D₃. Bone ash of hens on 0 IU dietary inclusion of vitamin D₃ (56.29) was higher than 52.94, 52.57 and 53.49 obtained at 1000, 2000 and 3000 IU inclusion of Vitamin D₃, respectively.

The relationship between tibia Ca and P with increased dietary supplement of vitamin C is shown in Figure 3 and explained by equations 3 and 4. There were significantly negative linear correlations of the variables.

$$Y = -0.000x + 13.9 \quad (R^2 = 0.067) \quad (3)$$

$$Y = -0.000x + 14.02 \quad (R^2 = 0.445) \quad (4)$$

The relationships between supplemental dietary vitamin C and tibia Mg and K is shown in Figure 4. The significantly negative linear relationships ($P < 0.05$) are represented in equations 5 and 6. The relationship among bone Ca and P compositions as affected by supplemental dietary vitamin D₃ are shown in Figure 5. The polynomial regressions were quadratic and are explained in equations 7 and 8. As shown in Figure 6 and equation 9, the relationship between increased inclusion levels of vitamin D₃ and bone Mg composition was quadratic and the level of vitamin D₃ for optimum Mg in the bone was 1000 IU. The significantly negative correlation of bone K composition with increased

supplementation of vitamin D₃ is shown in Figure 6 and the relationship is represented by equation 10.

$$Y = 6E-05x + 0.409 \quad (R^2=0.242) \quad (5)$$

$$Y = -8E-05x + 0.416 \quad (R^2=0.969) \quad (6)$$

$$Y = -4E-07x^2 + 0.000x + 13.86 \quad (R^2=0.76) \quad (7)$$

$$Y = -5E-07x^2 + 0.001x + 13.98 \quad (R^2=0.54) \quad (8)$$

$$Y = -3E-08x^2 + 0.000x + 0.415 \quad (R^2=0.61) \quad (9)$$

$$Y = -6E-09x^2 + 9E-06x + 0.401 \quad (R^2=0.12) \quad (10)$$

Table 5. Main effect of supplemental vitamin D₃ on bone chemical characteristics of laying hens

Parameters	Inclusion levels (IU)				SEM	P-value
	0	1000	2000	3000		
Calcium (%)	13.52 ^b	14.98 ^{ab}	14.44 ^{ab}	15.21 ^a	0.59	0.04
Phosphorus (%)	13.76 ^c	14.70 ^b	15.34 ^a	14.84 ^b	0.67	0.04
Magnesium (%)	0.42 ^b	0.48 ^a	0.44 ^b	0.44 ^b	0.03	0.04
Potassium (%)	0.39 ^a	0.33 ^b	0.38 ^a	0.37 ^a	0.02	0.01
Ash (%)	56.29 ^a	52.94 ^b	52.57 ^b	53.49 ^b	2.42	0.04

^{a,b,c} means on the same row with different superscripts are significantly different (P<0.05); SEM: Standard Error of Means




Figure 3. Relationship between supplemental vitamin C and laying hen tibia calcium and phosphorus




Figure 4. Relationship between supplemental vitamin C and laying hen tibia magnesium and potassium




Figure 5. Relationship between supplemental vitamin D₃ and laying hens tibia calcium and phosphorus




Figure 6. Relationship between supplemental vitamin C and laying hens tibia magnesium and potassium

The effect of interaction of supplemental vitamin C and D₃ on bone chemical characteristics in laying chickens is shown in Table 6. Among all the minerals evaluated, bone calcium only, differed significantly (P<0.05) across the treatments. Highest value of calcium (1683.33) was obtained in bones of hens on 3000 IU D₃+600 mg vitamin C and the least (1269.17) in those on 0 IU D₃+ 600 mg vitamin C.

Table 6. Effect of interaction of supplemental vitamin C and D₃ on tibia bone calcium of laying hens

Treatments		Tibia calcium (mg/dL)
Vitamin C (mg/kg)	Vitamin D (IU/kg)	
	0	13.70 ^{ab}
	1000	14.79 ^{ab}
0	2000	13.37 ^{ab}

	3000	12.73 ^b
	0	14.18 ^{ab}
300	1000	14.45 ^{ab}
	2000	14.52 ^{ab}
	3000	16.06 ^{ab}
	0	12.69 ^b
600	1000	15.69 ^{ab}
	2000	15.44 ^{ab}
	3000	16.83 ^a
	SEM	1.23
	P- value	<0.001

^{ab} means with different superscripts within the same row differ significantly (p<0.05). SEM: Standard Error of Means

4. Discussion

Bone mineralisation has been studied with respect to processes (chemical and physical) that improves bone matrix as mineralisation affects bone strength with an implication on the ability of the skeleton to withstand gravity and additional loading. Biochemical parameters are direct indicators of bone quality which include bone density [15] and bone flexibility and strength [16]. Rao *et al.* [17] reported that different measurements can be used to determine bone mineralisation, structure and health in poultry.

The improvements observed with respect to the bone parameters monitored in this study could be linked to the documented roles of both vitamins in bone mineralisation. Leeson and Summer [18] emphasized the crucial roles of vitamin C in the hydroxylation of proline residues necessary for synthesis of procollagen a precursor of bone formation. It has been recognized to play a role as a cofactor to several key enzymes involved in procollagen synthesis [19] and bioconversion of D₃ to its metabolic active form [20].

Similarly, the importance of vitamin D₃ in the absorption and utilization of calcium for bone development and strength have been reported [21, 22] reported improved bone resistance with vitamin C supplementation in chicks while Ogunwole [6] observed no beneficial effect of vitamin C on tibia bone physical parameters measured in broiler chickens. Conversely, Bruno *et al.* [7] and [5] observed decreased tibia and femur parameters with vitamin C supplementation in broiler chickens reared under heat stress. Fritts and Waldroup [10] reported increased bone weight while improvement in bone parameters was reported by Matilla *et al.* [23] with vitamin D supplementation in chickens. Sahin *et al.* [24] however observed no influence of vitamin D₃ on bone improvement in laying quails.

Higher bone weight/length index observed in this study with supplementation of vitamin C and D₃ corroborated earlier report [25] that increased bone weight/length index indicated more bone density and increased bone strength. Similar observation was reported for the effect of vitamin C and vitamin D on robusticity index of the birds as supplementation increased. Lower robusticity index has been associated with increased bone strength [26] thus emphasising that vitamin C and D supplementation in this study favoured formation of stronger bone.

Calcitriol the metabolically active form of vitamin D₃, is a secosteroid hormone involved in Ca and P metabolism [27] while vitamin C is a key player in the bioconversion of vitamin D to this active form [20]. This role of vitamin C and D could be attributed to the increased bone Ca and P observed in this study. High rate of bone resorption during lay resulted in bone weakness at the end of their production cycle but with the inclusion of vitamin D₃ there would be more calcium in the bone which reduced osteoporosis [28]. Although, there was increased calcium and phosphorus deposition in the tibia with the supplementation of either of the vitamins, magnesium and potassium content in the bone

did not follow a particular trend thus emphasizing the importance of Ca and P as the key players in bone mineralization.

The importance of vitamin D was further strengthened by Pereira [29] that vitamin D₃ deficiency could impair bone formation and as such ascribed vitamin D, calcium and phosphorus as extremely important for bone mobilization and increased bone strength. Tibia ash was reportedly increased with supplemental vitamin D₃ [10]. The effect of interaction of dietary vitamin C and D₃ supplementation on bone chemical characteristics of laying chickens was only significant for bone calcium. This further suggests the synergistic role of vitamin C and D₃ in improving calcium absorption and utilization. The involvement of vitamin C in the bioconversion of vitamin D had earlier been highlighted [20].

The influence of vitamin C and D₃ observed on increased tibia calcium deposition may be attributed to the several metabolic functions of calcium in poultry [30]. The high regression coefficients in Figures 2 and 3 suggest that robusticity index of the tibia was strongly dependent on supplemental vitamins C and D. This implied that 79.9 and 99.8% of the observed improvement in robusticity index were adduced to supplemental vitamin C and D₃, respectively. As shown in Figure 3, it seems vitamin C supplementation was more favourable to phosphorus deposition in the tibia compared to calcium as 44.5% of the deposition of P in the bone was ascribed to supplemental vitamin C. A very strong relationship between vitamin C supplementation and K deposition ($R^2=0.96$) compared with 0.24 for Mg deposition in the tibia suggests that increasing supplemental vitamin C would result in improved bone K deposition.

Supplemental vitamin D₃ on the other hand was more favourable to Ca deposition. Optimal tibia Ca and P were attained at 1400 IU supplementation of vitamin D₃. Tibia magnesium and potassium improved with supplemental vitamin D₃ as optimal tibia magnesium and potassium were attained at 1600 and 1000 IU of supplemental vitamin D₃, respectively. This implied that supplementation of vitamin D₃ beyond these levels would reduce deposition of these minerals in the bone of laying hens. Though tibia calcium in hens on the control was similar ($P>0.05$) to those of hens from other treatment combinations, hens on combination of 600 mg ascorbic acid and 3000 IU D₃ had higher tibia calcium than those on sole supplementation of 600 mg ascorbic acid and 3000 IU cholecalciferol, respectively. This observation suggests a synergy between these vitamins for tibia calcium deposition when combined at higher levels in the diets of hens.

5. Conclusions

In conclusion, supplemental vitamin C and D₃ improved physical and chemical characteristics of tibia bone in laying hens while the interaction of both vitamins in the diet of

laying hens increased tibia calcium deposition.

ACKNOWLEDGEMENTS

Authors appreciate the Tertiary Education Trust Fund (TETFUND) for their provision of research materials for this study.

REFERENCES

- [1] Abdul-Aziz, T. A. (1998). Cage layer fatigue is a complicated problem. *Word Poultry* 14: 56-58.
- [2] Gregory and Wilkins (1991). Broken bones in hens. *Veterinary Record* 129: 559.
- [3] Randall, C. J. and Duff, S. R. I. (1988). Avulsion of the patellar ligament in osteopenic laying fowl. *Vet. Rec.* 123: 439-441.
- [4] Newman, S. and Leeson, S. (1999). The effect of dietary supplementation with 1, 25-dihydroxycholecalciferol or vitamin C on the characteristics of the tibia of old layer hens. *Poultry Science.*, 78 (1) 85-90.
- [5] Sgavioli, S., Domingues, C. H. F., Santos, E. T., Quadros, T. C. O., deBorges, L. L., Garcia, R. G., Louzada, M. J. Q. L. and Boleli, I. C. (2016). Effect of In-Ovo Ascorbic Acid Injection on the Bone Development of Broiler Chickens Submitted to Heat Stress During Incubation and Rearing. *Revista Brasileira de Ciência Avícola*, 18 (1), 153-162.
- [6] Ogunwole, O. A. (2015). Bone characteristics of broiler chickens fed diets supplemented with graded levels of vitamin C. *Agric. Biol. J. Nutr. Am* 6: 30-33.
- [7] Bruno, L. D. G., Furlan, R. L. and Malheiros, E. B. (2000). Influence of early quantitative food restriction on long bone growth at different environmental temperatures in broiler chickens. *Br. Poult. Sci.* vol. 41 (pg. 389-394).
- [8] McCormack, H., McTeir, L., Fleming, R. and Whitehead, C. (2002). Vitamin D requirements of broilers at different dietary concentrations of calcium, phosphorus and vitamin A. *Arch. Gelflu'gelkd.* 66:71.
- [9] Whitehead, C. (2002). Influence of vitamins and minerals on bone formation and quality. *Arch. Gelflugelkd.* 66:45.
- [10] Fritts, C. A. and Waldroup, P. W. (2003). Effect of source and level of vitamin D on live performance and bone development in growing broilers. *J. Appl. Poult. Res.* 12:45-52.
- [11] Lohakare, J. D., Kim, J. K., Ryn, M. H., Hahn, T. W. and Chae, B. J. (2005). Effects of vitamin C and vitamin D interactions on the performance, immunity and bone characteristics of commercial broilers. *Journal of Applied Poultry Research.* 14 (4) pp. 670-678.
- [12] Adebisi, O. A., Sokunbi, O. A. and Ewuola, O. A. (2009). Performance evaluation and bone characteristics of growing cockerels fed diets containing different levels of diatomaceous earth. *Middle East Journal of Scientific Research* 4 (1): 36-39.
- [13] AOAC (1990). Association of Official Chemist, Official Methods of Analyses 15th Edition, Washington DC, USA Horwitz, W. Official methods of analysis of association of official chemist. 15th Ed. Washington: AOAC, P. 1278.
- [14] SAS (2003). SAS/STAT User's guide Version 8 for windows SAS Institute Inc. Cary NC.
- [15] Barreiro, F. R., Amaral, L. A., Shimano, A. C., Alva, J. C. R., Barbosa, J. C. and Baraldi-Artoni, S. M. (2011). Physiologic values of broiler femurs at different growth phases using boné densitometry and bone breaking strength. *International Journal of Poultry Science*; 10(7):530-533.
- [16] Reis, D. T. C., Torres, R. A., Barbosa, A. A., Rodrigues, C. S., Moraes, G. H. K. (2011). Efeito de linhagem e sexo nas características geométricas e biomecânicas de tibias de frangos de corte. *Acta Scientiarum: Animal Science*; 33(1): 101-108.
- [17] Rao, S. K., West, M. S., Frost, T. J., Orban, J. I., Bryant, M. M. and Roland Sr., D. A. (1993). Sample size required for various methods of assessing bone status in commercial leghorn hens. *Poultry Science* 72:229-235.
- [18] Leeson, S. and Summers, J. D. (2001). Scott's nutrition of the chicken, 4th edition. Nottingham, UK, Nottingham University Press.
- [19] Tillman, P. B. (1993). Vitamin C for laying hens: a review. Pages 1-31 in: Multi-State Poultry Feeding and Nutrition Conference, Indianapolis, IN.
- [20] McDowell, L. R. (2000). Vitamins in animal and human nutrition. Iowa State University Press, Ames, IA. 793 pp.
- [21] Akhter, S., Kutuzova, G. D., Christakos, S., DeLuca, H. F. (2007). Calbindin D9k is not required for 1,25-dihydroxyvitamin D3-mediated Ca²⁺ absorption in small intestine. *Archives of Biochemistry and Biophysics* RCS 7~305-313.
- [22] Bikle, D., Adams, J. and Christakos, S. (2008). Vitamin D: Production, Metabolism, Mechanism of Action, and Clinical Requirements. *American Society for Bone and Mineral Research.* 141-149.
- [23] Mattila, P., J. Valaja, L. Rossow, E. Venalainen and T. Tupasela. (2004). Effect of vitamin D-2- and D-3-enriched diets on egg vitamin D content, production, and bird condition during an entire production period. *Poultry Science* 83:433-440.
- [24] Sahin, N., Balci, T. A., Kucuk, O., Smith, M. O. and Sahin, K. (2009). Effects of 25-hydroxycholecalciferol and soy isoflavones supplementation on bone mineralisation of quail. *British Poultry Science* 50(6):709-715.
- [25] Monteagudo, M. D., Hernández, E. R., Seco, C., Gonzalez-Riola, J., Revilla, M., Villa, L. F. and Rico, H. (1997). Comparison of the bone robusticity index and bone weight/bone length index with the results of bone densitometry and bone histomorphometry in experimental studies. *Acta Anat (Basel).* 160(3):195-9.
- [26] Reisenfeld, A. (1972). Metatarsal robusticity in bipedal rats. *Am.J. Phys. Anthropol.*, 40: 229-234.
- [27] Veum, T. (2010). Phosphorus and Calcium Nutrition and Metabolism, In: Vitti, D. and Kebreab, E. eds. Phosphorus

and calcium utilization and requirements in farm animals. Wiltshire: CABI Publishing, pp.94–111.

- [28] Whitehead, C. C. (2004). Overview of bone biology in the egg-laying hen. *Poultry Science, Champaign*; 83 (1): 193-199.
- [29] Pereira, R. (2010). Universidade de São Paulo; Piracicaba, São Paulo: Eficiência de uma fitase bacteriana na liberação de fósforo fítico em dietas de frangos de corte. Msc. Dissertação.
- [30] Nunes, R. V., Pozza, P. C., Scherer, C., Campestrini, E., Rocha, L. D., Nunes, C. G. V. and Costa, F. G. P. (2006). Efeito dos teores de cálcio para poedeiras semipesadas durante a fase de pré-postura e no início da postura. *Revista Brasileira de Zootecnia*; 35(5).