

RESEARCH ARTICLE

Open Access

Frailty and falls among adult patients undergoing chronic hemodialysis: a prospective cohort study

Mara A McAdams-DeMarco^{1,2}, Sunitha Suresh³, Andrew Law^{1,2}, Megan L Salter^{1,2}, Luis F Gimenez^{4,5,6}, Bernard G Jaar^{2,4,5,6}, Jeremy D Walston⁷ and Dorry L Segev^{1,2,8*}

Abstract

Background: Patients undergoing hemodialysis are at high risk of falls, with subsequent complications including fractures, loss of independence, hospitalization, and institutionalization. Factors associated with falls are poorly understood in this population. We hypothesized that insights derived from studies of the elderly might apply to adults of all ages undergoing hemodialysis; we focused on frailty, a phenotype of physiological decline strongly associated with falls in the elderly.

Methods: In this prospective, longitudinal study of 95 patients undergoing hemodialysis (1/2009-3/2010), the association of frailty with future falls was explored using adjusted Poisson regression. Frailty was classified using the criteria established by Fried et al., as a combination of five components: shrinking, weakness, exhaustion, low activity, and slowed walking speed.

Results: Over a median 6.7-month period of longitudinal follow-up, 28.3% of study participants (25.9% of those under 65, 29.3% of those 65 and older) experienced a fall. After adjusting for age, sex, race, comorbidity, disability, number of medications, marital status, and education, frailty independently predicted a 3.09-fold (95% CI: 1.38-6.90, $P=0.006$) higher number of falls. This relationship between frailty and falls did not differ for younger and older adults ($P=0.57$).

Conclusions: Frailty, a validated construct in the elderly, was a strong and independent predictor of falls in adults undergoing hemodialysis, regardless of age. Our results may aid in identifying frail hemodialysis patients who could be targeted for multidimensional fall prevention strategies.

Keywords: Hemodialysis, Falls, Frailty

Background

Patients undergoing chronic hemodialysis are at high risk of falls: 13-25% experience a fall after hemodialysis initiation [1,2]. This population is also at high-risk for severe complications after a fall including fractures, loss of independence, hospitalization, institutionalization, and mortality [3-6]. One in seven patients undergoing chronic hemodialysis suffers a major bone fracture after a fall [7]. Additionally, fractures double the mortality risk in this population [5].

Unfortunately, falls are poorly understood in patients undergoing hemodialysis. Although some correlates have

been identified, such as age, comorbidities, and number of medication burden [2,3], the ability to predict which patients will suffer falls remains limited. Improved risk prediction would allow for better patient care through prehabilitation and protective living arrangements.

We hypothesized that insights into falls derived from studies of the elderly might inform risk prediction in adults of all ages undergoing chronic hemodialysis; briefly stated, we hypothesized that frailty would be predictive of falls in this population. Frailty is a phenotype of physiological decline strongly associated with falls in the elderly [8,9]. We have identified frailty as important risk factor for mortality and hospitalization in patients undergoing dialysis of all ages; frailty was independently associated with a 2.60-fold higher risk of death and 1.43-fold higher number of hospitalizations [10]. To our knowledge, there are no

* Correspondence: dorry@jhmi.edu

¹Department of Surgery, Johns Hopkins University School of Medicine, Baltimore, MD, USA

²Department of Epidemiology, Johns Hopkins School of Public Health, Baltimore, MD, USA

Full list of author information is available at the end of the article

studies assessing the association between frailty and falls in patients undergoing chronic hemodialysis. The primary goal of this study was to identify whether frailty was associated with falls independent of other risk factors for falls in patients undergoing hemodialysis of all ages.

Methods

Study design

This was a longitudinal study of 115 prevalent hemodialysis patients from a single dialysis center in Baltimore, Maryland, recruited between January 2009 and March 2010. Hemodialysis patients who were 18 years or older, spoke English, and agreed to participate were recruited into this study from a hospital-based outpatient dialysis unit. Frailty was measured at enrollment, and number of falls was self-reported at a follow-up visit in the 95 patients who survived at least 5 months. The follow-up visit was conducted at the dialysis center and was conducted on average within 6.7 months after enrollment. Demographics and comorbidities were obtained from medical record review at enrollment. The Johns Hopkins Institutional Review Board approved the study and all participants provided informed consent.

Frailty

Frailty was measured as defined and validated by Fried et al. [8,11-20], as a phenotype based on 5 components: shrinking (self-report of unintentional weight loss of more than 10 lbs. in the past year based on dry weight); weakness (grip-strength below an established cutoff based on gender and BMI) [8]; exhaustion (self-report); low activity (Kcals/week below an established cutoff) [8]; and slowed walking speed (walking time of 15 feet below an established cutoff by gender and height) [8]. The 5 components of frailty were measured using previously described methods [8]. A score of 1 was given to those with the presence of each measured component. The aggregate frailty score was calculated as the sum of the component scores (range 0-5) and categorized as nonfrail (0-1), intermediately frail (2), and frail (3-5), as we previously established in ESRD patients [10].

Statistical analysis

The association between frailty at enrollment and the number of falls reported during follow-up was estimated using Poisson regression. All adjusted models were adjusted for age, sex, and race. Additional models were also adjusted for comorbidity (as previously defined by Fried in the elderly and validated by our group in ESRD patients, namely 4 or more of the following conditions ascertained from medical records abstraction: peripheral vascular disease, rheumatoid arthritis, cancer, hypertension, COPD, diabetes, congestive heart failure, angina, and myocardial infarction) [8], disability in activities of

daily living [21] (also previously described by Fried in the elderly and validated by our group in ESRD patients) [8], number of medications, education, and marital status at enrollment. Interactions between frailty and age were tested using a Wald test. Additionally, as sensitivity analyses we tested whether the results were similar when 1) we used negative binomial regression and 2) adjusted for other factors including anemia, arrhythmia, and cognitive function. All analyses were performed using STATA 12.0/MP for Linux (College Station, Texas).

Results

Study population

Among 95 study participants, the mean age was 60.5 years (SD=12.6), 43.6% were over 65, 46.3% were female, and 81.1% were African American; 25.3% of participants were nonfrail, 28.4% were intermediately frail, and 46.3% were frail. The median follow-up time was 6.7 months (range 5.0-13.3), which did not differ by frailty (6.7 months vs. 6.7 months, $P=0.51$) or falls (6.7 months vs. 6.7 months, $P=0.98$). The median time on dialysis was 3.7 years.

Falls

During follow-up, 28.3% of participants had one or more falls. The total number of falls was 70 and maximum number of falls a single participant experienced was 8 (Figure 1). Participants with one or more falls did not differ from those without falls based on comorbidity, number of medications, or education (Table 1). However, there were notable but not statistically significant differences in the number of years on chronic hemodialysis (no falls: 6.0 vs. falls: 4.0 years, $P=0.17$), percent of participants with prevalent disability (no falls: 17.4% vs. falls: 38.5%, $P=0.054$), and percent of participants who were married/cohabitating (no falls: 27.5% vs. falls: 46.2%, $P=0.09$). Age was not statistically associated with falls, with 25.9% of younger adults and 29.3% of older adults experiencing one or more falls ($P=0.72$).

Frailty and falls

Frailty predicted a 3.55-fold (95% CI: 1.68-7.46, $P=0.001$) higher number of falls in the unadjusted model and predicted a 3.89-fold (95% CI: 1.78-8.49, $P=0.001$) higher number of falls after adjusting for age, sex, and race (Table 2). In a separate model that was additionally adjusted for comorbidity, disability, number of medications, education, and marital status, frailty predicted a 3.09-fold (95% CI: 1.38-6.90, $P=0.006$) higher number of falls. In all models, intermediately frailty was not itself associated with an increased number of falls, but there was a statistically significant trend in the risk of falls from nonfrail to intermediately frail to frail (all $P\leq 0.001$). The association of frailty and number of falls did not differ for younger and older adults ($P=0.57$ for interaction between age and frailty).




Figure 1 Distribution of number of falls in adults undergoing hemodialysis.

Table 1 Characteristics of the study population, by falls during longitudinal follow-up

	No falls (n=69)	Falls (n=26)	P value
Age (mean, SD)	59.9 [13.3]	62.2 [10.6]	0.44
Female (%)	44.9	50.0	0.82
Caucasian (%)	8.7	15.4	0.45
Enrollment body mass index (kg/m ²) (mean, SD)	28.8 [8.6]	29.8 [7.9]	0.59
Pre-dialysis body mass index (kg/m ²) (mean, SD)	29.9 [10.2]	32.0 [9.1]	0.37
Smoking history (%)	18.8	19.2	0.99
Years on hemodialysis (mean, SD)	6.0 [7.0]	4.0 [3.3]	0.17
Married/cohabitating (%)	27.5	46.2	0.09
Education (% high school or higher)	82.6	88.5	0.75
Comorbidities (%)			
Peripheral vascular disease	31.9	23.1	0.46
Rheumatoid arthritis	7.3	11.5	0.68
History of cancer	17.4	26.9	0.38
Hypertension	91.3	88.5	0.70
Chronic obstructive pulmonary disease	15.9	26.9	0.25
Diabetes	66.7	76.9	0.46
Congestive heart failure	29.1	42.3	0.82
Angina	5.8	7.7	0.66
Myocardial infarction	17.4	15.4	0.99
Number of medications (mean, SD)	8.7 [3.4]	8.1 [3.1]	0.42
Comorbidity (%) ¹	30.4	34.6	0.81
Disability (%) ²	17.4	38.5	0.05

¹Comorbidity = 4 or more of the following conditions: peripheral vascular disease, rheumatoid arthritis, cancer, hypertension, COPD, diabetes, congestive heart failure, angina, and myocardial infarction.

²Disability = Needing assistance on 3 or more of the following activities: feeding, dressing, ambulation, grooming, using a toilet, and bathing.

Sensitivity analyses

The association of frailty and falls was robust to the statistical methods used. Additionally, the adjustment for anemia (frailty: 2.64, 95% CI: 1.17-5.98), arrhythmias (frailty: 3.04, 95% CI: 1.36-6.80), and low cognitive

Table 2 Factors associated with falls in adults undergoing hemodialysis

	Unadjusted	Parsimonious model	Full model
Frailty status			
Nonfrail	Reference	Reference	Reference
Intermediately frail	1.11 (0.44, 2.82)	1.35 (0.52, 3.49)	1.19 (0.44, 3.24)
Frail	3.55 (1.68, 7.46)**	3.89 (1.78, 8.49)**	3.09 (1.38, 6.90)**
Age (in 10 years)	-	0.83 (0.68, 1.01)	0.90 (0.69, 1.17)
Female	-	1.68 (1.01, 2.79)*	1.84 (1.06, 3.19)*
Caucasian	-	0.98 (0.46, 2.12)	0.84 (0.37, 1.91)
Comorbidity ¹	-	-	0.88 (0.47, 1.65)
Disability ²	-	-	1.71 (0.97, 3.00)
Medication use ³	-	-	0.91 (0.85, 0.98)*
High school education or higher	-	-	3.63 (1.29, 10.24)*
Marital status			
Single	-	-	Reference
Married/cohabitating	-	-	1.71 (0.86, 3.39)
Separated/divorced	-	-	1.13 (0.49, 2.61)
Widowed	-	-	1.10 (0.43, 2.81)

¹Comorbidity = 4 or more of the following conditions: peripheral vascular disease, rheumatoid arthritis, cancer, hypertension, COPD, diabetes, congestive heart failure, angina, and myocardial infarction.

²Disability = Needing assistance on 3 or more of the following activities: feeding, dressing, ambulation, grooming, using a toilet, and bathing.

³Medication use reflects the association of a 1 medication increase and number of falls.

**P <0.01.

*P <0.05.

function (frailty: 3.02, 95% CI: 1.35-6.78) did not change the inferences on the association of frailty and number of falls.

Discussion

In this single-center study, 28.3% of prevalent hemodialysis patients of all ages had a fall. Frailty was a strong independent predictor of falls in this population. Frail patients undergoing chronic hemodialysis experienced 3-times as many falls as their non-frail counterparts. Frailty was a strong independent predictor of falls in patients undergoing hemodialysis, regardless of age.

Elderly hemodialysis patients have been found to be at increased risk of falls [22], and thus, most research has focused on falls in this vulnerable patient population [1,3,6]. Our incidence estimates of falls in patients undergoing hemodialysis are similar to previous studies [1], but we also present data that these estimates are similar even in younger hemodialysis patients. In addition to previously reported predictors of falls (age, comorbidity, and medication use) we have introduced frailty as a novel predictor of falls in both younger and older patients undergoing chronic hemodialysis. In fact, in our study frailty was one of the strongest predictors of falls in our adjusted model. This study not only confirms the association between frailty and falls observed in elderly adults [8,9,23] but extends the findings to patients undergoing hemodialysis of all ages. In older adults, frailty has been hypothesized to be a unique disorder that is independent of chronic conditions and disability. Our work confirms that frailty is an independent phenotype and predictive of falls, regardless of comorbidities and disability for patients undergoing hemodialysis.

Frailty, as defined by Fried et al., is a physiologic precursor and an etiologic factor in the onset of disability [8]. Frailty is thought to lead to disability and subsequent falls due to weakness and low endurance. Although frailty is a distinct phenotype from disability, frailty begins affecting mobility before clinically important outcomes like falls. Thus, the onset of frailty may be the optimal time to initiate interventions to prevent disabilities in mobility and prevent falls.

Strengths of this study were the direct measurement of a validated, objective frailty instrument, and granular ascertainment of comorbidities using medical records abstraction. Additionally, frailty was ascertained prior to the report of falls, so its predictive value could be assessed. The main limitation is the single-center study design, so direct inferences must be interpreted in the context of the demographics of our study population. While this study was conducted in a hospital-based outpatient dialysis unit in Baltimore, the findings are likely generalizable to patients undergoing hemodialysis in a community setting across the US. Additionally, the sample size of the cohort was small, which will not bias the observed associations, but did lead to low power to detect subgroup differences.

Although follow-up was limited, we were interested in ascertaining the number of falls within 6 months from the measurement of frailty, which does not require longer follow-up. We had limited ability to adjust for a number of factors that may be associated with falls (regardless of the association with frailty); however, the strength of the association between frailty and falls was similar when adjusted for anemia, arrhythmia, and cognitive function. Finally, survivor bias limits inferences from our finding that frailty was not associated with time on hemodialysis.

Conclusions

Frailty was identified as a novel risk factor for falls in adults of all ages undergoing hemodialysis. Additionally, the results suggest that among older and younger patients undergoing hemodialysis, frailty increases the short-term risk of experiencing a fall. Our results may aid in identifying frail hemodialysis patients who could be targeted for multidimensional fall prevention strategies.

Abbreviations

ESRD: End stage renal disease; COPD: Chronic obstructive pulmonary disease.

Competing interests

The authors declare that they have no competing interest.

Authors' contributions

All authors 1) have made substantial contributions to conception and design, or acquisition of data, or analysis and interpretation of data; 2) have been involved in drafting the manuscript or revising it critically for important intellectual content; and 3) have given final approval of the version to be published.

Acknowledgements

We thank the study participants, as well as the staff at the Renal Dialysis Program at Good Samaritan, including Colleen Reft, R.N. for their dedication to this study.

This study was supported by a Clinical Scientist Development Award from the Doris Duke Charitable Foundation and the National Institute on Aging (R01-AG042504) (Segev, PI). Mara McAdams-DeMarco was supported by the American Society of Nephrology Carl W. Gottschalk Research Scholar Grant and Johns Hopkins University Claude D. Pepper Older Americans Independence Center, National Institute on Aging (P30-AG021334). Megan Salter was supported by T32AG000247 from the National Institute on Aging.

Author details

¹Department of Surgery, Johns Hopkins University School of Medicine, Baltimore, MD, USA. ²Department of Epidemiology, Johns Hopkins School of Public Health, Baltimore, MD, USA. ³Johns Hopkins School of Medicine, Baltimore, MD, USA. ⁴Nephrology Center of Maryland, Baltimore, MD, USA. ⁵Department of Medicine, Division of Nephrology, Johns Hopkins University School of Medicine, Baltimore, MD, USA. ⁶Dialysis Program, Good Samaritan Hospital, Baltimore, MD, USA. ⁷Division of Geriatric Medicine and Gerontology, Johns Hopkins University School of Medicine, Baltimore, MD, USA. ⁸Clinical Research, Transplant Surgery, Johns Hopkins Medical Institutions, 720 Rutland Ave, Ross 771B, Baltimore, MD 21205, USA.

Received: 15 May 2013 Accepted: 4 October 2013

Published: 16 October 2013

References

1. Cook WL, Jassal SV: Prevalence of falls among seniors maintained on hemodialysis. *Int Urol Nephrol* 2005, **37**(3):649–652.
2. Desmet C, Beguin C, Swine C, Jadoul M: Falls in hemodialysis patients: prospective study of incidence, risk factors, and complications. *Am J Kidney Dis* 2005, **45**(1):148–153.

3. Cook WL, Tomlinson G, Donaldson M, Markowitz SN, Naglie G, Sobolev B, Jassal SV: **Falls and fall-related injuries in older dialysis patients.** *Clin J Am Soc Nephrol* 2006, **1**(6):1197–1204.
4. Leinau L, Perazella MA: **Hip fractures in end-stage renal disease patients: incidence, risk factors, and prevention.** *Semin Dial* 2006, **19**(1):75–79.
5. Mittalhenkle A, Gillen DL, Stehman-Breen CO: **Increased risk of mortality associated with hip fracture in the dialysis population.** *Am J Kidney Dis* 2004, **44**(4):672–679.
6. Li M, Tomlinson G, Naglie G, Cook WL, Jassal SV: **Geriatric comorbidities, such as falls, confer an independent mortality risk to elderly dialysis patients.** *Nephrol Dial Transplant* 2008, **23**(4):1396–1400.
7. Kohlmeier M, Saupe J, Schaefer K, Asmus G: **Bone fracture history and prospective bone fracture risk of hemodialysis patients are related to apolipoprotein E genotype.** *Calcif Tissue Int* 1998, **62**(3):278–281.
8. Fried LP, Tangen CM, Walston J, Newman AB, Hirsch C, Gottdiener J, Seeman T, Tracy R, Kop WJ, Burke G, et al: **Frailty in older adults: evidence for a phenotype.** *J Gerontol A Biol Sci Med Sci* 2001, **56**(3):M146–M156.
9. Ensrud KE, Ewing SK, Taylor BC, Fink HA, Stone KL, Cauley JA, Tracy JK, Hochberg MC, Rodondi N, Cawthon PM: **Frailty and risk of falls, fracture, and mortality in older women: the study of osteoporotic fractures.** *J Gerontol A Biol Sci Med Sci* 2007, **62**(7):744–751.
10. McAdams-DeMarco MA, Law A, Salter ML, Boyarsky B, Gimenez L, Jaar BG, Walston JD, Segev DL: **Frailty as a novel predictor of mortality and hospitalization in individuals of all ages undergoing hemodialysis.** *J Am Geriatr Soc* 2013, **61**(6):896–901.
11. Bandeen-Roche K, Xue QL, Ferrucci L, Walston J, Guralnik JM, Chaves P, Zeger SL, Fried LP: **Phenotype of frailty: characterization in the women's health and aging studies.** *J Gerontol A Biol Sci Med Sci* 2006, **61**(3):262–266.
12. Barzilay JI, Blaum C, Moore T, Xue QL, Hirsch CH, Walston JD, Fried LP: **Insulin resistance and inflammation as precursors of frailty: the Cardiovascular Health Study.** *Arch Intern Med* 2007, **167**(7):635–641.
13. Cappola AR, Xue QL, Fried LP: **Multiple hormonal deficiencies in anabolic hormones are found in frail older women: the Women's Health and Aging studies.** *J Gerontol A Biol Sci Med Sci* 2009, **64**(2):243–248.
14. Leng SX, Hung W, Cappola AR, Yu Q, Xue QL, Fried LP: **White blood cell counts, insulin-like growth factor-1 levels, and frailty in community-dwelling older women.** *J Gerontol A Biol Sci Med Sci* 2009, **64**(4):499–502.
15. Leng SX, Xue QL, Tian J, Walston JD, Fried LP: **Inflammation and frailty in older women.** *J Am Geriatr Soc* 2007, **55**(6):864–871.
16. Newman AB, Gottdiener JS, McBurnie MA, Hirsch CH, Kop WJ, Tracy R, Walston JD, Fried LP: **Associations of subclinical cardiovascular disease with frailty.** *J Gerontol A Biol Sci Med Sci* 2001, **56**(3):M158–M166.
17. Walston J, McBurnie MA, Newman A, Tracy RP, Kop WJ, Hirsch CH, Gottdiener J, Fried LP: **Frailty and activation of the inflammation and coagulation systems with and without clinical comorbidities: results from the Cardiovascular Health Study.** *Arch Intern Med* 2002, **162**(20):2333–2341.
18. Xue QL, Bandeen-Roche K, Varadhan R, Zhou J, Fried LP: **Initial manifestations of frailty criteria and the development of frailty phenotype in the Women's Health and Aging Study II.** *J Gerontol A Biol Sci Med Sci* 2008, **63**(9):984–990.
19. Chang SS, Weiss CO, Xue QL, Fried LP: **Association between inflammatory-related disease burden and frailty: results from the Women's Health and Aging Studies (WHAS) I and II.** *Arch Gerontol Geriatr* 2012, **54**(1):9–15.
20. Chang SS, Weiss CO, Xue QL, Fried LP: **Patterns of comorbid inflammatory diseases in frail older women: the Women's Health and Aging Studies I and II.** *J Gerontol A Biol Sci Med Sci* 2010, **65**(4):407–413.
21. Katz S, Akpom CA: **A measure of primary sociobiological functions.** *Int J Health Serv: Plann, Adm, Eval* 1976, **6**(3):493–508.
22. Roberts R, Jeffrey C, Carlisle G, Brierley E: **Prospective investigation of the incidence of falls, dizziness and syncope in haemodialysis patients.** *Int Urol Nephrol* 2007, **39**(1):275–279.
23. Shim EY, Ma SH, Hong SH, Lee YS, Paik WY, Seo DS, Yoo EY, Kim MY, Yoon JL: **Correlation between frailty level and adverse health-related outcomes of community-dwelling elderly, One year retrospective study.** *Korean J Fam Med* 2011, **32**(4):249–256.

doi:10.1186/1471-2369-14-224

Cite this article as: McAdams-DeMarco et al.: Frailty and falls among adult patients undergoing chronic hemodialysis: a prospective cohort study. *BMC Nephrology* 2013 **14**:224.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

