



US007531692B2

(12) **United States Patent**
Raveglia et al.

(10) **Patent No.:** **US 7,531,692 B2**
(45) **Date of Patent:** **May 12, 2009**

(54) **DERIVATIVES OF
1-PHENYLALKANECARBOXYLIC ACIDS
FOR THE TREATMENT OF
NEURODEGENERATIVE DISEASES**

(58) **Field of Classification Search** 564/155;
514/616
See application file for complete search history.

(75) Inventors: **Luca Raveglia**, Parma (IT); **Ilaria Peretto**, Parma (IT); **Stefano Radaelli**, Parma (IT); **Bruno Pietro Imbimbo**, Parma (IT); **Andrea Rizzi**, Parma (IT); **Gino Villetti**, Parma (IT)

(56) **References Cited**

U.S. PATENT DOCUMENTS

6,114,391	A	9/2000	Chiesi et al.
6,117,901	A	9/2000	Wu et al.
6,191,166	B1	2/2001	Audia et al.
6,476,263	B1	11/2002	Wu et al.
2002/0052322	A1	5/2002	Audia et al.
2007/0060752	A1	3/2007	Raveglia et al.

(73) Assignee: **Chiesi Farmaceutici S.p.A.**, Parma (IT)

FOREIGN PATENT DOCUMENTS

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

WO	99 41224	8/1999
WO	2004 009565	1/2004

(21) Appl. No.: **11/572,974**

OTHER PUBLICATIONS

(22) PCT Filed: **Jul. 26, 2005**

Tomiyaama et al, Oya Yakuri, 1980, 19(6), 895-900.*
Turbanti et al, Farmaco, Edizione Scientifica, 1960, 15, 406-13(Ab-
stract only).*
L. Turbanti, G. F. Di Paco, "Acidi α -Difenililettilacetamino- β -
Arlilacrilici A Prusunta Azione Coleretica E Ipoolesterioemica", Il
Farmaco-ED. Sc., vol. 15, No. 6, pp. 406-413, 1960.
U.S. Appl. No. 11/572,974, filed Jan. 30, 2007, Raveglia et al.

(86) PCT No.: **PCT/IB2005/002189**

§ 371 (c)(1),
(2), (4) Date: **Oct. 12, 2007**

(87) PCT Pub. No.: **WO2006/016219**

PCT Pub. Date: **Feb. 16, 2006**

* cited by examiner

(65) **Prior Publication Data**

US 2008/0096968 A1 Apr. 24, 2008

Primary Examiner—Shailendra Kumar
(74) *Attorney, Agent, or Firm*—Oblon, Spivak, McClelland,
Maier & Neustadt, P.C.

(30) **Foreign Application Priority Data**

Aug. 3, 2004 (EP) 04425604

(57) **ABSTRACT**

(51) **Int. Cl.**
C07C 233/05 (2006.01)
A61K 31/16 (2006.01)

The present invention concerns novel derivatives of 1-phenyl-
lalkanecarboxylic acids, pharmaceutical compositions
thereof, a process for their preparation and their use for the
treatment and/or prevention of neurodegenerative diseases
such as Alzheimer's disease.

(52) **U.S. Cl.** **564/155; 514/616**

21 Claims, No Drawings

**DERIVATIVES OF
1-PHENYLALKANECARBOXYLIC ACIDS
FOR THE TREATMENT OF
NEURODEGENERATIVE DISEASES**

CROSS REFERENCES TO RELATED
APPLICATIONS

This application is a 371 of International Patent Application No. PCT/IB05/02189, filed on Jul. 26, 2005, and claims priority to European Patent Application No. 04425604.8, filed on Aug. 3, 2004.

The present invention concerns novel derivatives of 1-phenylalkanecarboxylic acids, pharmaceutical compositions thereof, a process for their preparation and their use for the treatment of neurodegenerative diseases, in particular Alzheimer's disease.

INTRODUCTION

Alzheimer's disease is a neurodegenerative disorder of the Central Nervous System (C.N.S.) characterized, from the anatomical point of view, by atrophy of the cerebral cortex and by a massive loss of cortical neurons and cholinergic projections of the nuclei basalis towards the cortex. From the histopathologic point of view a diffuse presence of extracellular and perivascular neuritic plaques and intracellular neurofibrillary tangles in the cerebral parenchyma of the patients is observed.

Neuritic plaques are mainly composed of aggregates of a peptide with 39-43 amino acid residues known as β -amyloid (β A), and, depending on the numbers of amino acids, $A\beta_{39}$, $A\beta_{40}$, $A\beta_{42}$ and $A\beta_{43}$.

In addition to these histopathologic lesions, there is lack in some neurotransmitters, particularly acetylcholine, serotonin, noradrenalin, dopamine, glutamate and substance P. The pharmacological approaches aimed at increasing acetylcholine cerebral levels, mainly through acetylcholine-esterase inhibitors, attained poor results from the clinical standpoint, or anyhow results which cannot significantly prevent the progress of the disease. For this reason, in recent years the mechanisms of formation of the main pathologic lesions in the brain of the patients have been investigated, namely both neuritic plaques and neurofibrillary tangles, and more effective therapeutical approaches have been sought.

PRIOR ART

Epidemiological studies evidenced that chronic administration of non steroid anti-inflammatory drugs (NSAIDs) significantly decreases the risk of onset of Alzheimer's disease in the population regularly taking these drugs. The mechanism underlying such NSAIDs preventive action has not been fully elucidated yet, but in the earlier hypothesis, it was connected with their inherent anti-inflammatory activity, i.e. with their ability of inhibiting the cyclooxygenase (COX) enzyme.

In WO 99/41224 novel biaryl-acetic acid derivatives with anti-inflammatory activity as cyclooxygenase-2 inhibitors, useful for the treatment of a number of diseases, including Alzheimer's disease, are claimed.

More recently, a novel pharmacological action of some non steroid anti-inflammatory drugs (NSAIDs) has been described: indomethacin, sulindac, ibuprofen and flurbiprofen can selectively reduce the production of the most neurotoxic isoform of β -amyloid peptide in cell cultures, namely the form containing 42 amino acids ($A\beta_{42}$), thus favoring the release of a less harmful isoform, $A\beta_{38}$ (Weggen et al., Nature

2001; 414 (6860): 212-6). However, the inhibition of the production of $A\beta_{42}$, which can be ascribed to the interaction of these drugs with γ -secretase enzyme (a macromolecular/multiprotein complex with aspartyl-protease activity) has been observed in vitro at very high concentrations. Plasma and cerebral levels corresponding to the dosages used in the in vitro experimentation could significantly increase in treated patients the risk of side effects typical of COX inhibitors, such as gastrointestinal bleeding and perforating ulcers.

WO 01/78721 claims a method of preventing, delaying or reversing the progression of Alzheimer's disease by administering an $A\beta_{42}$ lowering agent, under conditions in which levels of $A\beta_{38}$ are increased and levels of $A\beta_{42}$ are left unchanged. Furthermore, methods and materials for identifying and developing $A\beta_{42}$ lowering agents and methods for identifying agents that increase the risk of developing, or hasten progression of Alzheimer's disease, are disclosed. The examples concern indomethacin and flufenamic acid derivatives, but no examples concerning flurbiprofen derivatives are reported.

Jantzen et al, *J Neurosci* 2002; 22: 2246-2254, described a flurbiprofen derivative capable of releasing nitric oxide. The paper generically states that flurbiprofen derivatives are apparently more efficacious than other NSAIDs in clearing β -amyloid deposits, but no mention concerning an $A\beta_{42}$ lowering selective activity is made.

In the co-pending International application No. PCT/EP2004/001596 the applicant claims 1-phenylalkanecarboxylic acids and their functional derivatives such as esters, amides, sulfonamides, and bioisosters as compounds provided with a more selective and more potent inhibitory activity on the peptide $A\beta_{42}$ than that on the cyclooxygenase enzymes.

On the other hand, drugs aimed at the treatment of C.N.S. diseases such as Alzheimer's disease, in order to efficaciously exercise their therapeutic activity, need to cross the blood-brain barrier. The passage and the distribution in the C.N.S. of polar drugs such the carboxylic acids and their derivatives are strongly limited by the presence of said barrier.

Therefore it would be highly advantageous to provide carrier molecules to link said 1-phenylalkanecarboxylic acids in such a way to obtain novel compounds acting as pro-drugs able of crossing more efficaciously the blood-brain barrier and then, releasing in situ the active moiety of the molecule in order to allow its distribution in the brain.

SUMMARY OF THE INVENTION

The present invention concerns novel derivatives of 1-phenylalkanecarboxylic acids, pharmaceutical compositions thereof, a process for their preparation and their use for the prevention or therapeutical treatment of neurodegenerative diseases connected with an increased production of the neurotoxic peptide $A\beta_{42}$, in particular Alzheimer's disease.


In particular the invention concerns derivatives of 1-phenylalkanecarboxylic acids wherein the carboxylic group is linked to a residue allowing the passage of the blood-brain barrier and the distribution of the active moiety in the brain.

In an embodiment of the invention, said residue is represented by the amide of an alpha-amino acid and preferably is glycylamide.

DETAILED DESCRIPTION OF THE INVENTION

The present invention is directed to compounds of general formula (I):

3



wherein:

B is H or the side chain of an alpha-amino acid;

R and R₁ are the same and are a linear or branched C₁-C₄ alkyl;

or they form a 3 to 6 carbon atoms ring with the carbon atom to which they are linked;

R₂ is H, CF₃, OCF₃ or a halogen selected from the group of F, Cl, Br, I, preferably fluorine.

Ar is phenyl substituted with one or more groups R₃ wherein R₃ represents:

halogen as previously defined; CF₃; C₃-C₈ cycloalkyl optionally substituted with one or more C₁-C₄ alkyl and/or oxo groups; CH=CH₂; NO₂; CH₂OH; CN; methylenedioxy; ethylenedioxy;

phenyl optionally substituted with one or more of the following groups: halogen as previously defined; CF₃, OCF₃, OH; linear or branched C₁-C₄ alkyl; a saturated heterocycle with at least 4 carbon atoms and at least 1 heteroatom; C₃-C₈ cycloalkyl optionally substituted with one or more linear or branched C₁-C₄ alkyl groups, CF₃ and/or OH;

OR₄ or NHCOR₄ wherein R₄ is CF₃, linear or branched C₂-C₆ alkenyl or alkynyl; benzyl; phenyl optionally substituted with one or more of the following groups: halogen as previously defined, CF₃, OCF₃, OH, linear or branched C₁-C₄ alkyl; a saturated heterocycle with at least 4 carbon atoms and at least 1 heteroatom; C₃-C₈ cycloalkyl optionally substituted with one or more linear or branched C₁-C₄ alkyl groups, CF₃ and/or OH;

SR₅, SO₂R₅ or COR₅ wherein R₅ is linear or branched C₁-C₆ alkyl;

or Ar is a heterocycle selected from the group consisting of pyrrole, pyrazole, furan, thiophene, indole, isoindole, benzofuran, benzothiophene, imidazole, oxazole, isoxazole, thiazole, benzoimidazole, benzoxazole, benzothiazole, pyrimidine, pyrazine, quinoline, isoquinoline, quinazoline, quinoxaline, dibenzofuran, dibenzothiophene, thianthrene, carbazole, pyridazine, cinnoline, phthalazine, 1,5-naphthiridine, 1,3-dioxole, 1,3-benzodioxole, said heterocycle being optionally substituted with one or more groups R₃ as defined above.

Amides of an alpha-amino acid representative of the invention are preferably selected from the group of glycylamide, alanylamide, serinamide, and valinamide, even more preferably in the levo form.

The preferred amide is glycylamide (H₂NCH₂CONH₂).

A first group of preferred compounds is that in which:

B is H;

R and R₁ form a 3 carbon atoms ring;

R₂ is fluorine;

Ar is phenyl as defined above.

A second group of preferred compounds is that in which:

B is H;

R and R₁ are both methyl;

4

R₂ is fluorine;

Ar is phenyl as defined above.

(I) A third group of preferred compounds is that in which:

B is H;

5 R and R₁ form a 3 carbon atoms ring;

R₂ is fluorine;

Ar is a heterocycle as defined above.

A fourth group of preferred compounds is that in which:

B is H;

R and R₁ are both methyl;

10 R₂ is fluorine;


Ar is a heterocycle as defined above.

The invention also includes the enantiomers, metal and organic salts and other esters pharmaceutically acceptable.

A further object of the present invention is the use of the aforementioned compounds for the therapeutical treatment and/or prevention of neurodegenerative diseases connected with an increased production of the neurotoxic peptide Aβ₄₂, such as Alzheimer's disease.

Still a further object of the invention are solid or liquid pharmaceutical compositions, preferably for the oral use, comprising at least one compound of formula (I) in admixture with pharmaceutically acceptable excipients and/or vehicles, for example those described in Remington's Pharmaceutical Sciences Handbook, XVII Ed., Mack Pub., N.Y., U.S.A.

20 The compounds of general formula (I) can be prepared according to methods known literature by conversion of an acid of formula (II)



(II)

in which R, R₁, R₂, and Ar are as defined above, into the corresponding acyl chloride, followed by reaction with the amide of the suitable alpha-amino acid.

Alternatively, said compounds can be prepared by direct reaction of the acid of formula (II) with the amide of the suitable alpha-amino acid in the presence of coupling agents such as dicyclohexylcarbodiimide (DCC), polymer supported-DCC, or N,N'-carbonyldiimidazole or treating a corresponding ester with the compound formed in situ by reacting trimethylaluminum and the amide of a suitable alpha-amino acid.

The acids of formula (II) can be prepared as described in the co-pending International application No. PCT/EP2004/001596.

EXAMPLES

The following Example illustrates the invention in a more detail.

Preparation of the glycylamide of the 1-[2-fluoro-4'-[[4-(trifluoromethyl)-cyclohexyl]oxy]-1,1'-biphenyl-4-yl]-cyclopropanecarboxylic acid

60 Preparation of the 1-(2-fluoro-4'-hydroxy-biphenyl-4-yl)-cyclopropanecarboxylic acid

To a solution of K₂CO₃ (447 mg, 3.2 mmoles) in dioxane (30 ml) and water (3 ml), 250 mg of 1-(4-bromo-3-fluorophenyl)-cyclopropanecarboxylic acid (1.0 mmole) and 265 mg of hydroxyphenylboronic acid (1.9 mmoles) are added under stirring. Under nitrogen atmosphere, 39 mg of [1,1'-bis

(diphenylphosphino)ferrocene]dichloropalladium (PdCl₂-DPPF) are added. The resulting mixture is heated at 65° C. for one hour, then the reaction is quenched by adding a 10% w/v HCl solution.

Extractions with ethyl acetate, washings with water and brine and evaporation afford the crude product, which is crystallized from ethyl acetate/hexane, to give the 1-(2-fluoro-4'-hydroxy-biphenyl-4-yl)-cyclopropanecarboxylic acid. Yield: 85%.

Preparation of the 1-[2-fluoro-4'-(4-trifluoromethyl-cyclohexyloxy)-biphenyl-4-yl]-cyclopropanecarboxylic acid

Diethyl azodicarboxylate (370 μL, 2.4 mmoles) is added to a solution of 200 mg of 1-(2-fluoro-4'-hydroxy-biphenyl-4-yl)-cyclopropanecarboxylic acid (0.8 mmoles), triphenyl phosphine (623 mg, 2.4 mmoles) and 4-trifluoromethyl cyclohexanol (400 mg, 2.4 mmoles) in dry THF (3 mL) at 0° C. under nitrogen atmosphere. The resulting clear orange solution is left under stirring overnight, then quenched with water and extracted with ethyl acetate. The organic phase is washed with 10% w/v HCl solution, dried on Na₂SO₄ and evaporated. Purification by chromatography on silica gel by eluting with methylene chloride:methanol 98:2 v/v furnishes 1-[2-fluoro-4'-(4-trifluoromethyl-cyclohexyloxy)-biphenyl-4-yl]-cyclopropanecarboxylic acid as a white solid.

Yield: 26%.

HPLC purity (254 nm): 98%

Preparation of the glycineamide of the 1-[2-fluoro-4'-(4-trifluoromethyl-cyclohexyloxy)-biphenyl-4-yl]-cyclopropanecarboxylic acid

Oxalyl chloride (530 μL, 6.0 mmoles) is added dropwise to a solution of 1-[2-fluoro-4'-(4-trifluoromethyl-cyclohexyloxy)-biphenyl-4-yl]-cyclopropane carboxylic acid (480 mg, 1.0 mmoles) in 5 ml of methylene chloride, under nitrogen. The resulting solution is stirred at room temperature for 2 hours, then evaporated to dryness and redissolved in methylene chloride. A suspension of glycineamide hydrochloride (133 mg, 1.3 mmoles) and triethylamine (3 ml) in 5 ml of methylene chloride is added to the solution and the resulting mixture is stirred for 2 hours at room temperature. After addition of water, the organic phase is washed with a 2N K₂CO₃ solution and brine, dried on Na₂SO₄ and evaporated. The solid obtained is purified by chromatography. After recrystallization from ethyl acetate/petroleum ether a white solid is obtained.

HPLC purity (254 nm): 98%.


MS (ESI⁺): 479.2 (MH⁺).

¹H NMR(CDCl₃): 7.47 (dd, 2H); 7.43 (dd, 1H); 7.25 (dd, 1H); 7.19 (dd, 1H); 6.97 (dd, 2H); 6.05 (t, 1H); 5.94 (s, 1H); 5.34 (s, 1H); 4.23 (m, 1H); 3.88 (d, 2H); 2.35-2.25 (m, 2H); 2.15-2.01 (m, 3H); 1.64 (m, 2H); 1.48 (dd, 4H); 1.14 (m, 2H).

Legend: dd=doublet of doublets; t=triplet; d=doublet; s=singlet; m=multiplet.

The invention claimed is:

1. A compound of formula (I):



wherein:

B is H or the side chain of an alpha-amino acid;

R and R₁ are the same and are linear or branched C₁-C₄ alkyl;

or together with the carbon atom to which they are linked form a ring containing 3 to 6 carbon atoms;

R₂ is H, CF₃, OCF₃ or a halogen selected from the group consisting of F, Cl, Br, and I; and

Ar is phenyl substituted with one or more groups R₃, wherein R₃ represents:

halogen selected from the group consisting of F, Cl, Br, and I; CF₃; C₃-C₈ cycloalkyl optionally substituted with one or more C₁-C₄ alkyl and/or oxo groups; CH=CH₂; NO₂; CH₂OH; CN; methylenedioxy; ethylenedioxy; phenyl optionally substituted with one or more of the following groups: halogen selected from the group consisting of F, Cl, Br, and I; CF₃; OCF₃; OH; linear or branched C₁-C₄ alkyl; a saturated heterocycle with at least 4 carbon atoms and at least 1 heteroatom; and C₃-C₈ cycloalkyl optionally substituted with one or more linear or branched C₁-C₄ alkyl groups, CF₃ and/or OH; OR₄ or NHCOR₄ wherein R₄ is CF₃, linear or branched C₂-C₆ alkenyl or alkynyl; benzyl; phenyl optionally substituted with one or more of the following groups: halogen as previously defined, CF₃, OCF₃, OH, linear or branched C₁-C₄ alkyl; a saturated heterocycle with at least 4 carbon atoms and at least 1 heteroatom; C₃-C₈ cycloalkyl optionally substituted with one or more linear or branched C₁-C₄ alkyl groups, CF₃ and/or OH; and SR₅, SO₂R₅ or COR₅ wherein R₅ is linear or branched C₁-C₆ alkyl; or

Ar is a heterocycle selected from the group consisting of pyrrole, pyrazole, furan, thiophene, indole, isoindole, benzofuran, benzothiophene, imidazole, oxazole, isoxazole, thiazole, benzoimidazole, benzoxazole, benzothiazole, pyrimidine, pyrazine, quinoline, isoquinoline, quinazoline, quinoxaline, dibenzofuran, dibenzothiophene, thianthrene, carbazole, pyridazine, cinnoline, phthalazine, 1,5-naphthiridine, 1,3-dioxole, and 1,3-benzodioxole, said heterocycle being optionally substituted with one or more groups R₃ as defined above;

or a pharmaceutically acceptable salt or ester thereof.

2. The compound or pharmaceutically acceptable salt or ester thereof as claimed in claim 1, wherein B is H.

3. The compound or pharmaceutically acceptable salt or ester thereof as claimed in claim 1, wherein B is H; R and R₁ together with the carbon atom to which they are linked form a ring containing 3 carbon atoms; R₂ is fluorine; and Ar is substituted phenyl.

4. The compound or pharmaceutically acceptable salt or ester thereof as claimed in claim 1, wherein B is H; R and R₁ are both CH₃; R₂ is fluorine; and Ar is substituted phenyl.

5. The compound or pharmaceutically acceptable salt or ester thereof as claimed in claim 1, wherein B is H; R and R₁ together with the carbon atom to which they are linked form a ring containing 3 carbon atoms; R₂ is fluorine; and Ar is a substituted or unsubstituted heterocycle.

6. The compound or pharmaceutically acceptable salt or ester thereof as claimed in claim 1, wherein B is H; R and R₁ are both CH₃; R₂ is fluorine; and Ar is a substituted or unsubstituted heterocycle.

7. A compound or pharmaceutically acceptable salt thereof as claimed in claim 1.

7

8. A pharmaceutical composition, comprising a compound of formula (I) or pharmaceutically acceptable salt or ester thereof as claimed in claim 1 in admixture with a pharmaceutically acceptable carrier.

9. The pharmaceutical composition as claimed in claim 8, which is in a form suitable for oral administration.

10. The compound or pharmaceutically acceptable salt or ester thereof according to claim 1, wherein R_2 is F.

11. A pharmaceutical composition, comprising a compound of formula (I) or pharmaceutically acceptable salt or ester thereof as claimed in claim 2 in admixture with a pharmaceutically acceptable carrier.

12. A pharmaceutical composition, comprising a compound of formula (I) or pharmaceutically acceptable salt or ester thereof as claimed in claim 3 in admixture with a pharmaceutically acceptable carrier.

13. A pharmaceutical composition, comprising a compound of formula (I) or pharmaceutically acceptable salt or ester thereof as claimed in claim 4 in admixture with a pharmaceutically acceptable carrier.

14. A pharmaceutical composition, comprising a compound of formula (I) or pharmaceutically acceptable salt or ester thereof as claimed in claim 5 in admixture with a pharmaceutically acceptable carrier.

15. A pharmaceutical composition, comprising a compound of formula (I) or pharmaceutically acceptable salt or

8

ester thereof as claimed in claim 6 in admixture with a pharmaceutically acceptable carrier.

16. A pharmaceutical composition, comprising a compound of formula (I) or pharmaceutically acceptable salt thereof as claimed in claim 1 in admixture with a pharmaceutically acceptable carrier.

17. A pharmaceutical composition, comprising a compound of formula (I) or pharmaceutically acceptable salt thereof as claimed in claim 2 in admixture with a pharmaceutically acceptable carrier.

18. A pharmaceutical composition, comprising a compound of formula (I) or pharmaceutically acceptable salt thereof as claimed in claim 3 in admixture with a pharmaceutically acceptable carrier.

19. A pharmaceutical composition, comprising a compound of formula (I) or pharmaceutically acceptable salt thereof as claimed in claim 4 in admixture with a pharmaceutically acceptable carrier.

20. A pharmaceutical composition, comprising a compound of formula (I) or pharmaceutically acceptable salt thereof as claimed in claim 5 in admixture with a pharmaceutically acceptable carrier.

21. A pharmaceutical composition, comprising a compound of formula (I) or pharmaceutically acceptable salt thereof as claimed in claim 6 in admixture with a pharmaceutically acceptable carrier.

* * * * *