

- arrhythmic outcome and sudden cardiac death in nonischemic dilated cardiomyopathy. *Heart Rhythm* 2014. doi:10.1016/j.hrthm.2014.01.014.
22. Kjaergaard J, Akkan D, Iversen KK, Kjoller E, Kober L, Torp-Pedersen C et al. Prognostic importance of pulmonary hypertension in patients with heart failure. *Am J Cardiol* 2007;**99**:1146–50.
23. Damy T, Goode KM, Kallvikbacka-Bennett A, Lewinter C, Hobkirk J, Nikitin NP et al. Determinants and prognostic value of pulmonary arterial pressure in patients with chronic heart failure. *Eur Heart J* 2010;**31**:2280–90.
24. Ghio S, Gavazzi A, Campana C, Inserra C, Klersy C, Sebastiani R et al. Independent and additive prognostic value of right ventricular systolic function and pulmonary artery pressure in patients with chronic heart failure. *J Am Coll Cardiol* 2001;**37**:183–88.
25. Bourantas CV, Loh HP, Bragadeesh T, Rigby AS, Lukaschuk E, Garg S et al. Relationship between right ventricular volumes measured by cardiac magnetic resonance imaging and prognosis in patients with chronic heart failure. *Eur Heart J* 2011;**13**:52–60.

IMAGE FOCUS

doi:10.1093/ehjci/jeu149

Online publish-ahead-of-print 7 August 2014

Isolated left ventricular apical hypoplasia

Carlos Galvão Braga^{1*}, Paulo Silva^{2,3}, Sónia Magalhães¹, Nuno Bettencourt^{4,5}, and Raquel Themudo^{4,6}

¹Department of Cardiology, Hospital de Braga, Braga, Sete Fontes – São Victor, 4710-243, Portugal; ²Clínica Jácome, Famalicão, Portugal; ³Hospital de Dia de Famalicão, Famalicão, Portugal; ⁴Hospital Privado de Alfena, Alfena, Portugal; ⁵Department of Cardiology, Centro Hospitalar de Vila Nova de Gaia-Espinho, VN Gaia, Portugal; and ⁶Department of Radiology, Karolinska University Hospital, Stockholm, Sweden

* Corresponding author. Tel: +351 915301245. E-mail: carlos.galvaobraga@gmail.com

A 66-year-old woman with hypertension, dyslipidaemia and history of stable angina was referred for cardiological assessment after a transthoracic echocardiogram raising the suspicion of right ventricular arrhythmogenic dysplasia. The patient reported atypical chest pain. The 12-lead electrocardiogram showed sinus rhythm with a complete left bundle branch block. The echocardiogram was repeated and demonstrated mild left ventricular (LV) systolic dysfunction with apical hypokinesia, abnormal interventricular septal motion and elongated right ventricle (RV). A myocardial perfusion scan showed a small fixed apical attenuation defect and absence of reversible ischaemia. To clarify diagnosis, cardiac MRI was performed and demonstrated the four phenotypical features of isolated LV apical hypoplasia: a bizarre LV with spherical configuration, truncated apex, interventricular septum bulging to the right and impaired systolic function (ejection fraction of 48%; Panel A and Supplementary data online *Video S1*); elongated RV wrapping around the deficient LV apex (*, Panel A; Supplementary data online *Video S1*); anomalous origin of anterior LV papillary muscle from the LV apex (*, Panel B); and replacement of the LV apex with fatty material [arrows, Panels B and C, which represent T1-weighted black-blood images without and with fat suppression (STIR) in four-chamber view, respectively], in this case with extension to the mesocardium of anterior, lateral and inferior midwall segments and continuity with the epicardial fat (arrow, Panel D, which represents cine-balanced steady-state-free precession in three-chamber view).

Isolated LV apical hypoplasia is a rare entity and MRI provides an excellent method to completely characterize the disease and evaluate the extent of myocardial fat replacement.

Supplementary data are available at *European Heart Journal – Cardiovascular Imaging* online.

