


Conformationally Restricted Criegee Intermediates: Evidence for Formation and Stereoelectronically Controlled Fragmentation

Sosale Chandrasekhar* and Chandra Deo Roy

Department of Organic Chemistry, Indian Institute of Science, Bangalore 560 012, India

The Baeyer–Villiger reaction of 2-(2-oxocyclohexyl)acetic acid occurs *via* a bicyclic Criegee intermediate, which fragments with stereoelectronic control, as evidenced by product analysis; the reaction of the but-2-yl ester and of 2-(2-oxocyclopentyl)acetic acid also show evidence of such stereoelectronic control, but less convincingly.


Stereoelectronic theory requires that the Criegee intermediates of the Baeyer–Villiger reaction fragment with the cleaving C–C and O–O bonds mutually antiperiplanar (**1**).^{1a} We have briefly



reported evidence for such stereoelectronic control,² and herein present a comprehensive study. This study is challenging as Criegee intermediates are not normally isolable, so that unmasking stereoelectronic effects needs a novel strategy. The one devised by us is a study of the intramolecular Baeyer–Villiger reaction, which produces a conformationally restricted Criegee intermediate whose mode of fragmentation may be traced by the nature of the final products. It is noteworthy that stereoelectronic theory has been vigorously debated,^{1b} and that the present results are amongst the least ambiguous in support of it.

Results and Discussion



With the aim of producing Criegee intermediate **3** by cyclisation, ketoacid **2a** was oxidised with peroxyacetic acid in




- 2a** R = H
b R = OH
c R = Bu^s

acetic acid, in order to produce keto-peroxy acid **2b**. The reaction produced 2-(4-acetoxybutyl)butanedioic acid (**5a**) in 62% yield (see Scheme 1); this could be hydrolysed to **5b**, or esterified to **5c** with diazomethane, both in good yield. Importantly, there was no trace of isomer **6a** in the oxidation; NMR of the product mixture showed no absorption at δ_{H} 4.3–5.5, as expected for AcOCH in **6a** and base treatment of the mixture did not form any octenedioic acid **7**. Apparently, a Baeyer–Villiger reaction takes place, but with the seven-membered ring lactone product (**4a**) cleaving further. Clearly, **5a** can form by stereoelectronically controlled fragmentation of **3**, by migration of a secondary centre (path A); the absence of **6a** suggests that migration of a tertiary centre (path B) is relatively disfavoured, *i.e.* the classical migratory preferences are overruled.

Evidence for the cyclisation of **2b** to **3** was obtained by treating but-2-yl ester **2c** with peroxyacetic acid, when both




4	R	R'
a	CH ₂ CO ₂ H	H
b	H	CH ₂ CO ₂ H
c	CH ₂ CO ₂ Bu ^s	H
d	H	CH ₂ CO ₂ Bu ^s



5	R	R'	R''
a	Ac	H	H
b	H	H	H
c	Ac	Me	Me
d	Ac	H	Bu ^s


possible Baeyer–Villiger oxidation products, **5d** and **6b** (approx. 2:1 ratio), were formed—presumably *via* lactones **4c** and **4d** respectively (see Scheme 1). Clearly, only intermolecular attack of peroxy acid on ketone is possible, so evidence of stereoelectronic control is lost. However, the relatively higher yield of **5d** could well be due to intramolecular assistance for O–O cleavage by the ester carbonyl as in **8**: the stereoelectronic situation would be similar to that in the cleavage of **3**. Note that the formation of Criegee intermediates is thermodynamically controlled,³ but intermediacy of **3** is not unequivocal *vis-à-vis* *cis*-fused epimer **9a, b** (different enantiomers are shown in **9a** and **9b** for convenience of representation): C-9-substituted decalins rather prefer *cis* ring fusion,⁴ and **9a** has the requisite stereoelectronic features to give the observed products.

Interesting results were obtained when the above oxidation strategy was applied to 2-(2-oxocyclopentyl)acetic acid (**10a**) and its but-2-yl ester **10b**. Ketoacid **10a** produced six-membered lactones **11a** (10%) and **11b** (22%), and the cleavage product of **11a, 12** (32%) (see Scheme 1). Thus, regioselectivity has been partially lost, suggesting the incursion of an intermolecular



pathway. Ester **10b** produced lactones **11c** (5%) and **11d** (35%), and the cleavage product of **11d**, **13** (20%), which constitutes nearly complete migratory aptitude control. All these results may be explained by invoking strain in the corresponding analogues of **3**, **8** and **9** (*i.e.* with the carbocyclic ring five-membered), without prejudice to stereoelectronic theory.

These results constitute evidence for stereoelectronic control in the fragmentation of Criegee intermediates. The study of stereoelectronic effects is almost always plagued by a fundamental problem, namely the impossibility of designing models



Scheme 1

which are conformationally rigid enough for stereoelectronic relationships to be well defined, and yet flexible enough to react. The above study is unique in being relatively free of this problem, apparently because of the structure of the cyclic peroxide **3**.

Experimental

IR spectra were recorded with a Perkin-Elmer 781 or Hitachi 270-50 spectrometer as thin films (liquids) or nujol mulls (solids). NMR spectra were recorded with a JEOL FX-90Q or a Varian T-60 spectrometer, at 90 MHz in CDCl₃ solution unless otherwise stated. EI mass spectra were recorded with a JEOL MS-DX 303 at 70 eV. Extracts of reaction mixtures were worked up by washing with H₂O, drying (MgSO₄ or Na₂SO₄) and evaporating the solvent *in vacuo*. Chromatography was carried out on silica gel columns. M.p.s are uncorrected. Ether refers to diethyl ether.

The methyl and ethyl esters of ketoacids **2a** and **10a** were easily hydrolysed under the reaction conditions, presumably by keto participation, but the but-2-yl esters were stable (but unstable to distillation, so did not provide satisfactory elemental analyses). Product mixtures were usually sensitive to chromatography and distillation, and were further derivatised; the yields shown are those of the isolated derivatives. Thus, mixture (**5d** + **6b**) gave, with alkali, hydroxydiacid **5b** (50%) and octenedioic acid **7** (30%), separated by crystallisation. This bears out the relative yields given in the Results and Discussion section, which are NMR estimates. Acid **7** was hydrogenated to octanedioic (suberic) acid. Neither elemental analysis nor high-resolution mass spectral data were obtained for lactone **11c**.

The products of the reaction of ketoacid **10a** were esterified with CH₂N₂ before resolution. In the reaction of ketoester **10b**, partial hydrolysis of the ester occurred, so the product mixture was heated with Bu^sOH/H⁺ to re-esterify the CO₂H groups; the above hydrolysis presumably occurred after the Baeyer-Villiger reaction, as the reaction of ketoacid **10a** showed a different 'product spread'.

Preparation of Keto-esters 2c and 10b.—The keto-acid⁵ (16 mmol) and conc. H₂SO₄ (0.07 cm³) were heated in butan-2-ol (10 cm³) for 7 h. After concentration, the mixture was extracted with CHCl₃, the extracts washed with aqueous NaHCO₃ and H₂O, and worked up. Chromatography (eluent: 0-1% EtOAc in hexane) gave the pure keto-esters (13.5 mmol, 85%) as liquids unstable to distillation.

Baeyer-Villiger Reactions: General Procedure.—A stirred solution of H₂O₂ (30%, 1.4 cm³; 23 mmol), acetic acid (4.2 cm³) and conc. H₂SO₄ (0.02 cm³) was treated with the substrate (6 mmol) at 25 °C for 15 h and 70 °C for 3 h. After concentration, the mixture was extracted with CH₂Cl₂ (100 cm³) and the extracts worked up.

The crude product of the reaction of keto-acid **2a** was

chromatographed (eluent: 5% MeOH in CHCl₃) to isolate **5a** as a viscous liquid (62%); attempted distillation seemed to form the anhydride, as shown by IR, so **5a** was hydrolysed to **5b** for characterisation, or esterified with CH₂N₂ (see below). In the case of keto-ester **2c**, the product mixture was also hydrolysed. In the case of keto-acid **10a**, the product mixture was esterified with CH₂N₂ in ether (as for **5a**), before chromatographic separation (eluent: 33% EtOAc in hexane) into the methyl esters of **11a** (10%), **11b** (22%) and **12** (32%). In the case of keto-ester **10b**, the product mixture was re-esterified with butan-2-ol and catalytic conc. H₂SO₄ (reflux 5 h); after work-up, the mixture was chromatographed (eluent: 33% EtOAc in hexane) to isolate **11c** (5%), **11d** (35%) and **13** (20%, as the dibut-2-yl ester).

Hydrolysis of 5a and (5d + 6b).—A solution of **5a** (0.334 g, 1.43 mmol) in 10% KOH in MeOH–H₂O (1:1, 8 cm³) was refluxed for 2 h. After concentration, followed by acidification, the mixture was extracted into EtOAc and the extracts worked up to obtain **5b** (0.15 g, 0.79 mmol, 50%), m.p. 110–112 °C (EtOAc–hexane).

Similarly, **(5d + 6b)** (2.1 g) in 10% KOH in MeOH–H₂O (1:1, 46 cm³), at reflux for 5 h, and work-up as above, gave a mixture (1.44 g) separated by fractional crystallisation (EtOAc) into **5b** (0.74 g, 3.9 mmol, 50%) and oct-2-enedioic acid (**7**) (0.40 g, 2.3 mmol, 30%), m.p. 155–160 °C (from H₂O). Compound **7** (0.05 g, 0.29 mmol) was catalytically (5% Pd/C) hydrogenated in EtOH–H₂O (1:1) to octanedioic (suberic) acid (0.045 g, 0.25 mmol, 86%), spectrally identical with an authentic sample, m.p. 139–140 °C (acetone), mixed m.p. 140–141 °C (lit.,⁶ 139–141 °C).

Esterification of 5a.—The diacid (0.232 g, 1.0 mmol) in ether (3 cm³) was slowly treated with diazomethane in ether⁷ at 0–5 °C, till a yellow colour persisted. After evaporation of volatiles, the residue was chromatographed (eluent: 4% EtOAc in hexane) to obtain pure triester **5c** (0.222 g, 0.9 mmol, 90%) as a liquid, unstable to distillation.

Characterising Data.—*Butyl ester 2c.* $\nu_{\max}/\text{cm}^{-1}$ 1730 (ester CO) and 1710 (ketone CO); δ_{H} (60 MHz; CCl₄) 4.70 (1 H, sextet, *J* 7, OCH) and 3.00–0.66 (19 H, m, CH); *m/z* 212 (M⁺), 156 (M⁺ – 56) and 139 (M⁺ – 73).

Acetate diacid 5a. $\nu_{\max}/\text{cm}^{-1}$ 3300–2300 (acid OH) and 1725 (CO); δ_{H} 11.20 (2 H, br s, CO₂H, D₂O exchangeable), 4.15 (2 H, m, AcOCH₂), 3.10–2.20 (3 H, m, COCH, COCH₂), 2.08 (3 H, s, CH₃) and 2.00–1.00 (6 H, m, CH₂); *m/z* 172 (M⁺ – 60), 154 (M⁺ – 78) and 136 (M⁺ – 96).

Alcohol diacid 5b. (Found: C, 50.8; H, 7.5. C₈H₁₄O₅ requires C, 50.5; H, 7.4%) $\nu_{\max}/\text{cm}^{-1}$ 3500–2300 (OH) and 1725 (CO); δ_{H} [(CD₃)₂CO–CDCl₃] 5.56 (3 H, br s, COOH + OH), 4.05 (2 H, m, HOCH₂), 2.40 (3 H, m, COCH, COCH₂) and 2.00–1.40 (6 H, m, CH₂).

Triester 5c. $\nu_{\max}/\text{cm}^{-1}$ 1740 (CO); δ_{H} 4.05 (2 H, m, AcOCH₂), 3.72 (3 H, s, CO₂Me), 3.68 (3 H, s, CO₂Me), 2.98–3.10 (3 H, m, COCH, COCH₂), 2.07 (3 H, s, OCOMe) and 2.00–1.00 (6 H, m, CH₂).

Mixture (5d + 6b). $\nu_{\max}/\text{cm}^{-1}$ 3650–2350 (OH) and 1720 (CO); δ_{H} (60 MHz; CCl₄) 8.96 (br s, CO₂H), 5.33–4.50 (m, AcOCH and OCH), 4.20–3.80 (m, AcOCH₂) and 2.90–0.65 (m, COCH and CH).

Octenedioic acid 7. (Found: C, 55.8; H, 7.3. C₈H₁₂O₄ requires C, 55.8; H, 7.0%) $\nu_{\max}/\text{cm}^{-1}$ 3350–2400 (OH), 1695 (CO) and 1650 (conj. CO); δ_{H} [(CD₃)₂CO–CDCl₃] 6.95 (1 H, m,

CH=CCO₂H), 5.80 (1 H, d, *J* 15, C=CHCO₂H), 5.35 (2 H, br s, CO₂H), 2.60–2.00 (4 H, m, HO₂CCH, C=CCH) and 1.80–1.00 (4 H, m, CH).

Butyl ester 10b. $\nu_{\max}/\text{cm}^{-1}$ 1740 (ester CO) and 1730 (ketone CO); δ_{H} 4.90 (1 H, sextet, *J* 7, CO₂CH), 3.00–1.40 (11 H, m, COCH and *s*-CH), 1.20 (3 H, d, *J* 7, CH₃CEt) and 0.90 (3 H, t, *J* 7, CH₂CH₃).

Lactone 11a (Me ester). $\nu_{\max}/\text{cm}^{-1}$ 1740 (CO); δ_{H} 4.30 (2 H, m, CO₂CH), 3.76 (3 H, s, CO₂Me), 3.00–2.10 (3 H, m, COCH) and 2.10–1.00 (4 H, m, CH) (Found: M⁺, 172.0738. C₈H₁₂O₄ requires *M*, 172.0736).

Lactone 11b (Me ester). $\nu_{\max}/\text{cm}^{-1}$ 1740 (CO); δ_{H} 4.80 (1 H, m, CO₂CH), 3.76 (3 H, s, CO₂Me), 2.70 (4 H, m, COCH) and 1.80 (4 H, m, CH) (Found: M⁺, 172.0735. C₈H₁₂O₄ requires *M*, 172.0736).

Lactone-ester 11c. $\nu_{\max}/\text{cm}^{-1}$ 1740 (CO); δ_{H} 4.90 (1 H, sextet, *J* 7, CO₂CH), 4.30 (2 H, m, CO₂CH₂), 3.00–1.40 (9 H, m, COCH, CH), 1.20 (3 H, d, *J* 7, CH₃CEt) and 0.90 (3 H, t, *J* 7, CH₂CH₃).

Lactone-ester 11d. $\nu_{\max}/\text{cm}^{-1}$ 1740 (CO); δ_{H} 4.90 (2 H, m, CO₂CH), 2.60 (4 H, m, COCH), 2.10–1.40 (6 H, m, CH), 1.20 (3 H, d, *J* 7, CHCH₃) and 0.90 (3 H, t, *J* 7, CH₂CH₃) (Found: M⁺, 214.1205. C₁₁H₁₈O₄ requires *M*, 214.1205).

Acetate 12 (dimethyl ester). (Found: C, 53.55; H, 7.5. C₁₁H₁₈O₆ requires C, 53.7; H, 7.3%) $\nu_{\max}/\text{cm}^{-1}$ 1740 (CO); δ_{H} (60 MHz; CCl₄) 3.95 (2 H, m, AcOCH), 3.78 (3 H, s, CO₂Me), 3.68 (3 H, s, CO₂Me), 2.80–2.00 (3 H, m, COCH), 2.07 (3 H, s, COMe) and 1.60 (4 H, m, CH); *m/z* 246 (M⁺), 187 (M⁺ – 59) and 173 (M⁺ – 73).

Alcohol ester-acid 13 (dibutyl ester). (Found: C, 62.0; H, 9.8. C₁₅H₂₈O₅ requires C, 62.5; H, 9.7%) $\nu_{\max}/\text{cm}^{-1}$ 3550 (OH) and 1740 (CO); δ_{H} 4.95 (2 H, m, CO₂CH), 4.03 (1 H, m, HOCH), 2.80 (1 H, s, OH, D₂O exchangeable), 2.40 (4 H, m, COCH), 2.00–1.40 (8 H, m, CH), 1.20 (6 H, 2 d, *J* 7, CH₃CEt) and 0.90 (6 H, t *J* 7, CH₂CH₃); *m/z* 288 (M⁺), 159 (M⁺ – 129), 141 (M⁺ – 147) and 99 (M⁺ – 189).

Acknowledgements

We thank CSIR for fellowship support to C. D. R.

References

- (a) P. Deslongchamps, *Stereoelectronic Effects in Organic Chemistry*, Pergamon, Oxford, 1983, pp. 163–208; (b) C. L. Perrin and J. D. Thornburn, *J. Am. Chem. Soc.*, 1993, **115**, 3140 and references therein.
- S. Chandrasekhar and C. D. Roy, *Tetrahedron Lett.*, 1987, **28**, 6371.
- J. March, *Advanced Organic Chemistry*, Wiley, New York, 1985, p. 990 and references therein; H. O. House, *Modern Synthetic Reactions*, Benjamin, Menlo Park, 1972, p. 321; G. R. Krow, *Tetrahedron*, 1981, **37**, 2697; C. H. Hassall, *Org. React. (N. Y.)*, 1957, **9**, 73.
- E. L. Eliel, *Stereochemistry of Carbon Compounds*, McGraw–Hill, New York, 1962, p. 279.
- G. Stork, A. Brizzolara, H. Landesman, J. Smuszkovicz and R. Terrell, *J. Am. Chem. Soc.*, 1963, **85**, 207; M. S. Newman and C. A. VanderWerf, *J. Am. Chem. Soc.*, 1945, **67**, 233; R. P. Linstead and E. M. Meade, *J. Chem. Soc.*, 1934, 941.
- J. Cason, L. Wallclave and C. N. Whiteside, *J. Org. Chem.*, 1949, **14**, 41.
- A. I. Vogel, *A Textbook of Practical Organic Chemistry*, Longmans, London, 1971, p. 969.

Paper 4/02516D

Received 27th April 1994

Accepted 31st May 1994