



# The Effect of Inhalation of Fumes from Esbiothrin Based Mosquito Coil on Some Renal Function Markers and Hematological Parameters in Male Wistar Rats

Okeke Chioma Blessing <sup>a\*</sup>, Oladejo Afees Adebayo <sup>a</sup>,  
Metu Mary Cynthia Dumebi <sup>a</sup>, Ndubueze Ifeanyi Kennedy <sup>a</sup>  
and Ayo Ebunolorun Ibukunoluwa <sup>b</sup>

<sup>a</sup> Department of Applied Biochemistry, Faculty of Biosciences, Nnamdi Azikiwe University, Awka, Anambra State, Nigeria.

<sup>b</sup> Department of Biochemistry, Faculty of Life Sciences, University of Ilorin, Ilorin, Nigeria.

## **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

## **Article Information**

### **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/100487>

**Original Research Article**

**Received: 22/03/2023**

**Accepted: 24/05/2023**

**Published: 05/07/2023**

## **ABSTRACT**

Esbiothrin is a synthetic pyrethroid with quick activity used in public health against mosquitoes, houseflies, ectoparasites and cockroaches as well as agricultural applications. These coils and the inhalation of their fumes have been reported and proved to cause grave consequences to the respiratory tract majorly. Hence, this current study was aimed at assessing the renal and hematological effect of inhalation of fumes from esbiothrin based mosquito coil using experimental

\*Corresponding author: E-mail: [ob.chioma@unizik.edu.ng](mailto:ob.chioma@unizik.edu.ng);

animal models. Ten (10) male wistar albino rats were divided into two (2) groups (A and F) of five rats each. Rats in group A were named control and were therefore not exposed to esbiothrin based mosquito coil fumes while group F animals were named test group and were exposed to esbiothrin based mosquito coil fumes for about 6 hours per day, five (5) days per week for four (4) weeks. At the end of the experimental period, blood was collected from each rat for the hematological analysis of Red Blood Cell (RBC), White Blood Cell (WBC), Packed Cell Volume and platelets Count using automated hematology analyzer and also for the analysis of some renal function markers (urea and creatinine). Result revealed that test group had a significant decrease ( $p>0.05$ ) in urea level while there was no significant difference in the creatinine levels. Results further showed a significant decrease ( $p>0.05$ ) in the WBC count, RBC count, PCV and PLT count. Results from this study indicates that esbiothrin based mosquito coil fumes, though considered the least toxic insecticide, are very capable of causing harm.

*Keywords: Esbiothrin; mosquito coil; hematology; renal function.*

## 1. INTRODUCTION

“Mosquitoes are group of very familiar insects that grow and pass their early stages of lives in a variety of aquatic habitats: permanent and transient. Permanent habitats are river, ponds, lakes and so on. and transient habitats include hollow trees, bromeliad tanks, and so on” [1]. “Several lines of evidence established that mosquitoes are main vectors of some catastrophic mammalian and animal diseases such as malaria, dengue, filariasis, chikungunya and West Nile virus” [2]. “As stated by the United States Environmental Protection Agency, mosquito bites can cause skin irritation through an allergic reaction to the mosquito's saliva - this is what causes the red bump and itching” [3].

“As reported by the National Pesticide Information Center, insecticides are pesticides that are formulated to kill, harm, repel or mitigate one or more species of insect” [4]. “However, the use of mosquito coil as a controlling measure against mosquitoes at household level is widespread in malaria endemic countries across Africa, Asia and South America” [5]. “Mosquito coils are smoldering insecticides used in close proximity to persons requiring protection against mosquitoes in order to prevent malaria, a disease of wide distribution” [6]. “Mosquito coil is a mosquito repelling incense-like product formulated from a paste of granulated insecticide and a filler, such as sawdust and other solid materials, and then extruded into a coiled shape” [5]. “Because of their cheapness and readily availability, mosquito coils are the preferred anti-mosquito materials in many developing countries especially Nigeria” [7].

According to the World Health Organization [8] and Midouhas et al [9], “air inside our homes can

be 5 times more toxic than air outside our homes depending on pollutant sources and nature of ventilation of the indoor environment. Mosquito repellents are used in most homes to drive away mosquitoes which are the principal vectors for malaria parasite. The use of these coils is a significant cause of indoor air pollution”. “Recent WHO figures estimated nearly 7 million annual premature deaths; 1 in 8 of the total deaths was due to exposure to air pollution” [10]. “This incredible figure makes air pollution one of the biggest health threats in the world. Most of the deaths from this conservative estimate come from combustion of biomass fuels and smoldering mosquito coils where half of the world's population in the developing countries relies on these crude fuels for multiple needs” [11,12]. “The disease burden in developing countries is relatively high, and malaria-a vector borne disease is no exception. Approximately 2 billion people worldwide are using mosquito coils” [13]. “These coils are usually made up active ingredients that could be any or a combination of pyrethroids such as metofluthrin, d-allethrin d-trans allethrin, and prallethrin and the percentage of the active ingredients usually ranged between 0.10% and 2.0%” [10,13]. These substances are low-toxicity insecticides.

“In Nigeria, mosquito coils have been the major mosquito repellents because they are not expensive and can easily be purchased in the neighborhood” [10]. “Mosquito coils contain insecticides that slowly vaporize into the air to provide mosquito protection to prevent malaria. Some of the toxic contaminants that may result from the burning of mosquito coils and related incense-like items are carbon monoxide (CO), sulfur dioxide (SO<sub>2</sub>), nitrogen dioxide (NO<sub>2</sub>), and particulate matters (PM)” [14].

This research focused on the investigation of the effect of inhaling mosquito coil fume (esbiothrin) on some renal function markers and on some hematological parameters. It may provide scientific rationale to either promote or discourage the combustion of this mosquito coil in a population.

## 2. MATERIALS AND METHODS

### 2.1 Mosquito Coil

The mosquito coil was purchased from a retail outlet in Awka, Anambra State, Nigeria.

### 2.2 Experimental Animals

Ten (10) male wistar albino rats were purchased from an animal farm in Mgbakwu, Anambra State. Prior to commencement of the study, the rats were acclimatized for two (2) weeks. The rats were housed in standard cages dressed with sawdust litter, at housing conditions of 12 light: 12 dark cycles in the animal house and fed growers mash and water for the entire duration of the study. All procedures were followed according to the guidelines of proper conduct of animal experimentation.

### 2.3 Design of Exposure

Inhalation of ablaze coil fumes was achieved by the construction of a microenvironment mimicking a room with a dimension of 50cm x 45cm x 20cm and two openings with a dimension of 10cm x 10cm (mimicking the windows) on opposite sides for cross ventilation. The rats were observed for any clinical signs associated with the exposure to the active ingredient from the fumes.

### 2.4 Research Plan

After acclimatization, the animals were weighed before the commencement of the experiment. They were then randomly grouped into two groups (the test and control group) with five rats per group.

**Group A:** Control group which were not exposed to the esbiothrin based mosquito coil.

**Group B:** Rats were exposed to esbiothrin based mosquito coil for six (6) hours of five days per week for four weeks.

### 2.5 Place of Study

This research was carried out at the laboratory of Applied Biochemistry, Nnamdi Azikiwe University, Awka.

### 2.6 Collection of Samples

After the last day of exposure, the rats were fasted overnight before being sacrificed. Blood collection was done by heart puncture with use of a syringe and into clean plain and EDTA bottles.

### 2.7 Biochemical Analysis

Blood Urea and creatinine contents were determined using commercial kits from Randox Laboratories and manufacturer's instructions were duly followed.

### 2.8 Hematological Analysis

The samples were then sent to the laboratory for the determination of the full blood count (packed cell volume, red blood cell, and white blood cell) using automated hematology analyzer (Minary Model BC 3200).

### 2.9 Data Analysis

Data was analyzed statistically using Statistical Package for the Social Sciences (SPSS). Statistical significance was evaluated by one-way analysis of variance (ANOVA) for renal function test and hematological analysis.

## 3. RESULTS

The result of this study is presented in Figs. 1 and 2 as well as in Table 1.

Fig. 1 showed the effect of esbiothrin based mosquito coil on the urea levels of experimental animals after exposure. Result showed a significant decrease ( $p>0.05$ ) in the urea levels of test group compared to the control group.

There was no significant difference in the serum creatinine levels of the test group ( $0.65\pm 0.12\text{mg/dl}$ ) and that of the control group ( $0.65\pm 0.08\text{mg/dl}$ ) (Fig. 2).

Effect of esbiothrin based mosquito coil on hematological parameters of experimental animals after exposure to esbiothrin is presented in Table 1 result showed a significant decrease ( $p>0.05$ ) in the levels of white blood cell count,

red blood cell count, packed cell volume and platelets count of the test group when compared with the control.

#### 4. DISCUSSION

“Mosquito coils are insecticides primarily manufactured from the pyrethrum plant. However, with increasing demand for this type of pest control, due to its cheapness and availability, synthetic pyrethroid mainly esters are produced” [15]. One of the pyrethroid esters is esbiothrin (the active ingredient of the mosquito coil used in this study).


“Assessment of renal function is important in the management of patients with kidney disease or pathologies affecting renal function. Tests of renal function have utility in identifying the presence of renal disease monitoring the response of kidneys to treatment, and determining the progression of renal disease” [16]. The result showing the effect of the fumes of esbiothrin based mosquito coil in urea level is presented in Fig 1. From this result, there was a significant decrease in the urea level of the rats exposed to the mosquito coil when compared to the control group. The control had a higher urea level ( $57.40 \pm 6.33 \text{ mg/dl}$ ) while the test group had a lower urea level ( $30.85 \pm 2.63 \text{ mg/dl}$ ). The findings of this research are in discordant with the findings of Abdrabouh, [17] who stated that there was a significant increase in serum levels of urea of young and adult rat.

Urea is major nitrogen containing substance that is cleared from the blood by the kidney in to the urine [18]. “It is useful in differential diagnosis of

acute renal failure and pre-renal condition where blood urea nitrogen–creatinine ratio is increased” [19]. According to Ambardekar, [20], the normal range of urea nitrogen in blood or serum ranges from 7 to 20 mg/dl. Since increased levels of serum urea are pointer to severe kidney damage, it could be inferred, from the result of this study that continuous inhalation of fumes from this coil might pose a great threat to the kidney.


The result showing the effect of the fumes of esbiothrin based mosquito coil in creatinine level is presented in Fig 2. There was no significant difference in the serum creatinine levels of the test group ( $0.65 \pm 0.12 \text{ mg/dl}$ ) and that of the control group ( $0.65 \pm 0.08 \text{ mg/dl}$ ). This was contrary with the findings of Hui et al., [21] that reported an increase in creatinine level of wistar rats exposed to pyrethroid based compound. This result showed a significant decrease in the level of creatinine of the test group when compared to the control group.

“Creatinine is an end product of muscle catabolism of creatine. For a given individual, creatinine production is relatively stable and mainly depends on muscle mass” [22]. “The creatinine clearance test is used to monitor the progression of renal disease. In chronic renal failure and uremia, an eventual reduction occurs in the excretion of creatinine by both the glomeruli and the tubules” [23]. The normal reference range for creatinine levels is 0.6 to 1.2 mg/dl in males and 0.5 to 1.1 mg/dl in females. Therefore, with the values gotten, it falls within the range of the standard reference range which might indicates no loss of kidney function.



**Fig. 1. Effect of esbiothrin based mosquito coil on the urea levels of experimental animals after exposure**

*The test group shows a significant decrease ( $p > 0.05$ ) in the urea levels*



**Fig. 2. Effect of esbiothrin based mosquito coil on the creatinine levels of experimental animals after exposure**

*The test group shows no significant difference in the creatinine levels*

**Table 1. Effect of esbiothrin based mosquito coil on hematological parameters of experimental animals after exposure**

	WBC	RBC	HGB	PCV	PLT
<b>Control</b>	13.01±1.72	6.97 ±0.20	15.52 ±0.20	46.92±0.73	717.40±50.45
<b>Esbiothrin</b>	12.89±0.23	5.02±0.88	10.00±1.77	28.94±6.91	642.60±41.72

*Result showed a significant decrease ( $p>0.05$ ) in the levels of white blood cell count, red blood cell count, packed cell volume and platelets count*

The white blood cell count (WBC) estimates the total number of white blood cells per liter in the blood [24]. The control group had the higher WBC ( $13.01\pm 1.72 \times 10^9/L$ ) while the test group had the lower WBC ( $12.89\pm 0.23 \times 10^9/L$ ) as presented in Table 1. When the control group was compared to the test group, there was a significant decrease ( $p>0.05$ ). This was at variance with the findings of Saka et al., [25] whose findings showed an increase in WBC after exposure to pyrethroid insecticides. This may be due to the fact that the authors carried out a minimal exposure of three (3) minutes daily for 3 weeks, a shorter duration of exposure. The normal reference range WBC is  $4.5$  to  $11.0 \times 10^9/L$ . The values gotten from the research surpasses the normal standard range. High white blood cell count could indicate infections, inflammation or immune system disorder [26].

A red blood cell count is a test that measures the number of red blood cells (RBCs) in the blood [27,28]. The control group had higher RBC ( $6.97\pm 0.20 \times 10^9/L$ ) while the test group had lower RBC ( $5.02\pm 0.88 \times 10^9/L$ ). When the control group was compared to the test group, there was a significant decrease ( $p>0.05$ ). The normal standard range for RBC is  $4.5$  to  $6.5 \times 10^{12}/L$  in males while  $3.8$  to  $5.8 \times 10^{12}/L$  [29]. The values

gotten from this study is a little lower than the normal standard reference range. A low RBC count could also indicate a vitamin B6, B12 or folate deficiency [26]. It may also signify kidney disease [30].

Packed Cell Volume (PCV) is the percentage of red blood cells in the total blood volume [28,31]. It is carried out to measure the volume of packed red blood cells relative to whole blood i.e. the proportion of red blood cells in the blood. The control group had the higher PCV ( $46.92\pm 0.73$  %) while the test group had the least ( $28.94\pm 6.91$  %). When the control group was compared to the test group, there was a significant decrease ( $p>0.05$ ). This result is in consonance with Kamal et al. [32] that also recorded a decrease in PCV levels but was in contrast with the findings of Saka et al. [25], whose report showed an increase in PCV levels. This however, could be due to minimal duration of exposure. A normal adult male shows a PCV of 40% to 54% and a female show 36% to 48% [33]. Increased PCV levels may be grounds to indicate dehydration.

The platelet count is a standard component in screening for coagulation abnormalities [34]. The control group has a higher PLT count ( $717.40\pm 50.45 \times 10^9/L$ ) while the test group has a

lower PLT count ( $642.60 \pm 41.7210^9/L$ ). Results from this study which presents platelets count decrease is in contrast with the previous by study Saka et al., [25], this might suggest that at minimal exposure, esbiothrin do not have a suppressive effect on the thrombopoietin, the regulator of platelets production. It also shows that pyrethroid cause structural or functional damage at high exposure. The normal platelet count in humans ranges from  $150 \times 10^9/L$  to  $400 \times 10^9/L$  [35]. High level of platelets count could be an indicator of hemostasis or lung disease.

## 5. CONCLUSION

The results of the present study suggest that despite being the least toxic insecticide safer for human beings than other insecticides such as dichlorodiphenyltrichloroethane (DDT), esbiothrin based mosquito coil still have harmful effects, as exposure to it can challenge the immune system and induce adverse changes on the kidney. However, based on the results from this work, studies aimed at producing alternative mosquito coils with minimal toxicity should be an area of practical interest.

## CONSENT

It is not applicable.

## ETHICAL APPROVAL

This research was cleared and approved by the Animal Research Ethics Committee (AREC) of Nnamdi Azikiwe University, Awka.

## CONFERENCE DISCLAIMER

Some part of this manuscript was previously presented in the conference: ISPEC 11th International Conference on Agriculture, Animal Sciences and Rural Development 03-05 March 2023-MUS, Turkey. Web Link of the proceeding: [https://en.ispeco.org/\\_files/ugd/614b1f\\_37d23320e66646bd901b58aeaf711c5c.pdf](https://en.ispeco.org/_files/ugd/614b1f_37d23320e66646bd901b58aeaf711c5c.pdf).

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Santos CF, Silva AC, Rodrigues RA, Sannidy J, Jesus RD, Borges M. Inventory of mosquitoes (diptera: Culicidae) in

- conservation units in Brazilian tropical dry forests. *Revista do Instituto de Medicina Tropical de Sao Paulo (Journal of the Institute of Tropical Medicine of Sao Paulo)*. 2015;57(3):227-232.
2. Garba S, Adelaiye A, Mshelia L. Histopathological and biochemical changes in the rat's kidney following exposure to a pyrethroid based mosquito coil. *The Journal of Applied Science Research*. 2007;3:1788-1793.
3. United States Environmental Protection Agency (EPA). General Information about mosquitoes; 2021. Available:<https://www.epa.gov/mosquitocontrol/general-information-about-mosquitoes>
4. National Pesticide Information Center (NPIC). Mosquitoes; 2019. Available:<http://npic.orst.edu/pest/mosquito/index.html>
5. Hogarh NJ, Thomas PA, Emmanuel DJ, Silas WA, Julius NF, Kwasi O. Environmental health risks and benefits of the use of mosquito coils as malaria prevention and control strategy. *Malaria Journal*. 2018;17:265.
6. Anyabolu AE, Ezejindu DN, Obinwa BN. Evaluation of toxic effect of D-allevrin based mosquito's coil on the lungs and selected haematological parameters of adult wistar rats. *Journal of Advances in Medical and Pharmaceutical Sciences*. 2021;23(4):9-19.
7. Agrawal A, Yadav AK, Singh R. Ameliorating effects of garlic oil against mosquito coil induced histopathological changes in rat kidney. *International Journal of Pharma and Biosciences*. 2013;4(2):1112-1116.
8. WHO. Health Aspects of Air Pollution With Particulate Matter, Ozone and Nitrogen Dioxide: Report on a WHO Working Group; 2003. Available:[http://www.euro.who.int/\\_data/assets/pdf\\_file/0005/112199/E79097](http://www.euro.who.int/_data/assets/pdf_file/0005/112199/E79097).
9. Midouhas E, Kokosi T, Flouri E. The quality of air outside and inside the home: Associations with emotional and behavioural problem scores in early childhood. *BMC Public Health*. 2019;19:406.
10. Elehinafe FB, Okedere OB, Adesanmi AJ, Jimoh EM. Assessment of indoor levels of carbon monoxide emission from smoldering mosquito coils used in Nigeria. *Environmental Health Insights*. 2022;16:1-6.

11. Taylor ET, Nakai S. The levels of toxic air pollutants in kitchens with traditional stoves in rural Sierra Leone. *Journal of Environmental Protection*. 2012;03:1353-1363.
12. Taylor ET, Wirmvem MJ, Sawyerr VH, Nakai S. Diurnal concentrations and variation of carbon monoxide in indoor and outdoor air of residential homes in Western Sierra Leone. *Environmental Pollution*. 2015;4:10-18.
13. Ogoma SB, Moore SJ, Maia MF. A systematic review of mosquito coils and passive emanators: Defining recommendations for spatial repellency testing methodologies. *Parasite Vectors*. 2012;5:287.
14. Huang Y, Wang J, Chen Y. Household PM<sub>2.5</sub> pollution in rural Chinese homes: Levels, dynamic characteristics and seasonal variations. *Science of the Total Environment*: 2022;817:153085.
15. Taylor ET, Beah J, Barrie M, James MS, Kaitibi D, Sannoh A. Characterizing emission of particulate matter from combusting different products of mosquito coils. *JSM Environmental Science & Ecology*. 2017;5(4):1054.
16. Damiati SA. A pilot study to assess kidney functions and toxic dimethylarginines as risk biomarkers in women with low vitamin D levels. *Journal of Medical Biochemistry*. 2019;38(2):145-152.
17. Abdrabouh EA. Susceptibility of young and adult rat kidney to impacts of mosquito coil fumes. *Egyptian Journal of Basic and Applied Sciences*. 2021;8(1):1-11.
18. Stoppler MC. Definition of urea; 2021. Available:<https://www.rxlist.com/urea/definition.html>
19. Shivaraj G, Prakash S, Sonal N, Vinayak K. Markers of renal function tests. *North American Journal of Medical Sciences*. 2010;2(4):170-173.
20. Ambardekar N. What is a blood urea? 2021. Available:<https://www.webmd.com/a-to-z-guides/blood-urea-nitrogen-test>
21. Hui L, Shaik AH, Daoud A, Suliman YA, Umeshareen S, Hamzah AHA, Narendra M. Induced alteration of rat erythrocyte membrane with effect of pyrethroid based compounds. *Saudi Journal of Biological Sciences*. 2020;27(12):3669-3675.
22. Delanaye P, Cavalier E, Maillard N, Krzesinski JM, Mariat C, Cristol JP, Pieroni L. Creatinine. *Annales de Biologie Clinique*, the official Journal of the French Society of Clinical Biology (SFBC). 2010;68:531-543.
23. Edmund L, David J. Kidney function tests. In: *Tietz Textbook of Clinical Chemistry and Molecular Diagnostic*. 2016:797-808.
24. Higuera V. White blood cell count; 2018. Available:<https://www.healthline.com/health/wbc-count> Access on 16<sup>th</sup> January, 2022
25. Saka WA, Akhigbe RE, Azeez OM, Babatunde TR. Effects of pyrethroid insecticide exposure on hematological and hemaostatic profiles in rats. *Pakistan Journal of Biological Sciences*. 2011;14(22):1024-1027.
26. Oladejo AA, Osukoya O. Hematological profiles of naturally infected pigs treated with *Bridelia ferruginea* leaf extracts. *Asian Hematology Research Journal*. 2021;4(2):1-10.
27. Cafasso J. Red blood cell count; 2018. Available:<https://www.healthline.com/health/rbc-count>
28. Ezeigwe OC, Oladejo AA, Iloanya EL. The impact of *Citrus aurantium* fruit juice on bodyweight and haematological parameters of wistar rats. *International Journal of Research and Reports in Hematology*. 2022;5(2):56-67.
29. Dean L. Blood groups and red cell antigens. National Center for Biotechnology Information (NCBI). Chapter 1: Blood and the cells it contains; 2016. Available:<https://www.ncbi.nlm.nih.gov/books/NBK2263/table/ch1.T1/>
30. Oladejo AA, Onwubuya EI, Okafor CS, Okeke CB, Ikimi CG, Okeke CM. Blood cells formative properties of *B. pinnatum* in chronic inflammatory disorders: an experience with wistar rats. *International Journal of Research and Reports in Hematology*. 2022;5(2):274-279.
31. Nall R. Hematocrit test. 2018. Available:<https://www.healthline.com/health/hematocrit>
32. Kamal SM, Khan A, Rizvi F, Sadeeq UR. Effect of cypermethrin on clinico-hematological parameters in rabbits. *Paulistsan Veterinary Journal*. 2007;27(4):171-175.
33. Billet H. Haemoglobin and haematocrit. In: *Clinical Methods: the physical and laboratory examinations*. 3<sup>rd</sup> edition. 2003:110.

34. Gropper MD. Thrombocytopenia. In: Miller's Science and Analysis. 2020:112.
35. Martina E. Platelets count. Haemtologica. 2011;96(1):10-13.

---

© 2023 Okeke et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*

*<https://www.sdiarticle5.com/review-history/100487>*