

# Use of stable isotopes to determine diets of living and extinct bears

G.V. Hilderbrand, S.D. Farley, C.T. Robbins, T.A. Hanley, K. Titus, and C. Servheen

**Abstract:** The potential use of stable-isotope analyses ( $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$ ) to estimate bear diets was assessed in 40-day feeding trials using American black bears (*Ursus americanus*). Bear plasma and red blood cells have half-lives of ~4 days and ~28 days, respectively. The isotopic signature of bear plasma is linearly related to that of the diet, and with the exception of adipose tissue, there is no isotopic fractionation across bear tissues. Isotopic analyses were used to estimate the diets of three bear populations: Pleistocene cave bears (*U. speleaus*) in Europe, grizzly bears (*Ursus arctos horribilis*) inhabiting the Columbia River drainage prior to 1931, and brown bears (*U. arctos*) of Chichagof and Admiralty islands, Alaska. Cave bears were omnivores with terrestrially produced meat contributing from 41 to 78% ( $58 \pm 14\%$ ) of their metabolized carbon and nitrogen. Salmon contributed from 33 to 90% ( $58 \pm 23\%$ ) of the metabolized carbon and nitrogen in grizzly bears from the Columbia River drainage. Finally, most brown bears on Chichagof and Admiralty islands feed upon salmon during the late summer and fall; however, a subpopulation of bears exists that does not utilize salmon.

**Résumé :** Le potentiel des analyses au moyen d'isotopes stables ( $\delta^{13}\text{C}$  et  $\delta^{15}\text{N}$ ) pour estimer le régime alimentaire des ours a été évalué au cours d'expériences d'alimentation d'une durée de 40 jours chez des Ours noirs (*Ursus americanus*). Le plasma des ours a une demi-vie de ~4 jours et leurs érythrocytes, une demi-vie de ~28 jours. La signature isotopique du plasma est en relation linéaire avec celle du régime alimentaire; de plus, à l'exception de ce qui prévaut dans le tissu adipeux, il n'y a pas de fractionnement isotopique dans les tissus de l'ours. Des analyses au moyen d'isotopes ont été utilisées pour étudier le régime alimentaire chez trois populations d'ours : des ours des cavernes du Pléistocène (*U. speleaus*) en Europe, des Grizzlis (*Ursus arctos horribilis*) du bassin du Columbia d'avant 1931 et des Ours bruns (*U. arctos*) des îles Chichagof et Admiralty, en Alaska. Les ours des cavernes étaient des omnivores et consommaient de la chair d'animaux terrestres leur fournissant de 41 à 78% ( $58 \pm 14\%$ ) de leur carbone et azote métabolisables. Les saumons fournissaient aux Grizzlis du bassin du Columbia de 33-90% de leur carbone et azote métabolisables. Enfin, la plupart des Ours bruns des îles Chichagof et Admiralty se nourrissent de saumon à la fin de l'été et à l'automne; cependant, une sous-population de ces ours survit sans consommer de saumons.

[Traduit par la Rédaction]

## Introduction

Food habits of bears have been studied by scat analyses and direct observation of foraging behavior. Scat analyses, even when corrected for differential disappearance, almost always estimate diet at the level of the population rather than at that of the individual or subpopulation (e.g., males versus females), cannot be used for foods that are completely digested and produce no quantifiable residues (e.g., meat),

and do not indicate the nutritional importance of different foods. Fecal correction factors used to convert scat analyses to actual food habits can be very difficult to estimate. For example, bears frequently avoid consuming the skin of large mammals, and other carnivores may use the same carcass. A correction factor for large-mammal residues in a bear consuming ungulates varies 18-fold when the consumption of skin and hair ranges from 5 to 100% (Hewitt 1989).

Stable-isotope analyses, e.g., carbon ( $\delta^{13}\text{C}$ ) and nitrogen ( $\delta^{15}\text{N}$ ), of bear tissues can complement fecal analyses by overcoming the above weaknesses, as well as indicate the historical diet of dead bears and the current and past diets of living bears. Carbon isotope analyses of animal tissue indicate the relative contributions of marine and terrestrial sources to the animal's carbon pool (DeNiro and Epstein 1978; Ramsey and Hobson 1991; Hobson and Welch 1992). In marine ecosystems, carbon enters as bicarbonate and  $^{13}\text{C}$  is enriched relative to terrestrial ecosystems where carbon enters as  $\text{CO}_2$  (Peterson and Fry 1987). Nitrogen isotope analyses indicate the trophic position of the consumer.  $^{15}\text{N}$  is enriched relative to  $^{14}\text{N}$  at each step in a food chain (DeNiro and Epstein 1980; Minagawa and Wada 1984; Hobson and Welch 1992), probably because of the preferen-

Received December 13, 1995. Accepted May 17, 1996.

**G.V. Hilderbrand and S.D. Farley.** Department of Zoology, Washington State University, Pullman, WA 99164, U.S.A.

**C.T. Robbins.** Departments of Natural Resource Sciences and Zoology, Washington State University, Pullman, WA 99164, U.S.A.

**T.A. Hanley.** United States Forest Service, Forestry Sciences Laboratory, 2700 Sherwood Lane, Suite 2A, Juneau, AK 99801, U.S.A.

**K. Titus.** Division of Wildlife, Alaska Department of Fish and Game, Douglas, AK 99824, U.S.A.

**C. Servheen.** United States Fish and Wildlife Service, Forestry Sciences Laboratory, University of Montana, Missoula, MT 59812, U.S.A.

tial excretion of the lighter  $^{14}\text{N}$  in urinary waste products (Peterson and Fry 1987).

Thus, the objectives of the experimental portion of this study were to determine (i) the relationship between carbon and nitrogen isotope ratios in bear tissues relative to ratios found in the diet, and (ii) the turnover time of isotopes in different bear tissues relative to dietary change. Finally, to illustrate the ecological applications of isotopic analyses, we determined the trophic level of extinct European cave bears (*Ursus spelaeus*), the importance of salmon to early grizzlies (*Ursus arctos horribilis*) of the Pacific Northwest, and the spring and summer diets of brown bears (*U. arctos*) on Chichagof and Admiralty islands, southeast Alaska.

### Cave bear trophic level

Cave bears were  $2\frac{1}{2}$  to 3 times larger than modern brown bears, presumably hibernated as do other temperate-zone bears, and, based on dental and jaw structure, are hypothesized to have been the most herbivorous of all bears (Kurten 1976). However, the herbivore hypothesis seems untenable, owing to the poor digestibility of vegetation by bears (Dierenfeld et al. 1982; Pritchard and Robbins 1990). Rates of gain by bears are lower on vegetation than on more digestible dietary items such as meat or berries (C.T. Robbins, unpublished data). Additionally, the small size of most wild fruits produces relatively small bite sizes and harvest rates, so the maximum bear size attainable on most wild fruits is only a fraction of the size of an adult cave bear (Welch et al.).<sup>1</sup> Thus, we hypothesize that cave bears were omnivores like most other temperate-zone bears, but because of their large size were specialized for feeding on megafauna, including carrion.

### Historical use of salmon by Pacific Northwest grizzly bears

Columbia River salmon populations have been reduced by hydroelectric dams and irrigation projects impeding or completely blocking their migration routes; by mining, road construction, and logging suffocating spawning streams with silt, debris, or toxic chemicals; and by overfishing (Lantz 1967; Fulton 1968, 1970; Platts 1970). Current research and management efforts are directed towards recovery of both salmon populations and their spawning areas (Volkman and McConaha 1993).

The grizzly bear, which is listed as threatened under the Endangered Species Act of 1975, was one of the organisms that historically fed on spawning salmon (Davis et al. 1986). Two of the six proposed grizzly bear recovery areas are the North Cascades of Washington and the Bitterroot Ecosystem of Idaho (Servheen 1984). The North Cascades' runs of chinook salmon (*Oncorhynchus tshawytscha*), sockeye salmon (*O. nerka*), coho salmon (*O. kisutch*), and steelhead trout (*Salmo gairdneri*) and the Bitterroot Ecosystem's runs of chinook salmon and steelhead trout have been depleted by human activities and development (Fulton 1968, 1970). Information on the historical importance of salmon in the diet of grizzly bears would benefit the drafting and implementation of grizzly bear recovery plans.

### Seasonal diets of brown bears on Chichagof and Admiralty islands

Telemetry data collected from brown bears on Chichagof and Admiralty islands suggest that a subpopulation exists which geographically has access to spawning salmon but does not utilize salmon as a food resource (K. Titus, unpublished data). However, because of the difficulty or expense of continuously monitoring bear locations, one can never be certain that a specific bear did not visit a stream containing salmon. Stable-isotope analyses of bear tissue can determine if a specific bear utilized salmon as a food resource.

## Methods

### Sample collection

Feeding trials were conducted at the Washington State University Bear Research, Conservation, and Education Center in 1994. Experimental groups consisting of three American black bears (*Ursus americanus*) were fed one of seven experimental diets (Table 1) for 40 days during which blood samples were collected every 10 days. The mixed diets were formulated to provide 35 and 65% of their digestible dry matter from either herring or beef, with vegetable matter providing the rest. Dry matter digestibilities of the mixed-diet ingredients were determined according to Pritchard and Robbins (1990). The plant matter in the mixed diets was derived from  $\text{C}_3$  plants, which are depleted in  $^{13}\text{C}$  relative to  $\text{C}_4$  plants. Temperate-zone and subarctic bears such as cave bears, American black bears, and brown or grizzly bears fed or feed almost exclusively on  $\text{C}_3$  plants, whereas  $\text{C}_4$  plants could contribute significantly to the diets of tropical bears such as the sloth bear and sun bear (Raven et al. 1986).


To assess isotopic fractionation between consumer tissues, plasma, red blood cells, hair, claws, adipose tissue, muscle, skin, and bone collagen were collected from three domestic rabbits (*Oryctolagus cuniculus*), three laboratory mice (*Mus musculus*), and three yearling American black bears. The rabbits and mice were raised on a single diet (R and R Rabbit Pellets 781 and Rat and Mouse Standard Expanded, Beekay Feeds) from birth to the time of sampling, while the bears were fed a single diet (Dealer's Pride, Purina) for the 7 months prior to sampling.

Bone samples from 10 cave bears were obtained from the American Museum of Natural History, New York, the National Museum of Natural History, Washington, D.C., and the Bulgarian Museum of Natural History, Sofia. Original collection sites included France, Czechoslovakia, Bulgaria, and Italy. Bone samples from four extinct, similarly aged herbivores (woolly rhinoceros (*Coelodonta antiquitatus*), woolly mammoth (*Mammuthus primigenius*), auroch (*Bos primigenius*), and horse/zebra (*Equus* sp.)) from Denmark, Hungary, and Italy were also obtained from collections at the same museums.

All federal, state, and university museums in the United States that house ursid collections were surveyed (Choate and Genoways 1975). Of the approximately 2500 grizzly specimens listed, 11 had been collected between 1890 and 1931 from the Columbia River drainage of Washington, Oregon, and Idaho (American Museum of Natural History, New York; National Museum of Natural History, Washington D.C.; University of Montana, Missoula). One of the 11 bears is a type specimen and therefore was not available for sampling. Samples of bone ( $n = 9$ ; cranial material) and hair ( $n = 3$ ) were collected from the 10 remaining specimens. Bone ( $n = 5$ ) and hair ( $n = 1$ ) samples were also obtained from grizzly bears that were collected in central Montana and eastern Wyoming between 1856 and 1888. The latter bears did not have access to salmon during their lifetime. Finally, bone samples ( $n = 5$ ) were obtained from southeastern Alaska coastal brown bears (1916–1921) that had access to abundant salmon.

<sup>1</sup> C.A. Welch, J. Keay, K.C. Kendall, and C.T. Robbins. Constraints to frugivory by bears. Submitted for publication.

**Fig. 1.** Representative isotope equilibration curves for plasma (a and b) and red blood cells (c and d) of American black bears when a dietary change occurs. Dietary changes were as follows: bear chow to 35% herring : 65% vegetation (a and b), salmon to apples (c), and 35% herring : 65% vegetation to mule deer (d).




**Table 1.** Diets fed to American black bears during 40-day feeding trials.

Diet	δ <sup>13</sup> C	δ <sup>15</sup> N
Salmon ( <i>Oncorhynchus tshawytscha</i> )	-20.1	11.2
Mule deer ( <i>Odocoileus hemionus</i> )	-26.2	6.4
Apples ( <i>Malus</i> sp.)	-24.1	2.3
46.6% alfalfa, 42.7% herring meal, 5.6% wheat, 5.0% honey, 0.1% vitamix	-21.8	9.5
72.4% alfalfa, 17.9% herring meal, 5.6% wheat, 5.0% honey, 0.1% vitamix	-22.4	5.8
53.3% alfalfa, 39.7% beef meal, 5.0% honey, 1.9% fat, 0.1% vitamix	-22.3	4.4
77.1% alfalfa, 15.9% beef meal, 5.0% honey, 1.9% fat, 0.1% vitamix	-22.1	2.9

Eleven blood and 7 hair samples were collected between June 24 and July 1, 1994, from 11 brown bears inhabiting Chichagof and Admiralty islands. Spawning salmon are abundant in the streams of

these islands during the late summer and fall, but the returning salmon had not yet arrived when these samples were collected. An additional five hair samples were obtained from bears captured on

**Fig. 2.** Relationship between  $^{13}\text{C}$  signatures of the diet and of equilibrated plasma in American black bears (●) and polar bears (▲; Hobson and Welch 1992). SEE, standard error of the estimate.



Admiralty Island between 1981 and 1992. The latter bears were suspected to be alpine bears that did not use salmon.

**Sample analyses**

Lipids were extracted from hair, skin, and muscle with a chloroform-ether solvent. Collagen was extracted from bone according to Chisholm et al. (1983). All samples were then freeze-dried and ground to <1 mm. For  $^{13}\text{C}$  analysis, 25 mg of sample and 0.5 g of  $\text{CuO}$  were placed in a 6-mm Vycor tube, flame-sealed under vacuum, and baked at  $550^\circ\text{C}$  for 6 h (Hobson and Clark 1992a). Carbon dioxide from the combusted sample was then collected and purified using successive dry ice - methanol and liquid nitrogen traps. To assess  $^{15}\text{N}$  content, 10 mg of sample was mixed with 4 g of  $\text{CuO}$ , 3 g of  $\text{Cu}$ , and 0.4 g of  $\text{CaO}$ , placed in a 9-mm Vycor tube, flame-sealed under vacuum, and baked at  $850^\circ\text{C}$  for 2 h (Fiedler and Proksch 1975). Collected samples were then analyzed for carbon and nitrogen isotope content using a Finnigan MAT  $\delta\text{S}$  mass spectrometer. Results are reported as ratios relative to PeeDee limestone ( $^{13}\text{C}$ ) or atmospheric nitrogen ( $^{15}\text{N}$ ) as follows:

$$\delta X = [(R_{\text{sample}}/R_{\text{standard}}) - 1] \times 10^3$$


where  $X$  is  $^{13}\text{C}$  or  $^{15}\text{N}$  and the  $R$  is  $^{13}\text{C}/^{12}\text{C}$  or  $^{15}\text{N}/^{14}\text{N}$  ratio (Peterson and Fry 1987). Mixed-diet isotopic signatures (Table 1) are weighted averages based on the product of the digestible dry matter and isotopic ratios of each ingredient.

**Results**

**Captive bear feeding trials**

Carbon and nitrogen isotope ratios in bear plasma equilibrated with the diet within 10 days, whereas the isotope ratios of red blood cells continued to change after 40 days (Fig. 1). The half-life of the plasma was less than 4 days, whereas the half-life of the red blood cells averaged 28 days. Carbon and nitrogen isotope ratios in bear plasma were linearly related to the carbon and nitrogen isotope ratios of single-item diets and the weighted, available dietary carbon

**Fig. 3.** Relationship between  $^{15}\text{N}$  signatures of the diet and that of equilibrated plasma in American black bears (●) and polar bears (▲; Hobson and Welch 1992). SEE, standard error of the estimate.



and nitrogen ratios in mixed diets (Figs. 2, 3). As hypothesized, plasma from bears consuming terrestrially produced plants or animals was much less enriched with  $^{13}\text{C}$  than that of bears consuming food from marine sources (Fig. 2). Similarly,  $^{15}\text{N}$  concentrations of bear plasma increased with increasing trophic level (Fig. 3). Finally, all tissues except fat (ANOVA,  $p < 0.0001$ ) have the same isotopic enrichment when animals are on a constant diet (Fig. 4). Thus, the isotopic relationships between diet and plasma (Figs. 2, 3) also apply to hair, bone collagen, and red blood cells.

The dietary sources of assimilated carbon and nitrogen can be estimated by first determining the magnitude of the marine component. The marine salmon component (percent digestible dry matter) of a bear's diet can be estimated as

$$[1] \quad M = \frac{^{13}\text{C} + 24.1}{4.0} \times 100$$


where  $^{13}\text{C}$  is the estimated isotope ratio of the diet when the equation in Fig. 2 is solved for  $X$ ,  $-24.1 \pm 1.3\text{‰}$  (reported as the mean  $\pm$  standard deviation) is the average temperate-zone plant and animal terrestrial signature (Chisholm and Schwartz 1982; Tieszen et al. 1983; this study), and  $4.0 \pm 0.5\text{‰}$  is the difference between the average terrestrial signature and the average signature ( $-20.1 \pm 1.0\text{‰}$ ) for four species of salmon (chinook salmon, coho salmon, chum salmon (*O. keta*), and pink salmon (*O. gorbuscha*)) (M. Ben-David, unpublished data; this study).

Once the marine salmon component has been determined, the dietary contribution of carbon and nitrogen absorbed from terrestrially produced meat ( $T_m$ , %) can be estimated by


$$[2] \quad T_m = \left[ \frac{^{15}\text{N} - (M \times 11.5) - 2.3}{4.1} \times (1 - M) \right] \times 100$$

where  $^{15}\text{N}$  is the estimated nitrogen enrichment of the diet when the equation in Fig. 3 is solved for the measured iso-

**Fig. 4.** Isotopic signatures of tissues collected from laboratory mice, domestic rabbits, and American black bears fed a constant diet for either 7 months (bears) or throughout life (rabbits and mice).



**Fig. 5.** Isotopic signatures of bone collagen extracted from cave bears ( $\blacktriangle$ ) and Pleistocene herbivores ( $\bullet$ ).



tope ratio of the bear's diet,  $M$  is the marine dietary contribution in decimal form,  $11.5 \pm 0.4\text{‰}$  is the average  $^{15}\text{N}$  signature of four species of salmon (Mathisen et al. 1988; Kline et al. 1990; M. Ben-David, unpublished data; this study),  $2.3 \pm 0.6\text{‰}$  is the average  $^{15}\text{N}$  signature of terrestrial plant matter (DeNiro and Epstein 1980; this study), and  $4.1 \pm 0.5\text{‰}$  is the average difference in  $^{15}\text{N}$  enrichment between primary producers (plants) and their consumers (Minagawa and Wada 1984; Sholto-Douglas et al. 1991; Hobson and Welch 1992; this study). Finally, the proportion of carbon and nitrogen absorbed from terrestrial plant matter is 100 minus the sum of the marine and terrestrially produced meat contributions.


#### Cave bear trophic level

Cave bear bones are significantly enriched ( $t$  test,  $p < 0.0001$ ) with  $^{15}\text{N}$  ( $\bar{x} = 8.4 \pm 0.5\text{‰}$ ) relative to the four herbivore bones ( $\bar{x} = 6.2 \pm 0.3\text{‰}$ ), but there is no difference in  $^{13}\text{C}$  (Fig. 5). When eq. 2 is used to predict the amount of meat in the cave bear's diet with the mean terrestrial plant signature reduced from  $2.3\text{‰}$  to  $1.6\text{‰}$ , the average based on solving the equation in Fig. 3 for the observed nitrogen enrichments in the four herbivores, meat provided  $58 \pm 14\%$  (range 41–78%) of all metabolized carbon and nitrogen absorbed by cave bears.

#### Historical use of salmon by Pacific Northwest grizzly bears

Grizzly bears collected from coastal Alaska and the Columbia River drainage were significantly enriched in  $^{13}\text{C}$  and  $^{15}\text{N}$  relative to those collected from eastern Wyoming and central Montana ( $t$  test,  $p < 0.0002$ ) (Fig. 6). Salmon contributed 33–90% ( $58 \pm 23\%$ ) of the carbon and nitrogen assimilated from the historical diet of grizzly bears living in the Columbia River drainage (Fig. 7). Terrestrially produced meat (e.g., ungulates, rodents, and insects) contributed 0–21% ( $8 \pm 11\%$ ) and plant matter contributed 10–64% ( $34 \pm 19\%$ ) of metabolized carbon and nitrogen. Grizzly

**Fig. 6.** Carbon and nitrogen enrichment of hair and bone collected from grizzly bears killed between 1856 and 1931 in eastern Wyoming and central Montana ( $\bullet$ ), the Columbia River drainage ( $\blacksquare$ ), and coastal southeast Alaska ( $\blacktriangle$ ).



bears from central Montana and eastern Wyoming obtained their assimilated carbon and nitrogen from terrestrially produced meat (13–54%,  $35 \pm 15\%$ ) and plant matter (46–87%,  $65 \pm 15\%$ ), while coastal Alaskan bears obtained virtually all of their carbon and nitrogen from salmon ( $94 \pm 9\%$ ).

#### Seasonal diets of brown bears on Chichagof and Admiralty islands


The carbon and nitrogen isotope signatures of red blood cells collected during the early summer prior to the annual return of spawning salmon are much less enriched than the hair that grows from late June through September, when the salmon are present (Fig. 8). The absorbed carbon and nitrogen used to produce the red blood cells was derived from terrestrial plants ( $85 \pm 15\%$ ), terrestrially produced meat ( $14 \pm 16\%$ ), and salmon ( $1 \pm 2\%$ ). The values contrast with the sources of the carbon and nitrogen in the hair ( $34 \pm 27\%$  from terrestrial plants,  $11 \pm 21\%$  from terrestrially produced meat,  $55 \pm 34\%$  from salmon) for those bears with significant  $^{13}\text{C}$  enrichment. However, as hypothesized, a subpopulation of bears does exist that maintained a year-long terrestrial diet.

## Discussion


#### Captive bear feeding trials

Owing to different turnover rates, the isotopic signatures of different consumer tissues serve as windows to different dietary periods. Plasma provides an indication of an animal's diet during the week prior to sampling, while red blood cells reflect the diet of the consumer over the past 2–3 months (Hobson and Clark 1992b; this study). Hair signatures are representative of the diet during the period of hair growth, which begins in midsummer and continues throughout the fall, while collagen, because of its long turnover rate, can be used to indicate a yearly or lifetime diet (e.g., van

**Fig. 7.** Locations and estimated contributions of salmon, terrestrially produced meat, and plant matter (values are given in this order) to the assimilated carbon and nitrogen pool of grizzly bears killed in the northwestern United States between 1856 and 1931. Values in parentheses are from hair samples; all other values are from bone collagen. Idaho's Bitterroot Ecosystem and Washington's North Cascades grizzly bear recovery areas are indicated.



**Fig. 8.** Carbon and nitrogen enrichment of hair ( $\blacktriangle$ ) and red blood cells ( $\bullet$ ) collected from brown bears on Chichagof and Admiralty islands, Alaska.



der Merwe and Vogel 1978; Stenhouse and Baxter 1979; Chisholm and Schwarcz 1982; Tieszen et al. 1989; Hobson and Montevecchi 1991; Loken et al. 1992).

Equilibrated bear plasma was enriched in  $^{13}\text{C}$  from 0.4 to 4.5 at dietary values of  $-18.5$  to  $-25.5$ ‰. Hobson and

Clark (1992a) reported enrichment values in various quail tissues ranging from  $0.2$ ‰ (liver) to  $2.7$ ‰ (collagen). An enrichment in  $^{15}\text{N}$  of  $4.1 \pm 0.8$ ‰ over the diet was present in equilibrated bear plasma. Similar  $^{15}\text{N}$  enrichments at each increase in trophic level have been reported by DeNiro and Epstein (1980), Minagawa and Wada (1984), and Hobson and Welch (1992). The depletion of  $^{13}\text{C}$  in adipose tissue relative to other tissues analyzed has been reported elsewhere (DeNiro and Epstein 1977; Tieszen et al. 1983; Hobson 1995).

#### Cave bear trophic level

Bocherens et al. (1994) concluded that cave bears were primarily herbivores, because of low  $^{15}\text{N}$  enrichments. While collagen  $^{15}\text{N}$  levels in Pleistocene herbivores reported by Bocherens et al. (1994) ( $6.6 \pm 0.7$ ‰) did not differ from those reported in this study ( $6.2 \pm 0.3$ ‰), their reported levels for cave bears spanned two entire trophic levels, with signatures ranging from that characteristic of a plant ( $3.4$ ‰) to that of a true carnivore ( $9.3$ ‰). Similarly, Bocherens et al.'s (1994)  $^{15}\text{N}$  levels for living ( $1.9$ – $5.9$ ‰) and extinct ( $3.8$ – $7.4$ ‰) herbivores were not sufficiently different from the lower limit of living ( $6.2$ – $9.9$ ‰) and extinct ( $8.0$ ‰) carnivores to have identified an omnivore. Known omnivores in their data set, such as brown and black bears, had nitrogen enrichments ( $3.8$ – $8.7$ ‰) that are very similar to the enrichments they reported for cave bears. Thus, owing to the variation in their  $^{15}\text{N}$  results, Bocherens et al.'s (1994) conclusion that cave bears were herbivores is untenable.

ble. Matheus (1995) and Bocherens et al. (1995) have constructed trophic models of bear populations based on isotopic signatures. When applied to the current data, both models show a clear distinction between herbivores and the cave bears in this study, and support the hypothesis that cave bears were omnivores.

Kurten (1976) concluded that the cause of cave bear extinction, which occurred 11 000 years ago (Brown 1993), is unknown. However, the extinction of the cave bear mirrored the extinction of large herbivores (Kurten 1968). The loss of this major source of dietary carbon and nitrogen would have been extremely detrimental to such a large omnivore and was likely the most important factor contributing to its extinction.

### Historical use of salmon by Pacific Northwest grizzly bears

While the recovery of historic spawning areas would benefit grizzly bear populations reintroduced to regions with reduced salmon runs, salmon are not essential for bear survival (Davis et al. 1986). In other ecosystems lacking salmon as a food resource, grizzly bears feed on vegetation, insects, ungulates, freshwater fish, and small mammals (Hamer et al. 1991; Mattson et al. 1991; Clevenger et al. 1992). However, studies have identified a strong positive correlation between the autumn mass of female bears and their reproductive success during hibernation (Rogers 1976; Stringham 1990). Bears' reproduction rate, one of the slowest of any terrestrial mammal (Bunnell and Tait 1981), is a critical factor in long-term population viability. Because spawning salmon are more nutrient dense than virtually any other food resource available to bears in the Pacific Northwest, grizzly bear recovery would be enhanced by simultaneous salmon recovery in the North Cascades and Bitterroot Ecosystem.

### Seasonal diets of brown bears on Chichagof and Admiralty islands

Brown bears of Chichagof and Admiralty islands feed exclusively on food from terrestrial sources during spring and early summer. Upon the arrival of spawning salmon in the late summer and fall, most of the brown bears shift their diet to include salmon. However, four bears analyzed did not utilize the returning salmon and may be representative of a subpopulation that survives on a wholly terrestrial diet.

In summary, stable-isotope analyses offer insight into the biology of both living and dead bears. The ability to determine the proportions of meat and (or) vegetation in a bear's diet during various time intervals from samples collected during a single capture can lead to a more comprehensive understanding of the nutritional ecology of individual bears and bear populations than is possible using more conventional scat analyses or direct observation alone.

### Acknowledgements

Financial support for this project was provided by the Inter-mountain Research Station (J. Lyon) and the Northwest Forest and Range Experiment Station of the United States Forest Service, the United States Fish and Wildlife Service, and the International Grizzly Fund. Peter Larson and Charles Knaack, Washington State University Department of Geology, provided the mass spectrometer and assisted with its use, and Randy George, National Hydrology Research Insti-

tute, Saskatoon, provided advice regarding  $^{15}\text{N}$  analyses. D. Dyer of the University of Montana, B. Mader of the American Museum of Natural History, R. Fisher of the National Museum of Natural History, and N. Spassov of the Bulgarian Museum of Natural History donated specimens for isotopic analyses. M. Evans, Washington State University Department of Statistics, provided advice regarding statistical analyses. Additionally, two anonymous reviewers provided valuable and constructive comments. Finally, thanks are extended to the many students who assisted in the care of the captive bears throughout this project.

### References

- Bocherens, H., Emslie, S.D., Billiou, D., and Mariotti, A. 1995. Stable isotopes ( $^{13}\text{C}$ ,  $^{15}\text{N}$ ) and paleodiet of the giant short-faced bear (*Arctodus simus*). *Paleontol. Geochim.* **320**: 779–784.
- Bocherens, H., Fizet, M., and Mariotti, A. 1994. Diet, physiology and ecology of fossil mammals as inferred from stable carbon and nitrogen isotope biogeochemistry: implications for Pleistocene bears. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **107**: 213–225.
- Brown, G. 1993. *The great bear almanac*. Lyons and Burford, Publishers, New York.
- Bunnell, F.L., and Tait, D.E.N. 1981. Population dynamics of bears—implications. *In* Dynamics of large mammal populations. Edited by C.W. Fowler and T.D. Smith. John Wiley and Sons, New York. pp. 75–98.
- Chisholm, B.S., and Schwarcz, H.P. 1982. Stable-carbon isotope ratios as a measure of marine versus terrestrial protein in ancient diets. *Science (Washington, D.C.)*, **216**: 1131–1132.
- Chisholm, B.S., Nelson, D.E., Hobson, K.A., Schwarcz, H.P., and Kynf, M. 1983. Carbon isotope measurement techniques for bone collagen: notes for the archaeologist. *J. Archaeol. Sci.* **10**: 355–360.
- Choate, J.R., and Genoways, H.H. 1975. Collections of recent mammals in North America. *J. Mammal.* **56**: 452–502.
- Clevenger, A.P., Purroy, F.J., and Pelton, M.R. 1992. Food habits of brown bears (*Ursus arctos*) in the Cantabrian Mountains, Spain. *J. Mammal.* **73**: 415–421.
- Davis, D.L., Melquist, W.E., and Graham, D. 1986. The Selway–Bitterroot Ecosystem as grizzly bear habitat. *U.S. For. Serv. Gen. Tech. Rep.* **207**. pp. 158–162.
- Dierenfeld, E.S., Hintz, H.F., Robertson, J.B., VanSoest, P.J., and Oftedahl, O.T. 1982. Utilization of bamboo by the giant panda. *J. Nutr.* **112**: 636–641.
- DeNiro, M.J., and Epstein, S. 1977. Mechanism of carbon fractionation associated with lipid synthesis. *Science (Washington, D.C.)*, **197**: 261–263.
- DeNiro, M.J., and Epstein, S. 1978. Influence of diet on the distribution of carbon isotopes in animals. *Geochim. Cosmochim. Acta*, **42**: 495–506.
- DeNiro, M.J., and Epstein, S. 1980. Influence of diet on the distribution of nitrogen isotopes in animals. *Geochim. Cosmochim. Acta*, **45**: 341–351.
- Fiedler, R., and Proksch, G. 1975. The determination of nitrogen-15 by emission and mass spectrometry in biochemical analysis: a review. *Anal. Chem. Acta*, **78**: 1–62.
- Fulton, L.A. 1968. Spawning areas and abundance of chinook salmon (*Oncorhynchus tshawytscha*) in the Columbia River Basin—past and present. *U.S. Fish. Wildl. Ser. Spec. Sci. Rep. Fish. No.* **571**. pp. 1–26.
- Fulton, L.A. 1970. Spawning areas and abundance of steelhead trout, coho, sockeye, and chum salmon in the Columbia River Basin—past and present. *Nat. Mar. Fish. Ser. Spec. Sci. Rep. Fish. No.* **618**. pp. 1–37.



- Hamer, D., Herrero, S., and Brady, K. 1991. Food and habitat used by grizzly bears, *Ursus arctos*, along the Continental Divide in Waterton Lakes National Park, Alberta. *Can. Field-Nat.* **105**: 325–329.
- Hewitt, D.G. 1989. Correcting grizzly bear fecal analysis to observed food habits. M.S. thesis, Washington State University, Pullman.
- Hobson, K.A. 1995. Reconstructing avian diets using stable-carbon and nitrogen isotope analysis of egg components: patterns of isotopic fractionation and turnover. *Condor*, **97**: 752–762.
- Hobson, K.A., and Clark, R.G. 1992a. Assessing avian diets using stable isotopes I: turnover of  $^{13}\text{C}$  in tissues. *Condor*, **94**: 181–188.
- Hobson, K.A., and Clark, R.G. 1992b. Turnover of  $^{13}\text{C}$  in cellular and plasma fractions of blood: implications for nondestructive sampling in avian dietary studies. *Auk*, **110**: 638–641.
- Hobson, K.A., and Welch, H.E. 1992. Determination of trophic relationships within a high Arctic marine food web using  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  analysis. *Mar. Ecol. Prog. Ser.* **84**: 9–18.
- Hobson, K.A., and Montevecchi, W.A. 1991. Stable isotopic determinations of trophic relationships of great auks. *Oecologia*, **87**: 528–531.
- Kline, T.C., Goering, J.J., Mathisen, O.A., and Poe, P.H. 1990. Recycling of elements transported upstream by runs of Pacific salmon: I.  $\delta^{15}\text{N}$  and  $\delta^{13}\text{C}$  evidence in Sashin Creek, south-eastern Alaska. *Can. J. Fish. Aquat. Sci.* **47**: 136–144.
- Kurten, B. 1968. Pleistocene mammals of Europe. Aldine Publishing Co., Chicago.
- Kurten, B. 1976. The cave bear story. Columbia University Press, New York.
- Lantz, R.L. 1967. An ecological study of the effects of logging on salmonids. *In* Proceedings of the 47th Annual Conference of the Western Association of State Game and Fish Commissioners, July 16–20, 1967, Honolulu, Hawaii. pp. 323–335.
- Loken, B., Rothenburger, D., and Tieszen, L.L. 1992. Inferences about diets based on  $\delta^{13}\text{C}$  analysis of collagen and other tissues from modern and early historic animals. *Proc. S.D. Acad. Sci.* **71**: 85–94.
- Matheus, P.E. 1995. Diet and co-ecology of Pleistocene short-faced bears and brown bears in eastern Beringia. *Quat. Res. (N.Y.)*, **44**: 447–453.
- Mathisen, O.A., Parker, P.L., Goering, J.J., Kline, T.C., Poe, P.H., and Scanlin, R.S. 1988. Recycling of marine elements transported into freshwater by anadromous salmon. *Verh. Int. Ver. Limnol.* **23**: 2249–2258.
- Mattson, D.J., Blanchard, B.M., and Knight, R.R. 1991. Food habits of Yellowstone grizzly bears, 1977–1987. *Can. J. Zool.* **69**: 1619–1629.
- Minagawa, M., and Wada, E. 1984. Stepwise enrichment of  $^{15}\text{N}$  along food chains: further evidence and the relation between  $\delta^{15}\text{N}$  and animal age. *Geochim. Cosmochim. Acta*, **48**: 1135–1140.
- Peterson, B.J., and Fry, B. 1987. Stable isotopes in ecosystem studies. *Annu. Rev. Ecol. Syst.* **18**: 293–320.
- Platts, W.S. 1970. The effects of logging and road construction on the aquatic habitat of the South Fork Salmon River, Idaho. *In* Proceedings of the 50th Annual Conference of the Western Association of State Game and Fish Commissioners, July 13–16, 1970, Victoria, B.C. pp. 182–185.
- Pritchard, G.T., and Robbins, C.T. 1990. Digestive and metabolic efficiencies of grizzly and black bears. *Can. J. Zool.* **68**: 1645–1651.
- Ramsey, M.A., and Hobson, K.A. 1991. Polar bears make little use of terrestrial food webs: evidence from stable-carbon isotope analysis. *Oecologia*, **86**: 598–600.
- Raven, P.H., Evert, R.F., and Eichorn, S.E. 1986. Biology of plants. Worth Publishers, New York.
- Rogers, L. 1976. Effects of mast and berry crop failure on survival, growth, and reproductive success in black bears. *Trans. North Am. Wildl. Nat. Resour. Conf.* **41**: 431–438.
- Servheen, C. 1984. The status of the grizzly bear and the inter-agency grizzly bear recovery effort. *In* Proceedings of the 64th Annual Conference of the Western Association of Fish and Wildlife Agencies, July 16–19, 1984, Victoria, B.C. pp. 227–234.
- Sholto-Douglas, A.D., Field, J.G., James, A.G., and van der Merwe, N.J. 1991.  $^{13}\text{C}/^{12}\text{C}$  and  $^{15}\text{N}/^{14}\text{N}$  isotope ratios in the Southern Benguela Ecosystem: indicators of food web relationships in different size-classes of plankton and pelagic fish: differences between fish muscle and bone collagen tissues. *Mar. Ecol. Prog. Ser.* **78**: 23–32.
- Stenhouse, M.J., and Baxter, M.S. 1979. The uptake of bomb  $^{14}\text{C}$ . *In* Radiocarbon dating. Edited by R. Burger and H.E. Suess. University of California Press, Berkeley. pp. 324–341.
- Stringham, S.F. 1990. Grizzly bear reproductive rate relative to body size. *In* Bears—Their Biology and Management: Proceedings of the Eighth International Conference on Bear Research and Management, 20–25 February 1989, Victoria, B.C. Edited by L.M. Darling and W.R. Archibald. International Association for Bear Research and Management. [Available from Michael R. Pelton, Department of Forestry, Wildlife, and Fisheries, The University of Tennessee, P.O. Box 1071, Knoxville, TN 37901-1071, U.S.A. pp. 433–443.
- Tieszen, L.L., Boutton, T.W., Otchichilo, W.K., Nelson, D.E., and Brandt, D.H. 1989. An assessment of long-term food habits of Tsavo elephants based on stable carbon and nitrogen isotope ratios of bone collagen. *Afr. J. Ecol.* **27**: 219–226.
- Tieszen, L.L., Boutton, T.W., Tesdahl, K.G., and Slade, N.A. 1983. Fractionation and turnover of stable carbon isotopes in animal tissues: implications  $\delta^{13}\text{C}$  analysis of diet. *Oecologia*, **57**: 32–37.
- van der Merwe, N.J., and Vogel, J.C. 1978.  $^{13}\text{C}$  content of human bone collagen as a measure of prehistoric diet in woodland North America. *Nature (Lond.)*, **276**: 815–816.
- Volkman, J.M., and McConnaha, W.E. 1993. Through a glass, darkly: Columbia River salmon, the Endangered Species Act, and adaptive management. *Environ. Law*, **23**: 1249–1272.

**This article has been cited by:**

1. Oliver E. Barker, Andrew E. Derocher, Mark A. Edwards. 2014. Use of Arctic ground squirrels (*Urocitellus parryii*) by brown bears (*Ursus arctos*). *Polar Biology* . [[CrossRef](#)]
2. Ricardo Fernandes, John Meadows, Alexander Dreves, Marie-Josée Nadeau, Pieter Grootes. 2014. A preliminary study on the influence of cooking on the C and N isotopic composition of multiple organic fractions of fish (mackerel and haddock). *Journal of Archaeological Science* **50**, 153-159. [[CrossRef](#)]
3. Phillips Donald L., Inger Richard, Bearhop Stuart, Jackson Andrew L., Moore Jonathan W., Parnell Andrew C., Semmens Brice X., Ward Eric J.. 2014. Best practices for use of stable isotope mixing models in food-web studies. *Canadian Journal of Zoology* **92**:10, 823-835. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
4. Jun Matsubayashi, Junko Morimoto, Tsutomu Mano, Achyut Aryal, Futoshi Nakamura. 2014. Using stable isotopes to understand the feeding ecology of the Hokkaido brown bear (*Ursus arctos*) in Japan. *Ursus* **25**, 87-97. [[CrossRef](#)]
5. Diana J. R. Lafferty, Jerrold L. Belant, Donald L. Phillips. 2014. Testing the niche variation hypothesis with a measure of body condition. *Oikos* n/a-n/a. [[CrossRef](#)]
6. Charles C. Schwartz, Justin E. Teisberg, Jennifer K. Fortin, Mark A. Haroldson, Christopher Servheen, Charles T. Robbins, Frank T. van Manen. 2014. Use of isotopic sulfur to determine whitebark pine consumption by Yellowstone bears: A reassessment. *Wildlife Society Bulletin* **38**:10.1002/wsb.v38.3, 664-670. [[CrossRef](#)]
7. Maciej T. Krajcarz, Magdalena Krajcarz. 2014. The 200,000 year long record of stable isotopes ( $\delta^{18}O$ ,  $\delta^{13}C$ ) of cave bear (*Ursus spelaeus*) teeth from Biśnik Cave, Poland. *Quaternary International* **339-340**, 119-130. [[CrossRef](#)]
8. Hervé Bocherens, Gennady Baryshnikov, Wim Van Neer. 2014. Were bears or lions involved in salmon accumulation in the Middle Palaeolithic of the Caucasus? An isotopic investigation in Kudaro 3 cave. *Quaternary International* **339-340**, 112-118. [[CrossRef](#)]
9. Anneke H. van Heteren, Ann MacLarnon, Christophe Soligo, Todd C. Rae. 2014. Functional morphology of the cave bear (*Ursus spelaeus*) cranium: a three-dimensional geometric morphometric analysis. *Quaternary International* **339-340**, 209-216. [[CrossRef](#)]
10. M. Cattet, B. J. Macbeth, D. M. Janz, A. Zedrosser, J. E. Swenson, M. Dumond, G. B. Stenhouse. 2014. Quantifying long-term stress in brown bears with the hair cortisol concentration: a biomarker that may be confounded by rapid changes in response to capture and handling. *Conservation Physiology* **2**, cou026-cou026. [[CrossRef](#)]
11. John B. Hopkins, Jesse Whittington, Anthony P. Clevenger, Michael A. Sawaya, Colleen Cassady St. Clair. 2014. Stable isotopes reveal rail-associated behavior in a threatened carnivore. *Isotopes in Environmental and Health Studies* **50**, 322-331. [[CrossRef](#)]
12. Jennapher L. Teunissen van Manen, Lisa I. Muller, Zheng-hua Li, Arnold M. Saxton, Michael R. Pelton. 2014. Using stable isotopes to assess dietary changes of American black bears from 1980 to 2001. *Isotopes in Environmental and Health Studies* **50**, 382-398. [[CrossRef](#)]
13. Carolyn M. Kurlle, Paul L. Koch, Bernie R. Tershy, Donald A. Croll. 2014. The effects of sex, tissue type, and dietary components on stable isotope discrimination factors ( $\Delta^{13}C$  and  $\Delta^{15}N$ ) in mammalian omnivores. *Isotopes in Environmental and Health Studies* **50**, 307-321. [[CrossRef](#)]
14. Andrew R. Thurber. 2014. Diet-dependent incorporation of biomarkers: implications for food-web studies using stable isotope and fatty acid analyses with special application to chemosynthetic environments. *Marine Ecology* n/a-n/a. [[CrossRef](#)]
15. H. M. Bryan, C. T. Darimont, P. C. Paquet, K. E. Wynne-Edwards, J. E. G. Smits. 2014. Stress and reproductive hormones reflect inter-specific social and nutritional conditions mediated by resource availability in a bear-salmon system. *Conservation Physiology* **2**, cou010-cou010. [[CrossRef](#)]
16. Torsten W. Bentzen, Richard T. Shideler, Todd M. O'Hara. 2014. Use of stable isotope analysis to identify food-conditioned grizzly bears on Alaska's North Slope. *Ursus* **25**, 14-23. [[CrossRef](#)]
17. Ashley K. McGrew, Lora R. Ballweber, Sara K. Moses, Craig A. Stricker, Kimberlee B. Beckmen, Mo D. Salman, Todd M. O'Hara. 2014. Mercury in gray wolves (*Canis lupus*) in Alaska: Increased exposure through consumption of marine prey. *Science of The Total Environment* **468-469**, 609-613. [[CrossRef](#)]
18. A.H. Thompson, A.S. Wilson, J.R. Ehleringer Hair as a Geochemical Recorder 371-393. [[CrossRef](#)]

19. Charles C. Schwartz, Jennifer K. Fortin, Justin E. Teisberg, Mark A. Haroldson, Christopher Servheen, Charles T. Robbins, Frank T. Van Manen. 2014. Body and diet composition of sympatric black and grizzly bears in the Greater Yellowstone Ecosystem. *The Journal of Wildlife Management* **78**:10.1002/jwmg.v78.1, 68-78. [[CrossRef](#)]
20. Alexander Braun, Karl Auerswald, Armin Vikari, Hans Schnyder. 2013. Dietary protein content affects isotopic carbon and nitrogen turnover. *Rapid Communications in Mass Spectrometry* **27**:10.1002/rcm.v27.23, 2676-2684. [[CrossRef](#)]
21. Maciej T. Krajcarz, Magdalena Krajcarz, Adrian Marciszak. 2013. Paleoecology of bears from MIS 8–MIS 3 deposits of Biśnik Cave based on stable isotopes ( $\delta^{13}\text{C}$ ,  $\delta^{18}\text{O}$ ) and dental cementum analyses. *Quaternary International* . [[CrossRef](#)]
22. K. J. Jenkins, N. D. Chelgren, K. A. Sager-Fradkin, P. J. Happe, M. J. Adams. 2013. OCCUPANCY PATTERNS OF MAMMALS AND LENTIC AMPHIBIANS IN THE ELWHA RIVER RIPARIAN ZONE BEFORE DAM REMOVAL. *River Research and Applications* n/a-n/a. [[CrossRef](#)]
23. Kimberly Asmus Hersey, Joseph D. Clark, James B. Layzer. 2013. Consumption of Freshwater Bivalves by Muskrats in the Green River, Kentucky. *The American Midland Naturalist* **170**, 248-259. [[CrossRef](#)]
24. Erik J. Blomberg, Simon R. Poulson, James S. Sedinger, Daniel Gibson. 2013. Prefledging diet is correlated with individual growth in Greater Sage-Grouse ( *Centrocercus urophasianus* ). *The Auk* **130**, 715-724. [[CrossRef](#)]
25. Seger R.L., Servello F.A., Cross R.A., Keisler D.H.. 2013. Body mass and mast abundance influence foraging ecology of the American black bear (*Ursus americanus*) in Maine. *Canadian Journal of Zoology* **91**:7, 512-522. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)] [[Supplemental Material](#)]
26. Luis Malpica-Cruz, Sharon Z. Herzka, Oscar Sosa-Nishizaki, Miguel A. Escobedo-Olvera. 2013. Tissue-specific stable isotope ratios of shortfin mako ( *Isurus oxyrinchus* ) and white ( *Carcharodon carcharias* ) sharks as indicators of size-based differences in foraging habitat and trophic level. *Fisheries Oceanography* n/a-n/a. [[CrossRef](#)]
27. Rumiko Nakashita, Yuzuru Hamada, Eishi Hirasaki, Juri Suzuki, Toru Oi. 2013. Characteristics of stable isotope signature of diet in tissues of captive Japanese macaques as revealed by controlled feeding. *Primates* **54**, 271-281. [[CrossRef](#)]
28. Andrew Robertson, Robbie A. McDonald, Richard J. Delahay, Simon D. Kelly, Stuart Bearhop. 2013. Whisker growth in wild Eurasian badgers *Meles meles*: implications for stable isotope and bait marking studies. *European Journal of Wildlife Research* **59**, 341-350. [[CrossRef](#)]
29. Robu Marius, Fortin Jennifer K., Richards Michael P., Schwartz Charles C., Wynn Jonathan G., Robbins Charles T., Trinkaus Erik. 2013. Isotopic evidence for dietary flexibility among European Late Pleistocene cave bears (*Ursus spelaeus*). *Canadian Journal of Zoology* **91**:4, 227-234. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)] [[Supplemental Material](#)]
30. L. Cheng, A. M. Nicol, R. J. Dewhurst, G. R. Edwards. 2013. The effects of dietary nitrogen to water-soluble carbohydrate ratio on isotopic fractionation and partitioning of nitrogen in non-lactating sheep. *animal* 1-6. [[CrossRef](#)]
31. Van Daele M.B., Robbins C.T., Semmens B.X., Ward E.J., Van Daele L.J., Leacock W.B.. 2013. Salmon consumption by Kodiak brown bears (*Ursus arctos middendorffi*) with ecosystem management implications. *Canadian Journal of Zoology* **91**:3, 164-174. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
32. Shannon Crowley, Chris J. Johnson, Dexter P. Hodder. 2013. Spatio-Temporal Variation in River Otter ( *Lontra canadensis* ) Diet and Latrine Site Activity. *Ecoscience* **20**, 28-39. [[CrossRef](#)]
33. M. Vašinová Galiová, M. Nývltová Fišáková, J. Kynický, L. Prokeš, H. Neff, A.Z. Mason, P. Gadas, J. Košler, V. Kanický. 2013. Elemental mapping in fossil tooth root section of *Ursus arctos* by laser ablation inductively coupled plasma mass spectrometry (LA-ICP-MS). *Talanta* **105**, 235-243. [[CrossRef](#)]
34. Brian Milakovic, Katherine L. Parker. 2013. Quantifying carnivory by grizzly bears in a multi-ungulate system. *The Journal of Wildlife Management* **77**:10.1002/jwmg.v77.1, 39-47. [[CrossRef](#)]
35. Ana C. Pinto-Llona. 2013. Macrowear and occlusal microwear on teeth of cave bears *Ursus spelaeus* and brown bears *Ursus arctos*: Inferences concerning diet. *Palaeogeography, Palaeoclimatology, Palaeoecology* **370**, 41-50. [[CrossRef](#)]
36. C. A. M. France, D. W. Owsley. 2013. Stable Carbon and Oxygen Isotope Spacing Between Bone and Tooth Collagen and Hydroxyapatite in Human Archaeological Remains. *International Journal of Osteoarchaeology* n/a-n/a. [[CrossRef](#)]
37. Ricardo Fernandes, Marie-Josée Nadeau, Pieter M. Grootes. 2012. Macronutrient-based model for dietary carbon routing in bone collagen and bioapatite. *Archaeological and Anthropological Sciences* **4**, 291-301. [[CrossRef](#)]

38. Silvina Botta, Aleta A. Hohn, Stephen A. Macko, Eduardo R. Secchi. 2012. Isotopic variation in delphinids from the subtropical western South Atlantic. *Journal of the Marine Biological Association of the United Kingdom* **92**, 1689-1698. [[CrossRef](#)]
39. T.C. O'Connell, C.J. Kneale, N. Tasevska, G.G.C. Kuhnle. 2012. The diet-body offset in human nitrogen isotopic values: A controlled dietary study. *American Journal of Physical Anthropology* **149**:10.1002/ajpa.v149.3, 426-434. [[CrossRef](#)]
40. Anthony J. Orr, Seth D. Newsome, Jeffrey L. Laake, Glenn R. VanBlaricom, Robert L. DeLong. 2012. Ontogenetic dietary information of the California sea lion (*Zalophus californianus*) assessed using stable isotope analysis. *Marine Mammal Science* **28**, 714-732. [[CrossRef](#)]
41. Sora Lee Kim, Dave R. Casper, Felipe Galván-Magaña, Ruth Ochoa-Díaz, Sandra Berenice Hernández-Aguilar, Paul L. Koch. 2012. Carbon and nitrogen discrimination factors for elasmobranch soft tissues based on a long-term controlled feeding study. *Environmental Biology of Fishes* **95**, 37-52. [[CrossRef](#)]
42. Daryl Codron, Matt Sponheimer, Jacqui Codron, Ian Newton, John L. Lanham, Marcus Clauss. 2012. The confounding effects of source isotopic heterogeneity on consumer-diet and tissue-tissue stable isotope relationships. *Oecologia* **169**, 939-953. [[CrossRef](#)]
43. Jonathan J. Derbridge, Paul R. Krausman, Chris T. Darimont. 2012. Using Bayesian stable isotope mixing models to estimate wolf diet in a multi-prey ecosystem. *The Journal of Wildlife Management* **76**:10.1002/jwmg.v76.6, 1277-1289. [[CrossRef](#)]
44. A. Borrell, N. Abad-Oliva, E. Gómez-Campos, J. Giménez, A. Aguilar. 2012. Discrimination of stable isotopes in fin whale tissues and application to diet assessment in cetaceans. *Rapid Communications in Mass Spectrometry* **26**, 1596-1602. [[CrossRef](#)]
45. Leslie R. Germain, Matthew D. McCarthy, Paul L. Koch, James T. Harvey. 2012. Stable carbon and nitrogen isotopes in multiple tissues of wild and captive harbor seals (*Phoca vitulina*) off the California coast. *Marine Mammal Science* **28**:10.1111/mms.2012.28.issue-3, 542-560. [[CrossRef](#)]
46. Donald L. Phillips. 2012. Converting isotope values to diet composition: the use of mixing models. *Journal of Mammalogy* **93**, 342-352. [[CrossRef](#)]
47. John P. Whiteman, Katie A. Greller, Henry J. Harlow, Laura A. Felicetti, Karyn D. Rode, Merav Ben-David. 2012. Carbon isotopes in exhaled breath track metabolic substrates in brown bears (*Ursus arctos*). *Journal of Mammalogy* **93**, 413-421. [[CrossRef](#)]
48. Merav Ben-David, Seth D. Newsome, John P. Whiteman. 2012. Lipid and amino acid composition influence incorporation and discrimination of  $^{13}\text{C}$  and  $^{15}\text{N}$  in mink. *Journal of Mammalogy* **93**, 399-412. [[CrossRef](#)]
49. Matthew J. Kohn, Moriah P. McKay. 2012. Paleoecology of late Pleistocene-Holocene faunas of eastern and central Wyoming, USA, with implications for LGM climate models. *Palaeogeography, Palaeoclimatology, Palaeoecology* **326-328**, 42-53. [[CrossRef](#)]
50. Jason N. Waite, Stephen J. Trumble, Vladimir N. Burkanov, Russel D. Andrews. 2012. Resource partitioning by sympatric Steller sea lions and northern fur seals as revealed by biochemical dietary analyses and satellite telemetry. *Journal of Experimental Marine Biology and Ecology* **416-417**, 41-54. [[CrossRef](#)]
51. Gary A. King. 2012. Isotopes as palaeoeconomic indicators: new applications in archaeoentomology. *Journal of Archaeological Science* **39**, 511-520. [[CrossRef](#)]
52. M. Pérez-Rama, D. Fernández-Mosquera, A. Grandal-d'Anglade. 2011. Recognizing growth patterns and maternal strategies in extinct species using stable isotopes: The case of the cave bear *Ursus spelaeus* ROSENMÜLLER. *Quaternary International* **245**, 302-306. [[CrossRef](#)]
53. J. J. Duda, H. J. Coe, S. A. Morley, K. K. Kloehn. 2011. Establishing spatial trends in water chemistry and stable isotopes ( $\delta^{15}\text{N}$  and  $\delta^{13}\text{C}$ ) in the Elwha River prior to dam removal and salmon recolonization. *River Research and Applications* **27**:10.1002/rra.v27.10, 1169-1181. [[CrossRef](#)]
54. Christine A.M. France, Jennifer A. Giaccai, Nadia Cano. 2011. The effects of PVAc treatment and organic solvent removal on  $\delta^{13}\text{C}$ ,  $\delta^{15}\text{N}$ , and  $\delta^{18}\text{O}$  values of collagen and hydroxyapatite in a modern bone. *Journal of Archaeological Science* **38**, 3387-3393. [[CrossRef](#)]

55. Hervé Bocherens, Mathias Stiller, Keith A. Hobson, Martina Pacher, Gernot Rabeder, James A. Burns, Thomas Tütken, Michael Hofreiter. 2011. Niche partitioning between two sympatric genetically distinct cave bears (*Ursus spelaeus* and *Ursus ingressus*) and brown bear (*Ursus arctos*) from Austria: Isotopic evidence from fossil bones. *Quaternary International* **245**, 238-248. [[CrossRef](#)]
56. KnottKatrina K., BoydDaryle, YlitaloGina M., O'HaraTodd M.. 2011. Concentrations of mercury and polychlorinated biphenyls in blood of Southern Beaufort Sea polar bears (*Ursus maritimus*) during spring: variations with lipids and stable isotope ( $\delta^{15}\text{N}$ ,  $\delta^{13}\text{C}$ ) values. *Canadian Journal of Zoology* **89**:11, 999-1012. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
57. K. A. Hatch, B. L. Roeder, R. S. Buckman, B. H. Gale, S. T. Bunnell, D. L. Eggett, J. Auger, L. A. Felicetti, G. V. Hilderbrand. 2011. Isotopic and gross fecal analysis of American black bear scats. *Ursus* **22**, 133-140. [[CrossRef](#)]
58. Andrew D. Lowther, S. D. Goldsworthy. 2011. Detecting alternate foraging ecotypes in Australian sea lion (*Neophoca cinerea*) colonies using stable isotope analysis. *Marine Mammal Science* **27**:10.1111/mms.2011.27.issue-3, 567-586. [[CrossRef](#)]
59. Scott T. Florin, Laura A. Felicetti, Charles T. Robbins. 2011. The biological basis for understanding and predicting dietary-induced variation in nitrogen and sulphur isotope ratio discrimination. *Functional Ecology* **25**:10.1111/fec.2011.25.issue-3, 519-526. [[CrossRef](#)]
60. Seth G. Cherry, Andrew E. Derocher, Keith A. Hobson, Ian Stirling, Gregory W. Thiemann. 2011. Quantifying dietary pathways of proteins and lipids to tissues of a marine predator. *Journal of Applied Ecology* **48**:10.1111/jpe.2011.48.issue-2, 373-381. [[CrossRef](#)]
61. Mark A. Edwards, Andrew E. Derocher, Keith A. Hobson, Marsha Branigan, John A. Nagy. 2011. Fast carnivores and slow herbivores: differential foraging strategies among grizzly bears in the Canadian Arctic. *Oecologia* **165**, 877-889. [[CrossRef](#)]
62. Ryo Narita, Tsutomu Mano, Reo Yokoyama, Atsushi Takayanagi. 2011. Variation in Maize Consumption by Brown Bears (*Ursus arctos*) in Two Coastal Areas of Hokkaido, Japan. *Mammal Study* **36**, 33-39. [[CrossRef](#)]
63. F. Brischox, X. Bonnet, Y. Cherel, R. Shine. 2011. Isotopic signatures, foraging habitats and trophic relationships between fish and seasnakes on the coral reefs of New Caledonia. *Coral Reefs* **30**, 155-165. [[CrossRef](#)]
64. A. J. Orr, J. Orr, G. R. VanBlaricom, G. R. VanBlaricom, R. L. DeLong, R. L. DeLong, V. H. Cruz-Escalona, V. H. Cruz-Escalona, S. D. Newsome, S. D. Newsome. 2011. Intraspecific comparison of diet of California sea lions (*Zalophus californianus*) assessed using fecal and stable isotope analyses. *Canadian Journal of Zoology* **89**:2, 109-122. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)] [[Supplemental Material](#)]
65. JPY Arnould, Y Cherel, J Gibbens, JG White, CL Littnan. 2011. Stable isotopes reveal inter-annual and inter-individual variation in the diet of female Australian fur seals. *Marine Ecology Progress Series* **422**, 291-302. [[CrossRef](#)]
66. Emanuel Joaquim Victória Monteiro Oliveira, Léa Silvia Sant'Ana, Carlos Ducatti, Juliana Célia Denadai, Cibele Regina Souza Kruliski. 2011. The use of stable isotopes for authentication of gadoid fish species. *European Food Research and Technology* **232**, 97-101. [[CrossRef](#)]
67. Audrey Jaeger, Vincent J. Lecomte, Henri Weimerskirch, Pierre Richard, Yves Cherel. 2010. Seabird satellite tracking validates the use of latitudinal isoscapes to depict predators' foraging areas in the Southern Ocean. *Rapid Communications in Mass Spectrometry* **24**, 3456-3460. [[CrossRef](#)]
68. Sarah Habran, Cathy Debier, Daniel E. Crocker, Dorian S. Houser, Gilles Lepoint, Jean-Marie Bouquegneau, Krishna Das. 2010. Assessment of gestation, lactation and fasting on stable isotope ratios in northern elephant seals (*Mirounga angustirostris*). *Marine Mammal Science* **26**:10.1111/mms.2010.26.issue-4, 880-895. [[CrossRef](#)]
69. D. J. Halley, M. Minagawa, M. Nieminen, E. Gaare. 2010. Diet: tissue stable isotope fractionation of carbon and nitrogen in blood plasma and whole blood of male reindeer *Rangifer tarandus*. *Polar Biology* **33**, 1303-1309. [[CrossRef](#)]
70. Frank J. Wessels, Daniel A. Hahn. 2010. Carbon 13 discrimination during lipid biosynthesis varies with dietary concentration of stable isotopes: implications for stable isotope analyses. *Functional Ecology* **24**, 1017-1022. [[CrossRef](#)]
71. Frédéric Bailleul, Matthieu Authier, Simon Ducatez, Fabien Roquet, Jean-Benoît Charrassin, Yves Cherel, Christophe Guinet. 2010. Looking at the unseen: combining animal bio-logging and stable isotopes to reveal a shift in the ecological niche of a deep diving predator. *Ecography* **33**:10.1111/eco.2010.33.issue-4, 709-719. [[CrossRef](#)]

72. Massimiliano Drago, Luis Cardona, Enrique A. Crespo, Néstor García, Santiago Ameghino, Alex Aguilar. 2010. Change in the foraging strategy of female South American sea lions (Carnivora: Pinnipedia) after parturition. *Scientia Marina* **74**:10.3989/scimar.2010.74n3, 589-598. [[CrossRef](#)]
73. Stephane Caut, Elena Angulo, Franck Courchamp, Jordi Figuerola. 2010. Trophic experiments to estimate isotope discrimination factors. *Journal of Applied Ecology* **47**:10.1111/jpe.2010.47.issue-4, 948-954. [[CrossRef](#)]
74. Karl Auerswald, Maximilian H. O. M. Wittmer, Antoine Zazzo, Rudi Schäufele, Hans Schnyder. 2010. Biases in the analysis of stable isotope discrimination in food webs. *Journal of Applied Ecology* **47**:10.1111/jpe.2010.47.issue-4, 936-941. [[CrossRef](#)]
75. Rebecca C. Redfern, Christine Hamlin, Nancy Beavan Athfield. 2010. Temporal changes in diet: a stable isotope analysis of late Iron Age and Roman Dorset, Britain. *Journal of Archaeological Science* **37**, 1149-1160. [[CrossRef](#)]
76. Massimiliano Drago, Luis Cardona, Alex Aguilar, Enrique A. Crespo, Santiago Ameghino, Néstor García. 2010. Diet of lactating South American sea lions, as inferred from stable isotopes, influences pup growth. *Marine Mammal Science* **26**:10.1111/mms.2010.26.issue-2, 309-323. [[CrossRef](#)]
77. Charles T. Robbins, Laura A. Felicetti, Scott T. Florin. 2010. The impact of protein quality on stable nitrogen isotope ratio discrimination and assimilated diet estimation. *Oecologia* **162**, 571-579. [[CrossRef](#)]
78. Borja Figueirido, Juan A. Pérez-Claros, Vanessa Torregrosa, Alberto Martín-Serra, Paul Palmqvist. 2010. Demythologizing *Arctodus simus*, the 'short-faced' long-legged and predaceous bear that never was. *Journal of Vertebrate Paleontology* **30**, 262-275. [[CrossRef](#)]
79. Layne G. Adams, Sean D. Farley, Craig A. Stricker, Dominic J. Demma, Gretchen H. Roffler, Dennis C. Miller, Robert O. Rye. 2010. Are inland wolf-ungulate systems influenced by marine subsidies of Pacific salmon?. *Ecological Applications* **20**, 251-262. [[CrossRef](#)]
80. Egle Vulla, Keith A. Hobson, Marju Korsten, Malle Leht, Ants-Johannes Martin, Ave Lind, Peep Männil, Harri Valdmann, Urmas Saarma. 2009. Carnivory is Positively Correlated with Latitude among Omnivorous Mammals: Evidence from Brown Bears, Badgers and Pine Martens. *Annales Zoologici Fennici* **46**, 395-415. [[CrossRef](#)]
81. Travis W. Horton, Joel D. Blum, Zhouqing Xie, Michael Hren, C. Page Chamberlain. 2009. Stable isotope food-web analysis and mercury biomagnification in polar bears (*Ursus maritimus*). *Polar Research* **28**:10.1111/por.2009.28.issue-3, 443-454. [[CrossRef](#)]
82. S. Peigne, C. Goillot, M. Germonpre, C. Blondel, O. Bignon, G. Merceron. 2009. Predormancy omnivory in European cave bears evidenced by a dental microwear analysis of *Ursus spelaeus* from Goyet, Belgium. *Proceedings of the National Academy of Sciences* **106**, 15390-15393. [[CrossRef](#)]
83. Naho Mitani, Shozo Mihara, Nobuo Ishii, Hiroko Koike. 2009. Clues to the cause of the Tsushima leopard cat (*Prionailurus bengalensis euptilura*) decline from isotopic measurements in three species of Carnivora. *Ecological Research* **24**, 897-908. [[CrossRef](#)]
84. MARTINA PACHER, ANTHONY J. STUART. 2009. Extinction chronology and palaeobiology of the cave bear (*Ursus spelaeus*). *Boreas* **38**:10.1111/bor.2009.38.issue-2, 189-206. [[CrossRef](#)]
85. Hayley C. Norman, Matt G. Wilmot, Dean T. Thomas, David G. Masters, Dean K. Revell. 2009. Stable carbon isotopes accurately predict diet selection by sheep fed mixtures of C3 annual pastures and saltbush or C4 perennial grasses. *Livestock Science* **121**, 162-172. [[CrossRef](#)]
86. Stéphane Caut, Elena Angulo, Franck Courchamp. 2009. Variation in discrimination factors ( $\Delta 15\text{ N}$  and  $\Delta 13\text{ C}$ ): the effect of diet isotopic values and applications for diet reconstruction. *Journal of Applied Ecology* **46**:10.1111/jpe.2009.46.issue-2, 443-453. [[CrossRef](#)]
87. Eva M. Krümmel, Michael Scheer, Irene Gregory-Eaves, Robie W. Macdonald, Lynda E. Kimpe, John P. Smol, Bruce Finney, Jules M. Blais. 2009. Historical analysis of salmon-derived polychlorinated biphenyls (PCBs) in lake sediments. *Science of The Total Environment* **407**, 1977-1989. [[CrossRef](#)]
88. D. R. Klinka, T. E. Reimchen. 2009. Darkness, Twilight, and Daylight Foraging Success of Bears (*Ursus americanus*) on Salmon in Coastal British Columbia. *Journal of Mammalogy* **90**, 144-149. [[CrossRef](#)]
89. Toru Oi, Hitomi Furusawa. 2008. Nutritional condition and dietary profile of Japanese black bear (*Ursus thibetanus japonicus*) killed in western Japan in autumn 2004. *Mammal Study* **33**, 163-171. [[CrossRef](#)]

90. Simon Ducatez, Sébastien Dalloyau, Pierre Richard, Christophe Guinet, Yves Cherel. 2008. Stable isotopes document winter trophic ecology and maternal investment of adult female southern elephant seals (*Mirounga leonina*) breeding at the Kerguelen Islands. *Marine Biology* **155**, 413-420. [[CrossRef](#)]
91. Heidi Porras-Peters, David Auriolos-Gamboa, Victor Hugo Cruz-Escalona, Paul L. Koch. 2008. Trophic level and overlap of sea lions (*Zalophus californianus*) in the Gulf of California, Mexico. *Marine Mammal Science* **24**:10.1111/mms.2008.24.issue-3, 554-576. [[CrossRef](#)]
92. Stéphane Caut, Elena Angulo, Franck Courchamp. 2008. Caution on isotopic model use for analyses of consumer diet. *Canadian Journal of Zoology* **86**:5, 438-445. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
93. Gregory W. Thiemann. 2008. Using fatty acid signatures to study bear foraging: Technical considerations and future applications. *Ursus* **19**, 59-72. [[CrossRef](#)]
94. Kena Fox-Dobbs, Jennifer A. Leonard, Paul L. Koch. 2008. Pleistocene megafauna from eastern Beringia: Paleocological and paleoenvironmental interpretations of stable carbon and nitrogen isotope and radiocarbon records. *Palaeogeography, Palaeoclimatology, Palaeoecology* **261**, 30-46. [[CrossRef](#)]
95. Stéphane Caut, Elena Angulo, Franck Courchamp. 2008. Discrimination factors ( $\Delta 15\text{ N}$  and  $\Delta 13\text{ C}$ ) in an omnivorous consumer: effect of diet isotopic ratio. *Functional Ecology* **22**, 255-263. [[CrossRef](#)]
96. T. W. Bentzen, E. H. Follmann, S. C. Amstrup, G. S. York, M. J. Wooller, D. C. G. Muir, T. M. O'Hara. 2008. Dietary biomagnification of organochlorine contaminants in Alaskan polar bears. *Canadian Journal of Zoology* **86**:3, 177-191. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
97. Nancy Beavan Athfield, Roger C. Green, Jacqueline Craig, Bruce McFadgen, Simon Bickler. 2008. Influence of marine sources on 14 C ages: Isotopic data from Watom Island, Papua New Guinea inhumations and pig teeth in light of new dietary standards. *Journal of the Royal Society of New Zealand* **38**, 1-23. [[CrossRef](#)]
98. M. P. Richards, M. Pacher, M. Stiller, J. Quiles, M. Hofreiter, S. Constantin, J. Zilhao, E. Trinkaus. 2008. Isotopic evidence for omnivory among European cave bears: Late Pleistocene *Ursus spelaeus* from the Pestera cu Oase, Romania. *Proceedings of the National Academy of Sciences* **105**, 600-604. [[CrossRef](#)]
99. Vicki K. Stegall, Sean D. Farley, Lorrie D. Rea, Kenneth W. Pitcher, Robert O. Rye, Cynthia L. Kester, Craig A. Stricker, Carleton R. Bern. 2008. Discrimination of carbon and nitrogen isotopes from milk to serum and vibrissae in Alaska Steller sea lions (*Eumetopias jubatus*). *Canadian Journal of Zoology* **86**:1, 17-23. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
100. Tuula Sinisalo, Roger I. Jones, Eero Helle, E. Tellervo Valtonen. 2008. Changes in diets of individual Baltic ringed seals (*Phoca hispida botnica*) during their breeding season inferred from stable isotope analysis of multiple tissues. *Marine Mammal Science* **24**:10.1111/mms.2008.24.issue-1, 159-170. [[CrossRef](#)]
101. KERRY CRAWFORD, ROBBIE A. MCDONALD, STUART BEARHOP. 2008. Applications of stable isotope techniques to the ecology of mammals. *Mammal Review* **38**, 87-107. [[CrossRef](#)]
102. J. K. Bump, K. Fox-Dobbs, J. L. Bada, P. L. Koch, R. O. Peterson, J. A. Vucetich. 2007. Stable isotopes, ecological integration and environmental change: wolves record atmospheric carbon isotope trend better than tree rings. *Proceedings of the Royal Society B: Biological Sciences* **274**, 2471-2480. [[CrossRef](#)]
103. MARK D. SCHEUERELL, JONATHAN W. MOORE, DANIEL E. SCHINDLER, CHRIS J. HARVEY. 2007. Varying effects of anadromous sockeye salmon on the trophic ecology of two species of resident salmonids in southwest Alaska. *Freshwater Biology* **52**:10.1111/fwb.2007.52.issue-10, 1944-1956. [[CrossRef](#)]
104. Jeffrey A. Seminoff, Karen A. Bjorndal, Alan B. Bolten. 2007. Stable Carbon and Nitrogen Isotope Discrimination and Turnover in Pond Sliders *Trachemys Scripta*: Insights for Trophic Study of Freshwater Turtles. *Copeia* **2007**, 534-542. [[CrossRef](#)]
105. T. W. Bentzen, E. H. Follmann, S. C. Amstrup, G. S. York, M. J. Wooller, T. M. O'Hara. 2007. Variation in winter diet of southern Beaufort Sea polar bears inferred from stable isotope analysis. *Canadian Journal of Zoology* **85**:5, 596-608. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
106. K. Fox-Dobbs, J. K. Bump, R. O. Peterson, D. L. Fox, P. L. Koch. 2007. Carnivore-specific stable isotope variables and variation in the foraging ecology of modern and ancient

- wolf populations: case studies from Isle Royale, Minnesota, and La Brea. *Canadian Journal of Zoology* **85**:4, 458-471. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)] [[Supplemental Material](#)]
107. Jennifer K. Fortin, Sean D. Farley, Karyn D. Rode, Charles T. Robbins. 2007. Dietary and spatial overlap between sympatric ursids relative to salmon use. *Ursus* **18**, 19-29. [[CrossRef](#)]
  108. Jerrold L. Belant, Knut Kielland, Erich H. Follmann, Layne G. Adams. 2006. INTERSPECIFIC RESOURCE PARTITIONING IN SYMPATRIC URSIDS. *Ecological Applications* **16**, 2333-2343. [[CrossRef](#)]
  109. Ryo Narita, Atsuko Sugimoto, Atsushi Takayanagi. 2006. Animal components in the diet of Japanese black bears *Ursus thibetanus japonicus* in the Kyoto area, Japan. *Wildlife Biology* **12**, 375-384. [[CrossRef](#)]
  110. BRETT P. MURPHY, DAVID M. J. S. BOWMAN. 2006. Kangaroo metabolism does not cause the relationship between bone collagen  $\delta^{15}N$  and water availability. *Functional Ecology* **20**:10.1111/fec.2006.20.issue-6, 1062-1069. [[CrossRef](#)]
  111. Stéphane Caut, Gary W. Roemer, C. Josh Donlan, Franck Courchamp. 2006. COUPLING STABLE ISOTOPES WITH BIOENERGETICS TO ESTIMATE INTERSPECIFIC INTERACTIONS. *Ecological Applications* **16**, 1893-1900. [[CrossRef](#)]
  112. Ryo Narita, Kazuyoshi Sasaki, Katsumi Goda, Naoko Maeda, Atsushi Takayanagi. 2006. Turnover of stable isotopes in Hokkaido brown bear (*Ursus arctos yesoyensis*). *Mammal Study* **31**, 59-63. [[CrossRef](#)]
  113. Katherine W. McFadden, Raymond N. Sambrotto, Rodrigo A. Medellín, Matthew E. Gompper. 2006. FEEDING HABITS OF ENDANGERED PYGMY RACCOONS (*PROCYON PYGMAEUS*) BASED ON STABLE ISOTOPE AND FECAL ANALYSES. *Journal of Mammalogy* **87**, 501-509. [[CrossRef](#)]
  114. Liying Zhao, Donald M. Schell, Michael A. Castellini. 2006. Dietary macronutrients influence  $^{13}C$  and  $^{15}N$  signatures of pinnipeds: Captive feeding studies with harbor seals (*Phoca vitulina*). *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology* **143**, 469-478. [[CrossRef](#)]
  115. Garth Mowat, Douglas C Heard. 2006. Major components of grizzly bear diet across North America. *Canadian Journal of Zoology* **84**:3, 473-489. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]
  116. Hervé Bocherens, Dorothee G. Drucker, Daniel Billiou, Jean-Michel Geneste, Johannes van der Plicht. 2006. Bears and humans in Chauvet Cave (Vallon-Pont-d'Arc, Ardèche, France): Insights from stable isotopes and radiocarbon dating of bone collagen. *Journal of Human Evolution* **50**, 370-376. [[CrossRef](#)]
  117. Donald L. Phillips, Peter M. Eldridge. 2006. Estimating the timing of diet shifts using stable isotopes. *Oecologia* **147**, 195-203. [[CrossRef](#)]
  118. Vincent Balter, Laurent Simon, H el ene Fouillet, Christophe L ecuyer. 2006. Box-modeling of  $^{15}N/^{14}N$  in mammals. *Oecologia* **147**, 212-222. [[CrossRef](#)]
  119. Xianfeng Yi, Yueqin Yang, Xiaoi Zhang. 2006. Modeling trophic positions of the alpine meadow ecosystem combining stable carbon and nitrogen isotope ratios. *Ecological Modelling* **193**, 801-808. [[CrossRef](#)]
  120. James M. Helfield, Robert J. Naiman. 2006. Keystone Interactions: Salmon and Bear in Riparian Forests of Alaska. *Ecosystems* **9**, 167-180. [[CrossRef](#)]
  121. Andrea Ballinger, P. S. Lake. 2006. Energy and nutrient fluxes from rivers and streams into terrestrial food webs. *Marine and Freshwater Research* **57**, 15. [[CrossRef](#)]
  122. B. Sorkin. 2006. Ecomorphology of the giant short-faced bears *Agriotherium* and *Arctodus*. *Historical Biology* **18**, 1-20. [[CrossRef](#)]
  123. Yves Cherel, Keith A. Hobson, Fr ed eric Bailleul, Ren e Groscolas. 2005. NUTRITION, PHYSIOLOGY, AND STABLE ISOTOPES: NEW INFORMATION FROM FASTING AND MOLTING PENGUINS. *Ecology* **86**, 2881-2888. [[CrossRef](#)]
  124. Monika Winder, Daniel E Schindler, Jonathan W Moore, Susan P Johnson, Wendy J Palen. 2005. Do bears facilitate transfer of salmon resources to aquatic macroinvertebrates?. *Canadian Journal of Fisheries and Aquatic Sciences* **62**:10, 2285-2293. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]
  125. M O Hammill, V Lesage, P Carter. 2005. What do harp seals eat? Comparing diet composition from different compartments of the digestive tract with diets estimated from stable-isotope ratios. *Canadian Journal of Zoology* **83**:10, 1365-1372. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]



126. Y. Cherel, K. A. Hobson, H. Weimerskirch. 2005. Using stable isotopes to study resource acquisition and allocation in procellariiform seabirds. *Oecologia* **145**, 533-540. [[CrossRef](#)]
127. Erin J. M. Urton, Keith A. Hobson. 2005. Intrapopulation variation in gray wolf isotope ( $\delta^{15}\text{N}$  and  $\delta^{13}\text{C}$ ) profiles: implications for the ecology of individuals. *Oecologia* **145**, 316-325. [[CrossRef](#)]
128. Charles T. Robbins, Laura A. Felicetti, Matt Sponheimer. 2005. The effect of dietary protein quality on nitrogen isotope discrimination in mammals and birds. *Oecologia* **144**, 534-540. [[CrossRef](#)]
129. F. Dalerum, A. Angerbjörn. 2005. Resolving temporal variation in vertebrate diets using naturally occurring stable isotopes. *Oecologia* **144**, 647-658. [[CrossRef](#)]
130. Seiji Yanai, Kaori Kochi. 2005. Effects of salmon carcasses on experimental stream ecosystems in Hokkaido, Japan. *Ecological Research* **20**, 471-480. [[CrossRef](#)]
131. Owen T. Nevin, Barrie K. Gilbert. 2005. Measuring the cost of risk avoidance in brown bears: Further evidence of positive impacts of ecotourism. *Biological Conservation* **123**, 453-460. [[CrossRef](#)]
132. V.C. Cruz, C. Ducatti, A.C. Pezzato, D.F. Pinheiro, J.R. Sartori, J.C. Gonçaves, A.S. Carrijo. 2005. Influence of diet on assimilation and turnover of  $^{13}\text{C}$  in the tissues of broiler chickens. *British Poultry Science* **46**, 382-389. [[CrossRef](#)]
133. David T.J. Sare, John S. Millar, Frederick J. Longstaffe. 2005. Tracing dietary protein in red-backed voles (*Clethrionomys gapperi*) using stable isotopes of nitrogen and carbon. *Canadian Journal of Zoology* **83**:5, 717-725. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]
134. Rumiko N. Mizukami, Mitsuki Goto, Shigeyuki Izumiya, Hidetake Hayashi, Muneoki Yoh. 2005. Estimation of feeding history by measuring carbon and nitrogen stable isotope ratios in hair of Asiatic black bears. *Ursus* **16**, 93-101. [[CrossRef](#)]
135. R. N. Mizukami, M. Goto, S. Izumiya, M. Yoh, N. Ogura, H. Hayashi. 2005. Temporal diet changes recorded by stable isotopes in Asiatic black bear (*Ursus thibetanus*) hair. *Isotopes in Environmental and Health Studies* **41**, 87-94. [[CrossRef](#)]
136. Garth Mowat, Douglas C. Heard, Dale R. Seip, Kim G. Poole, Gord Stenhouse, David W. Paetkau. 2005. Grizzly *Ursus arctos* and black bear *U. americanus* densities in the interior mountains of North America. *Wildlife Biology* **11**, 31-48. [[CrossRef](#)]
137. David W. Podlesak, Scott R. McWilliams, Kent A. Hatch. 2005. Stable isotopes in breath, blood, feces and feathers can indicate intra-individual changes in the diet of migratory songbirds. *Oecologia* **142**, 501-510. [[CrossRef](#)]
138. Michele Thums, Marcel Klaassen, Ian D. Hume. 2005. Seasonal changes in the diet of the long-nosed bandicoot (*Perameles nasuta*) assessed by analysis of faecal scats and of stable isotopes in blood. *Australian Journal of Zoology* **53**, 87. [[CrossRef](#)]
139. Jianzhu Wang, Guanghui Lin, Jianhui Huang, Xingguo Han. 2004. Applications of stable isotopes to study plant-animal relationships in terrestrial ecosystems. *Chinese Science Bulletin* **49**, 2339-2347. [[CrossRef](#)]
140. Charles T. Robbins, Charles C. Schwartz, Laura A. Felicetti. 2004. Nutritional ecology of ursids: a review of newer methods and management implications. *Ursus* **15**, 161-171. [[CrossRef](#)]
141. Jonathan W. Moore, Daniel E. Schindler. 2004. Nutrient export from freshwater ecosystems by anadromous sockeye salmon (*Oncorhynchus nerka*). *Canadian Journal of Fisheries and Aquatic Sciences* **61**:9, 1582-1589. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]
142. STUART BEARHOP, COLIN E. ADAMS, SUSAN WALDRON, RICHARD A. FULLER, HAZEL MACLEOD. 2004. Determining trophic niche width: a novel approach using stable isotope analysis. *Journal of Animal Ecology* **73**, 1007-1012. [[CrossRef](#)]
143. Hervé Bocherens, Alain Argant, Jacqueline Argant, Daniel Billiou, Evelyne Crégut-Bonnoure, Bernadette Donat-Ayache, Michel Philippe, Michel Thionon. 2004. Diet reconstruction of ancient brown bears (*Ursus arctos*) from Mont Ventoux (France) using bone collagen stable isotope biogeochemistry ( $^{13}\text{C}$ ,  $^{15}\text{N}$ ). *Canadian Journal of Zoology* **82**:4, 576-586. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]
144. Grant V. Hilderbrand, Sean D. Farley, Charles C. Schwartz, Charles T. Robbins. 2004. Importance of salmon to wildlife: Implications for integrated management. *Ursus* **15**, 1-9. [[CrossRef](#)]
145. Danielle K. Greaves, Michael O. Hammill, Jim D. Eddington, Darcy Pettipas, Jason F. Schreer. 2004. GROWTH RATE AND SHEDDING OF VIBRISSAE IN THE GRAY SEAL, HALICHOERUS GRYPUS:

A CAUTIONARY NOTE FOR STABLE ISOTOPE DIET ANALYSIS. *Marine Mammal Science* 20:10.1111/mms.2004.20.issue-2, 296-304. [[CrossRef](#)]

146. S M Gende, T P Quinn. 2004. The relative importance of prey density and social dominance in determining energy intake by bears feeding on Pacific salmon. *Canadian Journal of Zoology* 82:1, 75-85. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]
147. Kaneaki Edo, Kenichi Suzuki. 2003. Preferable summering habitat of returning adult masu salmon in the natal stream. *Ecological Research* 18:10.1111/ere.2003.18.issue-6, 783-791. [[CrossRef](#)]
148. Shana R Lavin, Timothy R Van Deelen, Patrick W Brown, Richard E Warner, Stanley H Ambrose. 2003. Prey use by red foxes (*Vulpes vulpes*) in urban and rural areas of Illinois. *Canadian Journal of Zoology* 81:6, 1070-1082. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]
149. Laura A Felicetti, Charles C Schwartz, Robert O Rye, Mark A Haroldson, Kerry A Gunther, Donald L Phillips, Charles T Robbins. 2003. Use of sulfur and nitrogen stable isotopes to determine the importance of whitebark pine nuts to Yellowstone grizzly bears. *Canadian Journal of Zoology* 81:5, 763-770. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]
150. H. Bocherens, D. Drucker. 2003. Trophic level isotopic enrichment of carbon and nitrogen in bone collagen: case studies from recent and ancient terrestrial ecosystems. *International Journal of Osteoarchaeology* 13:10.1002/oa.v13:1/2, 46-53. [[CrossRef](#)]
151. Deanne C. Drake, Robert J. Naiman, James M. Helfield. 2002. RECONSTRUCTING SALMON ABUNDANCE IN RIVERS: AN INITIAL DENDROCHRONOLOGICAL EVALUATION. *Ecology* 83, 2971-2977. [[CrossRef](#)]
152. C T Darimont, T E Reimchen. 2002. Intra-hair stable isotope analysis implies seasonal shift to salmon in gray wolf diet. *Canadian Journal of Zoology* 80:9, 1638-1642. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]
153. Charles T. Robbins, Grant V. Hilderbrand, Sean D. Farley. 2002. Incorporating concentration dependence in stable isotope mixing models: a response to Phillips and Koch (2002). *Oecologia* 133, 10-13. [[CrossRef](#)]
154. Paul L. Koch, Donald L. Phillips. 2002. Incorporating concentration dependence in stable isotope mixing models: a reply to Robbins, Hilderbrand and Farley (2002). *Oecologia* 133, 14-18. [[CrossRef](#)]
155. David J. Mattson, Troy Merrill. 2002. Extirpations of Grizzly Bears in the Contiguous United States, 1850 -2000. *Conservation Biology* 16, 1123-1136. [[CrossRef](#)]
156. G. R. Graves, C. S. Romanek, A. Rodriguez Navarro. 2002. Stable isotope signature of philopatry and dispersal in a migratory songbird. *Proceedings of the National Academy of Sciences* 99, 8096-8100. [[CrossRef](#)]
157. Carolyn M Kurlle. 2002. Stable-isotope ratios of blood components from captive northern fur seals (*Callorhinus ursinus*) and their diet: applications for studying the foraging ecology of wild otariids. *Canadian Journal of Zoology* 80:5, 902-909. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]
158. VERONIQUE Lesage, Mike O. Hammill, Kit M. Kovacs. 2002. DIET-TISSUE FRACTIONATION OF STABLE CARBON AND NITROGEN ISOTOPES IN PHOCID SEALS. *Marine Mammal Science* 18:10.1111/mms.2002.18.issue-1, 182-193. [[CrossRef](#)]
159. Pablo Sabat, Carlos Martinez del Rio. 2002. Inter- and intraspecific variation in the use of marine food resources by three Cinclodes (Furnariidae, Aves) species: carbon isotopes and osmoregulatory physiology. *Zoology* 105, 247-256. [[CrossRef](#)]
160. S J Iverson, J E McDonald, Jr., L K Smith. 2001. Changes in the diet of free-ranging black bears in years of contrasting food availability revealed through milk fatty acids. *Canadian Journal of Zoology* 79:12, 2268-2279. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]
161. Jose M. Fedriani, Michael H. Kohn. 2001. Genotyping faeces links individuals to their diet. *Ecology Letters* 4, 477-483. [[CrossRef](#)]
162. C. L. Kester, R. O. Rye, C. A. Johnson, Ch. Schwartz, Ch. Holmes. 2001. On-line Sulfur Isotope Analysis of Organic Material by Direct Combustion: Preliminary Results and Potential Applications. *Isotopes in Environmental and Health Studies* 37, 53-65. [[CrossRef](#)]
163. M. Ben-David, D. M. Schell. 2001. Mixing models in analyses of diet using multiple stable isotopes: a response. *Oecologia* 127, 180-184. [[CrossRef](#)]
164. S C Polischuk, K A Hobson, M A Ramsay. 2001. Use of stable-carbon and -nitrogen isotopes to assess weaning and fasting in female polar bears and their cubs. *Canadian Journal of Zoology* 79:3, 499-511. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]

165. Serge Larivière. 2001. *Ursus americanus*. *Mammalian Species* **647**, 1-11. [[CrossRef](#)]
166. M C Drever, L K Blight, K A Hobson, D F Bertram. 2000. Predation on seabird eggs by Keen's mice (*Peromyscus keeni*): using stable isotopes to decipher the diet of a terrestrial omnivore on a remote offshore island. *Canadian Journal of Zoology* **78**:11, 2010-2018. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]
167. Keith A Hobson, Bruce N McLellan, John G Woods. 2000. Using stable carbon ( $\delta^{13}\text{C}$ ) and nitrogen ( $\delta^{15}\text{N}$ ) isotopes to infer trophic relationships among black and grizzly bears in the upper Columbia River basin, British Columbia. *Canadian Journal of Zoology* **78**:8, 1332-1339. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]
168. James D Roth, Keith A Hobson. 2000. Stable carbon and nitrogen isotopic fractionation between diet and tissue of captive red fox: implications for dietary reconstruction. *Canadian Journal of Zoology* **78**:5, 848-852. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]
169. T E Reimchen. 2000. Some ecological and evolutionary aspects of bear-salmon interactions in coastal British Columbia. *Canadian Journal of Zoology* **78**:3, 448-457. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]
170. Jeffrey F Kelly. 2000. Stable isotopes of carbon and nitrogen in the study of avian and mammalian trophic ecology. *Canadian Journal of Zoology* **78**:1, 1-27. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]
171. G V Hilderbrand, S G Jenkins, C C Schwartz, T A Hanley, C T Robbins. 1999. Effect of seasonal differences in dietary meat intake on changes in body mass and composition in wild and captive brown bears. *Canadian Journal of Zoology* **77**:10, 1623-1630. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]
172. R R Doucett, R K Booth, G Power, R S McKinley. 1999. Effects of the spawning migration on the nutritional status of anadromous Atlantic salmon (*Salmo salar*): insights from stable-isotope analysis. *Canadian Journal of Fisheries and Aquatic Sciences* **56**:11, 2172-2180. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]
173. Sergine Ponsard, Pierre Averbuch. 1999. Should growing and adult animals fed on the same diet show different  $\delta^{15}\text{N}$  values?. *Rapid Communications in Mass Spectrometry* **13**:10.1002/(SICI)1097-0231(19990715)13:13<1305::AID-RM1305>3.0.CO;2-C, 1305-1310. [[CrossRef](#)]
174. G V Hilderbrand, C C Schwartz, C T Robbins, M E Jacoby, T A Hanley, S M Arthur, C Servheen. 1999. The importance of meat, particularly salmon, to body size, population productivity, and conservation of North American brown bears. *Canadian Journal of Zoology* **77**:1, 132-138. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]
175. Troy Merrill, David J Mattson, R.Gerald Wright, Howard B Quigley. 1999. Defining landscapes suitable for restoration of grizzly bears *Ursus arctos* in Idaho. *Biological Conservation* **87**, 231-248. [[CrossRef](#)]
176. Hervé Bocherens. 1998. Comment: Use of stable isotopes to determine diets of living and extinct bears. *Canadian Journal of Zoology* **76**:12, 2299-2300. [[Citation](#)] [[PDF](#)] [[PDF Plus](#)]
177. G V Hilderbrand, C T Robbins, S D Farley. 1998. Response: Use of stable isotopes to determine diets of living and extinct bears. *Canadian Journal of Zoology* **76**:12, 2301-2303. [[Citation](#)] [[PDF](#)] [[PDF Plus](#)]
178. Mary F. Willson, Scott M. Gende, Brian H. Marston. 1998. Fishes and the Forest. *BioScience* **48**, 455-462. [[CrossRef](#)]
179. Leonard Z. Gannes, Carlos Martínez del Rio, Paul Koch. 1998. Natural Abundance Variations in Stable Isotopes and their Potential Uses in Animal Physiological Ecology. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology* **119**, 725-737. [[CrossRef](#)]
180. M. Ben-David, T. A. Hanley, D. R. Klein, D. M. Schell. 1997. Seasonal changes in diets of coastal and riverine mink: the role of spawning Pacific salmon. *Canadian Journal of Zoology* **75**:5, 803-811. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]
181. Rebecca Spindler, Joanna Wiszniewski, David Slip The potential role of zoos in climate change research and mitigation 107-113. [[CrossRef](#)]