

Left Generalized Derivations on Prime Γ -Rings

C. Jaya Subba Reddy¹, K. Nagesh² and A. Sivakameshwara Kumar³

¹Department of Mathematics, Sri Venkateswara University
Tirupati-517502, India. E-mail: cjsreddysvu@gmail.com

²Department of Mathematics, Rayalaseema University
Kurnool, Andhrapradesh, India. E-mail: nagesh.kunda100@gmail.com

³Department of Mathematics, Rayalaseema University
Kurnool, Andhrapradesh, India. E-mail: kamesh1069@gmail.com

Received 6 December 2017; accepted 26 December 2017

Abstract. Let M be a prime Γ -ring with 2-torsion free, I a nonzero ideal of M and $f: M \rightarrow M$ a left generalized derivation of M , with associated nonzero derivation d on M . If $f(x) \in Z(M)$ for all $x \in I$, then M is a commutative Γ -ring.

Keywords: Gamma ring, prime gamma ring, derivation, generalized derivation, left generalized derivation, commutators.

AMS Mathematics Subject Classification (2010): 06F25

1. Introduction

The notion of Γ -ring was first introduction by Nobusawa [9] and also shown that Γ -ring, more general than rings. Barnes [1] slightly weakened the conditions in the definitions of a Γ -rings in the sense of Nobusawa. After the study of Γ -rings by Nobusawa [9] and Barnes [1], many researchers have a done lot of work and have obtained some generalizations of the corresponding results in ring theory [6][8]. Barnes [1] and kyuno [8] studied the structure of Γ -ring and obtained various generalizations of the corresponding results of ring theory. Hvala [4] introduced the concept of Generalized derivations in rings. Dey, Paul and Rakhimov [3] discussed some properties of Generalized derivations in semiprime gamma rings Bresar [2] studied on the distance of the composition of two derivations to the generalized derivations. Jaya Subba Reddy. et al. [5] studied centralizing and commuting left generalized derivation on prime ring is commutative. Jaya Subba Reddy et al. [12] studied some results of symmetric reverse bi-derivations on prime rings, Ozturk et al. [10] studied on derivations of prime gamma rings. Khan et al. [6,7] studied on derivations and generalized derivations on prime Γ -rings is a commutative. In this paper we extended some results on left generalized derivations on prime Γ -ring is a commutative.

2. Preliminaries

If M and Γ are additive abelian groups and there exists a mapping $M \times \Gamma \times M \rightarrow M$ which satisfies the following conditions:

For all $a, b \in M$ and $\alpha, \beta \in \Gamma$,

C. Jaya Subba Reddy, K. Nagesh and A. Sivakameshwara Kumar

- (i) (a, β, b) , denoted by $a\beta b$, is an element of M
- (ii) $(a + b)\beta c = a\beta c + b\beta c$, $a(\alpha + \beta)b = a\alpha b + a\beta b$, $a\beta(b + c) = a\beta b + a\beta c$
- (iii) $(aab)\beta c = a\alpha(b\beta c)$

then M is called a Γ -ring [1]. It is known that from (i), (iii) the following follows:

$$0\beta b = a0b = a\beta 0 = 0 \quad (\text{A})$$

for all a and b in M and all β in Γ [1].

Every ring is a Γ -ring with $M = \Gamma$. However a Γ -ring need not be a ring. Let M be a Γ -ring, then M is called a prime Γ -ring, if $a\Gamma M\Gamma b = 0$ implies $a = 0$ or $b = 0$, for all $a, b \in M$ and M is called a semiprime Γ -ring, if $a\Gamma M\Gamma a = 0$ implies $a = 0$, for all $a \in M$. Every prime Γ -ring is obviously semiprime. If M is a Γ -ring, then M is said to be 2-torsion free if $2x = 0$ implies $x = 0$, for all $x \in M$. An additive subgroup I of M is called a left (right) ideal of M if $M\Gamma I \subseteq I$ ($I\Gamma M \subseteq I$). If I is both left and right ideal of M , then we say I is an ideal of M . Moreover, the set $Z(M) = \{x \in M : x\beta y = y\beta x \ \forall \beta \in \Gamma, y \in M\}$ is called the centre of the Γ -ring M . We shall write $[x, y]_\beta = x\beta y - y\beta x$, for all $x, y \in M$ and $\beta \in \Gamma$. We shall make use of the basic commutator identities: $[x\beta y, z]_\alpha = [x, z]_\alpha \beta y + x\beta [y, z]_\alpha$ and $[x, y\beta z]_\alpha = [x, y]_\alpha \beta z + y\beta [x, z]_\alpha$, for all $x, y, z \in M$ and $\alpha, \beta \in \Gamma$. If Γ -ring satisfies the assumption (B) $aab\beta c = a\beta bac$, for all $a, b, c \in M$ and $\alpha, \beta \in \Gamma$. Let M be a Γ -ring. An additive mapping $d: M \rightarrow M$ is called a derivation on M if $d(x\gamma y) = d(x)\gamma y + x\gamma d(y)$ holds for all $x, y \in M$ and $\gamma \in \Gamma$. An additive mapping $f: M \rightarrow M$ is called a generalized derivation if there exists a derivation $d: M \rightarrow M$ such that $f(x\gamma y) = f(x)\gamma y + x\gamma d(y)$ holds for all $x, y \in M$ and $\gamma \in \Gamma$. An additive mapping $f: M \rightarrow M$ is called a left generalized derivation if there exists a derivation $d: M \rightarrow M$ such that $f(x\gamma y) = x\gamma f(y) + d(x)\gamma y$ holds for all $x, y \in M$ and $\gamma \in \Gamma$. A derivation of the form $x \rightarrow aax + xab$ where a, b are fixed elements of M and $\alpha \in \Gamma$ is called generalized inner derivation. An additive mapping $T: M \rightarrow M$ is called a left (right) centralizer if $T(x\alpha y) = T(x)\alpha y$ ($T(x\alpha y) = x\alpha T(y)$) for all $x, y \in M$ and $\alpha \in \Gamma$.

Lemma 2.1. Let M be a prime Γ -ring with 2-torsion free and I a nonzero ideal of M . Let $f: M \rightarrow M$ be a left generalized derivation of M , associated with derivation d . If $f(y) = 0$, for all $y \in I$, then $f = 0$.

Proof: For all $x, y \in I$ and $\beta \in \Gamma$, $f(x\beta y) = 0$. That is, $x\beta f(y) + d(x)\beta y = 0$, which implies $d(x)\beta y = 0$. Let $z \in M, \alpha \in \Gamma$. The last relation along with (A) gives, $d(x)\beta z\alpha y = 0$. Since M is prime Γ -ring and I is a nonzero ideal, so $d(x) = 0$, for all $x \in I$. Hence, by hypothesis, $f(x\beta r) = 0$, for all $x \in I$, and $\beta \in \Gamma$, and $r \in M$. That is, $x\beta f(r) + d(x)\beta r = 0$, which gives $x\beta f(r) = 0$. Let $w \in M, \gamma \in \Gamma$. The last relation along with (A), implies $x\gamma w\beta f(r) = 0$. Since I is nonzero and primeness of M , gives $f = 0$.

Lemma 2.2. Let I be a nonzero ideal of a prime Γ -ring M , $a \in M$ and $f \neq 0$ is a left generalized derivation of M , with associated nonzero derivation d , then

- (i) If $f(y)\beta a = 0$ for all $y \in I$ and $\beta \in \Gamma$, then $a = 0$,
- (ii) If $a\beta f(y) = 0$ for all $y \in I$ and $\beta \in \Gamma$, then $a = 0$.

Proof:

Left Generalized Derivations on Prime Γ -Rings

- (i) For any $y \in I, r \in M$ and $\alpha, \beta \in \Gamma, f(r\alpha y)\beta a = 0$. That is,
 $r\alpha f(y)\beta a + d(r)\alpha y\beta a = 0$ Which implies, $d(r)\alpha y\beta a = 0$. Since I is a nonzero ideal of M and $d \neq 0$, we get $a = 0$.
- (ii) Proof is similar to (i).

3. Main results

Theorem 3.1. Let M be a prime Γ -ring with 2-torsion free and I a nonzero ideal of M . Let $f: M \rightarrow M$ be a left generalized derivation of M , with associated nonzero derivation d on M . If $f(x) \in Z(M)$ for all $x \in I$, then M is a commutative Γ -ring.

Proof: Using hypothesis, we have $[f(x\beta y), x]_\alpha = 0$, for all $x, y \in I, \alpha, \beta \in \Gamma$, which gives $[x\beta f(y) + d(x)\beta y, x]_\alpha = 0$

$$x\beta[f(y), x]_\alpha + [x, x]_\alpha\beta f(y) + d(x)\beta[y, x]_\alpha + [d(x), x]_\alpha\beta y = 0$$

Using hypothesis, we get

$$d(x)\beta[y, x]_\alpha + [d(x), x]_\alpha\beta y = 0$$

$$d(x)\beta y\alpha x - d(x)\beta x\alpha y + d(x)\alpha x\beta y - xad(x)\beta y = 0$$

Using (B), from the last equation we get

$$d(x)\beta y\alpha x - d(x)\alpha x\beta y + d(x)\alpha x\beta y - xad(x)\beta y = 0$$

$$d(x)\beta y\alpha x - xad(x)\beta y = 0, \text{ for all } y \in I, \alpha, \beta \in \Gamma. \quad (1)$$

Let $z \in I$. Replacing y by $z\beta y$ in equation (1), we get

$$d(x)\beta z\beta y\alpha x - xad(x)\beta z\beta y = 0$$

Which along with equation (1) and (B) gives ,

$$d(x)\beta z\beta y\alpha x - d(x)\beta z\beta x\alpha y = 0$$

$$d(x)\beta z\beta[y, x]_\alpha = 0, \text{ for all } x, y \in I \text{ and } \alpha, \beta \in \Gamma.$$

Since I is a nonzero ideal of M and $d \neq 0$, therefore M is a commutative Γ -ring.

Theorem 3.2. Let M be a prime Γ -ring with 2-torsion free and I a nonzero ideal of M . Let $f: M \rightarrow M$ be a generalized derivation and left generalized derivation of M , with associated derivation d on M . If $a \in M$ and $[f(x), a]_\alpha = 0$, for all $x \in I, \alpha \in \Gamma$, then either $a \in Z(M)$ or $d(a) = 0$.

Proof: Using hypothesis, we have

$$[f(x\beta y), a]_\alpha = 0, \text{ for any } x \in M, y \in I \text{ and } \alpha, \beta \in \Gamma.$$

This gives $[x\beta d(y) + f(x)\beta y, a]_\alpha = 0$

$$[x\beta d(y), a]_\alpha + [f(x)\beta y, a]_\alpha = 0$$

The last equation gives

$$x\beta[d(y), a]_\alpha + [x, a]_\alpha\beta d(y) + f(x)\beta[y, a]_\alpha + [f(x), a]_\alpha\beta y = 0$$

Using hypothesis, from the last equation we get

$$x\beta[d(y), a]_\alpha + [x, a]_\alpha\beta d(y) + f(x)\beta[y, a]_\alpha = 0$$

$$x\beta d(y)aa - x\beta aad(y) + xaa\beta d(y) - aax\beta d(y) + f(x)\beta yaa - f(x)\beta aay = 0$$

Using (B), from the last equation we get

$$x\beta d(y)aa - xaa\beta d(y) + xaa\beta d(y) - aax\beta d(y) + f(x)\beta yaa - f(x)\beta aay = 0$$

$$x\beta d(y)aa - aax\beta d(y) + f(x)\beta yaa - f(x)\beta aay = 0 \quad (2)$$

Let $z \in M$. Replacing x by $z\gamma x$ in equation (2), we get

$$z\gamma x\beta d(y)aa - aaz\gamma x\beta d(y) + f(z\gamma x)\beta yaa - f(z\gamma x)\beta aay = 0$$

$$z\gamma x\beta d(y)aa - aaz\gamma x\beta d(y) + z\gamma(f(x)\beta yaa - f(x)\beta aay) + d(z)\gamma x\beta(yaa - aay)$$

$$= 0$$

Using equation (2), from the last equation we get

C. Jaya Subba Reddy, K. Nagesh and A. Sivakameshwara Kumar

$$z\gamma x\beta d(y)\alpha\alpha - a\alpha z\gamma x\beta d(y) + z\gamma(a\alpha x\beta d(y) - x\beta d(y)\alpha\alpha) + d(z)\gamma x\beta[y, a]_\alpha = 0$$

$$z\gamma x\beta d(y)\alpha\alpha - a\alpha z\gamma x\beta d(y) + z\gamma a\alpha x\beta d(y) - z\gamma x\beta d(y)\alpha\alpha + d(z)\gamma x\beta[y, a]_\alpha = 0$$

Using (B), from the last equation we get

$$-a\alpha z\gamma x\beta d(y) + z\alpha a\gamma x\beta d(y) + d(z)\gamma x\beta[y, a]_\alpha = 0$$

$$[z, a]_\alpha \gamma x\beta d(y) + d(z)\gamma x\beta[y, a]_\alpha = 0$$

Replacing y by a from the last equation, we get

$$[z, a]_\alpha \gamma x\beta d(a) + d(z)\gamma x\beta[a, a]_\alpha = 0$$

$$[z, a]_\alpha \gamma x\beta d(a) = 0, \text{ for all } x \in I, z \in M \text{ and } \alpha, \beta, \gamma \in \Gamma.$$

Since I is nonzero ideal of prime Γ -ring M , therefore either $d(a) = 0$ or $a \in Z(M)$.

Corollary 3.2.1. Let M be a prime Γ -ring with 2-torsion free and I a nonzero ideal of M . Let $f: M \rightarrow M$ be a left generalized derivation of M , with associated derivation d on M . If $[f(x), f(y)]_\beta = 0$, for all $x, y \in I, \beta \in \Gamma$, then M is a commutative Γ -ring.

Proof: Using Theorem 3.2, we have $f(I) \subset Z(M)$, we get the corollary 3.2.1 proof.

Theorem 3.3. Let M be a prime Γ -ring with 2-torsion free and I a nonzero ideal of M . Let $f: M \rightarrow M$ be a left generalized derivation of M , with associated derivation d on M . If $f(x\beta y) = f(x)\beta f(y)$, for all $x, y \in I, \beta \in \Gamma$, then $d = 0$.

Proof: $f(x\beta y) = x\beta f(y) + d(x)\beta y$, for any $x, y \in I, \beta \in \Gamma$.

$$f(x)\beta f(y) = x\beta f(y) + d(x)\beta y \quad (3)$$

Let $w \in I, \gamma \in \Gamma$. Then replacing y by $w\gamma y$ in equation (3), we get

$$f(x)\beta f(w\gamma y) = x\beta f(w\gamma y) + d(x)\beta w\gamma y$$

$$d(x)\beta w\gamma (f(y) - y) = 0, \text{ for all } x, y \in I, \text{ and } \gamma, \beta \in \Gamma.$$

Since I is a nonzero ideal of the prime Γ -ring M , therefore either $f(y) - y = 0$ for all $y \in I$ or $d(x) = 0$ for all $x \in I$. If $f(y) - y = 0$, then $f(y) = y$ for all $y \in I$. Replacing y by $y\beta x$ in the last equation, we get $f(y\beta x) = y\beta x$, which implies $y\beta f(x) + d(y)\beta x = y\beta x$, which gives $y\beta x + d(y)\beta x = y\beta x$. That is $d(y)\beta x = 0$, for all $x, y \in I, \beta \in \Gamma$. Thus $d(y) = 0$ for all $y \in I$ for both cases. So $d = 0$.

Theorem 3.4. Let M be a prime Γ -ring with 2-torsion free and I a nonzero ideal of M . Let $f: M \rightarrow M$ be a left generalized derivation of M , with associated derivation d on M . If $f(x\beta y) = f(y)\beta f(x)$, for all $x, y \in I, \beta \in \Gamma$, then $d = 0$.

Proof: $f(x\beta y) = x\beta f(y) + d(x)\beta y$, for all $x, y \in I, \beta \in \Gamma$.

$$f(y)\beta f(x) = x\beta f(y) + d(x)\beta y \quad (4)$$

Let $x \in I, \gamma \in \Gamma$. Replacing y by $x\gamma y$ in equation (4), we get

$$f(x\gamma y)\beta f(x) = x\beta f(x\gamma y) + d(x)\beta x\gamma y$$

$$x\gamma f(y)\beta f(x) + d(x)\gamma y\beta f(x) = x\beta f(y)\gamma f(x) + d(x)\beta x\gamma y$$

Using (B), from the last equation we get

$$x\beta f(y)\gamma f(x) + d(x)\gamma y\beta f(x) = x\beta f(y)\gamma f(x) + d(x)\beta x\gamma y$$

$$d(x)\gamma y\beta f(x) = d(x)\beta x\gamma y \quad (5)$$

Let $w \in I, \alpha \in \Gamma$. Then replacing y by $y\alpha w$, we get

$$d(x)\gamma y\alpha w\beta f(x) = d(x)\beta x\gamma y\alpha w$$

Using equation (5) in above equation, we get

$$d(x)\gamma y\alpha w\beta f(x) = d(x)\gamma y\beta f(x)\alpha w$$

Using (B), from the last equation we get

Left Generalized Derivations on Prime Γ -Rings

$$d(x)\gamma y\alpha f(x)\beta w - d(x)\gamma y\alpha w\beta f(x) = 0$$

$$d(x)\gamma y\alpha [f(x), w]_{\beta} = 0$$

Since I is a nonzero ideal of the prime Γ -ring M . Therefore either $d(x) = 0$ for all $x \in I$ or $[f(x), w]_{\beta} = 0$ for all $x, w \in I$ and $\beta \in \Gamma$. Let $A = \{x \in I: d(x) = 0\}$ and $B = \{x \in I: [f(x), w]_{\beta} = 0, \forall w \in I\}$. Obviously A and B are additive subgroups of I . Moreover I is the set theoretic union of A and B . But a group cannot be set theoretic union of two proper sub groups. Hence either $A = I$ or $B = I$. If $A = I$, we have $d(R) = 0$, which completes the proof. If $B = I$, then $0 = [f(x), w]_{\beta} = w\alpha[f(x), r]_{\beta}$ for all $x, w \in I$, $r \in M$ and $\alpha, \beta \in \Gamma$. Thus, we obtain $f(I) \subset Z(M)$, Using Theorem 3.2, we get $d = 0$.

Acknowledgements. We are thankful to the reviewers for their comments to improve the presentation of the paper.

REFERENCES

1. W.E.Barnes, On the Γ -rings of Nobusawa, *Pacific J. Math.*, 18 (3) (1966) 411-422.
2. M.Bresar, On the distance of the composition of two derivations to the generalized Derivations, *Glasgow Math. J.*, 33 (1) (1991) 89-93.
3. K.K.Dey, A.C.Paul and I.S.Rakhimov, Generalized derivations in semiprime gamma rings, *IJMMS*, 2012.
4. B.Hvala, Generalized derivations in rings, *Comm. Algebra*, 26 (4) (1998) 1147-1166.
5. C.Jaya Subba Reddy, S.Mallikarjuna Rao and V.Vijaya Kumar, Centralizing and commuting left generalized derivations on prime rings, *Bulletin of Mathematical Science and Applications*, 11 (2015) 1-3.
6. A.R.Khan, M.Anwar Chaudhry and Imran Javaid, Generalized derivations on prime Γ -rings, *World Applied Sciences Journal*, 23 (12) (2013) 59-64.
7. A.R.Khan, I.Javaid and M.Anwar Chaudhry: Derivations on semiprime Γ -rings, *Utilitas Mathematica*, 90 (2013) 171-185.
8. S.Kyuno, On prime gamma rings, *Pacific J. Math.*, 75 (1) (1978) 185-190.
9. N.Nobusawa, On the generalization of the ring theory, *Osaka J. Math.*, 1 (1964) 81-89.
10. M.A.Ozturk, Y.B.Jun and K.H.Kim, On derivations of prime gamma rings. *Turk. J. Math.*, 26 (2002) 317-327.
11. C.Yilmaz, and M.A.Ozturk, On Jordan generalized derivations in gamma rings, *Hacet. J. Math. Stat.*, 33 (2004) 11-14.
12. C. Jaya Subba Reddy, A. Sivakameshwara Kumar and B. Ramoorthy Reddy, Results of Symmetric Reverse bi-derivations on Prime Rings, *Ann. Pure and Applied Math*, 16(1)(2018) 1-6.
13. A.K.Kadhim, H.Sulaiman and A.R.H.Majeed, Gamma * derivation pair and jordan gamma*- derivation pair on gamma-ring M with involution, *Ann. Pure and Applied Math*, 10(2) (2015) 169-177.