

# Abrupt climate changes recorded in loess sequences

Denis-Didier Rousseau<sup>1,2</sup> and Adriana Sima<sup>1</sup>


The glacial-interglacial and orbital-timescale oscillations imprinted in loess records have been studied for a long time. Researchers have recently started to investigate millennial and sub-millennial variations by means of high-resolution field studies and data analyses combined with modeling experiments.

The terrestrial sediments known as “loess” represent an important archive of paleoclimatic variability. Mineral material mobilized by wind at the ground surface was transported over distances from a few hundred to thousands of kilometers and deposited back on the ground where it underwent a gradual transformation into loess sediment. The largest loess deposits are found in the Northern Hemisphere, primarily because the continental surface is much larger than in the Southern Hemisphere. Depending on their location on the globe (Fig. 1), the deposits have formed under the influence of different climate factors, and contain more or less detailed records of regional climate change. Here we review the current knowledge about the three main loess regions of the Northern Hemisphere. The many underlying references are provided in an extensive online archive of references, listed by regions.

## European loess

An almost continuous loess band stretches, along approximately 50°N, from western Europe to the Dnieper valley in Ukraine. In this band, loess sequences mainly record the North-Atlantic millennial-scale climate changes of the last glaciation (between ca. 110-15 ka BP): the Dansgaard-Oeschger (DO) events and Heinrich stadials (Rousseau et al. 2007, 2011). The DO events correspond in the Northern Hemisphere to abrupt warmings, of about 10°C in Greenland within 50-100 years, and are followed by a more gradual return to cold (stadial) conditions. The Heinrich stadials represent particularly cold climate episodes caused by massive iceberg discharges from the Northern Hemisphere ice sheets known as Heinrich events.

Multidisciplinary loess studies and detailed chronological analyses indicate a strong correlation between these climate variations and European loess sedimentation. In the 50°N European loess band, the alternating warm and cold episodes are clearly imprinted in the stratigraphy as a succession of doublets of paleosol-loess units, in particular during marine isotope stages 3 and 2 (between approx. 60-15 ka BP). The soils were formed during the relatively warm North-Atlantic phases (Greenland interstadials) associated with DO events, when the eolian sedimentation was reduced or even absent. The degree



**Figure 1:** The Northern Hemisphere ice-sheets (in light blue) and glaciers (dark blue dots) at the Last Glacial Maximum (map compiled by Jürgen Ehlers, available at [www.qpg.geog.cam.ac.uk/lgmextent.html](http://www.qpg.geog.cam.ac.uk/lgmextent.html)), the schematic location of the polar jet stream (blue arrows), wind patterns (black arrows) generated by the presence of the ice sheets (Kutzbach 1987), and (labeled) the loess regions discussed in the text.

of development of each soil depended on the duration of the corresponding warm episode. The loess units correspond to the cold North-Atlantic phases (Greenland stadials and Heinrich stadials), when the dust cycle was very active and the sedimentation rates were high. Other loess deposits can be found in Europe at lower latitudes, but the millennial climate variations represented by the alternation of paleosols and loess units are absent in the stratigraphy, suggesting drier conditions than in the 50°N loess band.

Recent studies have employed numerical modeling to investigate the mechanistic link between the North-Atlantic millennial climate

signal and the loess sedimentation variations in Europe around 50°N (Sima et al. 2009, 2013). These studies focused on variations in dust mobilization, reasonably assuming that the deposition fluxes strongly depended on the emission fluxes. An important aspect of modeling the formation of loess deposits in Europe is to identify the geographical areas that were potentially subject to deflation during glacial times, but are not anymore under present-day climate conditions.

For Western Europe, data-based studies have shown that the continental shelf that emerged during glacials (when sea-level dropped), especially in the English Channel and the

North Sea, used to be a strong dust deflation area. Transported from there by westerly winds, fine material must have reached loess deposits located many hundreds of kilometers away, together with coarser material from nearer sources, such as large river valleys or periglacial outwash plains. Modeling results (Sima et al. 2009) point to vegetation changes in response to the millennial climate variations as a key factor in modulating dust emission, and hence ultimately also in controlling loess sedimentation (Fig. 2). They also show the strong seasonality of the dust cycle. Dust emission was mainly active in spring-time when the snow cover had melted, the topsoil layer had begun to thaw, the surface winds were still strong (even though weaker than in winter), and the vegetation had not yet developed enough to completely protect the surface from wind erosion. The colder the climate, the later the emission season started, and the later it ended. Numerical experiments generated about one month of delay for a given region between the warmest ("Greenland interstadial") and the coldest ("H stadial") simulated climate state.

For Eastern Europe, previous data-based studies only suggested the general direction in which the source areas should have been located with respect to the investigated loess deposits. Numerical modeling results (Sima et al. 2013) now show the likely dust source areas (Fig. 2). They also suggest that Heinrich stadials, even though generally colder and drier than the other stadials, were not necessarily dustier. Dust storms appeared to be generally less frequent during Heinrich stadials, but individual events could become stronger than during non-Heinrich stadials. These modeling results are in agreement with findings of relatively coarser-grained layers in loess sequences interpreted as Heinrich-Stadial signature.

### Asian loess

In the eastern part of Eurasia, the sedimentation in the Chinese Loess Plateau (Fig. 1), which commenced about 22 Ma ago, strongly depended on fluctuations of the Southeast Asian monsoon. Initial investigations on loess sequences from this area have identified orbital timescale variations related to changes in the three astronomical parameters: eccentricity, obliquity, and precession. Some series from the north of the Chinese Loess Plateau also exhibit particle-size variations with millennial frequency, similar to the DO oscillations observed in North Atlantic records. A correlation, mainly based on luminescence dates, has been established between intervals of coarse loess layers and Greenland stadials (without particularly distinguishing the Heinrich stadials), while intervals of fine-grained loess were associated with the DO warming events. The grain size changes are interpreted as reflecting variations in the strength of the atmospheric circulation during the SE Asian winter monsoon over the deserts of Mongolia and northern China, the main dust suppliers to the Chinese Loess Plateau.

The Chinese loess sequences do not contain any paleosols related to DO events, because the corresponding climate is too dry to allow

pedogenesis. Alternating paleosol-loess units similar to those observed in Europe along 50°N can, however, be found in Siberian loess deposits north of Lake Baikal. The available radiocarbon dates allow the Siberian paleosols to be correlated with the DO events and suggest that the impact of the North-Atlantic millennial climate variations reached at least as far east as Lake Baikal.

### North American loess

The North American loess deposits have recorded various influences, mostly related to climate changes originating in the North Atlantic and Pacific regions. According to climate simulations run with an Earth

system model of intermediate complexity, the eastern part of North America should have been affected by the DO and Heinrich abrupt climate changes, but very few loess deposits can be found there to verify this model result. On the other hand, modeling results indicate that much of the Great Plains, where the thickest North American loess deposits are located, and the western part of the continent were not influenced by DO and Heinrich events. Indeed, the succession of paleosol-loess units in sediments in the state of Illinois, 120 km south of the Laurentide ice sheet margin at its maximum extent, is interpreted (using radiocarbon dates) as reflecting sub-millennial timescale inflow of air from the Gulf of Mexico, with a periodicity of about  $450 \pm 100$  yr. Loess sequences located further west, mainly in the state of Nebraska, show bedded mineral layers corresponding to even finer timescales. These were formed as a result of the transport of material during the Last Glacial Termination (~23-12 ka BP) from nearby areas, including the outwash areas of the southwest margin of the Laurentide ice sheet and the east margin of the Cordilleran ice sheet.

### Outlook

Many loess deposits, especially across the Northern Hemisphere, have recorded past climate changes at millennial and sub-millennial timescales. Some loess sequences also contain evidence of particularly strong dust events with a much shorter characteristic timescale. Thus, during Marine Isotopic Stage 5, European sequences (especially from Central Europe) have recorded dust storms of continental magnitude in response to atmospheric blocking episodes (Rousseau et al. 2013). More effort is needed, on the data side, to better quantify the past dust-cycle variations at these different timescales, and on the modeling side, to more realistically simulate them.

### ACKNOWLEDGEMENTS

The authors were supported by the ANR ACTES project (ANR-08-BLAN-0227). DDR was supported by the Alliance program for a Visiting Professorship at Columbia University. This is LDEO contribution 2014-7803.

### AFFILIATIONS


<sup>1</sup>Laboratoire de Météorologie Dynamique & CERES-ERTI, CNRS - Ecole Normale Supérieure, Paris, France  
<sup>2</sup>Lamont-Doherty Earth Observatory of Columbia University, Palisades, USA

### CONTACT

Denis-Didier Rousseau: denis.rousseau@lmd.ens.fr

### REFERENCES

- Full reference list under:  
[www.pages-igbp.org/products/magazine/ref2014\\_2.pdf](http://www.pages-igbp.org/products/magazine/ref2014_2.pdf)  
Rousseau DD et al. (2007) *Geophys Res Lett* 34, doi:10.1029/2007gl031716  
Rousseau DD et al. (2011) *Clim Past* 7: 221-234  
Rousseau DD et al. (2013) *Clim Past* 9: 2213-2230  
Sima A et al. (2009) *Quat Sci Rev* 28: 2851-2866  
Sima A et al. (2013) *Clim Past* 9: 1385-1402



**Figure 2:** (A) Simulated dust emission fluxes for Western Europe for a "Greenland interstadial" (GIS) are relatively close to those for a "Greenland stadial" (GS) when the GIS and GS fluxes are computed without taking the vegetation effect into account (GIS/GS flux ratio between 70-130%); (B) the flux ratio GIS/GS strongly decreases when the vegetation effect of inhibiting eolian erosion is taken into account (modified from Sima et al. 2009). (C) Simulated dust emission fluxes for Eastern Europe for a GIS, superimposed on a topographic map; the most active spots (indicated with dotted lines) have probably been important sources for some Eastern European loess deposits (in yellow, in background), such as those around Stayky (white dot) in NW Ukraine (modified from Sima et al. 2013).

Springer, in press

- Pigati JS et al. (2013) Radiocarbon dating late Quaternary loess deposits using small terrestrial gastropods. *Quat Sci Rev* 76: 114-128
- Prins MA, Vriend M (2007) Glacial and interglacial eolian dust dispersal patterns across the Chinese Loess Plateau inferred from decomposed grain-size records. *Geochem Geophys Geosys* 8, doi:10.1029/2006GC001563

30 [Abrupt climate changes recorded in loess sequences](#)  
[Denis-Didier Rousseau and Adriana Sima](#)

*Complete set of references*

- Kutzbach JE (1987) Model simulations of the climatic patterns during the deglaciation of North America, In: Ruddiman WF, Wright HE (Eds.) *North America and Adjacent Oceans during the Last Deglaciation - The Geology of North America*. The Geological Society of America, Boulder, 425-446
- Rousseau DD et al. (2007) European loess records, In: Elias S (Ed) *Encyclopedia of Quaternary Science*, Elsevier, 1440-1456

*European loess*

- Antoine P et al. (2001) High-resolution record of the last interglacial-glacial cycle in the loess palaeosol sequences of Nussloch (Rhine Valley-Germany), *Quat Int* 76/77: 211-229
- Antoine P et al. (2009) High-resolution record of the last climatic cycle in the southern Carpathian Basin (Surduk, Vojvodina, Serbia), *Quat Int* 198: 19-36
- Antoine P et al. (2013) High-resolution record of the environmental response to climatic variations during the Last Interglacial-Glacial cycle in Central Europe: The loess-palaeosol sequence of Dolni- Vestonice (Czech Republic), *Quat Sci Rev* 67: 17-38
- Bond G et al. (1992) Evidence for massive discharges of icebergs into the North Atlantic Ocean during the last glacial period, *Nature*: 360: 245-249
- Bond GC, Lotti R (1995) Iceberg discharges into the North Atlantic on millennial time scales during the last glaciation, *Science* 267: 1005-1010
- Dansgaard W (1993) Evidence for general instability of past climate from a 250-kyr ice-core record, *Nature* 364: 218-220
- Hatté C, Guiot J (2005) Paleoprecipitation reconstruction by inverse modelling using the isotopic signal of loess organic matter: Application to the Nussloch loess sequence (Rhine Valley, Germany), *Clim Dyn* 25: 315-327
- Hatté C et al. (2013) Excursions to C4 vegetation recorded in the upper Pleistocene loess of surduk (Northern Serbia): An organic isotope geochemistry study, *Clim Past* 9: 1001-1014
- Hemming SR (2004) Heinrich events: Massive late Pleistocene detritus layers of the North Atlantic and their global climate imprint, *Rev Geophys* 42, doi:10.1029/2003RG000128.
- Kindler P et al. (2013) NGRIP temperature reconstruction from 10 to 120 kyr b2K, *Clim Past* 9: 4099-4143
- Lang A et al. (2003) High-resolution chronologies for loess: comparing AMS14C and optical dating results, *Quat Sci Rev* 22: 953-959
- Lautridou JP (1985) Le cycle périglaciaire Pléistocène en Europe du Nord-Ouest et plus particulièrement en Normandie, *Centre Géomorphologie Caen, Université Caen*, 908 pp
- Masson-Delmotte V et al. (2005) GRIP deuterium excess reveals rapid and orbital-scale changes in Greenland moisture origin, *Science* 309: 118-121
- Rousseau DD et al. (1998) Late Pleistocene climatic variations at Achenheim, France based on a magnetic susceptibility and TL chronology of loess, *Quat Res* 49: 255-263
- Rousseau DD et al. (2002) Abrupt millennial climatic changes from Nussloch (Germany) Upper Weichselian eolian records during the Last Glaciation, *Quat Sci Rev* 21: 1577-1582
- Rousseau DD et al. (2007) Link between European and North Atlantic abrupt climate changes over the last glaciation, *Geophys Res Lett* 34, doi:10.1029/2007gl031716
- Rousseau DD et al. (2011) North Atlantic abrupt climatic events of the last glacial period recorded in Ukrainian loess deposits, *Clim Past* 7: 221-234
- Rousseau DD et al. (2013) Major dust events in Europe during marine isotope stage 5 (130-74 ka): A climatic interpretation of the "markers", *Clim Past* 9: 2213-2230
- Sima A et al. (2009) Imprint of North-Atlantic abrupt climate changes on western European loess deposits as viewed in a dust emission model, *Quat Sci Rev* 28: 2851-2866
- Sima A et al. (2013) Modeling dust emission response to North Atlantic millennial-scale climate variations from the perspective of East European MIS3 loess deposits, *Clim Past* 9: 1385-1402

*Asian loess*

- Ding ZL et al. (1995) Ice-Volume Forcing of East Asian Winter Monsoon Variations in the Past 800,000 Years, *Quat Res* 44: 149-159
- Ding ZL et al. (2002) Stacked 2.6-Ma grain size record from the Chinese loess based on five sections and correlation with the deep-sea d18O record, *Paleoceanography* 17: 5.1-5.21
- Guo ZT et al. (2002) Onset of Asian desertification by 22 Myr ago inferred from loess deposits in China, *Nature* 416: 159-163
- Haesaerts P et al. (2005) The loess-palaeosol succession of Kurtak (Yenisei Basin, Siberia): a reference record for the Karga stage (MIS 3), *Quaternaire* 16: 3-24
- Kukla G, An ZS (1989) Loess stratigraphy in central China, *Palaeogeog Palaeoclimatol Palaeoecol* 72: 203-225
- Liu TS et al. (1985) *Loess and the environment*, China Ocean Press, 251 pp
- Sun Y et al. (2012) Influence of Atlantic meridional overturning circulation on the East Asian winter monsoon, *Nature Geosci* 5: 46-49

- Wang YJ et al. (2001) A high-resolution absolute-dated late Pleistocene monsoon record from Hulu Cave, China, *Science* 294: 2345-2348
- Wang Y et al. (2008) Millennial- and orbital-scale changes in the East Asian monsoon over the past 224,000 years, *Science* 451: 1090-1093

#### *North American loess*

- Bettis III EA et al. (2003) Cenozoic eolian sedimentary systems of the USA midcontinent, *Quaternary geology of the United States*, INQUA 2003 Field Guide Volume, Reno, 195-218
- Bettis III EA et al. (2003) Last Glacial loess in the conterminous USA, *Quat Sci Rev* 22: 1907-1946
- Ganopolski A, Rahmstorf S (2001) Rapid changes of glacial climate simulated in a coupled climate model, *Nature* 409: 153-158
- Muhs DR et al. (1999) Late Quaternary loess in northeastern Colorado: Part I-Age and paleoclimatic significance, *Geol Soc Am Bull* 111: 1861-1875
- Muhs DR, Bettis III EA (2000) Geochemical variations in Peoria loess of western Iowa indicate paleowinds of midcontinental North America during last glaciation, *Quat Res* 53: 49-61
- Muhs DR, Bettis III EA (2003) Quaternary loess-paleosol sequences as examples of climate-driven sedimentary extremes, in: *Extreme depositional environments: Mega end members in Geologic time*, Geological Society of America Special papers, The Geological Society of America, 53-74
- Muhs DR, Zarate MA (2001) Late Quaternary eolian records of the Americas and their significance, In: Markgraf V (Ed) *Interhemispheric climate linkages*, Academic Press, 183-216
- Rousseau DD et al. (2007) Evidence of cyclic dust deposition in the US Great plains during the last deglaciation from the high-resolution analysis of the Peoria Loess in the Eustis sequence (Nebraska, USA), *Earth Planet Sci Lett* 262: 159-174
- Sweeney MT et al. (2004) Glacial anticyclone recorded in Palouse loess of northwestern United States, *Geology* 32: 705-708
- Sweeney MR et al. (2005) Topographic and climatic influences on accelerated loess accumulation since the last glacial maximum in the Palouse, Pacific Northwest, USA, *Quat Res* 63: 261
- Wang H et al. (2000) Isotope evidence of paleo-El Nino-Southern Oscillation cycles in loess-paleosol record in the central United States, *Geology* 28: 771-774

#### 32 [Extracting paleodust information from peat geochemistry](#)

[François De Vleeschouwer, M. Ferrat, H. McGowan, H. Vanneste and D. Weiss](#)

- Allan M et al. (2013) Mid and late Holocene dust deposition in western Europe: the Misten peat bog (Hautes Fagnes – Belgium). *Clim Past* 9: 2285-2298
- Blaauw M, Christen JA (2011) Flexible paleoclimate age-depth models using an autoregressive gamma process. *Bayesian Anal* 6: 457-474
- De Vleeschouwer F et al. (2012) Multiproxy paleoenvironmental study in the Misten bog (East Belgium) during the last millenium. *Quat Int* 268: 44-57
- De Vleeschouwer F et al. (2009) Multiproxy evidence of “Little Ice Age” palaeoenvironmental changes in a peat bog from northern Poland. *Holocene* 19: 625-637
- Ferrat M et al. (2013) Numerical simulations of dust fluxes to the eastern Qinghai-Tibetan Plateau: Comparison of model results with a Holocene peat record of dust deposition. *J Geophys Res Atmos* 118: 4597-4609
- Ferrat M et al. (2012) The inorganic geochemistry of a peat deposit on the eastern Qinghai-Tibetan Plateau and insights into changing atmospheric circulation in central Asia during the Holocene, *Geochim Cosmochim Acta* 91: 7-31
- Ferrat M et al. (2011) Improved provenance tracing of Asian dust sources using rare earth elements and selected trace elements for palaeomonsoon studies on the eastern Tibetan Plateau, *Geochim Cosmochim Acta* 75: 6374-6399
- Gallagher K et al. (2011) Inference of abrupt changes in noisy geochemical records using transdimensional changepoint models, *Earth Planet Sci Lett* 311: 182-194
- Harrison SP et al. (2001) The role of dust in climate changes today, at the last glacial maximum and in the future. *Earth Sci Rev* 54: 43-80
- Kamber BS et al. (2010) Comment on "Lead isotopic evidence for an Australian source of aeolian dust to Antarctica at times over the last 170,000 years". *Palaeogeog Palaeoclimatol Palaeoecol* 298: 432-436
- Kohfeld KE, Tegen I (2007) The record of soil dust aerosols and their role in the Earth system, In: Holland HD, Turekian KK (Eds) *Treatise on Geochemistry Update 1*, Elsevier, 1-26
- Kohfeld KE, Harrison SP (2001) DIRTMAP: the geological record of dust. *Earth Sci Rev* 54: 81-114
- Kylander ME et al. (2013) A novel geochemical approach to paleorecords of dust deposition and effective humidity: 8500 years of peat accumulation at Store Mosse ("Big Bog"), Sweden. *Quat Sci Rev* 69: 69-82
- Kylander ME et al. (2007) Rare Earth Element and Pb isotope variations in a 52 kyr peat core from Lynch's Crater (NE Queensland, Australia): Proxy development and application to paleoclimate in the Southern Hemisphere. *Geochim Cosmochim Acta* 71: 942-960
- Kylander ME et al. (2005) Refining the pre-industrial atmospheric Pb isotope evolution curve in Europe using an 8000 year old peat core from NW Spain. *Earth Planet Sci Lett* 240: 467-485
- Le Roux G et al. (2012) Volcano- and climate-driven changes in atmospheric dust sources and fluxes since the Late Glacial in Central Europe. *Geology* 40: 335-338
- Lee JA, Tallis JH (1973) Regional and historical aspects of lead pollution in Britain. *Nature* 245: 216-218
- Mahowald NM et al. (2006) Change in atmospheric mineral aerosols in response to climate: last glacial period pre-industrial, modern and doubled carbon dioxide climates. *J Geophys Res Atmos* 111, doi:10.1029/2005JD006653
- Marx SK et al. (2011) Holocene dust deposition rates in Australia's Murray-Darling Basin, record the interplay between aridity, position of the mid-latitude westerlies, *Quat Sci Rev* 30: 3290-3305
- Marx SK et al. (2009) Long-range dust transport from eastern Australia: a proxy for Holocene aridity and ENSO induced climate