

Exercise Causes Muscle GLUT4 Translocation in an Insulin-Independent Manner

Giovanni Messina^{1,4}, Filomena Palmieri¹, Vincenzo Monda¹, Antonietta Messina¹, Carmine Dalia¹, Andrea Viggiano², Domenico Tafuri³, Antonietta Messina¹, Fiorenzo Moscatelli^{4,5}, Anna Valenzano⁴, Giuseppe Cibelli⁴, Sergio Chieffi¹ and Marcellino Monda^{1*}

¹Department of Experimental Medicine, Section of Human Physiology and Clinical Dietetic Service, Second University of Naples, Via Costantinopoli 16, 80138 Naples, Italy

²Faculty of Medicine, University of Salerno, Salerno, Italy

³Department of Motor Sciences and Wellness, University of Naples "Parthenope", Naples, Italy

⁴Department of Clinical and Experimental Medicine University of Foggia, Foggia, Italy

⁵Department of Movement, Human and Health Sciences, University of Rome "Foro Italico", Rome, Italy

*Corresponding author: Marcellino Monda, Department of Experimental Medicine, Section of Human Physiology and Clinical Dietetic Service, Second University of Naples, Via Costantinopoli 16, 80138 Naples, Italy, Tel. +39 +81 566 5804; Fax +39 +81 5665844; E-mail: marcellino.monda@unina2.it

Received date: May 14, 2015, Accepted date: July 27, 2015, Published date: August 3, 2015

Copyright: ©2015 Monda M. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Glucose uptake in skeletal muscle is dependent on the translocation of GLUT4 glucose transporters to the plasma membrane. The most important stimulators of glucose transport in skeletal muscle are insulin and exercise. Glucose uptake in skeletal muscle during exercise induces acceleration of many processes compared to the resting state. The scientific literature does not underline the role played by muscle contraction to increase glucose uptake with insulin-independent mechanisms. Search on Pub Med (May 05, 2015) using the key words "contraction and glucose uptake and muscle" gives 717 reports, while a search using the key words "insulin and glucose uptake and muscle" cites 5676 publications. The present paper describes the role of exercise in the muscle glucose uptake. Contraction of muscle induces GLUT4 translocation in the absence of insulin. There are different intracellular "pools" of GLUT4, one stimulated by insulin and another one stimulated by exercise. The roles exerted by AMPK, AICAR, calcium, NO, glycogen and hypoxia in the glucose uptake during exercise are emphasized. The effects of these phenomena on human wellness are reported.

Keywords: Glucose; GLUT4; Muscle; Exercise

Introduction

The population statistics of most countries of the world are indicating that industrialization and computerization have been associated with an increase in sedentary behavior and more recently with a significant shift from healthy weight to overweight. In general, this change in the overweight/obesity prevalence is attributed by health professionals to suboptimal diet and physical activity practices.

The beneficial effects of physical exercise on the decreased insulin sensitivity caused by detrimental lifestyle have been demonstrated by experimental evidences. In epidemiological studies, disease prevention has been considered at three levels: primary (avoiding the occurrence of disease), secondary (early detection and reversal), and tertiary (prevention or delay of complications) [1-5]. The major purpose of physical exercise for primary prevention and treatment of lifestyle-related diseases is to improve insulin sensitivity [6-9]. It is known that, during physical exercise, glucose uptake by the working muscles rises 7 to 20 times over the basal level, depending on the intensity of the work performed [10]. However, intense exercise provokes the release of insulin-counter regulatory hormones such as glucagon and catecholamines, which ultimately counter-balance insulin action [11-16]. Continued physical training improves the reduced peripheral tissue sensitivity to insulin in impaired glucose tolerance and Type II diabetes, along with regularization of abnormal lipid metabolism. Furthermore, combination of salt intake restriction and

physical training ameliorates hypertension. In practical terms, before diabetic patients undertake any program of physical exercise, various medical examinations are needed to determine whether they have good glycemic control and are without progressive complications. Because the effect of exercise that is manifested in improved insulin sensitivity decreases within 3 days after exercise and is no longer apparent after 1 week, a continued program is needed.

Muscle glucose uptake [17] can be separated into three sequential steps, i.e. delivery of glucose from the blood to the muscle, transport across the sarcolemma by a GLUT, and irreversible phosphorylation to glucose-6-phosphate by an HK isozyme. Each of these steps can serve as a barrier to MGU and, thus, are important in regulating glucose influx. During resting conditions, the transport step exerts the most control in regulating MGU, as GLUT1 (1-4) or GLUT4 [18,19] overexpression augments basal MGU. Previous work suggests that normal GLUT4 content is sufficient for increases in MGU during exercise, because GLUT4 overexpression alone does not further increase exercise-stimulated MGU [20]. Instead of glucose transport, glucose phosphorylation is a primary limitation of exercise-stimulated MGU [21,22]. Heterozygous GLUT4 knock-out mice serve as a useful tool for examining the impact of reductions in glucose transport capacity on MGU [23-26].

Insulin stimulation and physical exercise are the most physiologically relevant stimulators of glucose transport in skeletal muscle [27,28] and interestingly in patients with Type 2 diabetes,

insulin- but not contraction-stimulated glucose transport is impaired [29].

Both insulin and exercise/muscle contraction increase skeletal muscle glucose uptake by translocation of glucose transporters from an intracellular location to the plasma membrane and t-tubules. GLUT4 is the predominant glucose transporter isoform expressed in skeletal muscle. Early studies have demonstrated that there are distinct proximal signaling mechanisms responsible for the stimulation of GLUT4 translocation and glucose transport by insulin and exercise. Insulin signaling involves the rapid phosphorylation of the insulin receptor, insulin receptor substrate-1/2 (IRS-1/2) on tyrosine residues, and the activation of phosphatidylinositol 3-kinase (PI3-K)[30,31]. In contrast, exercise and muscle contraction have no effect on insulin receptor and IRS-1 phosphorylation or on PI3-K activity[32], and muscle-specific knockout of the insulin receptor does not impair contraction-stimulated glucose transport [33,34]. Clearly, these data demonstrate that the initiating signals that lead to GLUT4 translocation by insulin and exercise in skeletal muscle are distinct.

Stimulation of insulin secretion by glucose involves the enhanced synthesis of ATP by mitochondria [35] and closure of ATP-sensitive K⁺ channels [36]. Subsequent depolarization of the plasma membrane [37] then opens voltage-sensitive (L-type) Ca²⁺ channels [38] causing insulin containing vesicles to fuse at the plasma membrane[39]. The mechanisms that link changes in glucose concentration to the regulation of B-cell genes are less well understood[40,41]. Recently, we have demonstrated that AMP-activated protein kinase (AMPK) is involved in the regulation of gene expression by glucose in this cell type[42]. AMPK is a multisubstrate, heterotrimeric serine/threonine protein kinase, consisting of a catalytic α -subunit and non-catalytic β - and γ -subunits [43].

Regular physical activity leads to a number of adaptations in skeletal muscle that allow the muscle to more efficiently utilize substrates for ATP production and thus become more resistant to fatigue. The three major adaptations to exercise training are: 1) muscle fiber type transformations as defined by the expression of specific contractile proteins (myosin heavy chain isoforms), 2) increases in mitochondrial activity and content, and 3) increases in GLUT4 protein expression.

Adaptations, which might underlie the increased insulin sensitivity in trained individuals, include increases in levels of the glucose transporter protein GLUT-4 and in muscle glycogen synthase activity, a decrease in the serum triglyceride concentration and, possibly, an increase in the muscle capillary network.

In post absorptive humans, there are 100g of glycogen in the liver and 400g of glycogen in muscle. Carbohydrate oxidation by the working muscle can go up by 10-fold with exercise, and yet after 1 h, blood glucose is maintained at 4g. Blood glucose is preserved at the expense of liver and muscle glycogen. Liver glycogen breakdown protects blood glucose as the glucose moieties that comprise it are released into the blood. Muscle glycogen breakdown impedes the removal of glucose from the blood by increasing glucose 6-phosphate (G6P), which inhibits the hexokinase (HK) reaction, and by providing a source of fuel that diminishes the need for blood glucose. The amount of glucose in the blood can still be constant after 2 h of exercise in well-nourished subjects. Blood glucose is protected by liver gluconeogenesis after glycogen stores become critically low. Only after extremely prolonged exercise does blood glucose fall to concentrations that result in hypoglycemia severe enough to cause neuroglycopenia[44].

Glucagon is the primary controller of hepatic glucose production in the sedentary state [45]. Exercise is a robust challenge of the processes involved because of the high rates of glucose production necessary to maintain blood glucose[44]. Glucagon secretion from the pancreatic α cell increases during exercise, whereas insulin secretion from the pancreatic β cell declines. The decline in insulin secretion potentiates the actions of glucagon [46–48]. Studies in animals [49–51] and humans [46,52] demonstrate that the increase in glucagon is the primary stimulator of hepatic glucose production during exercise. The powerful effect of glucagon on hepatic glucose production was recently demonstrated by Berglund et al. [53]. This study showed that increasing glucagon in sedentary mice to levels similar to those seen during exercise causes a marked discharge of hepatic energy stores so that the adenosine monophosphate (AMP) to adenosine triphosphate (ATP) ratio is increased. This increase in the AMP: ATP ratio, through allosteric mechanisms, facilitates the glucagon-induced breakdown of glycogen and the oxidation of fat in the liver [51,54].

The translocation of GLUT4 from an intracellular vesicle to the plasma membrane and T tubules is a major mechanism through which both insulin and exercise increase skeletal muscle glucose transport. Contractile activity can stimulate GLUT4 translocation in the absence of insulin, and some studies suggest there are different intracellular "pools" of GLUT4, one stimulated by insulin and one stimulated by exercise [55]. These findings have provided the basis for our understanding of the glucose transport system with exercise in skeletal muscle.

Conclusion

Systematic review of literature indicates that regular participation in moderately intense physical activity is associated with a substantially lower risk of type 2 diabetes. The association was partly independent of BMI, suggesting that moderate-intensity physical activity can reduce the risk of type 2 diabetes even in those who do not achieve weight loss [56–58]. Findings from several prospective studies [59–62] indicate that 30 min or more of daily moderate-intensity activity, as recommended in multiple guidelines [63,64] can substantially reduce the risk of type 2 diabetes as compared with being sedentary. Moderately intense activity as defined in guidelines (3.0–6.0 MET h) includes walking at brisk pace but not walking at an easy or casual pace [65] and walking at brisk pace also seems preferable for the prevention of type 2 diabetes [61,66–70]. Further studies are needed to define more specifically what combinations of duration and pace are optimal for reducing the risk of type 2 diabetes. However, given that only a low percentage adults in the industrialized countries currently meet the general physical activity recommendations, efforts to prevent type 2 diabetes should strongly emphasize the benefit of moderately intense physical activities and encourage wider participation in these activities.

References:

1. King H, Kriska AM (1992) Prevention of type II diabetes by physical training. *Epidemiological considerations and study methods*. *Diabetes Care* 15: 1794-1799.
2. Messina G, Dalia C, Tafuri D, Monda V, Palmieri F, et al. (2014) Orexin-A controls sympathetic activity and eating behavior. *Front Psychol* 5: 997.
3. Monda M, Messina G, Scognamiglio I, Lombardi A, Martin GA, et al. (2014) Short-Term Diet and Moderate Exercise in Young Overweight

- Men Modulate Cardiocyte and Hepatocarcinoma Survival by Oxidative Stress. *Oxid Med Cell Longev* 1-7.
4. Di Bernardo G, Messina G, Capasso S, Del Gaudio S, Cipollaro M, et al. (2014) Sera of overweight people promote in vitro adipocyte differentiation of bone marrow stromal cells. *Stem Cell Res Ther* 5: 4.
 5. Esposito M, Serpe FP, Diletti G, Messina G, Scortichini G, et al. (2014) Serum levels of polychlorinated dibenzo-p-dioxins, polychlorinated dibenzofurans and polychlorinated biphenyls in a population living in the Naples area, southern Italy. *Chemosphere* 94: 62-69.
 6. Viggiano A, Chieffi S, Tafuri D, Messina G, Monda M, et al. (2014) Laterality of a second player position affects lateral deviation of basketball shooting. *J Sports Sci* 32: 46-52.
 7. Messina G, De Luca V, Viggiano A, Ascione A, Iannaccone T, et al. (2013) Autonomic nervous system in the control of energy balance and body weight: personal contributions. *Neurol Res Int* 2013: 639280.
 8. Messina G, Vicidomini C, Viggiano A, Tafuri D, Cozza V, et al. (2012) Enhanced parasympathetic activity of sportive women is paradoxically associated to enhanced resting energy expenditure. *Auton Neurosci Basic Clin* 169: 102-106.
 9. Wagenmakers AJ, Strauss JA, Shepherd SO, Keske MA, Cocks M (2015) Increased muscle blood supply and transendothelial nutrient and insulin transport induced by food intake and exercise: effect of obesity and ageing. *J Physiol*.
 10. Yamanouchi K, Shinozaki T, Chikada K, Nishikawa T, Ito K, et al. (1995) Daily walking combined with diet therapy is a useful means for obese NIDDM patients not only to reduce body weight but also to improve insulin sensitivity. *Diabetes Care* 18: 775-778.
 11. Monda M, Viggiano A, Viggiano A, Viggiano E, Messina G, et al. (2007) Sympathetic and hyperthermic reactions by orexin A: role of cerebral catecholaminergic neurons. *Regul Pept* 139: 39-44.
 12. Monda M, Amaro S, Sullo A, De Luca B (1994) Posterior hypothalamic activity and cortical control during the PGE1 hyperthermia. *Neuroreport* 6: 135-139.
 13. Monda M, Pittman QJ (1993) Cortical spreading depression blocks prostaglandin E1 and endotoxin fever in rats. *Am J Physiol* 264: R456-459.
 14. Monda M, Amaro S, Sullo A, De Luca B (1995) Injection of muscimol in the posterior hypothalamus reduces the PGE1-hyperthermia in the rat. *Brain Res Bull* 37: 575-580.
 15. Monda M, Viggiano A, Viggiano A, Viggiano E, Messina G, et al. (2006) Quetiapine lowers sympathetic and hyperthermic reactions due to cerebral injection of orexin A. *Neuropeptides* 40: 357-363.
 16. Monda M, Viggiano A, Viggiano A, Fuccio F, De Luca V (2004) Injection of orexin A into the diagonal band of Broca induces sympathetic and hyperthermic reactions. *Brain Res* 1018: 265-271.
 17. Hansen PA, Marshall BA, Chen M, Holloszy JO, Mueckler M (2000) Transgenic overexpression of hexokinase II in skeletal muscle does not increase glucose disposal in wild-type or Glut1-overexpressing mice. *J Biol Chem* 275: 22381-22386.
 18. Hansen PA, Gulve EA, Marshall BA, Gao J, Pessin JE, et al. (1995) Skeletal muscle glucose transport and metabolism are enhanced in transgenic mice overexpressing the Glut4 glucose transporter. *J Biol Chem* 270: 1679-1684.
 19. Treadway JL, Hargrove DM, Nardone NA, McPherson RK, Russo JF, et al. (1994) Enhanced peripheral glucose utilization in transgenic mice expressing the human GLUT4 gene. *J Biol Chem* 269: 29956-29961.
 20. Fueger PT, Bracy DP, Malabanan CM, Pencek RR, Granner DK, et al. (2004) Hexokinase II overexpression improves exercise-stimulated but not insulin-stimulated muscle glucose uptake in high-fat-fed C57BL/6j mice. *Diabetes* 53: 306-314.
 21. Halseth AE, Bracy DP, Wasserman DH (1998) Limitations to exercise- and maximal insulin-stimulated muscle glucose uptake. *J Appl Physiol* (1985) 85: 2305-2313.
 22. Halseth AE, Bracy DP, Wasserman DH (1999) Overexpression of hexokinase II increases insulin and exercise-stimulated muscle glucose uptake in vivo. *Am J Physiol* 276: E70-77.
 23. Cline GW, Petersen KF, Krssak M, Shen J, Hundal RS, et al. (1999) Impaired glucose transport as a cause of decreased insulin-stimulated muscle glycogen synthesis in type 2 diabetes. *N Engl J Med* 341: 240-246.
 24. Kubo K, Foley JE (1986) Rate-limiting steps for insulin-mediated glucose uptake into perfused rat hindlimb. *Am J Physiol* 250: E100-102.
 25. Alvim RO, Cheuhen MR, Machado SR, Sousa AG, Santos PC (2015) General aspects of muscle glucose uptake. *An Acad Bras Cienc* 87: 351-368.
 26. Stenbit AE, Tsao TS, Li J, Burcelin R, Geenen DL, et al. (1997) GLUT4 heterozygous knockout mice develop muscle insulin resistance and diabetes. *Nat Med* 3: 1096-1101.
 27. Goodyear LJ, Kahn BB (1998) Exercise, glucose transport, and insulin sensitivity. *Annu Rev Med* 49: 235-261.
 28. Hayashi T, Wojtaszewski JF, Goodyear LJ (1997) Exercise regulation of glucose transport in skeletal muscle. *Am J Physiol* 273: E1039-1051.
 29. Kennedy JW, Hirshman MF, Gervino EV, Ocel JV, Forse RA, et al. (1999) Acute exercise induces GLUT4 translocation in skeletal muscle of normal human subjects and subjects with type 2 diabetes. *Diabetes* 48: 1192-1197.
 30. Folli F, Saad MJ, Backer JM, Kahn CR (1992) Insulin stimulation of phosphatidylinositol 3-kinase activity and association with insulin receptor substrate 1 in liver and muscle of the intact rat. *J Biol Chem* 267: 22171-22177.
 31. Goodyear LJ, Giorgino F, Balon TW, Condorelli G, Smith RJ (1995) Effects of contractile activity on tyrosine phosphoproteins and PI 3-kinase activity in rat skeletal muscle. *Am J Physiol* 268: E987-995.
 32. Treadway JL, James DE, Burcel E, Ruderman NB (1989) Effect of exercise on insulin receptor binding and kinase activity in skeletal muscle. *Am J Physiol* 256: E138-144.
 33. Wojtaszewski JF, Higaki Y, Hirshman MF, Michael MD, Dufresne SD, et al. (1999) Exercise modulates postreceptor insulin signaling and glucose transport in muscle-specific insulin receptor knockout mice. *J Clin Invest* 104: 1257-1264.
 34. Govers R (2014) Cellular regulation of glucose uptake by glucose transporter GLUT4. *Adv Clin Chem* 66: 173-240.
 35. Kennedy HJ, Pouli AE, Ainscow EK, Jouaville LS, Rizzuto R, et al. (1999) Glucose generates sub-plasma membrane ATP microdomains in single islet beta-cells. Potential role for strategically located mitochondria. *J Biol Chem* 274: 13281-13291.
 36. Bryan J, Aguilar-Bryan L (1997) The ABCs of ATP-sensitive potassium channels: more pieces of the puzzle. *Curr Opin Cell Biol* 9: 553-559.
 37. Soriano S, Gonzalez A, Marroquí L, Tudurí E, Vieira E, et al. (2010) Reduced insulin secretion in protein malnourished mice is associated with multiple changes in the beta-cell stimulus-secretion coupling. *Endocrinology* 151: 3543-3554.
 38. Safayhi H, Haase H, Kramer U, Bihlmayer A, Roenfeldt M, et al. (1997) L-type calcium channels in insulin-secreting cells: biochemical characterization and phosphorylation in RINm5F cells. *Mol Endocrinol* 11: 619-629.
 39. Rutter GA (2001) Nutrient-secretion coupling in the pancreatic islet beta-cell: recent advances. *Mol Aspects Med* 22: 247-284.
 40. Melloul D, Marshak S, Cerasi E (2002) Regulation of insulin gene transcription. *Diabetologia* 45: 309-326.
 41. Leclerc I, da Silva Xavier G, Rutter GA (2002) AMP- and stress-activated protein kinases: key regulators of glucose-dependent gene transcription in mammalian cells? *Prog Nucleic Acid Res Mol Biol* 71: 69-90.
 42. da Silva Xavier G, Leclerc I, Salt IP, Doiron B, Hardie DG, et al. (2000) Role of AMP-activated protein kinase in the regulation by glucose of islet beta cell gene expression. *Proc Natl Acad Sci U S A* 97: 4023-4028.
 43. Stapleton D, Mitchelhill KI, Gao G, Widmer J, Michell BJ, et al. (1996) Mammalian AMP-activated protein kinase subfamily. *J Biol Chem* 271: 611-614.

44. Wasserman DH (2009) Four grams of glucose. *Am J Physiol Endocrinol Metab* 296: E11–E21.
45. Liljenquist JE, Mueller GL, Cherrington AD, Keller U, Chiasson J-L, et al. (1977) Evidence for an important role of glucagon in the regulation of hepatic glucose production in normal man. *J Clin Invest* 59: 369–374.
46. Lavoie C, Ducros F, Bourque J, Langelier H, Chiasson JL (1997) Glucose metabolism during exercise in man: the role of insulin and glucagon in the regulation of hepatic glucose production and gluconeogenesis. *Can J Physiol Pharmacol* 75: 26–35.
47. Lins PE, Wajngot A, Adamson U, Vranic M, Efendić S (1983) Minimal increases in glucagon levels enhance glucose production in man with partial hypoinsulinemia. *Diabetes* 32: 633–636.
48. Wasserman DH, Williams PE, Lacy DB, Goldstein RE, Cherrington AD (1989) Exercise-induced fall in insulin and hepatic carbohydrate metabolism during muscular work. *Am J Physiol* 256: E500–509.
49. Wasserman DH, Lickley HL, Vranic M (1984) Interactions between glucagon and other counterregulatory hormones during normoglycemic and hypoglycemic exercise in dogs. *J Clin Invest* 74: 1404–1413.
50. Wasserman DH, Lickley HL, Vranic M (1985) Important role of glucagon during exercise in diabetic dogs. *J Appl Physiol* (1985) 59: 1272–1281.
51. Wasserman DH, Spalding JA, Lacy DB, Colburn CA, Goldstein RE, et al. (1989) Glucagon is a primary controller of hepatic glycogenolysis and gluconeogenesis during muscular work. *Am J Physiol* 257: E108–117.
52. Hirsch IB, Marker JC, Smith LJ, Spina RJ, Parvin CA, et al. (1991) Insulin and glucagon in prevention of hypoglycemia during exercise in humans. *Am J Physiol* 260: E695–E704.
53. Berglund ED, Lee-Young RS, Lustig DG, Lynes SE, Donahue EP, et al. (2009) Hepatic energy state is regulated by glucagon receptor signaling in mice. *J Clin Invest* 119: 2412–2422.
54. Wasserman DH, Spalding JA, Bracy D, Lacy DB, Cherrington AD (1989) Exercise-induced rise in glucagon and ketogenesis during prolonged muscular work. *Diabetes* 38: 799–807.
55. Roy D, Marette A (1996) Exercise induces the translocation of GLUT4 to transverse tubules from an intracellular pool in rat skeletal muscle. *Biochem Biophys Res Commun* 223: 147–152.
56. Monda M, Messina G, Mangoni C, De Luca B (2008) Resting energy expenditure and fat-free mass do not decline during aging in severely obese women. *Clin Nutr* 27: 657–659.
57. Viggiano A, Vicidomini C, Monda M, Carleo D, Carleo R, et al. (2009) Fast and low-cost analysis of heart rate variability reveals vegetative alterations in non-complicated diabetic patients. *J Diabetes Complications* 23: 119–123.
58. Monda M, Messina G, Vicidomini C, Viggiano A, Mangoni C, et al. (2006) Activity of autonomic nervous system is related to body weight in pre-menopausal, but not in post-menopausal women. *Nutr Neurosci* 9: 141–145.
59. Hsia J, Wu L, Allen C, Oberman A, Lawson WE, et al. (2005) Physical activity and diabetes risk in postmenopausal women. *Am J Prev Med* 28: 19–25.
60. Hu FB, Sigal RJ, Rich-Edwards JW, Colditz GA, Solomon CG, et al. (1999) Walking compared with vigorous physical activity and risk of type 2 diabetes in women: a prospective study. *JAMA* 282: 1433–1439.
61. Laaksonen DE, Lindström J, Lakka TA, Eriksson JG, Niskanen L, et al. (2005) Physical activity in the prevention of type 2 diabetes: The finnish diabetes prevention study. *Diabetes* 54: 158–165.
62. Weinstein AR, Sesso HD, Lee IM, Cook NR, Manson JE, et al. (2004) Relationship of physical activity vs body mass index with type 2 diabetes in women. *JAMA* 292: 1188–1194.
63. Haskell WL, Lee I-M, Pate RR, Powell KE, Blair SN, et al. (2007) Physical activity and public health: updated recommendation for adults from the American College of Sports Medicine and the American Heart Association. *Med Sci Sports Exerc* 39: 1423–1434.
64. Eyre H, Kahn R, Robertson RM (2004) Preventing cancer, cardiovascular disease, and diabetes: A common agenda for the American Cancer Society, the American Diabetes Association, and the American Heart Association. *Diabetes Care* 27: 1812–1824.
65. Ainsworth BE, Haskell WL, Whitt MC, Irwin ML, Swartz AM, et al. (2000) Compendium of physical activities: an update of activity codes and MET intensities. *Med Sci Sports Exerc* 32: S498–504.
66. Hu FB, Leitzmann MF, Stampfer MJ, Colditz GA, Willett WC, et al. (2001) Physical activity and television watching in relation to risk for type 2 diabetes mellitus in men. *Arch Intern Med* 161: 1542–1548.
67. Messina G, Viggiano A, De Luca V, Messina A, Chieffi S, et al. (2013) Hormonal changes in menopause and orexin-a action. *Obstet Gynecol Int* 2013: 209812.
68. De Luca V, Viggiano E, Messina G, Viggiano A, Borlido C, et al. (2008) Peripheral amino Acid levels in schizophrenia and antipsychotic treatment. *Psychiatry Investig* 5: 203–208.
69. Viggiano A, Nicodemo U, Viggiano E, Messina G, Viggiano A, et al. (2010) Mastication overload causes an increase in O₂ - production into the subnucleus oralis of the spinal trigeminal nucleus. *Neuroscience* 166: 416–421.
70. Monda M, Viggiano A, Viggiano A, Mondola R, Viggiano E, et al. (2008) Olanzapine blocks the sympathetic and hyperthermic reactions due to cerebral injection of orexin A. *Peptides* 29: 120–126.