

Plumbosolvency exacerbated by Water Fluoridation

Geoff Pain

September 2015

Abstract

Plumbosolvency, the dissolution of metallic Lead, which results in the contamination of drinking water and consequent damage to human health, is recognized as a major problem wherever Lead pipes, solder or brass fittings are present in the supply route. Plumbosolvency is exacerbated by Fluoridation of drinking water. The use of Phosphate in an attempt to reduce the dissolution rate leads to increased costs, waste of a scarce natural resource and environmental degradation. Immediate cessation of neurotoxic Fluoridation to reduce plumbosolvency makes more economic sense. Provision of communal water supplies from point of collection reverse osmosis filters can bring an immediate end to Lead exposure while sources of Lead contamination are gradually removed from the supply network, creating thousands of person years employment in a depressed economy.

Keywords: Crime, Delinquency, Fluoride, Fluoridation, Heart, Hexafluorosilicate, IQ, Lead, Neurotoxin, Phosphate, Plumbosolvency, Stroke

Introduction

The historical use of the malleable metal Lead in plumbing has left a toxic legacy that will cost many nations billions of dollars to remove. In the presence of oxygen, Lead will dissolve slowly with the initial formation of Lead Hydroxide, which readily reacts with other ions. With Carbonate or Chloride, slightly soluble salts form a crust on the metal. The following Table of Solubility shows relative solubilities for some Lead compounds [CRC].

Compound	Formula	Solubility in Cold (20°C) Water Gram per litre
Lead Hexafluorosilicate	PbSiF ₆	"soluble"
Lead Chloride	PbCl ₂	0.99
Lead Fluoride	PbF ₂	0.64
Lead Fluorochloride	PbFCl	0.37
Lead Oxychloride, the mineral Matlockite	PbCl ₂ ·Pb(OH) ₂	0.0095
Lead Carbonate, the mineral Cerussite	PbCO ₃	0.0011
Lead Orthophosphate	Pb ₃ (PO ₄) ₂	0.00014
Basic Lead Carbonate, White Lead	2PbCO ₃ ·Pb(OH) ₂	"insoluble"
EU 2013 Target		0.00001
USEPA Minimal Risk Level		0.0000001
EU "Goal"		0

The addition of Chlorine to water could in theory reduce leaching by formation of Matlockite, but is more likely to increase leaching of Lead by forming Lead Chloride. Chlorine also leads to formation of Chloramines which enhance leaching of brass, releasing more Lead [Miranda 2006, Switzer 2006, Maas 2007].

Governments until quite recently have been content to “do nothing” about plumbosolvency, however with the advent of the European Union, multinational standards for water safety have emerged and member nations have found themselves confronted with the obligation to try to meet those standards.

The EU set a target of less than 10 micrograms of Lead per litre (10 ppb) to be reached by 2013. The optimum level is Zero ppm.

Adverse Effects of Lead

It took decades of consumer activism to eliminate tetraethyllead from petrol despite decades of scientific proof of harm. Like Fluoride, Lead is bio-accumulative, both being recognized neurotoxins that permanently damage the brain, reducing IQ [Xiang 2003] and can cause behavioural problems. Mullinex [1995] found fluoride more potent than lead in damage to behaviour of experimental animals. Therefore the apparent concern over Lead contamination tends to obscure the neurotoxic hazard of Fluoride in public discussions of water contamination [Xiang 2003]. Recently the United States halved the market for Fluoride by reducing the concentration of Fluoride in US water supplies to 0.7 ppm [Gooch 2015].

Lead increases the severity of dental fluorosis. Increased body load of Lead reduces IQ [Masters 2004, Lamphear], decreases learning ability [Niu 2008, 2009], increases juvenile delinquency, and increases crime rates [Mann 2000]. Fetal death and reduced birth rates are associated with exposure to lead-contaminated drinking water [Edwards 2014].

Lead is also associated with high blood pressure, stroke and heart attacks [Pocock 1998].

Hexafluorosilicate increases Lead leaching

Hydrofluorosilicic acid and its salts are waste products from the phosphate fertilizer industry that cannot be legally dumped on land, surface water, groundwater or the sea. The compound reversibly hydrolyses forming oligomers. Presence of excess silica stabilizes hexafluorosilicate ion and its partial hydrolysis products [Borodin 1974].

Hydrofluorosilicic acid (H_2SiF_6) doubles the number of children with blood Lead (PbB) > 10 mg/dL [Coplan 2007].

The addition of Hydrofluorosilicic acid has been demonstrated to increase the dissolution rate and hence the concentration of Lead in drinking water supplies and this translates directly to higher Lead blood levels and associated human damage [Masters 2000, Allegood 2005, Clabby 2006 and Miranda 2006, discussed in Maas 2007].

This occurs because Hexafluorosilicate can form complex ions with Lead ions (coordination number up to nine) via bridging Fluorine atoms in solution [Bonomi 2001, Burt 2015, Cole 1981]. Stable Hexafluorosilicate compounds are readily obtained from aqueous solution [Conley 2002, Gelmboldt 2007, Burt 2015].

Rapid ion exchange causes shifts in ^{19}F NMR measurements and has led some researchers to mistakenly state that hydrolysis of hexafluorosilicate is “complete” in water. However the fact that Si-F coupling constants are observed at low temperature confirms attachment of F to Si under mild conditions [Borodin 1974, Conley 2002].

There is also a little studied Pentafluorosilicate ion SiF_5^- that is present at pH below 3.5 [Finney 2006]. The hydrated ion $[\text{SiF}_5(\text{H}_2\text{O})]^-$ and the neutral intermediate *trans*- $[\text{SiF}_4(\text{H}_2\text{O})_2]$ have been successfully isolated in stable solids [Gelmboldt 2007]. The stability of complex fluoride ions of silicon is enhanced by the fact that the Si-F bond is much stronger than the Si-O bond [CRC]. The toxicology of these fluorosilicate species has not been widely reported [Rice 2014], and perhaps even suppressed. According to one study “No carcinogenicity studies have been conducted using

(hydro)fluorosilicic acid, sodium silicofluoride, disodium hexafluorosilicate or hexafluorosilicate or hexafluorosilicic acid.” [European Commission 2010].

Hydrofluorosilicic acid is a more powerful inhibitor of acetylcholinesterase than Sodium Fluoride (NaF) [Westendorf 1975, cited in Coplan 2007].

Use of Phosphate in attempts to reduce Plumbosolvency

Some water suppliers have added Orthophosphate in an attempt to counter the leaching by formation of a thin coating of Lead Orthophosphate inside the pipes [Comber 2011]. However reference to the Table of Solubility above shows that, assuming saturated conditions, the EU target will clearly not be met by this treatment because any Lead Phosphate in the pipes has plenty of time to dissolve to reach saturation when the water is not flowing.

Phosphorus is essential to life but is a rapidly diminishing natural resource and should therefore be conserved for food production. Given that 99% of the added Phosphate will not be consumed by humans, the economic and environmental folly of this approach is exposed.

When excess Phosphorus is released to the environment, algal blooms can occur that damage the ecology and dosing concentrations commonly used in attempts to reduce plumbosolvency are up to 30 times those allowed in UK rivers [Goody 2015].

Phosphate sources commonly used are Phosphoric Acid or Monosodium dihydrogenphosphate (about seven times the cost of the acid). Zinc phosphates would be too toxic to contemplate. It has been found that some Polyphosphate chemicals actually increase Lead mobilization [Edwards and McNiell 2002 cited in Maas 2007].

All Phosphate sources are contaminated with toxic and carcinogenic elements including Arsenic, Cadmium, Chromium, Copper, Lead, Mercury, and Zinc. Clearly the addition of Phosphate increases risk to human health.

In a 2003 study of orthophosphate dosing [Jackson 2003], there was a statistically marginal reduction of measured Lead in only 3 out of 5 properties tested, however the fundamental flaw of that study was failure to monitor temperature on sample collection. Following poor results with orthophosphate a 2011 report [UK Water 2011] discussed the options for finding alternative strategies and briefly mentioned introduction of pipe liners. The economic model included an estimate of extra costs at waste water treatment plants to cope with the excess phosphate but failed to include environmental impact and associated costs of dumping phosphate in rivers.

A recent study of orthophosphate dosing performed for the Government of Alberta, Canada, concluded that the most cost-effective approach for Calgary is an accelerated programme of total pipe replacement, while orthophosphate will delay the inevitable need to replace pipes in Edmonton [Hayes 2014].

Plumbosolvency Lead Hazard Reduction by Cessation of Fluoridation

Studies in two US cities, Tacoma, Washington and Thurmont, Maryland, found an immediate reduction of Lead concentration when Fluoridation was stopped. In Tacoma the level dropped from 32 parts per billion (ppb) to 17 ppb and in Thurmont from 30 ppb to 7 ppb, that is below the 2013 EU target level [cited in Maas 2007].

Recommended actions and alternative Strategy

Considering the available information, the following actions could partially ameliorate the Plumbosolvency threat from supply networks:

- Immediate cessation of Fluoridation
- no addition of Phosphates
- minimal use of Chlorine
- use Ozone instead of Chlorine
- adjustment of pH with the least contaminating alkali available
- acceleration of Lead pipe replacement

Replacement of piping in the distribution network and customer premises will take time, generating much new employment and net economic growth. The improvement in human health could be assigned a monetary value.

In order to more rapidly remove this dangerous Lead exposure from the entire affected population, namely those who drink distributed system water, I suggest an alternative strategy.

As 99% of the system water is not consumed, it would be economically feasible to deliver reverse osmosis pure water at community taps so that each family could collect a few litres of water per day for drinking and cooking.

The installation requires a small footprint, could be powered by renewable energy and would need a small holding tank while producing pure water by pressure sensing pump on demand. Citizens would be advised to refill their own approved container, avoiding plastic waste.

This strategy has been implemented in New Zealand by communities that reject deliberate poisoning of water with Fluoride, enabling them to opt out at almost no cost apart from transport to the supply point. People are happy to meet at the clean water tap, similar to distribution of hot water in poorer areas of Scotland, prior to the Thatcher era.

It would be worthwhile if all water providers immediately installed a number of these facilities for a demonstration and trial of community acceptance while explaining the real and present danger of Lead poisoning wherever plumbosolvency occurs.

Hopefully the international attention currently being brought to plumbosolvency will see more governments making the decision to cease fluoridation of water (as did Israel in 2014) to remove another neurotoxic hazard.

Selected References

Allegood, J. 2005. Water treatment process called potential risk. Chemicals' mix with plumbing could put lead in tap water. *The News & Observer*. May 18, 2005 [online]. Available:

http://www.newsobserver.com/news/health_science/story/2417101p-8794959c.html

Auckland Healthcare Services. 2008. What is Plumbosolvency and are there any Health Risks?

Bonomi A, et al. 2001. Selective deleading process and bath for plumbing components made of a copper alloy. US Patent 6,284,053; September 4, 2001.

Borodin, PM, Zao, NGK and NS Petrov. 1974. Stability of the hexafluorosilicate ion in aqueous solutions of fluorosilicic acid, from ^{19}F NMR data *Journal of Structural Chemistry* 14(4) 564-568

Burt, J, Grantham, W, Levason, W, Light, M E and Reid, G. 2015. Hexafluorosilicate and tetrafluoroborate coordination to lead(II) di- and tri-imine complexes – Unusual fluoroanion coordination modes. *Polyhedron*, 85, 530-36

Burt, J, Grantham, W, Levason, W and Reid. 2015. Lead(II) nitrate and hexafluorosilicate complexes with neutral diphosphine coordination *Dalton Transactions* , 244, 11533

Clabby, C. 2005. Water tests to plumb extent of lead problem. Durham among cities where chemical could cause toxic taint in tap water. *The News and Observer*, Raleigh, NC, September 3 [online]. Available:

<http://www.newsobserver.com>

Cole ER Jr., et al. 1981. Electrowinning of lead from H_2SiF_6 solution. US Patent 4,272,340; June 9

Coplan MJ, Patch SC, Masters RD, Bachman MS. 2007. Confirmation of and explanations for elevated blood lead and other disorders in children exposed to water disinfection and fluoridation chemicals. *Neurotoxicology*. Sep; 28(5):1032-42. Epub Mar 1

Comber, S, Cassé, F, Brown, B, Martin, J, Hillis, P and Gardner, M. 2011. Phosphate treatment to reduce plumbosolvency of drinking water also reduces discharges of copper into environmental surface waters. *Water Environ. J.* 25, 266–270.

Conley, DC, Yearwood BC, Parkin S and Atwood DA. 2002. Ammonium hexafluorosilicate salts. *Journal of Fluorine Chemistry* 115, 155-160

CRC Handbook of Chemistry and Physics 57th Edition.

Dabeka, RW, Karpinski, KF McKenzie, AD and CD Bajdik. 1986. Survey of lead, cadmium and fluoride in human milk and correlation of levels with environmental and food factors. *Food Chem. Toxicol.* 24(9):913-921.

Dabeka, RW, and AD McKenzie. 1987. Lead, cadmium, and fluoride levels in market milk and infant formulas in Canada. *J. Assoc. Off. Anal. Chem.* 70(4):754-757.

Dabeka, RW, and AD McKenzie. 1995. Survey of lead, cadmium, fluoride, nickel, and cobalt in food composites and estimation on dietary intakes of these elements by Canadians in 1986-1988. *J. AOAC Int.* 78(4):897-909.

Dudi A, Schock M, Murray N, Edwards M. 2005. Lead leaching from inline brass devices: a critical evaluation of the existing standard. *J Am Water Works Assoc* 97(8):66–78.

Edwards M, McNeill LS. 2002. Effect of phosphate inhibitors on lead release from pipes. *J Am Water Works Assoc*;94(1):79–92

Edwards, M, Triantafyllidou, S, Best, D. 2009. Elevated blood lead in young children due to lead-containing drinking water. *Environ. Sci. Technol.* 43, 1618–1623.

Edwards, M. 2014. Fetal death and reduced birth rates associated with exposure to lead-contaminated drinking water. *Environ. Sci. Technol.* 48, 739–746.

European Commission. 2010. Critical review of any new evidence on the hazard profile, health effects, and human exposure to fluoride and the fluoridating agents of drinking water.

Finney WF, Wilson E, Callender A, Morris MD, Beck LW, 2006, Reexamination of Hexafluorosilicate Hydrolysis by ¹⁹F NMR and pH Measurement, *Environ. Sci. Technol.*, 40 (8), pp 2572–2577

Glemboldt, VO, Ganin VE, Fonaria MS, Simonov, YA, Koroeva, YA, Enna AA, Basok SS, Shova, S, Kahlig, H, Arion, VB and Keppler BK. 2007. Two new “onium” fluorosilicates, the products of interaction of fluorosilicic acid with 12-membered macrocycles: structures and spectroscopic properties. *Dalton Transactions.* 2915-2924

Gooch, BF. 2015. U.S. Public Health Service Recommendation for Fluoride Concentration in Drinking Water for the Prevention of Dental Caries. *Public Health Reports / July–August 2015 / Volume 130, 1-14*

Goody, Daren C, Lapworth, Dan J, Ascott, Matthew J, Bennett, Sarah A, Heaton Timothy H E, and Ben W J Surridge 2015. Isotopic Fingerprint for Phosphorus in Drinking Water Supplies *Environ. Sci. Technol.* 49, 9020–9028

Hac, E, Czarnowski, Gos, T and J Krechniak. 1997. Lead and fluoride content in human bone and hair in the Gdansk region. *Sci. Total Environ.* 206(2-3):249-254.

Hayes, CR and Croft, TN. 2014. Optimisation of plumbosolvency control using computational modelling techniques: a demonstration project for the Government of Alberta, working with the city of Calgary and EPCOR (Edmonton).

Holman, I P, Whelan, M J, Howden, N J K, Bellamy, P H, Wilby, N J, Rivas-Casado, M, McConvey, P. 2008. Phosphorus in groundwater - An overlooked contributor to eutrophication. *Hydrol. Processes* 22, 5121–5127.

Jackson PJ and JC Ellis. 2003. Report DWI 6173 Demonstration of Optimisation of Plumbosolvency Treatment and Control Measures. Final Report to the Drinking Water Inspectorate

Kennedy, David C. D.D.S 2003 IAOMT Policy position on ingested fluoride and fluoridation

Langphear, B P, Hornung, R, Khoury, J, Yolton, K, Baghurst, P, Bellinger, D C, Canfield, R L, Dietrich, K N, Bornschein, R, Greene, T, Rothenberg, S J, Needleman, H L, Schnaas, L, Wasserman, G, Graziano, J, Roberts, R. 2005. Low-level environmental lead exposure and children’s intellectual function: An international pooled analysis. *Environ. Health Perspect.* 113, 894–899.

Leite GAS, Sawan RMM, Teofilo JM, Porto IM, Sousa FB, Gerlach RF. 2011. Exposure to lead exacerbates dental fluorosis. *Arch Oral Biol* 56(7):695-702.

Luo G, Niu R, Sun Z, Zhang J, Wang J, Wang C, Wang J. 2011. Reduction of CAMKII expression in the hippocampus of rats from ingestion of fluoride and/or lead. *Fluoride April-June* 44(2):63–69

Maas, RP, Patch, SC and AM Smith. 2005. Effects of Fluorides and Chloramines on Lead Leaching from Leaded-Brass Surfaces. Technical Report 05-142. Environmental Quality Institute, University of North Carolina, Asheville, NC. June

Maas, R P, Patch, S C, Christian, A and Coplan, M J. 2007 Effects of fluoridation and disinfection agent combinations on lead leaching from leaded-brass parts *NeuroToxicology* 28, 1023–1031

Macek, MD, Matte, TD Sinks, T and DM Malvitz. 2006. Blood lead concentrations in children and method of water fluoridation in the United States, 1988-1994. *Environ. Health Perspect.* 114(1):130-134.

- Mahaffey, KR et al 1976. Effect of High Fluorine (F) intake on tissue lead (Pb) Concentrations. Fed. Proc., 35, 256.
- Mann, Judy. 2000. Chemicals and Crime: A Truly Toxic Effect by, May 26, page C 11, Washington Post
- Masters, R D and M J Coplan. 1999. Water Treatment With Silicofluorides and Lead Toxicity, by International Journal of Environmental Studies, 56. pp. 435-449
- Masters, R D. 2004. The Hidden Handicap: Lead, Brain Chemistry, and Educational Failure. Paper presented to 2004 Annual Meeting of the American Political Science Association, Chicago, IL., Sept. 3.
- Masters RD, Coplan MJ, Hone BT, Dykes JE. 2000. Association of silicofluoride treated water with elevated blood lead. Neurotoxicology 21(6):1091-100.
- Miranda ML, Kim D, Hull AP, Paul CJ, Galeano MAO. 2006 Changes in blood lead levels associated with use of chloramines in water treatment systems. Environ. Health Perspect. <http://dx.doi.org/>; online November 7.
- Mullenix, P et al. 1995. Neurotoxicity of Sodium Fluoride in Rats, Neurotoxicology and Teratology, 17, No. 2, pp. 169-177
- Niu, R, Sun, Z, Cheng, Z, Liu, H and Chen H. 2008. Effects of fluoride and lead on N-methyl-D-aspartate receptor 1 expression in the hippocampus of offspring rat pups Fluoride April-June 41(2):101-110
- Niu, R, Sun, Z, Wang, J, Cheng, Z and Wang J. 2008. Effects of fluoride and lead on locomotor behavior and expression of nissl body in brain of adult rats. Fluoride October-December 41(4):276-282
- Niu, R, Sun, Z, Cheng, Z, Li, Z and Wang J. 2009. Decreased learning ability and low hippocampus glutamate in offspring rats exposed to fluoride and lead. Environmental Toxicology and Pharmacology September 28(2):254-8.
- Pocock, S J, Shaper, A G, Ashby, H T, Dewes, H T and Clayton, B E. 1988. The relationship between blood lead, blood pressure, stroke and heart attacks in middle-aged British men. Environ. Health Perspect. 78, 23-30.
- Rice JR, Boyd WA, Chandra D, Smith MV, Den Besten PK, Freedman JH. 2014. Comparison of the toxicity of fluoridation compounds in the nematode *Caenorhabditis elegans* Environmental Toxicology and Chemistry January 33(1):82-8.
- Switzer JA, Rajasekharan VV, Boonsalee S, Kulp EA, Bohannon EW. 2006. Evidence that monochloramine disinfectant could lead to elevated Pb Levels in drinking water. Environ Sci Technol 40:3384-7.
- UK Water Industry Research Limited. 2011. Alternatives to Phosphate for Plumbosolvency Control
- Xiang Q, et al. 2003a. Effect of fluoride in drinking water on children's intelligence. Fluoride 36: 84-94.
- Xiang Q, et al. 2003b. Blood lead of children in Wamiao-Xinhuai intelligence study. Fluoride 36: 198-199.