

MESMAP-5 PROCEEDINGS BOOK
April 24th - 26th, 2019 / Cappadocia – Turkey

**The Fifth International Mediterranean Symposium on
Medicinal and Aromatic Plants**

MESMAP – 5
PROCEEDING BOOK
ABSTRACTS & FULL PAPERS

April 24th – 26th, 2019

Cappadocia – Turkey

ISBN: (PDF)

EDITORS

Prof. Dr. Nazım ŞEKEROĞLU

Assist. Prof. Dr. Sevgi GEZİCİ

Cover Design by

Mustafa ÖZYAĞLI

(Graphic Designer, Kilis-Turkey)

MESMAP – 5 Logo Designed by

Abdulgani BARUTÇU

(Kaju Digital, Kilis-Turkey)

CHAIR OF MESMAP-5

Prof. Dr. Nazım ŞEKEROĞLU

Department of Food Engineering, Faculty of Engineering and
Architecture,
Kilis 7 Aralık University, Kilis, TURKEY
President of AMAPMED
General Coordinator of GOFMAP

CO-CHAIRS OF MESMAP-5

Prof. Dr. İlkay ERDOĞAN ORHAN

Department of Pharmacognosy, Faculty of Pharmacy,
Gazi University, Ankara, TURKEY

Prof. Dr. Anake KIJJOA

Instituto de Ciências Biomédicas Abel Salazar & Interdisciplinary
Centre of Marine and Environmental Research (CIIMAR),
Universidade do Porto, PORTUGAL

SYMPOSIUM SECRETARY

Dr. Sevgi GEZİCİ

Department of Molecular Biology and Genetics,
Kilis 7 Aralık University, TURKEY

HONORARY BOARD

Prof. Dr. Mustafa Doğan KARACOŞKUN, Turkey

Prof. Dr. Sumon SAKOLCHAI, Thailand

Prof. Dr. Ricardo LOURO BERBARA, Brazil

Prof. Dr. Hirofumi KAI, Japan

Prof. Dr. Masahiko KIKUCHI, Japan

Prof. Dr. Paiboon DAOSODSAI, Thailand

Prof. Dr. M. Hakkı ALMA, Turkey

Bekir KARACABEY, Turkey

INTERNATIONAL ORGANIZING COMMITTEE

PROF. DR. ILKAY ERDOGAN ORHAN (TURKEY)
PROF. DR. ANAKE KIJOA (PORTUGAL)
PROF. DR. ERDEN BANOĞLU (TURKEY)
PROF. DR. UFUK KOCA CALISKAN (TURKEY)
PROF. DR. TAKASHI WATANABE (JAPAN)
PROF. DR. MONICA HANCIANU (ROMANIA)
PROF. DR. MASAMI OTSUKA (JAPAN)
PROF. DR. MARIA DAGLIA (ITALY)
PROF. DR. ÇİMEN KARASU (TURKEY)
PROF. DR. ANA PAULA DE ALMEIDA (BRAZIL)
PROF. DR. RAMAN DANG (INDIA)
PROF. DR. FRANCESCO EPIFANO (ITALY)
PROF. DR. KHAETTHAREEYA SUTTHANUT (THAILAND)
PROF. DR. MURAT TUNÇTÜRK (TURKEY)
ASSOC. PROF. DR. MIKIYO WADA (JAPAN)
ASSOC. PROF. DR. KOJI SUGIMURA (JAPAN)
ASSOC. PROF. DR. KOULA DOUKANI (ALGERIA)
ASSOC. PROF. DR. MIKAKO FUJITA (JAPAN)
ASSOC. PROF. DR. F. SEZER ŞENOL (TURKEY)
ASSIST. PROF. DR. SEVGI GEZICI (TURKEY)
DR. LAMIA HAMROUNI (TUNISIA)
DR. AHMAD ALI (INDIA)
DR. FARUK KARAHAN (TURKEY)
MSc. DIANA MIHAELA DUMITRASCU (ROMANIA)
MSc. HASNA BOUHENNI (ALGERIA)
MSc. LÜTFİ NOHUTÇU (TURKEY)
MSc. DECHA KUMLA (PORTUGAL)

INTERNATIONAL SCIENTIFIC COMMITTEE

NAME – SURNAME

COUNTRY

A.K. TRIPATHY	INDIA
AHMET ÖZDEMİR	TURKEY
ÁKOS MÁTHÉ	HUNGARY
ALBAN IBRALIU	ALBENIA
ANA PAULA DE ALMEIDA	BRAZIL
ANAKE KIJJOA	PORTUGAL
AYLA KAYA	TURKEY
BERAAT ÖZÇELİK	TURKEY
BILJIANA BAUER	MACEDONIA
CHARAFEDDINE JAMA	FRANCE
DOUGLAS CHAVES	BRAZIL
DURMUŞ ALPASLAN KAYA	TURKEY
ELACHOURI MOSTAFA	MOROCCO
EMİNE BAYRAM	TURKEY
ESVET AKBAŞ	TURKEY
FİLİZ AYANOĞLU	TURKEY
FİLİZ MERİÇLİ	TURKEY
GÜLSÜM YALDIZ	TURKEY
HAMDİ TEMEL	TURKEY
HARI PRASAD DEVKOTA	JAPAN
HUMA NAZ	INDIA
HYTHAM AHMED	EGYPT
IBRAHIM TUMEN	TURKEY
ILKAY ERDOGAN ORHAN	TURKEY
IRFAN ALI KHAN	INDIA
IVAN SALAMON	SLOVAKIA
JACKSON ROBERTO GUEDES DA SILVA ALMEIDA	BRAZIL
JANAR JENIS	KAZAKHSTAN
JIANBO XIAO	CHINA
JUTTA LUDWIG-MULLER	GERMANY
KENSAKU ANRAKU	JAPAN
KHAETTHAREEYA SUTTHANUT	THAILAND
KOULA DOUKANI	ALGERIA
KRYSTYNA SKALICKA-WOZNIAK	POLAND
LAMIA HAMROUNI	TUNISIA
MADALENA MARIA DE MAGALHÃES PINTO	PORTUGAL

NAME – SURNAME

COUNTRY

MARINA SPÎNU	ROMANIA
MARYNA KRYVTSOVA	UKRAINE
MASAKI FUJITA	JAPAN
MEHLİKA DİLEK ALTINTOP	TURKEY
MENŞURE ÖZGÜVEN	TURKEY
MIRIAM DE BARCELLOS FALKENBERG	BRAZIL
MOHAMMAD RAIS MUSTAFA	MALAYSIA
MOHAMMAD SANAD ABU- DARWISH	JORDAN
MOHAMMED HMAMOUHI	MOROCCO
MONICA HANCIANU	ROMANIA
MURAT KARTAL	TURKEY
NAZLI ARDA	TURKEY
NİLGÜN GÖKTÜRK BAYDAR	TURKEY
NİLGÜN ÖZTÜRK	TURKEY
NOUREDDINE DJEBLI	ALGERIA
NUTTINEE TEERAKULKITTIPONG	THAILAND
RACHID BELHATTAB	ALGERIA
RÜVEYDE TUNÇTÜRK	TURKEY
S.V. SHUKLA	INDIA
SAFIYE EMİİRDAĞ ÖZTÜRK	TURKEY
SALAH AKKAL	ALGERIA
SEVGI GEZICI	TURKEY
SIVARAM VENKATARAMANA GOWDA	INDIA
SONAL DUBEY	INDIA
SURESH KUMAR MALHOTRA	INDIA
TAKASHI WATANABE	JAPAN
TANWEER ALAM	INDIA
TETSUO YAMASAKI	JAPAN
THEERA RITTIROD	THAILAND
TIDA DETHOUP	THAILAND
TULAY ILTER BAKIREL	TURKEY
YAOWERED CHULIKHIT	THAILAND
YAVUZ BAĞCI	TURKEY
YOSHINARI OKAMOTO	JAPAN
YUSUF BARAN	TURKEY
YÜKSEL KAN	TURKEY
ZORA DAJIC STEVANOVIC	SERBIA

***Alphabetically ordered**

Dear colleagues,



Having respected scientific board and organizing committee members from all over the world, MESMAP Symposium series started in 2013. The first Mediterranean Symposium on Medicinal and Aromatic Plants (MESMAP-2013) was held on April 17-20, 2013 in Gazimagosa (Famagusta), Turkish Republic of Northern Cyprus (TRNC), which was organized by Faculty of Pharmacy, Eastern Mediterranean University (EMU) joint with AMAPMED (Association of Medicinal and Aromatic Plants of the Mediterranean).

MESMAP-2 Symposium was held on April 22-25, 2015 in Antalya – TURKEY, which was organized by academicians from Gazi University (TURKEY), Gaziantep University (TURKEY), Kilis 7 Aralık University (TURKEY), Yüzüncü Yıl University (TURKEY), Association of Pharmaceutical Teachers of India (APTI – INDIA) joint with AMAPMED (Association of Medicinal and Aromatic Plants of the Mediterranean). INDUSTRIAL CROPS AND PRODUCTS JOURNAL with high impact factor from ELSEVIER group published a special issue covering some of the full papers selected after scientific evaluation.

MESMAP-3 Symposium which was held on April 13-16, 2017 in Girne (Kryneia) – Turkish Republic of Northern Cyprus (TRNC), was the third event of MESMAP symposium series on Medicinal and Aromatic Plants. After scientific evaluation selected full papers published in Indian Journal of Pharmaceutical Education and Research (IJPER), indexed with THOMSON REUTERS. MESMAP-4 Symposium, which was held on April 18-22, 2018 in Sherwood Breezes Resort Hotel Antalya – Turkey, was the fourth event of MESMAP symposium series on Medicinal and Aromatic Plants. MESMAP Symposiums provide a platform for herbal medicines, botany, plant biotechnology, ethnobotany, phytopharmacology, pharmacognosy, food, agriculture and forestry, plant biology, phytochemistry and aromatherapy.

That symposium was the fifth meeting series of MESMAP, and you can find abstracts of all the scientific works presented in MESMAP-5 in this PROCEEDING BOOK. We are proud to announce that selected full papers will be published in the contracted journals of the symposium after scientific evaluation. We are happy to invite MESMAP-5 participants to submit their full papers which were presented at the symposium ‘Molecules’, ‘Annals of Phytomedicine’, International Journal of Agriculture, Environment and Food Sciences’, ‘Journal of Institute of Science and Technology-Iğdır University’, and ‘Current Perspectives on Medicinal and Aromatic Plants (CUPMAP)’.

We would like to thank for their sincere supports of Turkish General Directorate of Forestry, TURKISH AIRLINES, Austrian Drug Screening Institute, Kilis 7 Aralık University, Khon Kaen University, Kumamoto University, Iğdır University, Rural Federal University of Rio de Janeiro (UFRRJ)-Brazil, AMAPMED, Association of Pharmaceutical Teachers of India, Cosmetic Producers and Researchers Associations, Talya Herbal Company, and MASMANA Olive Oils and all the other supporters.

Organizing Committee hope that MESMAP-5 Symposium participants would have an amazing experience and unforgettable memories to take back their homes, and would like to thank for all MESMAP-5 participants for their valuable contributions. We would like to remind you that MESMAP Symposium series will be organized every year. Hope to meet you in the sixth meeting series of MESMAP-5 in 2020 spring.

Sincerely,
Symposium Chairman

Prof. Dr. Nazım ŞEKEROĞLU

Department of Food Engineering, Faculty of Engineering and Architecture,
Kilis 7 Aralık University, Kilis, TURKEY

President of AMAPMED

General Coordinator of GOFMAP

www.nazimsekeroglu.com ; www.mesmap.org

MESMAP - 5 SUPPORTERS & SPONSORS

Organizing committee would like to thank sincerely to the sponsors for their valuable support;

- General Directorate of Forestry, Republic of Turkey
- Kilis 7 Aralık University, Turkey
- Kumamoto University, Japan
- Khon Kaen University, Thailand
- Iğdır University, Turkey
- Rural Federal University of Rio de Janeiro (UFRRJ), Brazil
- Instituto Idehia, Portugal
- KUAD – Cosmetic Producers and Researchers Associations
- TURKISH AIRLINES
- ADSI – Austrian Drug Screening Institute
- MUSIAD – Independent Industrialists and Businessmen's Association, Japan
- APTI – Association of Pharmaceutical Teachers of India
- AMAPMED – Association of Medicinal and Aromatic Plants of Mediterranean
- AMAPSEEC – Association for Medicinal and Aromatic Plants of Southeast European Countries
- SILAE – Società Italo-Latinoamericana di Etnomedicina
- CTFC – Centre Forestal Centre Tecnològic Forestal de Catalunya
- INRGREF – National Research institute of Rural Engineering, Water and Forests
- FIARNS09 – Free international Association of Researchers on Natural Substances 2009
- ESCORENA – The European System of Cooperative Research Networks in Agriculture
- Societa Botanica Italiano
- Iranian Medicinal Plants Society
- Isık Spices and Herbs Company
- Talya Herbal Company
- Kilizi Organic Olive Oil
- MASMANA Olive Oils

**‘Selected full papers will be published in the official journals of
MESMAP- 5 after the symposium’**

Molecules

(Publication fee
will be covered by
authors)

<https://www.mdpi.com/journal/molecules>

**Annals of
Phytomedicine**

(Publication fee
will be covered by
authors)

<http://ukaazpublications.com/publications/index.php>

**International
Journal of
Agriculture,
Environment and
Food Sciences**

(Free)

<http://dergipark.gov.tr/jaefs>

**Journal of
Institute of
Science and
Technology, Iğdır
University**

(Free)

<http://fbed.igdir.edu.tr/>

**Current
Perspectives on
Medicinal and
Aromatic Plants
(CUPMAP)**

(Free)

<http://www.cupmap.org/>

Contents

International Organizing Committee.....	IV
International Scientific Committee	V
Welcome Speech from Chair of Symposium	VII
MESMAP-5 Supporters & Sponsors.....	VIII
Contracted Journals	IX
Contents.....	X

PLENARY & INVITED LECTURES

Planery Lecturer: Prof. Dr. Gülaçtı TOPÇU

Faculty of Pharmacy, Bezmialem Vakif University, İstanbul – TURKEY

Title: "Chemical Diversity of Bioactive Diterpenoids in Endemic Anatolian Lamiaceae Plants" 2

Planery Lecturer: Prof. Dr. Mohammad Rais MUSTAFA

Department of Pharmacology, Faculty of Medicine, University of Malaya, Kuala Lumpur – Malaysia

Title: "Elephant Trunk, Birds Nest, Cat Whiskers and Fatima's Dagger: Can Natural Products from Malaysia Treat Diabetes Mellitus?" 3

Invited Leturer: Dr. Thomas JAKSCHITZ

Senior Scientist and Head of Laboratory, Adsi (Austrian Drug Screening Institute Gmbh), Innsbruck – AUSTRIA

Title: "Combination Is the Key – Analytical and Biological Investigations in Phytopharmacy, Phytocosmetics and Phytonutrition" 4

Invited Lecturer: Prof. Dr. Filiz MERİÇLİ

Department Pharmacognosy, Faculty of Pharmacy, Near-East University, Nicosia – TURKISH REPUBLIC OF NORTHERN CYPRUS

Title: "Some Investigations on the Medicinal Plants Growing in Northern Cyprus" 5

Invited Lecturer: Prof. Dr. Alvaro VILJOEN

Department of Pharmaceutical Sciences, Tshwane University of Technology, Pretoria – SOUTH AFRICA

Title: "Evidence-Based African Natural Medicines - Exploring the Synergy between Ancient Wisdom and Modern Science" 6

Invited Lecturer: Prof. Dr. Monica HANCIANU

Department of Pharmacognosy-Phytotherapy, Faculty of Pharmacy, University of Medicine and Pharmacy, Iasi – ROMANIA

Title: "Neuroprotective Behavior for A Healthy Brain" 7

Invited Lecturer: Prof. Dr. Nouredine DJEBLI

Faculty of SNV, Pharmacognosy and Api-phytotherapy Laboratory, Mostaganem University, Mostaganem – ALGERIA

Title: "Can we Protect our Brain from Neurotoxicity by Natural Products" 8

Invited Lecturer: Prof. Dr. Zahia KABOUCHE

Université frères Mentouri-Constantine 1, Département de chimie, Laboratoire d'Obtention de Substances Thérapeutiques (LOST), Constantine – ALGERIA

Title: "Biological Potential of Compounds Isolated from Algerian Plants"..... 9

Invited Lecturer: Prof. Dr. Ufuk KOCA ÇALIŞKAN

Department Pharmacognosy, Faculty of Pharmacy, Gazi University, Ankara – TURKEY

Title: "Synthesis of Plant Based Hybrid Nanoflowers and Exploration in Novel Drug Discovery" 10

Invited Lecturer: Dr. Lamia HAMROUNI

National Research Institute of Rural Engineering, Water and Forestry (INRGREF), Institution of Agricultural Research and Higher Education (IRESA) University of Carthage – TUNISIA

Title: "Propagation, Improvement, Preservation and Enhancement of Forest Species in Tunisia" 11

ORAL PRESENTATIONS

Marina Spînu, Oana Pop, Mihaela Niculae, Elena Andruța Mureșan, Aurel Vasiiu, Adrian Potârniche, Constantin Cerbu, Carmen Dana Șandru, Silvana Popescu, Eموke Pall

In Vitro Salvia glutinosa Biological Effects Depend on Extraction Procedure and Cell Type ... 13

Cornelia Mircea, Denisa Batîr-Marin, andreia Corciovă, Cristina Iancu, Oana Cioanca, Flavia Burlec, Catalina Stan, Silvia Robu, Monica Hăncianu

In Vitro Antioxidant Properties of Extracts and Nanoparticles Obtained from *Equisetum* sp..... 14

Susmita Gudulkar, Usha Mukundan

Ethno-Botanical Studies of Plants Used by Tribes in Maharashtra for Treatment of Kidney Stones 15

Diana Ivanova, Alexander Tashev, Paraskev Nedialkov, Yana Ilieva, Galina Kalotova, George Angelov, Teodora Atanassova

Conifers in the Conquest of Cancer 16

Nazim Sekeroglu, Sevgi Gezici

Curcumin, a Polyphenol Derived from *Curcuma longa*: Clinical Evidence and Molecular Mechanisms in Prevention and Management of Cancer 17

Ivan Salamon, Anton Bubnov, Marianna Kharabara

Blue Essential Oil of Chamomile Variety “Lianka” With the Highest Content of α -Bisabolol and Its Importance to New Pharmaceutical Formulations 18

Zakia Boubechiche, Mounira Ouar-Korichi, Rabah Boukherroub, Alexandre Barras, Patrick Daubias, Nour-Eddine Chihib, Amina Hellal, Charafeddine Jama Chemical Composition and Biological Activities of Essential Oils of Bulbs and Bulbils from <i>Allium sativum</i> L.	19
<u>Safive Emirdağ-Öztürk</u> Semi-Synthesis of Bioactive Gypsogenin Derivatives.....	20
<u>Ciprian Ilașcă, Mihaela Niculae, Elena andruța Mureșan, Aurel VasIU, Eموke Pall, Raluca Seicheli, Constantin Cerbu, Carmen Dana Șandru, Silvana Popescu, Marina Spînu</u> Influence of an Aqueous Mint (<i>Mentha piperita</i>) Extract on Health Status and Immune Response in Young Calves.....	21
<u>Sevgi Gezici</u> Investigation of the Cytotoxic, Anticancer and Apoptotic DNA Fragmentation Properties of <i>Bongardia chrysogonum</i> L. Extracts on Brain Glioma and Lung Carcinoma Cancer Cells	22
<u>Mebarka Bouziane, Naima Hellali, Ferdous Cherifi, Bakir Abanou, Mahfoud Hadj Mahammed</u> Compounds from an Aromatic Annual Plant Growing in Arid Zone of Algeria and Screening of Its Antibacterial Activity.....	23
<u>Belgin Coşge Şenkal, Tansu Uskutođlu, Arzu Türker, Hülya Dođan</u> Impact of Different Cutting Stages on the Content, Chemical Composition and Antibacterial Activite of Essential Oils from <i>Salvia officinalis</i> L.	24
<u>Hilal Zaid, Sleman Kadan, Najlaa Bassalat, Raghad Albzoor</u> Medicinal Plants and Phytochemicals: A Source for GLUT4 Stimulators and Digestive Enzymes Inhibitors.....	25
<u>Gülsüm Yaldiz, Mahmut Çamlica, Abdurrahman Başol, Halit Aşkin</u> Morphology, Yield and Quality Characterizations of <i>Nigella sativa</i> L. and <i>Nigella damascena</i> L.	27
<u>Bushrah Karsou, Rana Samara</u> Natural Induced Resistance in Crop Plants by Palestinian Indigenous Plant Extracts Against Pests and Diseases.....	28
<u>Tuba Kuşman, Abdullah Dalar, Ömer Akgül, Gülhan Bora</u> Investigation of Antimicrobial Activity of Some Plants against <i>E.coli</i> Isolated from Blood Cultures	29
<u>Muhammad Zafar</u> The Role of Medicinal Plants Used in Traditional Medicine for Drug Discovery and Their Conservation in Pakistan.....	30

- Gülsüm Yıldız, Yavuz Bülent Köse, Mine Kürkçüoğlu**
Essential Oil Composition of *Centranthus longiflorus steven* subsp. *longiflorus* from Turkey 31
- Carmen Dana Sandru, Silvana Popescu, Mihaela Niculae, Elena andruța Mureșan, Brudașcă Gheorghe Florinel, Eموke Pall, Adrian Potarniche, Constantin Cerbu, Aurel Vasiu, Marina Spînu**
Influence of Antioxidant Capacity of Plant Extracts on Enhancement of Adaptive Cell-Mediated Response in Herbivores..... 32
- Amruta Kothare, Vinal Pardhi, Prajкта Chivte, Sakharam Muley, Tejashree Shanbhag**
Bromelain: A Possible Phytoremedy towards Breast Cancer 33
- Zglinska Klara, Matusiewicz Magdalena, Daniluk Karolina, Balaban Jasmina, Niemiec Tomasz**
Chemical Composition, Antioxidant Activity and Effect on Human Fibroblast Visibility of Extracts from Oleaster (*Elaeagnus umbellate*) Fruit Collected in Poland 34
- Fatma Güneş**
Wild Plants Used as Food in Edirne Central District and Villages..... 35
- Naima Hellali, Mebarka Bouziane, Mahfoud Hadj Mahammed**
Correlation between Chemical Composition and Antioxidant Activity of Essential Oils from Six Aromatic Medicinal Plants Growing in Illizi and Ghardaïa (Southern Algeria)..... 36
- Oya Ustuner, Ceren Anlas, Tulay Bakirel, Fulya Ustun Alkan, Belgi Diren Sigirci, Seyyal Ak, Huseyin Askin Akpulat**
Thymus sipyleus Boiss. Subsp. *rosulans* (Borbas) J alas Extracts as Alternative therapeutic Agents for the Treatment of Skin Injury 37
- Reem Hassan Ahmed, El khatim Hassan Abdelgadir**
Acute and Dermal Toxicity Studies of *Adansonia digitata* L. Seeds Oil on Wistar Albino Rats 38
- Taner Özcan**
Determination of Hybridization in *Nepeta X Tmolea* Boiss. Using Different Approaches..... 39
- Oufkir Sara, Ait Laaradia Mehdi, Laadrawi Jawad, Aboufatima Rachida, Chait Abderrahmane, Sokar Zahra**
Anti-Fertility and Abortifacient Effect of Aqueous Sprouted Seeds Extract of *Trigonella foenum graecum* (L.) in Mice..... 40
- Ceylan Dönmez, Ufuk Koca Çalışkan, Petek Korkusuz, Özge Boyacıoğlu**
From A Traditional Remedy to Modern Therapy; *In Vivo* Antihemorrhoidal Study of *Malva sylvestris* L. 41
- Nazim Sekeroglu, Sevgi Gezici, Ceren Serin Tanrıöver, Fatih Yayla**
Anticancer, Antiproliferative and Cellular DNA Fragmentation Activities of *Astragalus elongatus* subsp. *nucleiferus* on Human Cancer Cells 42

<u>Sevgi Gezici, Murat Tuncturk, Lütfi Nohutcu, Nazim Sekeroglu</u> Assessment of Cholinesterase and Tyrosinase Enzymes Inhibitory, Phytochemical Composition and Antioxidant Properties of <i>Mentha spicata</i> L.	43
<u>Zubeyir Elmazoglu, Edgar Rangel-López, Abel Santamaria, Cimen Karasu</u> Neuroprotective Effects of Active Phytoconstituents against A β ₁₋₄₂ -Mediated Injury in High Glucose-Induced Primary Hippocampal Neurotoxicity: <i>In Vitro</i> Hyperglycemic Alzheimer's Disease Model	44
<u>Marina Spînu, Laura andreea Rusu, Mihaela Niculae, Elena Andruța Mureșan, Jovan Bojkovski, Aurel Vasii, Eموke Pall, Diana Ioana Olah, Constantin Cerbu, Carmen Dana Șandru</u> Sea Buckthorn Treatment Influence on the Multi-Effector Immune Response in Swine Exposed to Potentially Pathogenic Microbiome	45
<u>Ayşe Esra Karadağ, Ayşegül Çaskurlu, Fatma Tosun</u> <i>In Vitro</i> Anti- <i>Helicobacter pylori</i> and Antimycobacterial Activity Evaluation of Plants from Turkey	46
<u>Argentieri Maria Pia, D'Addabbo Trifone, Avato Pinarosa</u> Nematicidal Activity of Essential Oils from Lamiaceae.....	47
<u>N. Gökçe Ulusoy, Safiye Emirdağ-Öztürk</u> New Bioactive Gypsogenin Compounds	48
<u>Elsadig A. Eltayeb</u> Levels (Production) and Biological Activity of Solamargine and Solasonine from <i>Solanum incanum</i>	49
<u>Emrah Şirin</u> Karyomorphology of Some Medicinal <i>Lepidium</i> Species (Brassicaceae) from Turkey.....	50
<u>Bouamama H., Nouaa S., Benharref A.</u> Chemical and Biological Valorisation of <i>Cedrus atlantica</i>	51
<u>Antoaneta Trendafilova, Suleyman Yur, Fatih Goger, Miroslav Rangelov, Milka Todorova, Ina Aneva, Temel Ozek, Gulmira Ozek</u> Phytochemical Study and Evaluation of Inhibitory Potential of <i>Angelica panicicii</i> Vandas. Essential Oils and Hexane Extracts against Enzymes Involved in Alzheimer Disease and Type II Diabetes	52
<u>Muhammad Ajmal Shah, Wantana Reanmongkol, Pharkphoom Panichayupakaranant</u> <i>In Vivo</i> Antidiabetic Activity of Rhinacanthins Enriched Extract Prepared by Microwave assisted Green Extraction Process.....	53
<u>Sultan Pekacar, Hasya Nazlı Ekin, Nilüfer Orhan, Burçin Özüpek, Didem Deliorman Orhan</u> Enzyme Inhibitory, Antioxidant Activities and Phytochemical Studies on <i>Viscum album</i> Samples Collected from Different Host Plants	54

Enkelejda Goci, Entela Haloci, Jona Berberi

Encapsulation of *Capsicum frutescens* Extracts With β – Cyclodextrin and 2-Hydroxypropyl – β – Cyclodextrin 55

Tansu Uskutođlu, Belgin Coşge Şenkal, Cüneyt Cesur, Hülya Dođan

Investigation of the Antimicrobial Activity of Extracts Obtained from Tesbi (*Styrax officinalis* L.) 56

Biljana Bauer

Ethnomedicinal Study of *Cornus mas* L. from the Folk and Traditional Medicine of Republic of Macedonia..... 57

Saloni Jain, Tabassum Khan

Cytotoxic Flavonoids from the Ethyl Acetate Extract of the Aerial Roots of *Ficus benghalensis* Linn 58

Sagarika Damle, Archana Thite

Seasonal Edible Plants of Maharashtra (India) - A Potential Medicinal Treasury..... 59

Eda Şensu, Kadriye Nur Kasapođlu, Mine Gültekin-Özğüven, Evren Demircan, Sevcan Erşan, Ayla Arslaner, Beraat Özçelik

Berberis Plants With Orange, Red and Purple Fruits: Antioxidant and Anti-Hypertensive Activities..... 60

Ezgi Şenol, Beraat Özçelik

Evaluation of the Reliability of Plants Usage in Foods: A Review 61

Büşra Karpuz, Fatma Ayaz, Esra Emerce, Bilge Şener, İhsan Çalıř, Hayri Duman, M. Iqbal Choudhary, Nurgün Küçükboyacı

In Vitro Cytotoxicity and *in Silico* Mutagenicity Assessments of the Isolated Compounds from *Sideritis germanicopolitana* Bornm. subsp. *germanicopolitana* 62

Fatma Ayaz, Mert İlhan, Esra Küpeli Akkol, Nezhun Gören, İhsan Çalıř, Muhammad Iqbal Choudhary, Nurgün Küçükboyacı

Evaluation of the *In Vitro* Wound Healing Activity of Isolated Constituents from *Chrysophthalmum montanum* (Dc.) Boiss. 63

Bouhenni Hasna, Doukani Koula, Tabak Souhila

Anti-Helicobacter Pylori Activity of Two Medicinal Plants; Fenugreek (*Trigonella foenum-graecum* L.) and Cumin (*Cuminum cyminum* L.) 64

Muhammad Zafar, Moona Nazish

Ethnomedicinal and Aromatic Salt Tolerant Plants Use Value in Khewra Salt Mine, Punjab, Pakistan 65

Zine El Abidine Ababsa, Wwahiba Kara Ali, Kamel Medjroubi, Salah Akkal, Nacira Abidli

Chemical Characterization and Biological Study of the Species *Senecio cineraria*..... 66

Djamila Benouchene, Inès Bellil, Douadi Khelifi Phenolic Contents and Anti-Oxidant Activities of <i>Abies numidica</i> Leaves Extracts	67
Koula Doukani, Brahim Kamel Louacini Physicochemical and Phytochemical Characterization and Antibacterial Activity of Spineless Cactus (<i>Opuntia ficus indica</i> F. <i>inermis</i>) Cladodes in the Region of Tiaret (Algeria)	68
Asih Triastuti, Oktavia Indrati, Farida Hayati Development of Microemulsion Containing <i>Plantago major</i> Extracts: Formulation and Evaluation of Topical Anti-Inflammatory Activities	69
Albert Katchborian Neto, Wanderleya Toledo Dos Santos, Ana Cláudia Chagas De Paula, Michael Murgu, Marisi Gomes Soares, Daniela Aparecida Chagas-Paula Untarget Metabolomics Analyses of the Neuroprotective Potential of Ayahuasca on Human Neuroblastoma Cells	70
Iram Nizam, M.Mushfiq Antioxidant Activity of the Fruit Extracts of <i>Withania somnifera</i> and <i>Solanum nigrum</i>	71
Mehtap Kılıç, Erdal Kaya, Bilge Şener Evaluation of Some Biological Activities of the Bulbs of <i>Corydalis solida</i> (L.) Clairv. ssp. <i>incisa</i> Liéden Species Growing in Turkey	72
Ceren Anlas Medicinal Plants Used in the Treatment of Skin Disorders in Turkish Veterinary Medicine	73
Faruk Karahan, Cumhuri Avşar Leaves Antimicrobial Activity of Some <i>Glycyrrhiza</i> L. Species	74
Ufuk Koca-Çalışkan, Elif Burcu Bali, Demet Erdönmez, Müjgan Özfenerci Quorum Sensing Inhibition Activity of Volatile Oils Applied in Aromatherapy	75
Pamela Jha, Renitta Jobby, Sonit Kumari, Rasika Ghodmare, Nitin Desai, Ahmad Ali Antioxidant, Antiglycation and Antidiabetic Potential of a Natural Sweetener, <i>Stevia rebaudiana</i>	76
Grozi Delchev Changes in the Weed Species in Fields of Milk Thistle, Coriander, Chickpea and Forage Pea Which Are Sown on Damaged By Frost Areas of Winter Oilseed Canola.....	77
Nihal Karakaş, Mehmet Evren Okur, Ayşe Esra Karadağ, Şükran Kültür, Betül Demirci Investigation of Volatile Components and Various <i>in Vitro</i> Biological Activities of <i>Tilia tomentosa</i> (Silver Linden) Moench Flowers	78
Ahmad Ali, Additiya Paramanya Antidiabetic and Antiglycating Properties of <i>Spirulina platensis</i>	79

<u>Aizhamal Baiseitova, Zhamilya Abilova, Janar Jenis</u> Phytochemical Profiles and Biological Activities of Endemic Medicinal Plants from Kazakhstan	80
<u>Tuğba Günbatan, Hasya Nazlı Ekin, Burçin Özüpek, Sultan Pekacar, İlhan Gürbüz, Didem Deliorman Orhan</u> Investigation of the Effect of Seven Plant Species on Enzymes That associated With Diabetes and Obesity.....	81
<u>Ferit Özen, Gülsüm Yaldiz, Mahmut Çamlıca</u> The Effects of Nitrogen and Farm Manure Applications on Essential Oil Content of Dill (<i>Anethum graveolens</i> L.)	82
<u>Archana Thite, Sagarika Damle</u> Elemental Analysis of Leafy Vegetables as Nutritional & Medicinal Benefactors	83
<u>Rüveyde Tunçtürk, Lütfi Nohutçu, Ünal Karik, Murat Tunçtürk</u> Chemical Analysis of Wild Plant <i>Inula helenium</i> subsp. <i>pseudohelenium</i> Grierson Growing in Van Lake Around	84
<u>Vikneswaran Murugaiyah, Ning Tiang, Zurina Hassan</u> Neuroprotective and Cognitive Enhancement Potentials of Xanthone-Enriched Fraction of <i>Garcinia mangostana</i> and A-Mangostin.....	85
<u>Ataman Bilge Sari</u> Medicinal Plants as Immunostimulants in Fish Farming	86
<u>Ahmet Yenikalayci, Kemal Gül, Mahmut Güneş</u> Studies on Medicinal and Aromatic Plants under the Farmers Conditions in the Province of Kayseri	87
<u>Aurel VasIU, Mihaela Niculae, Elena Andruța Mureșan, Brudașcă Gheorghe Florinel, Eموke Pall, Constantin Cerbu, Marina Spînu, Silvana Popescu, Carmen Dana Șandru</u> Solvent Depending Immunological Activity of Nettle Plant Extracts in Sheep with Contagious Ecthyma.....	88
<u>Kazunori Shimagaki, Ryoko Koga, Haruna Fujino, Halil Ibrahim Ciftci, Minami Yamamoto, Hiroshi Tateishi, Masami Otsuka, Yoshihiro Yamaguchi, Mikako Fujita</u> HIV-2 VPX/VPR Regulate their Stability Using Two Motifs.....	89
<u>Demush Bajraktari, Biljana Bauer, Aneta Dimitrovska, Zoran Kavrakovski, Lulzimzeneli</u> Analysis of Correlation between Heavy and Essential Metals in <i>Salix alba</i> L. (Fam. Salicaceae) in Highly Polluted Area in Kosovo	90
<u>Priya Sundarrajan, Sangeetha Shetty</u> Synthesis and Characterization of Silver Nanoparticles Using Various Plant Extracts and their Various Environmental Applications	91

<u>Lalita Piarasd, Panomkorn Kwakhong, Rachadaporn Benchawattananon, Kanyarat Holasut</u> Microwave Hydro-Diffusion and Gravity of Essential Oil from <i>Zingiber officinale</i> Rosc.	92
<u>Zehra Aydin Bek, Zubeyir Elmazoglu, Berna Goker, Berivan Bitik, Cem Nuri Aktekin, Cimen Karasu</u> Therapeutic Potential of Olive Leaf Polyphenols (ZeyEX®) in Human Osteoarthritic Chondrocytes <i>In Vitro</i>	93
<u>Nuttinee Teerakulkittipong, Yanisa Chanthophat, Papawarin Han-gla, Phatcharat Lueangworachartkul, Suthabodee Muongmee, Monsiri Chetkhtkran, Suchada Sukontanon</u> Efficacy and Safety Study of the New Formulation of Herbal Coffee on Weight Control	94
<u>Mihaela Niculae, Daniela Hanganu, Ilioara Oniga, Daniela Benedec, Radu Giupana, Carmen Dana Sandru, Marina Spinu</u> Phytochemical Profile and Antimicrobial Potential of Extracts Obtained from <i>Thymus marschallianus</i> Willd	95
<u>Tuna Uysal, Ela Nur Şimşek Sezer, Meryem Bozkurt</u> Headspace Solid Phase Microextraction (Hs-Spme) Analysis of <i>Sideritis ozturkii</i> Aytac & Aksoy	96
<u>Nabila Slougui, Ahlem Tlili, Hadjer Bentayeb, Mahfoud Hadj Mahammed</u> Antibacterial Activity and Composition of Essential Oils of Leaves and Basil Flowers of Three Different Growing Seasons under Arid Climate.....	97
<u>S. Ayşe Celik, İrem Ayran, Sefa Akbulut, Rabia Serpil Günhan, Asuman Kan, Yüksel Kan</u> The Essential Oil Yield and Compositions of Tarragon (<i>Artemisia dracuncululus</i> L.) Cultivated in Bayburt.....	98
<u>Rachadaporn Benchawattananon, Kanyarat Holasut</u> The Effectiveness of Thai Traditional Herbal Decoction Consumed in the Northeast of Thailand.....	99
<u>Thatsanand Xayavong, Theera Rittirod</u> Stability Study of Oseltamivir Extemporaneous Suspension.....	100
<u>Ebru Yilmaz, Deniz Çoban, Murat Er</u> Effects of Olive Leaf (<i>Olea europea</i> L.) Extract on the Growth Performance and Some Blood Parameters of Common Carp (<i>Cyprinus carpio</i>)	101
<u>Narin Sadıkoğlu, Mehmet Efe</u> Leaf Anatomy of the Genus <i>Origanum</i> L. (Labiatae) in Turkey	102
<u>Ela Nur Şimşek Sezer, Meryem Bozkurt, Nihal İlhan, Tuna Uysal</u> Characterization of Volatile Compounds of <i>Mentha pulegium</i> via HS-SPME.....	103

- Awatif Boumaza, Messaouda Khallaf, Mouna Cherairia**
Cytotoxic and Genotoxic Effects of Water Extracts of *Zygophyllum cornutum* and *Atractylis gummifera* Using *Allium cepa* assay 104
- Yigit Gunes, Tulay Bakirel, Oya Ustuner**
Adverse Effects of Conventional Drug-Medicinal Plant Interactions in Veterinary Medicine ... 105
- Meryem Bozkurt, Ela Nur Şimşek Sezer, Nihal İlhan, Tuna Uysal**
Determination of Volatile and Semi-Volatile Oil Content of *Isatis tinctoria* L. via SPME-GC-MS 106
- Murat Tuncurk, Tamer Eryigit, Ruveyde Tuncurk, Lütfi Nohutçu**
The Chemical Composition of *Delphinium szowitsianum* Boiss. Species Spreading in Van Region 107
- Emine Saliha Toprak, Hasan Hüseyin Yiğitbaşı, Himmet Özcan, İrem Ayran, Sadiye Ayşe Çelik, Hayal Akyıldırım Beğen**
The Essential Oil Yield and Compositions of Summer Savory (*Satureja hortensis* L.) Collected in Turkey 108
- İrem Ayran, S.Ayşe Çelik, Asuman Kan, Yüksel Kan**
The Essential Oil Yield and Compositions of *Nigella sativa* L. and *Nigella damascena* L. Cultivated in Konya Ecological Conditions 109
- Rukiye Özkaya, Hakan Çetinkaya, İbrahim Samet Gökçen, Mehmet Koç**
The Role and Use of Olive Leaves as a Functional Food 110
- İbrahim Samet Gökçen, Hakan Çetinkaya, Birhan Kunter, Nurhan Keskin**
As A Good Source of Melatonin: Grape and Grape Products 111

POSTER PRESENTATIONS

- Višnja Oreščanin, Štefica Findri Guštek, Josipa Rodić**
The Efficacy of New Bioapifit® Anti-Hemorrhoidal Ointment Versus Standard Treatment Protocol 113
- Nadjia Hamlat, A. Hassani, S. Boubekur**
Antioxidant Activity and Free Aglycone Compounds in Aerial Parts of Three Species of *Pistacia* Growing in Algeria 114
- Ana Flavia Burlec, Łukasz Pecio, Mariusz Kowalczyk, Cornelia Mircea, Oana Cioancă, Andreia Corciovă, Alina Nicolescu, Cătălina Stan, Wiesław Oleszek, Monica Hăncianu**
Chemical Profile and Antioxidant Activity of New Selective *Zinnia elegans* Fractions..... 115
- Zine El Abidine Ababsa, Mohamed Tahar Derouiche, Jean Pierre Mahy, Kamel Medjroubi, Salah Akkal**
In-Vivo Anti-ulcerative and Antidiarrheal Activities of the *Tetraclinis articulata* Species of the Cupressaceae family 116

Cristina Iancu, <u>Cornelia Mircea</u>, Oana Cioanca, Catalina Stan, Nina Filip, Cristinel Stan, Gabriela Tataringa, Monica Hăncianu <i>In Vivo</i> Anti-Inflammatory Effects of Some Extracts from <i>Pelargonium</i> sp.	117
<u>Ece Sözer</u>, Safiye Emirdağ-Öztürk, N. Gökçe Ulusoy, Nurettin Yaylı, Halil Ciftci, Mohamed O. Radwan, Mikako Fujita, Masami Otsuka New Gypsogenin-Amine Compounds	118
<u>Amantay A.</u>, Dyusebaeva M., Jenis J. Chemical Constituents of <i>Artemisia albicerata</i>	119
Nurlybekova A.K., <u>Amantay A.</u>, Ye Yang, Abilov Zh.A., Jenis J. Liposoluble Constituents from the Aerial Part of <i>Ligularia narynensis</i>	120
Izabela Korona-Glowniak, Diana Ivanova, Anna Malm, <u>Alexander Tashev</u>, Paraskev Nedialkov, Galina Kalotova, George Angelov Antibacterial Activity of Conifers	121
<u>Alexander Tashev</u>, Violeta Dimitrova Medicinal Plants of Bulgaria.....	122
<u>Kabouche A</u>, Bouratoua A, Kabouche Z, Al Abdulmagid A, Voutquenne-Nazabadioko L, Morjani H. Polyphenols from <i>Athamanta sicula</i> L.	123
<u>Elif Güzel</u>, Hüseyin Servi, Cansu Vatansever Antibacterial Activity of <i>Peganum harmala</i> Seed Extracts	124
<u>Elif Güzel</u>, Cansu Vatansever, Hüseyin Servi Determination of Antibacterial Activity from Seed Extracts of <i>Pistacia terebinthus</i>	125
<u>İrem Sengür</u>, Cansu Vatansever, Hüseyin Servi Antibacterial Activity of <i>Sambucus nigra</i> Seed Extracts	126
<u>İrem Sengür</u>, Hüseyin Servi, Cansu Vatansever Determination of Antibacterial Activity of <i>Linum usitatissimum</i> Seed Extracts	127
<u>Irmak Alev</u>, Cansu Vatansever, Hüseyin Servi The Antibacterial Activity of <i>Elaeagnus angustifolia</i> L. Fruit and Leaf Extracts	128
<u>Awatif Boumaza</u>, Messaouda Khallaf, Mouna Cherairia Mutagenic <i>In Vitro</i> Activity and Genotoxic Effect of <i>Zygophyllum cornutum</i> Methanolic Extract	129
<u>Boubakeur B.</u>, Khadem H., Drabo M. S., Tir Touil Meddah A. Biological activities of <i>Thymus fontanesii</i> Polyphenolic Extract: <i>In Vitro</i> and <i>In Vivo</i> Effect on Bacterial Adhesion and Colonisation	130

- Samukelisiwe Clerance Ngubane, Helene De Wet, Sandy Van Vuuren**
In Vitro Toxicological Evaluation of Some Medicinal Plants Used By Women in Northern Maputaland, South Africa 131
- Saiqa Ishtiaq, Rabia Arshad, Sairah Hafeez Kamran**
In-vitro Susceptibility Testing of Tea Tree Oil Acne Cream in Combination with Salicylic Acid and Nicotinamide..... 132
- Grozi Delchev**
Impact of Some Herbicides and Herbicide Combinations on Sowing Characteristics of Coriander (*Coriandrum sativum* L.) 133
- Fatma Ayaz, Nuraniye Eruygur, Yavuz Bağcı, H. Filiz Ayyıldız**
Profiling of Phenolic Compounds by HPLC–DAD, Antioxidant and Enzyme Inhibitory Activities of *Marrubium vulgare* L. and *Marrubium parviflorum* Fisch. & Mey. subsp. *parviflorum*..... 134
- Mahmoudi Kheira, Cheriti A., Didi Oueld El Hadj M.**
Preliminary Phytochemical Screening of Four Saharan Plants used in South West of Algeria .. 135
- Tabassum Khan, Anju Kashyap**
Phytochemistry and Cytotoxic Potential of *Litsea deccanensis* bark. 136
- Gabriela Tataringa, Ana Maria Zbancioc, Catalina Stan, Cornelia Mircea, Monica Hancianu**
In Vitro Antioxidant Prospection of Onion Oil from Romanian Source..... 137
- F. Sezer Senol Deniz, Ilkay Erdogan Orhan, İhsan Çalıř**
Profiling Enzyme Inhibitory Potential of Cycloartane Type of Triterpene Glycosides from the Roots of Three *Astragalus* Species 138
- Habiba Daroui-Mokaddem, Assia Zeghib, Fouad Zeghib, Linda Achouri, Sara Abbas, Khouloud Fradj, Amel Moussaoui, Soraya Hioun, Zahia Kabouche, Belgacem Djabri**
Antioxidant Activity and Phenolic Content of *Peganum harmala* Used in Algerian Folk Medicine..... 139
- Assia Zeghib, Habiba Daroui-Mokaddem, Fouad Zeghib, Soraya Hioun, Khouloud Fradj, Zahia Kabouche, Belgacem Djabri**
Phytochemical Screening and Antibacterial Activity of *Pituranthos scoparius* Aerial Parts Harvested from the Area of Tébessa (Algeria) 140
- Nabila Slougui, Naima Draoui, Cherifa Khaddar, Mahfoud Hadj Mahammed**
Study of the Varietal Behavior of 6 Basil Varieties (*Ocimum basilicum* L) Grown for the First Time in Algeria under Arid Climate 141
- Salima Douichene, Nessrine Ammari, Nouredine Djebli**
Neuroprotective Effect of *Hypericum thymopsis* against Chronic Exposure to Aluminum Chloride and Alzheimer's Disease..... 142

<u>Tuğba Günbatan, Mert İlhan, Esra Küpeli Akkol, İlhan Gürbüz, Hayri Duman, Ceyda Sibel Kiliç</u> <i>In Vitro</i> Enzyme Inhibitory Activity of Three <i>Ferulago</i> Species Growing in Turkey	143
<u>T.S.Mammadov, A.G.Rasulova, S.B.Bagirova</u> Comparative Analysis of Rare and Endangered Plants of Hirkan dendroflora.....	144
<u>Blagica Jovanova, Svetlana Kulevanova, Tatjana Kadifkova Panovska</u> ROS Scavenging and Reducing Capacity of Aromatic Herbs	145
<u>Rafik Bensegueni, Mounia Guergouri, Chawki Bensouici, Mustapha Bencharif</u> Antioxidant Activity of Four Nitrogen Organic Compounds: Correlation between Theoretical and Experimental Data	146
<u>Meliani Samiha</u> Study of the Hypglycemic Activity of the Common Figs of the Region of Ain Defla Algeria...	147
<u>Antoaneta Trendafilova, Miroslav Rangelov, Milka Todorova</u> Qualitative and Quantitative Determination of Mono- and Dicafeoylquinic Acids in <i>Inula</i> Species from Bulgaria	148
<u>Ilhem Semra, Mehmet Öztürk, Ahmed Kabouche, Zahia Kabouche</u> Total Phenolic Content and Antioxidant Activity of the n-Butanol Extract of <i>Mentha rotundifolia</i> from Algeria	149
<u>Inès Bellil, Djamila Benouchenne, Douadi Khelifi</u> Radical Scavenging and Anti-Bacterial Activities of Methanolic Extract from Algerian Fir (<i>Abies numidica</i>) Leaves	150
<u>Naima Hellali, Mahfoud Hadj Mahammed, Mebarka Bouziane</u> Chemical composition and Biological Activities of the Essential Oils from Two Chemotypes of <i>Cymbopogon schoenanthus</i> (L.) spreng. Growing in Southern Algeria.....	151
<u>Mounira Benteldjoune, Tarek Boudiar, Abdelhakim Bakhouch, María Del Mar Contreras, Jesús Lozano-Sánchez, Chawki Bensouici, Antonio Segura-Carretero, Zahia Kabouche</u> Evaluation of the Antioxidant Potential, Phenolic and Flavonoid Contents of the Methanol Extract of <i>Ammoides atlantica</i>	152
<u>Susmita Gudulkar, Usha Mukundan</u> Extraction of Natural Product from Plant by Using a Microwave	153
<u>Nidhi Pathak, Himanshu Dawda, Usha Mukundan</u> Ursolic Acid as a Biochemical Marker in Family <i>Myrtaceae</i>	154
<u>Orçun Cınar, Murat Tunçtürk, Tuba Seçmen, Ayşe Erdoğan, Ünal Karık</u> Determination of Essential Oil Composition, Phenolic and Flavanoid Contents of <i>Phlomis longifolia</i> var. <i>longifolia</i> Collected from Hatay District of Turkey	155

- Zehra Aydın Bek, Zubeyir Elmazoglu, Berna Goker, Berivan Bitik, Cem Nuri Aktekin, Cimen Karasu**
Effects of Mulberry (*Morus alba* L.) Leaf Extract (Immunflex®) on MMP-9, BMP-2 and COMP in Human Osteoarthritic Chondrocytes *In Vitro* 156
- Zubeyir Elmazoglu, Zeynep Inan, Cansel Keskin, Naciye Bozkurt, İhsan Calis, Cimen Karasu**
Astragaloside IV and Forsythoside B Modulate Hyperglycemia-Mediated Neurotoxicity in Primary Hippocampal Neuronal Cells 157
- M. Muharrem Özcan, Ayşegül Kırılı, Özbay Dede, İrem Ayran, S.Ayşe Çelik, Yüksel Kan**
The Essential Oil Yield and Compositions of Lemon Verbana (*Lippia citrodara* Kunth.) Cultivated in Ordu..... 158
- İrem Ayran, S.Ayşe Çelik, Asuman Kan, Yüksel Kan**
The Essential Oil Yield and Compositions of Laurier (*Laurus nobilis* L.) Cultivated in Samsun 159
- S.Ayşe Çelik, İrem Ayran, Asuman Kan, Emine Bilginoğlu, Yüksel Kan**
A Study on Essential Oil Yield and Components of Lemon Balm (*Melissa officinalis* L.) Cultivated Different Provinces in Turkey 160
- S.Ayşe Çelik, İrem Ayran, Asuman Kan, Yüksel Kan**
The Essential Oil Yield and Compositions of *Lavandula angustifolia* Miller and *Lavandula intermedia* Emeric ex Loisel. Cultivated in Bayburt Ecological Conditions 161
- Halil İbrahim Polat, Esra İnfal, Emine Saliha Toprak, S.Ayşe Çelik, İrem Ayran**
The Essential Oil Yield and Compositions of Fennel (*Foeniculum vulgare* Mill.) Cultivated in Konya, Turkey..... 162
- Hallal Nouria, Benyettou Imène, Kharoubi Omar**
The Polyphenols Extracted from the *Artemisia absinthium* L. Plant Combine Brain Oxidative Stress, Neuro-Inflammation and Neurodegeneration Induced by Heavy Metals (Aluminum and Mercury) 163
- FULL-TEXT PAPERS**
- Ahmet Yenikalayci, Kemal Gül, Mahmut Güneş**
Studies on Medicinal and Aromatic Plants under the Farmers Conditions in the Province of Kayseri 165
- Nazim Sekeroglu, Sevgi Gezici**
Curcumin, a Polyphenol Derived from *Curcuma longa*: Clinical Evidence and Molecular Mechanisms in Prevention and Management of Cancer 171
- Ferit Özen, Gülsüm Yaldiz, Mahmut Çamlica**

The Effects of Nitrogen and Farm Manure Applications on Essential Oil Content of Dill (<i>Anethum graveolens</i> L.)	177
<u>Mebarka Bouziane, Naima Hellali, Ferdous Cherifi, Bakir Abanou, Mahfoud Hadj Mahammed</u> Compounds from an Aromatic Annual Plant Growing in Arid Zone of Algeria and Screening of Its Antibacterial Activity.....	181
<u>Ela Nur Şimşek Sezer, Meryem Bozkurt, Nihal İlhan, Tuna Uysal</u> Characterization of Volatile Compounds of <i>Mentha pulegium</i> via HS-SPME.....	186
<u>Ebru Yilmaz, Deniz Çoban, Murat Er</u> Effects of Olive Leaf (<i>Olea europea</i> L.) Extract on the Growth Performance and Some Blood Parameters of Common Carp (<i>Cyprinus carpio</i>)	191
<u>Ezgi Şenol, Beraat Özçelik</u> Evaluation of the Reliability of Plants Usage in Foods: A Review	196
<u>Nabila Slougui, Naima Draoui, Cherifa Khaddar, Mahfoud Hadj Mahammed</u> Study of the Varietal Behavior of 6 Basil Varieties (<i>Ocimum basilicum</i> L.) Grown for the First Time in Algeria under Arid Climate	201
<u>Biljana Bauer</u> Ethnomedicinal Study of <i>Cornus mas</i> L. from the Folk and Traditional Medicine of Republic of Macedonia.....	205

- Asih Triastuti, Oktavia Indrati, Farida Hayati**
Development of Microemulsion Containing *Plantago major* Extracts: Formulation and Evaluation of Topical Anti-Inflammatory Activities 208
- Nabila Slougui, Ahlem Tlili, Hadjer Bentayeb, Mahfoud Hadj Mahammed**
Antibacterial Activity and Composition of Essential Oils of Leaves and Basil Flowers of Three Different Growing Seasons under Arid Climate..... 213
- Grozi Delchev**
Impact of Some Herbicides and Herbicide Combinations on Sowing Characteristics of Coriander (*Coriandrum sativum* L.)..... 217
- Grozi Delchev**
Changes in the Weed Species in Fields of Milk Thistle, Coriander, Chickpea and Forage Pea Which Are Sown on Damaged By Frost Areas of Winter Oilseed Canola..... 223
- Naima Hellali, Mebarka Bouziane, Mahfoud Hadj Mahammed**
Correlation between Chemical Composition and Antioxidant Activity of Essential Oils from Six Aromatic Medicinal Plants Growing in Illizi and Ghardaïa (Southern Algeria) 230
- Naima Hellali, Mahfoud Hadj Mahammed, Mebarka Bouziane**
Chemical composition and Biological Activities of the Essential Oils from Two Chemotypes of *Cymbopogon schoenanthus* (L.) spreng. Growing in Southern Algeria 235
- Sevgi Gezici, Murat Tuncturk, Lütfi Nohutcu, Nazim Sekeroglu**
Assessment of Cholinesterase and Tyrosinase Enzymes Inhibitory, Phytochemical Composition and Antioxidant Properties of *Mentha spicata* L. 239
- Meryem Bozkurt, Ela Nur Şimşek Sezer, Nihal İlhan, Tuna Uysal**
Determination of Volatile and Semi-Volatile Oil Content of *Isatis tinctoria* L. via SPME-GC-MS 245
- Saiqa Ishtiaq, Rabia Arshad, Sairah Hafeez Kamran**
In-vitro Susceptibility Testing of Tea Tree Oil Acne Cream in Combination with Salicylic Acid and Nicotinamide..... 250
- Tuna Uysal, Ela Nur Şimşek Sezer, Meryem Bozkurt**
Headspace Solid Phase Microextraction (Hs-Spme) Analysis of *Sideritis ozturkii* Aytac & Aksoy 253
- Ceylan Dönmez, Ufuk Koca Çalışkan, Petek Korkusuz, Özge Boyacıoğlu**
From A Traditional Remedy to Modern Therapy; *In Vivo* Antihemorrhoidal Study of *Malva sylvestris* L. 257

Bellil Inès, Benouchenne Djamila, Khelifi Douadi

Radical Scavenging and Anti-Bacterial Activities of Methanolic Extract from Algerian Fir
(*Abies numidica*) Leaves 263

Muhammad Zafar, Moona Nazish

Ethnomedicinal and Aromatic Salt Tolerant Plants Use Value in Khewra Salt Mine, Punjab,
Pakistan 267

Author List..... 287

MESMAP-6 & ISPBS-6 292

Current Perspectives on Medicinal and Aromatic Plants (CUPMAP) 293

**PLENARY
&
INVITED
LECTURES**

MESMAP-5 Draft Proceedings Book

PLENARY LECTURE

**CHEMICAL DIVERSITY OF BIOACTIVE DITERPENOIDS IN ENDEMIC
ANATOLIAN LAMIACEAE PLANTS**

Gülaçtı Topçu

Department of Pharmacognosy, Faculty of Pharmacy, Bezmialem Vakif University, Fatih-İstanbul, TURKEY

MESMAP-5 Draft Proceeding Book

PLENARY LECTURE

**ELEPHANT TRUNK, BIRDS NEST, CAT WHISKERS AND FATÍMA’S DAGGER:
CAN NATURAL PRODUCTS FROM MALAYSIA TREAT DIABETES MELLITUS?**

Mohammad Rais Mustafa

Department of Pharmacology, Faculty of Medicine and Centre for Natural Products Research and Drug Discovery (CENAR), University of Malaya, 50603, Kuala Lumpur, Malaysia.

Natural products are naturally derived metabolites and/or by-products from microorganisms, plants, or animals. These products have been exploited for human use for thousands of years continuously providing a broad and structurally diverse arsenal of pharmacologically active compounds that have been effectively utilised in the treatment of numerous diseases. More than 45 % of all marketed drugs in the world are natural products or their derivatives. Medicinal use of natural products in the treatment and prevention of diseases including diabetes has a long history compared to conventional medicine. Diabetes mellitus is one of the major public health concerns all over the world including Malaysia. Malaysia has the highest rate of diabetes in Asia and this prevalence is projected to rise to 21.6% by the year 2020. Most of diabetic patients are affected by type 2 diabetes with hyperglycemia, insulin resistance and insulin secretion defect. Despite the greater awareness and increasing number of drugs available, diabetes continues to be poorly treated with many patients succumbing to serious complications such as kidney failure, vision loss, heart diseases, and foot amputations. Natural products including herbal medicines are commonly used in Malaysia as a form of complementary and alternative therapy for the treatment of diabetes and its complications. This is attributed to Malaysia being one of the ten mega biodiversity countries in the world and having a mixed population of Malays, Chinese, Indians and indigenous tribes, each with its own ethnic health system. This talk will highlight local practices using natural products including elephant trunk (*Clinacanthus nutans*), birds nest (*Aerodramus fuciphagus*), cat whiskers (*Orthosiphon stamineus*) and Fatima’s dagger (*Marantodes pumilum*) to treat diabetes and similar scientific efforts being undertaken by researchers from the Centre for Natural Products Research and Drug Discovery (CENAR) at the University of Malaya.

Keywords: Natural products, diabetes, *Clinacanthus nutans*, *Aerodramus fuciphagus*, *Orthosiphon stamineus*, *Marantodes pumilum*

INVITED LECTURE

**COMBINATION IS THE KEY – ANALYTICAL AND BIOLOGICAL
INVESTIGATIONS IN PHYTOPHARMACY, PHYTCOSMETICS AND
PHYTONUTRITION**

Thomas A.E. Jakschitz¹; Günther K. Bonn^{1,2}

¹ *ADSI - Austrian Drug Screening Institute GmbH, Innrain 66a, 6020 Innsbruck, Austria*

² *Institute of Analytical Chemistry and Radiochemistry, University of Innsbruck, Innrain 80-82, 6020 Innsbruck, Austria*

Contact: Thomas.Jakschitz@adsi.ac.at

Phyto-sciences are playing an increasingly important role, not only because of scientific interest, but also because the strongly growing demand for nature-derived products in pharma, cosmetics and nutrition.

Compared with synthetic formulations, it is a big issue to generate effective and safe products as one has to deal with hundreds of different compounds in highly diverse concentrations. For identification of active compounds in these multi-component mixtures multiple different extracts have to be investigated by analytical technologies and biological assays to enable bioinformatic investigations with meaningful results. Once the active compounds are identified, the extraction procedure has to be optimized in terms of increasing the yield of active ingredients and eliminating the yield of disturbing/toxic compounds. The next hurdle is the production of sufficient quantities of plant material of appropriate quality. Especially in wild collections, the concentrations of the ingredients usually vary greatly, making the production of high-quality products significantly more difficult. The solution to this problem is controlled cultivation of optimized seedlings under optimized conditions like soil-composition, temperature and altitude.

This brief overview clearly figures out the importance of combined technologies at the very different development stages from an idea to an effective and safe product.

INVITED LECTURE

**SOME INVESTIGATIONS ON THE MEDICINAL PLANTS GROWING IN
NORTHERN CYPRUS**

Filiz Meriçli

*Department of Phytotherapy, Faculty of Pharmacy, Near East University, Nicosia, Turkish Republic of Cyprus,
filiz.mericli@neu.edu.tr*

Objective / Purpose: Cyprus, the third largest island of the Mediterranean Sea; is located at the crossroads between Asia, Africa and Europe. More than 1800 plants are growing wildly in Northern Cyprus, and 22 of them are endemic plants. Folk remedies and medicinal plants of Northern Cyprus are investigated by the scientists working in Faculty of Pharmacy at Near East University. The purpose of these researchs is to evaluate the medicinal plants grown on Northern Cyprus in pharmaceutical sciences, to develop the medicinal products to be patented and to gain economic value.

Material and Methods: Fatty oils of seeds of *Prunus dulcis* and *Opuntia ficus indica* were collected from different place of Northern Cyprus; obtained using supercritical CO₂ extraction method, and fatty acids were determined by GC-MS. Alkaloid content of *Colchicum pusillum* samples roots and flowers were determined by HPLC. Cladodes and fruits of *Opuntia ficus indica* were collected from Güzelyurt and phenolic compounds were analyzed by HPLC. Anticancer activities of these plants were investigated on human colon carcinoma or breast cancer cell lines. Volatile oil of *Crataegus azarolus* were investigated by GC-MS; phenolic compounds were determined by HPLC.

Results: Our results indicated that all of these plant materials can be evaluated in pharmaceutical sciences for human health; further investigations on developing the medicinal products are still going on.

Conclusion / Discussion: According to the results to develop the medicinal products to be patented has been stil going on.

Keywords: *Prunus dulcis*, *Opuntia ficus indica*, *Colchicum pusillum*, *Crataegus azarolus*, Northern Cyprus, anticancer activity.

References:

[1] Eda Becer, Duygu Yigit Hanoğlu, Hilal Kabadayı et al, (2018). The effect of *Colchicum pusillum* in humon colon cancer cells via Wnt/ β -catenin pathway, *Gene*, 686, 213-219.

INVITED LECTURE

**EVIDENCE-BASED AFRICAN NATURAL MEDICINES - EXPLORING THE
SYNERGY BETWEEN ANCIENT WISDOM AND MODERN SCIENCE**

Alvaro Viljoen

*Department of Pharmaceutical Sciences and SAMRC Herbal Drugs Research Unit, Tshwane University of
Technology, Pretoria, 175 Nelson Mandela Drive, Private Bag X680, Pretoria, 0001, South Africa*

Southern Africa harbours an impressive floral diversity and ranks as one of the most biodiverse countries in the world. Interweaved within this botanical tapestry is a cultural heritage characterised by rich indigenous knowledge systems (IKS) which have moulded one of the oldest healing modalities, African Traditional Medicines (ATM). This unique blend of medicinal plant use and IKS has created a unique research opportunity in ethnopharmacology. Over the past 20 years our group has endeavoured to provide a scientific rationale for medicinal plant use through an evidence-based research approach of traditional medicines. Several examples will be presented to demonstrate the challenging yet rewarding workflow to explore the chemistry and biological properties of the ethnomedicinal flora of South Africa. Using various *in vitro* and *in vivo* approaches, complemented by analytical methods and multivariate data analysis we aim to contribute to the fundamental research base required to convert these botanical assets into tangible consumer products. The various challenges facing translation research and the standardisation of ATMs will be highlighted.

MESMAP-5 Draft P1

INVITED LECTURE

NEUROPROTECTIVE BEHAVIOR FOR A HEALTHY BRAIN

Monica Hancianu, Oana Cioanca, Ursula Stanescu

Department of Pharmacognosy, Faculty of Pharmacy, "Grigore T. Popa" University of Medicine and Pharmacy, Iasi, Romania

Objective: Reactive oxygen species (ROS) and oxidative stress represent key factor for multiple health problems such as metabolic syndrome, neurodegeneration and cancer. The negative impact is expressed as lipid peroxidation, protein misfolding, DNA damage and mitochondrial dysfunction which lead to cell death. Neuroprotection is a major strategy for brain health and may include natural molecules with antioxidant potential. Often, neuronal alteration, subtle cognitive and behavioral changes occur long before the clinical stage of dementia symptoms. Newer trends in neuroscience imply that depression and dementia might have a common cause or they might be linked. It is thought that earlier-onset depressive symptoms are an indicator for higher risk of developing dementia later in life.

Material and Methods: Our study briefly presents the neuroprotective potential of several selective extracts and herbal compounds against ROS and on scopolamine-induced anxiety and depression animal models.

Results: Our results indicated that all of the extracts had a notable antidepressant and anxiolytic activity which lead to better memory and learning abilities for the tested rats. The intensity of the behavior changes and the antioxidant value depends on the concentration and the type of the substances included in the extracts.

Conclusion: All in all, the obtained results are suggesting the importance of herbs for neuroprotection; a moderate daily amount could possible help against oxidative damage. Diet comes most at hand when the most risk factors cannot be influenced (genes, age, sex), by adopting neuroprotective behaviors we can maintain the neuronal plasticity of our brains.

Keywords: Neuroprotection, depression, anxiety, oxidative stress

References:

[1] Singh-Manoux, A., Dugravot, A., Fournier, A., Abell, J., Ebmeier, K., Kivimaki, M., & Sabia, S. (2017). Trajectories of depressive symptoms before diagnosis of dementia: a 28-year follow-up study. *JAMA Psychiatry*, 74(7), 712-718.

INVITED LECTURE

CAN WE PROTECT OUR BRAIN FROM NEUROTOXICITY BY NATURAL PRODUCTS?

Noureddine Djebli

Laboratory of Pharmacognosy and Api-Phytotherapy (LPAP, University of Mostaganem. ALGERIA.
Djebli_n@yahoo.fr. Tel: (00213)772451866

Neurotoxicity is the damage to the brain and/or the peripheral nervous system from toxic chemicals. These chemicals include insecticides, solvents, lead, mercury, cadmium, car exhaust, formaldehyde, chlorine, phenol and thousands of others. Our environment has become polluted with large doses of these. It is difficult for the individual to avoid exposure. Symptoms for the brain toxicity are short term memory loss, loss of circulation, imbalance, and flu-like symptoms.

Researchers believe that there may be a link between neurotoxicity and progressive neurodegenerative disorders such as dementia, Parkinson's disease, amyotrophic lateral sclerosis, and multiple sclerosis. There's no doubt that we are living in a sea of neurotoxins, and it's not possible to avoid them all. But what you can do instead is reduce the total load — the total number of neurotoxic "burdens." The attentions of researchers have been inclined towards the phytochemicals, many of which have minimal side effects. Phytochemicals are promising therapeutic agents because many phytochemicals have anti-inflammatory, antioxidative as well as anticholinesterase activities. Various drugs of either synthetic or natural origin applied in the treatment of brain disorders

The aim of this lecture is to evaluate some of the medicinal plants and their active constituents that have been used in different methods induced neurotoxicity.

Keywords: Neurotoxic, heavy metals, neurodegenerative diseases, medicinal plants, mice.

INVITED LECTURE

BIOLOGICAL POTENTIAL OF COMPOUNDS ISOLATED FROM ALGERIAN PLANTS

Zahia Kabouche

Université frères Mentouri-Constantine 1, Département de chimie, Laboratoire d'Obtention de Substances Thérapeutiques (LOST), 25000 Constantine, Algeria
E-mail : zahiakabouche@gmail.com

Algeria is characterized by a rich flora. Caryophyllaceae, Amaranthaceae, Fabaceae, Euphorbiaceae and Pinaceae have been the subject of our studies. Flavonoids and terpenoids were the principal secondary metabolites isolated from the selected species. Antioxidant, antimicrobial, anti-inflammatory, anti-tyrosinase and cytotoxic activities of the isolated compounds have been established.

Keywords: Caryophyllaceae, Amaranthaceae, Fabaceae, Euphorbiaceae, Flavonoids, Terpenoids, Biological activities.

INVITED LECTURE

**SYNTHESIS OF PLANT BASED HYBRID NANOFLOWERS AND EXPLORATION
IN NOVEL DRUG DISCOVERY**

Ufuk Koca-Çalışkan

*Department of Pharmacognosy, Faculty of Pharmacy, Gazi University, Ankara, Turkey,
E-mail: ukoca@gazi.edu.tr*

Objective/Purpose: Organic–inorganic hybrid nanoflowers have received much attention due to their simple, fast and green synthesis, large surface-to-volume ratio, high efficiency, and enzyme stabilizing abilities [1-2]. Zare and co-workers reported protein-inorganic hybrid nanoflowers initially that was made of proteins (BSA, laccase etc.) and $\text{Cu}_3(\text{PO}_4)_2 \cdot 3\text{H}_2\text{O}$ [1], further, different kinds of organic molecules (such as protein, DNA, alginate etc.,) and ions (Cu^{2+} , Zn^{2+} , Mn^{2+} , Co^{2+} , etc.,) containing hybrid nanoflowers have been subsequently reported. In this review study, synthesis of novel hybrid nanoflowers exploiting plant extracts and copper (II) ions in order apply as enzyme inhibitor agents were given.

Material and Methods: Medicinal plants (*Achillea wilhelmsii*, *Ajuga chamaepitys*, *Phlomis grandiflora*, *Rosa canina*, *Solanum melongena*) that are grown in Turkey, have traditional applications against different diseases were collected and extracted according to their ethnopharmacological usage. The dried plant extracts in different concentrations were used as an organic component to synthesis hybrid nanoflowers All the organic-inorganic hybrid structures were synthesized with the collaboration of Dr. Nalan Özdemir laboratory (Erciyes University Kayseri/Turkey). The morphologies of the synthesized nanoflowers were examined using Scanning Electron Microscopy (SEM), the chemical and crystal structures of them were characterized using Fourier Transform Infrared Spectroscopy (FTIR,) and X-Ray Diffraction (XRD) analysis.

Results: Synthesized nanoflowers were analyzed with different physicochemical techniques. According to analyzed SEM images the concentration of the best shaped nanoflowers were selected for further antioxidant and enzyme inhibitory analysis. The data demonstrated that some of the hybrid nanoflowers (*Achillea wilhelmsii*, *Rosa canina*) have better antioxidant activity than the plain extracts. Moreover, some of the hybrid nanoflowers (such as *Achillea wilhelmsii*, *Ajuga chamaepitys*, *Phlomis grandiflora*) demonstrated better alfa-glicosidase inhibition activity, whereas some nanoflowers (such as *Ajuga chamaepitys*, *Phlomis grandiflora* and *Rosa canina*) alfa-amylase enzyme inhibitory activity compared to plain extracts.

Conclusion/Discussion: The synthesis of hybrid nanoflowers containing plant extracts is an important factor considering its suitability for commercial applications in pharmacy, cosmetics and various applications in industry.

Keywords: Nano flowers, hybrid nanoflowers, plant extracts, enzyme inhibition,

References

- [1] Jun Ge, Jiandu Lei & Richard N. Zare, Nature Nanotechnology volume 7, pages 428–432 (2012) Protein–inorganic hybrid nanoflowers
- [2] C. Altinkaynak, S. Tavlasoglu, N. Özdemir, I. Ocsoy, Enzyme MicrobTechnol, 93-94, (2016) 105-112.

INVITED LECTURE

PROPAGATION, IMPROVEMENT, PRESERVATION AND ENHANCEMENT OF FOREST SPECIES IN TUNISIA

Lamia Hamrouni

*National Research Institute of Rural Engineering, Water and Forestry (INRGREF)
Institution of Agricultural Research and Higher Education (IRESA) University of Carthage
Email: hamrounilam@yahoo.fr*

Abstract

The Republic of Tunisia, located in northern Africa, faces various environmental challenges caused by anthropogenic practices such as overgrazing, deforestation, and desertification. The conversion of natural ecosystems is the major cause of plant biodiversity loss.

Because of this great environmental diversity, there are distinctive vegetation and various genetic resources in Tunisia. However, these genetic resources are endangered and the risk of degradation due to various socio-economic and industrial (deforestation, wood industries and paper, ...) as well as environmental and climatic stresses (drought, salinity, erosion, ...) and it is therefore essential, to preserve in order to exploit them effectively and sustainably. The conservation and exploitation of these resources must be done in a rational and long term. This research was conducted to investigate propagation, improvement and preservation of forest species and aromatic and medicinal plants. The objective of this work is to achieve:

- The conservation of forest species and medicinal plants by *in vitro* and *ex vitro* techniques. This is to prevent them of degradation and erosion of forest genetic resources but also to maintain the adaptive potential of species or populations in long term.
- The recovery of forest species and medicinal use and exploitation by their essential oil to identify the bioactive substances of agronomic interest (ie biological herbicides and pesticides) and pharmaceuticals (phytotherapy, synthesis of new antibiotics and anticancer drugs ...).

The results show that the techniques of *in vitro* culture improve the rate of propagation of threatened forest species and also allow their conservation and solve the problem of genetic conformity. Similarly, the use of essential oils and fixed natural resources could contribute economically to solve several problems

The results show that the techniques of *in vitro* culture significantly improve the rate of propagation of the forest species (*Arbustus*, *argania spinosa*, *cupresus*, *pinus*, ...) and also allow their conservation and solve the problem of genetic conformity. Similarly, the use of essential oils and fixed natural resources could contribute economically to solve several problems.

Keywords: Propagation, improvement, preservation, enhancement, forest species, *in vitro* culture, essential oil, antibacterial activity.

ORAL PRESENTATIONS

MESMAP-5 Draft Proceeding Book

ORAL PRESENTATION

**IN VITRO SALVIA GLUTINOSA BIOLOGICAL EFFECTS DEPEND ON
EXTRACTION PROCEDURE AND CELL TYPE**

Marina Spînu¹, Oana Pop¹, Mihaela Niculae¹, Elena Andruța Mureșan², Aurel Vasîu¹, Adrian Potârniche¹, Constantin Cerbu¹, Carmen Dana Șandru¹, Silvana Popescu¹, Eموke Pall¹

¹*Department of Clinical Sciences, Division of Infectious Diseases, University of Agricultural Sciences and Veterinary Medicine, Faculty of Veterinary Medicine, Cluj-Napoca, Romania*

²*Food Engineering Department, Faculty of Food Science and Technology, University of Agricultural Sciences and Veterinary Medicine Cluj-Napoca, Romania*

Objective/Purpose: Cytotoxicity, antibacterial and *in vitro* immune effects were investigated for aqueous (tea and cooked) and methanol extracts of *Salvia glutinosa* in three different *in vitro* cellular systems: fibroblast, tumor and stem cell lines, bacteria and mononuclear blood cells.

Material and Methods: The herb of *Salvia glutinosa* was subject to extraction with cold and hot water and methanol, respectively. The extracts were tested for cytotoxicity in HS fibroblasts cell line and B16F10 cancer cell line systems and palatal stem cells, in ATCC and clinical *E. coli* and *Pseudomonas* isolates and also in whole blood cultures from vaccinated bovine. The antioxidant capacity was determined by assessing free radical scavenging effect over 1,1-diphenyl 1-2-picrylhydrazyl (DPPH) radical for all extracts in duplicate.

Results: The antioxidant activity was the highest in cooked extract (87.402%), while the methanolic extracts were less active (83.64%). The cytotoxicity was the lowest in the tea treated cells, with a VTI% of 101.853 at a dosage of 0.95 μ l on HS cells and of 102.580 at a dosage of 7.5 μ l on B16F10 cell line. No antibacterial effects were observed in any of the extracts against the used bacterial strains. While the methanolic sage extract was the most active in bovine blood cultures at the beginning of the experiment, the aqueous extracts were highly immune stimulating for the mononuclear blood cells 14 days post vaccination.

Conclusion / Discussion: The *in vitro* biological effects of *Salvia glutinosa* were dependant on the extraction procedure, radical scavenging activity and solvent, but also the cell type used, showing no cytotoxicity and antibacterial activities but with strong immune enhancing effect.

Keywords: *Salvia glutinosa* extracts, cell-lines, antioxidant, immune stimulating, antibacterial

References:

[1] Veličković D.T., Karabegović I. T., Stojičević S.S., Lazić M.L., Marinković V.D., Veljković V.B. (2011) Comparison of antioxidant and antimicrobial activities of extracts obtained from *Salvia glutinosa* L. and *Salvia officinalis* L. Hem. Ind. 65 (5) :599–605

ORAL PRESENTATION

IN VITRO ANTIOXIDANT PROPERTIES OF EXTRACTS AND NANOPARTICLES OBTAINED FROM *EQUISETUM SP.*

Cornelia Mircea^{1*}, Denisa Bafîr-Marin², Andreia Corciovã³, Cristina Iancu¹, Oana Cioanca², Flavia Burlec³, Catalina Stan⁴, Silvia Robu⁵, Monica Hâncianu²

¹Department of Pharmaceutical Biochemistry and Clinical Laboratory, Faculty of Pharmacy, "Grigore T. Popa" University of Medicine and Pharmacy, Iași, Romania

²Department of Pharmacognosy, Faculty of Pharmacy, "Grigore T. Popa" University of Medicine and Pharmacy, Iași, Romania

³Department of Drug Analysis, Faculty of Pharmacy, "Grigore T. Popa" University of Medicine and Pharmacy, Iași, Romania

⁴Department of Drugs Industry and Pharmaceutical Biotechnology, Faculty of Pharmacy, "Grigore T. Popa" University of Medicine and Pharmacy, Iași, Romania

⁵Department of Pharmacognosy, Faculty of Medicine and Pharmacy, "Dunarea de Jos" University, Galati, Romania

*corresponding author: corneliamircea@yahoo.com

Objective: Vegetal extracts contain antioxidant compounds able to block reactive oxygen species, metals involved in oxidative processes or enzymes that catalyze oxidative processes. In our study we evaluated the antioxidant properties of extracts from *Equisetum species* and for silver nanoparticles obtained from these.

Material and Methods: We analysed ethanol and water extracts obtained from *Equisetum pratense*, *Equisetum sylvaticum* and *Equisetum telmateia*. *In vitro* antioxidant properties have been evaluated by lipoxigenase inhibition test, iron chelating test, hydroxyl scavenging and superoxide scavenging tests [2]. Silver nanoparticles (AgNPs) have been prepared using extracts and AgNO₃ (1:10) [1]. The synthesis of AgNPs has been evaluated by UV-Vis spectrophotometry.

Results: Organic compounds especially polyphenols from extracts react with silver to form AgNPs that modify the wavelength of maximum absorption to 420 nm. Antioxidant properties of extracts and nanoparticles depend on the chemical composition of these. On lipoxigenase inhibition test ethanol extracts were more active than water extracts. The *Equisetum sylvaticum* ethanol extract was the most active (EC₅₀ = 19.17±0.80 µg/mL). AgNPs are at least twice more active than extracts, so on lipoxigenase inhibition test for *E. sylvaticum*-AgNPs the value of EC₅₀ decreased to 8.98±0.49 µg/mL.

Conclusion / Discussion: Polyphenols presented in water and methanol extracts react with silver and form AgNPs that improves the biological properties of extracts. Excepting superoxide scavenging test, extracts from *E. sylvaticum* presented the most important antioxidant properties.

Keywords: antioxidant, *Equisetum species*, nanoparticles, silver

References:

- [1]. Corciova, A., Ivanescu, B., Tuchilus, C., Fifere, A., Doroftei, F., Lungoci, A.L., Marangoci, N., Mircea, C. (2018). Biosynthesis of silver nanoparticles (AgNPs) using *Tilia cordata* flowers extracts and evaluation of some biological activities. Environmental Engineering and Management Journal, 17(12), 2957-2968.
- [2]. Mircea, C., Cioanca, O., Iancu, C., Tatarina, G., Hancianu M. (2015). *In vitro* antioxidant activity of some extracts obtained from *Agaricus bisporus* brown, *Pleurotus ostreatus* and *Fomes fomentarius*. Farmacia, 63(6), 927-933.

ORAL PRESENTATION

**ETHNO-BOTANICAL STUDIES OF PLANTS USED BY TRIBES IN
MAHARASHTRA FOR TREATMENT OF KIDNEY STONES**

Susmita Gudulkar & Usha Mukundan

*Department of Botany, Ramniranjan Jhunjhunwala College (Autonomous), Affiliated to University of Mumbai,
Ghatkopar (W), Mumbai 400 086, India.*

Authors email address: - umukundan@hotmail.com

Objective/Purpose: A renal calculi or kidney stones are becoming one of the most prevalent life styles diseases globally. Treatment of kidney stones is expensive and there is also recurrence of the same. Kidney being a complex organ needs to be treated carefully otherwise it results in irreversible kidney damage. A survey in Raigad region of Maharashtra revealed that the tribal communities have been successfully using plants and their extracts for dissolution of the kidney stones. A survey was conducted to collect information on the plants being used by these tribes and the tribal medical Practitioners.

Material and Methods: A detailed survey was conducted in Raigad district of Maharashtra by meeting and observing the tribal community and local medical practitioners to learn about the plants and plant parts which were being used by them to treat patients with kidney stones. Patients who were treated and were relieved of kidney stones were also interviewed. A plant based system using cystoliths of *Ficus elastica* leaves and raphides of *Colocasia* petiole has been developed to demonstrate the efficacy of the plants used by the tribals and the local medical practitioners. Phytochemical analysis of the extracts of *Tectona grandis* fruits and *Bauhinia purpurea* leaves was done using standard methods.

Results: The tribal community in Raigad used several plants for successful treatment of kidney stones. In this research phytochemical analysis of extracts of fruits of *Tectona grandis* and *Bauhinia pupurea* leaves revealed that major components were flavonoids, saponins and phytosterols. These extract had effective lithotropic activity as revealed by the test developed in our laboratory.

Conclusion: Aqueous extracts of *Tectona grandis* fruits and *Bauhinia purpurea* leaves were effective in dissolving calcium oxalate and calcium carbonate crystals.

Keywords: Kidney stones, Lithotropic, *Tectona grandis* fruits, *Bauhinia purpurea* leaves.

References: Choubey, A., Parasar,A., Choubay, A., Iyer, D., Pawar,R.S., & Patil U., K. (2010). Potential of medicinal plants in kidney, gall and urinary stones. International Journal of Drug Development and Research 2 (2), 431-447.

ORAL PRESENTATION

CONIFERS IN THE CONQUEST OF CANCER

Diana Ivanova¹, Alexander Tashev², Paraskev Nedialkov³, Yana Ilieva³, Galina Kalotova¹,
George Angelov¹, Teodora Atanassova³

¹ Institute of Chemical Engineering, Bulgarian Academy of Sciences, 1113 Sofia, Bulgaria, dianadoc@abv.bg

² Department of Dendrology, University of Forestry, 1756 Sofia, Bulgaria, altashev@abv.bg, altashev@mail.ru

³ Faculty of Pharmacy, Medical University, 1000 Sofia, Bulgaria

Objective / Purpose: Cancer is a global disease, causing millions of deaths every year. Along with the existing anticancer drugs, identification of new antiproliferative agents is necessary for patients in poor general condition who cannot tolerate conventional cytostatic drugs or in cases of resistance to the available chemotherapeutics. This study is aimed at identification of plant species, producing metabolites with high antiproliferative activities and with potential application as sources of lead compounds for drug discovery.

Material and Methods: *Juniperus sabina* var. *balkanensis* was delivered from its natural habitat in the Rhodope Mountains.¹ Other conifer representatives and *Buxus sempervirens* were from the Arboretum of the University of Forestry, Sofia, Bulgaria. *Sequoiadendron giganteum*, *Ruta graveolens*, *Betula pendula*, *Inula helenium* were collected from the Botanic garden area of the Bulgarian Academy of Sciences. *Rhodiola rosea* was of commercial origin. Antiproliferative activities of the extracts were evaluated by their half-maximum growth-inhibitory concentrations (IC₅₀), derived from the dose-response curves of the MTT-assay after treatment of NB4 cells with the corresponding plant extracts, as it was described previously.

Results: In the group of studied plant species, the best antiproliferative activities were determined after treatment of NB4 (acute promyelocytic leukemia) cells with the leaves extracts of *Juniperus sabina* var. *balkanensis* R. P. Adams & A. N. Tashev (IC₅₀=0.3±0.03 microgram/ml) and *Sequoiadendron giganteum* (Lindl.) J. Buchh. (IC₅₀=0.9±0.1 microgram/ml) in comparison with less active leaves extracts of other conifers *Taxus baccata* L., *Cupressus arizonica* Greene, *Chamaecyparis lawsoniana* Parl., *Chamaecyparis pisifera* (Siebold & Zucc.) Endl, and other medicinal plant species *Buxus sempervirens* L. (leaves), *Ruta graveolens* L. (leaves), *Inula helenium* L. (roots), *Betula pendula* Roth (cortex), *Rhodiola rosea* L. (rhizomes). Podophyllotoxin and other lignans were identified in the *J. sabina* var. *balkanensis* extract by UPLC/HRMS. Identification of *S. giganteum* metabolites is in progress. While *S. giganteum* (giant sequoia) belongs to the endangered plant species and its metabolites are useful as lead compounds for design of anticancer chemotypes, *J. sabina* varieties are easy for cultivation and feasible for industrial exploitation.

Conclusion / Discussion: The leaves extracts of *Juniperus sabina* var. *balkanensis* and *Sequoiadendron giganteum* were identified as new natural agents with superior anticancer activities in leukemia cells. Extension of their antiproliferative properties in other types of cancer cells is envisaged in the near future.

Keywords: Anticancer conifers, *Juniperus sabina* var. *balkanensis*, *Sequoiadendron giganteum*

Acknowledgements: The work is performed within the Bulgarian Science Fund contract DN 07/25 (2016) and project “Development of green eco-technologies” of the Bulgarian Academy of Sciences.

ORAL PRESENTATION

CURCUMIN, A POLYPHENOL DERIVED FROM *CURCUMA LONGA*: CLINICAL EVIDENCE AND MOLECULAR MECHANISMS IN PREVENTION AND MANAGEMENT OF CANCER

Nazim Sekeroglu^{a*} & Sevgi Gezici^b

^a *Department of Food Engineering, Faculty of Engineering and Architecture, Advanced Technology Application and Research Center, Kilis 7 Aralik University, 79000 Kilis, Turkey*

E-mail: nsekeroglu@gmail.com, www.nazimsekeroglu.com

^b *Department of Molecular Biology and Genetics, Faculty of Science and Literature; Advanced Technology Application and Research Center, Kilis 7 Aralik University, 79000 Kilis, Turkey*

E-mail: drsevgigezici@gmail.com, www.sevgigezici.com

Background and Purpose: Cancer is the second leading cause of death globally, and it has been a major health problem in the past decades. Several strategies are involved in treatment and prevention of cancer cases such as chemotherapy, immune therapy, radiation, hormone therapy, targeted therapy and herbal therapy. Among them, herbal therapy with medicinal and aromatic plants has been gained great importance in recent years. Medicinal plants and their bioactive compounds may offer clinically useful anticancer agents, due to their rich anticarcinogenic and chemoprotective potentials.

Curcumin, the rhizome of *Curcuma longa*, is widely used in therapeutic purposes because of its several biological properties such as anti-inflammatory, anticancer, antioxidant, and etc. From this point of view, this review is aimed to comprehensively highlight the effects of curcumin for the prevention and treatment of cancer. Methods: This report reviews the pharmacological and anticancer effects of curcumin with a focus on its molecular mechanisms in prevention and management of various cancer cases. For this purpose, electronic databases, including PubMed, Scopus, ScienceDirect, Cochrane library, Google Scholar and MedlinePlus, etc. were searched to summarize *in vitro*, *in vivo* and clinical studies on anticancer effects of *Curcuma longa* and curcumin, polyphenolic bioactive compound of the plant up-to-date.

Results and Conclusion: To sum up, curcumin, the active ingredient of turmeric and exerts remarkable therapeutic potential for the treatment of various types of cancers. Curcumin is strong anti-oxidant and anti-inflammatory properties and therefore it possesses anticancer potentials through various molecular mechanisms. Based on documented data this natural compound could be a key source of anticancer agent in modern anticancer therapy after clinical confirmation.

Key Words: *Curcuma longa*; curcumin; anti-cancer agents; molecular mechanism; natural products; medicinal plants.

ORAL PRESENTATION

BLUE ESSENTIAL OIL OF CHAMOMILE VARIETY “LIANKA” WITH THE HIGHEST CONTENT OF γ -BISABOLOL AND ITS IMPORTANCE TO NEW PHARMACEUTICAL FORMULATIONS

Ivan Salamon¹, Anton Bubnov¹, Marianna Kharabara²

¹ Department of Ecology, Faculty of Humanities and Natural Sciences, University of Presov, 01, 17th November St., SK-081 16 Presov, Slovak Republic

² Department of Biology and Pharmaceutical Botany, Medical University of Lodz, Muszynskiego 1, PL-90-151 Lodz, Poland

Objective/Purpose: Based on the study of pharmacodynamics properties, the chamomile variety “LIANKA” was bred at the University of Presov, Slovakia, between the years 2008 – 2013. Currently, this variety obtains the Certificate by the Community Plant Variety Office (CPVO) in Angers, France (in 2018).

Material and Methods: 5 samples of German Chamomile, diploid variety “LIANKA”, from the small-cultivation in Trebisov, the Eastern Slovakia (2018), with weight of 10 g was grounded in a blender. The essential oil was prepared by hydro-distillation in Clevenger-type apparatus according to the EU Pharm. The GC/MS analyze were carried out on a Varian 450-GC connected with a Varian 220-MS. Identification of components was made by comparison of their mass spectra with those stored in NIST 14. Kovat's samples (C-5 to C-22 alkane mixture) injected during sample analyze and Kovat's indices calculated from retention time using a third order polynomial. 40 reference authentic compounds (Extrasynthese, Merck, Sigma and Roth) were purchased. Many authenticated from MS data or retention index were compared by literature (Adams, 2007).

Results: The GC/MS results confirm earlier reports (Salamon et al., 2016) that major volatile constituents obtained from the chamomile antheridia were γ -bisabolol and chamazulene. The selected variety, environmental conditions, agronomic management practice and effective distillation (Salamon, 2007) were obtained of substantial contents of γ -bisabolol (69.87 \pm 2.1%) and chamazulene (10.69 \pm 0.5%) and the low contents of γ -bisabololoxides A and B (< 3 %). The blue essential oil content was ranged from 0.65 to 0.85 % in order to time of single flower harvests. γ -bisabolol is the most important component of the blue chamomile essential oil, also known as levomenol, chemically is a natural monocyclic sesquiterpene alcohol. The chemist Sorm and his research team isolated and discovered the γ -bisabolol in chamomile essential oil in 1951. The isopropylindene structure of natural γ -bisabolol was proved through infrared (IR) and nuclear magnetic resonance spectra (Franke and Schilcher, 2005).

Conclusion/Discussion: Formulation several liquid and ointment preparations (Schilcher, 2004) were standardized to contain essential oil, high contents of γ -bisabolol and chamazulene from flowers. These formulations are used on inflammations of the gums and oral cavity, rectum and urogenital region, and other mucous membranes. The formulations are also used for skin baths and enemas and as aides in curing bronchitis, stomach ulcers, digestive disorders, wounds, and dermatitis. The chamomile variety “LIANKA” and its large-scale cultivation, extraction and distillation is very forward raw-material to the pharmaceutical industry in the world.

References:

- ADAMS, R.P. (2007). *Identification of Essential Oil Components by Gas Chromatography/ Mass Spectrometry*. 4th Eds., Carol Stream, IL, USA: Allured Publishing Corporation, 804 p., ISBN: 1932633219
- FRANKE, R. – SCHILCHER, H. (2005). *Chamomile – Industrial Profiles*. 1st Eds., CRC Taylor & Francis, Boca Raton, London, New York, Singapore, 289 p., ISBN: 0415334632
- SALAMON, I. – FEJER, J. – GRULOVA, D. (2016). Breeding of German Chamomile, *Chamomila recutita* L., with the Highest Content of γ -Bisabolol. 1st Eds (V.D. Jeliakov – Ch.L. Cantrell). The book “Medicinal and Aromatic Crops: Production, Phytochemistry, and Utilization”, American Chemical Society, Washington, D.C., USA, 288 p., ISBN13: 9780841231276, eISBN: 9780841231269
- SALAMON, I. (2007). Effect of the Internal and External Factors on Yield and Qualitative-quantitative Characteristics of Chamomile Essential Oil. *Acta Horticulturae*, No. 749, p. 45-64, ISSN: 05677572
- SCHILCHER, H. (2004). *Wirkungsweise und Anwendungsformen der Kamillenblüten – Handbuch für Apotheker, Ärzte, Heilpraktiker und weitere Heilberufe sowie für Medizin-, Pharmazie- und Biologiestudenten*. BMV Berliner Medizinische Verlagsanstalt GmbH, 160 s., ISBN: 3880400067

ORAL PRESENTATION

CHEMICAL COMPOSITION AND BIOLOGICAL ACTIVITIES OF ESSENTIAL OILS OF BULBS AND BULBILS FROM *ALLIUM SATIVUM* L.

Zakia Boubechiche^{1,2}, Mounira Ouar-Korichi³, Rabah Boukherroub⁴, Alexandre Barras⁴, Patrick Daubias⁵, Nour-Eddine Chihib⁶, Amina Hellal⁷, Charafeddine Jama⁵

¹Ecole Nationale Supérieure Agronomique, 16000 Alger, Algérie, e-mail : boubechichez@yahoo.fr

²Ecole Nationale Supérieure des Sciences de la Mer et l'Aménagement du Littoral, 16000 Alger, ALGÉRIE

³Laboratoire des Entérobactéries, Institut Pasteur d'Alger, 16000 Alger, ALGÉRIE

⁴IEMN, CNRS-8520, Cité Scientifique, Avenue Poincaré, B.P. 60069, 59652 Villeneuve d'Ascq, FRANCE

⁵UMET, CNRS, ENSCL, UMR 8207, Avenue Dimitri Mendeleïev, 59652 Villeneuve d'Ascq, FRANCE

⁶UMET-PIHM, Université de Lille, CNRS, INRA, UMR 8207, CS 20039, 59651 Villeneuve d'Ascq, FRANCE

⁷Laboratoire des Sciences et Techniques de l'Environnement, Ecole Nationale Polytechnique, Alger, ALGÉRIE

Objective: This work aims to study in vitro, antioxidant, antibacterial, antifungal and cytotoxic activities of essential oils of bulbs and bulbils from garlic (*Allium sativum*).

Material and Methods: The chemical composition of these essential oils extracted by conventional hydrodistillation method was analysed using GC/MS. The antimicrobial activity was assessed against bacterial and fungal strains clinically significant and the antioxidant activity by using 2,2-diphenyl-1-picrylhydrazyl (DPPH) method. Also, the in vitro cytotoxicity of the bulbs essential oil was evaluated on human cancer cell lines, the HeLa cell line derived from cervical carcinoma from a 31-year-old female [ATCC® CCL-2™, ECACC, Sigma Aldrich, Saint-Quentin Fallavier, France] and the MDA-MB-231 cell line derived from human breast adenocarcinoma [ATCC® HTB-26™, ECACC, Sigma Aldrich, Saint-Quentin Fallavier, France] by the UptiBlue Viable Cell Counting Assay.

Results: A total of twenty-three compounds were identified and the diallyl disulfide, methyl allyl trisulfide and diallyl trisulfide were found to be the predominant compounds for both essential oils. These oils showed respectively potential DPPH radical scavenging activity with IC₅₀ (concentration providing 50% inhibition) values of 1.87 and 2.85 mg/ml for bulbs and bulbils. Similarly, antimicrobial activity showed moderate to potent broad spectrum activities. However, the strains *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Trichophyton violaceum* and *Epidermophyton floccosum* proved to be the most sensitive. Moreover, after 48 hours of incubation, the oil tested proved to be potent against both cell lines with a mortality of almost 100% at a concentration of 0.05%, the HeLa cell line appears more sensitive with a much lower viability at 50% for a concentration of 0.005%.

Conclusion: Based on these results, further experiments will be carried out on this golden vegetable in order to use it as an alternative product in food and pharmaceutical proposes.

Keywords: Garlic essential oil, GC-MS, Antioxidant activity, Antimicrobial activity, Cytotoxicity.

ORAL PRESENTATION

SEMI-SYNTHESIS OF BIOACTIVE GYPSOGENIN DERIVATIVES

Safiye Emirdağ-Öztürk¹

¹ Ege University, Faculty of Science, Department of Chemistry, Bornova, Izmir, Turkey

Objective / Purpose: Gypsophila species have been used for the traditional Chinese medicine. Gypsophila is well known to contain saponins. Saponins are glycosides which are found in many plants. These saponins have variety of biological activities including, antimicrobial, antifungal, antitumor, antioxidant, cytotoxic activity. In this study, gypsogenin aglycone which is a natural saponin was obtained from the roots of *Gypsophila arostii*. Gypsogenin and synthesised compounds were evaluated for activity. We synthesized some bioactive compounds.

Material and Method: We synthesized new gypsogenin derivatives (**1-9**). Purification has been carried out using chromatographic methods (Column Chromatography).


Results: Synthesized compounds (**1-9**) was determined by IR, UV, ¹H NMR, APT and LCMS analysis.

Conclusion / Discussion: Gypsogenin and synthesised compounds were evaluated for antimicrobial and cytotoxic activities.

Keywords: Gypsogenin, semi synthesis.

References:

- [1] The Chinese Medicine Dictionary, Shanghai People's Publishing House, Shanghai, 1977, 2170.
[2] Öztürk S., Karayildirim T., Capaci-Karagoz A., Alankus-Caliskan O., Ozmen A., Poyrazoglu-Coban E. (2014). Synthesis, antimicrobial and cytotoxic activities, and structure-activity relationships of gypsogenin derivatives against human cancer cells, *European Journal Of Medicinal Chemistry*, vol.82 565-573.



	R	R ¹	R ²
Gypsogenin	-OH	-COH	-COOH
Compound 1	-OH	-CH=N-OH	-COOH
Compound 2	-OCO-CH ₃	-CHO	-COOH
Compound 3	-OH	-CHO	-CO-OCH ₂ -Ph
Compound 4	-OCO-CH ₃	-CH=N-OH	-COOH
Compound 5	-OCO-CH ₃	-CH=N-NH-CS-NH ₂	-COOH
Compound 6	-OH	-CH=N-OH	-CO-OCH ₂ -Ph
Compound 7	-OH	-CH=N-NH-CS-NH ₂	-CO-OCH ₂ -Ph
Compound 8	-OCO-CH ₃	-CHO	-CO-OCH ₂ -Ph
Compound 9	-OCO-CH ₃	-CH=N-OH	-CO-OCH ₂ -Ph

ORAL PRESENTATION

INFLUENCE OF AN AQUEOUS MINT (*MENTHA PIPERITA*) EXTRACT ON HEALTH STATUS AND IMMUNE RESPONSE IN YOUNG CALVES

Ciprian Ilașcă¹, Mihaela Niculae¹, Elena Andruța Mureșan³, Aurel Vasiiu¹, Eموke Pall¹, Raluca Seicheli¹, Constantin Cerbu¹, Carmen Dana Șandru¹, Silvana Popescu², Marina Spînu¹

¹Department of Clinical Sciences, Division of Infectious Diseases, University of Agricultural Sciences and Veterinary Medicine, Faculty of Veterinary Medicine, Cluj-Napoca, Romania

²Department of Animal production, Division of Hygiene and Welfare, University of Agricultural Sciences and Veterinary Medicine, Faculty of Veterinary Medicine, Cluj-Napoca, Romania

³Food Engineering Department, Faculty of Food Science and Technology, University of Agricultural Sciences and Veterinary Medicine Cluj-Napoca, Romania

Objective/Purpose: The susceptibility to diseases in calves is increased during the neonatal period, thus natural products, such as plant extracts with increased bioavailability and less side effects could help in boosting immunity. *Mentha piperita*, an ubiquitous plant, with considerable antimicrobial activity, could serve for this purpose.

Material and Methods: Three equal groups of calves (n=7), with homogenous age within the group (15-25 days), were subjected to plane milk (I) or milk plus 200 ml(II) or 400 ml (III) of mint tea treatment for 10 days. The clinical health status of animals was monitored daily and blood was sampled on heparin (50 IU) on days 0 and 10. Blast transformation of mononuclear cells was quantified by a micromethod. Cultural variants included treatment with alcoholic extracts of *M. piperita*, *Echinacea angustifolia*, *E. purpurea*, *Calendula officinalis*, *Vaccinium myrtillus*, *Thymus vulgaris*, *Symphitum officinale* and propolis (1μl/well). The antioxidant activity of the aqueous extract of mint was determined by assessing free radical scavenging effect over DPPH (%) radical. Excel program was used to evaluate the statistical significance of the treatments' effects by group.

Results: All calves stayed clinically health during the entire experiment. The antioxidant activity of plants rendered from *E. purpurea* being the most to *C. officinalis* the least active antioxidant. The mint tea antioxidant activity was of 76.2%. The blast transformation indices were high in all groups and did not differ significantly, but with higher values for all plant extract treatments in the 200 ml tea treated group.

Conclusion / Discussion: Mint tea treatment somewhat increased the spontaneous blast transformation capacity of mononuclear cells but did not significantly influence the *in vitro* response to other plant extracts.

Keywords: calves, mint tea, antioxidant, alcoholic plant extracts, blast transformation

References:

[1] Bombik T., Bombik E., Frankowska A., Trawińska B., Saba L. (2012) Effect of herbal extracts on some haematological parameters of calves during rearing, Bull Vet Inst Pulawy, 56: 655-658.

ORAL PRESENTATION

INVESTIGATION OF THE CYTOTOXIC, ANTICANCER AND APOPTOTIC DNA FRAGMENTATION PROPERTIES OF *BONGARDIA CHRYSOGONUM* L. EXTRACTS ON BRAIN GLIOMA AND LUNG CARCINOMA CANCER CELLS

Sevgi Gezici*

Department of Molecular Biology and Genetics, Faculty of Science and Literature; Advanced Technology Application and Research Center, Kilis 7 Aralik University, 79000 Kilis, Turkey

E-mail: drsevgigezici@gmail.com, sevgigezici@kilis.edu.tr

Background and Objective: *Bongardia chrysogonum* L. is commonly known as ‘Uruf-el-Deek’ and locally known as ‘Catlak Otu’ in the Anatolia, where the tubers of this plant have been used for the treatment of prostate hypertrophy, hypercholesterolemia, hemorrhoids, diabetes, epilepsy and hematological malignancies in folk remedy. **Material and Methods:** The purpose of the current study was to investigate potential anticancer and antiproliferative activities of the tubers of *B. chrysogonum* L., extracted with water, ethanol and chloroform solvents, against human lung carcinoma (A549), non-small lung cancer (H1299), glioma (C6) cancer cells, and non-tumorous human umbilical vein endothelial cells (HUVECs). In this context, cell proliferation and cell viability of the cells treated with the plant extracts at various concentration ranged from 10, 25, 50, 100 to 200 µg/mL were determined by MTT and trypan blue exclusion assays. The dose response curve was used to generate the IC₅₀ (µg/mL) values for each cell line. ANOVA (one way) was used for evaluation statistical differences between the groups, and the experiments were carried out at least triplicate. Based on their cytotoxicity against A549, H1299 and C6 cancer cells, the concentrations of the extracts were selected for *in vitro* micronucleus (MN) assay. Apoptotic activity induced in the plant extracts treated and untreated cancer cell lines were analyzed for DNA fragmentation using the ‘Apoptotic DNA-Ladder Kit’ (QIAamp DNA mini kit, Qiagen), and the release of the enzyme lactate dehydrogenase (LDH) activity was also performed for detection of necrosis in the cultured cells. **Results:** The tuber extracts of *B. chrysogonum* L. exerted significant anticancer and antiproliferative activity towards all treated cancer cells with the IC₅₀ values ranged from 8.25±0.01 to 42.05±0.03 µg/mL. The methanolic-tuber extract seem to be possessed the strongest anticancer and antiproliferative activities on A549 cells (IC₅₀=8.25±0.01 µg/mL at 50 µg/mL concentration), whilst chloroform extract of the tuber was found to have the weakest ones on C6 cells. With regard to LDH activity, it was determined that the methanolic-tuber extract lead to higher leakage activity towards C6 cells as compared to that of lung cancer cell lines, in dosage and time dependently. In addition, the tuber extracts of the plant-induced apoptosis was determined in the cancer cells, and multinucleated cells treated with the extracts at different concentration were distinguished as apoptotic or necrotic cells under the microscope. **Conclusions:** As can be seen from the literature survey, the results presented herein could be considered as the first report on probable anticancer and antiproliferative effects of the tuber of *B. chrysogonum* L. However, the data obtained from the current research would be provided valuable information to discover plant-based chemotherapeutics for the cure of cancers, the extracts should be analyzed by using further *in vitro* and *in vivo* techniques for specify the bioactive phytochemicals responsible for this excellent anticancer potential.

Keywords: *Bongardia chrysogonum* L., anticancer, apoptosis, necrosis, DNA fragmentation, micronucleus

Acknowledgments: The author would like to thank to taxonomist Res. Assist. Fatih Yayla for his kind contribution of collecting and identifying the plant material, and Prof. Dr. Nazim Sekeroglu for his kind scientific contribution to the research.

References

- [1] Gezici S, Sekeroglu N, Kijjoo A. 2017. *In vitro* Anticancer Activity and Antioxidant Properties of Essential Oils from *Populus alba* L. and *Rosmarinus officinalis* L. from South Eastern Anatolia of Turkey. Indian Journal of Pharmaceutical Education and Research, 51(3):498-503.
- [2] Gezici, S. 2018. Promising anticancer activity of lavender (*Lavandula angustifolia* Mill.) essential oil through induction of both apoptosis and necrosis. Annals of Phytomedicine, 7(2), 38-45.
- [3] Gezici S. 2018. Molecular Mechanisms and Targeting Pathways of Plant Derived Bioactive Compounds in Cancer Prevention and Therapy. The 4th International Symposium on Pharmaceutical and Biomedical Sciences. March 17-19, 2018, Kumamoto, Japan. Abstract Book, Invited Speech S-13 p:20.
- [4] Gezici, S., Sekeroglu, N. 2019. Current perspectives in the application of medicinal plants against cancer: novel therapeutic agents. Anti-Cancer Agents in Medicinal Chemistry, 19, 1-11.

ORAL PRESENTATION

COMPOUNDS FROM AN AROMATIC ANNUAL PLANT GROWING IN ARID ZONE OF ALGERIA AND SCREENING OF ITS ANTIBACTERIAL ACTIVITY

Mebarka Bouziane^{1,2*}, Naima Hellali¹, Ferdous Cherifi^{1,2}, Bakir Abanou^{1,2},
Mahfoud hadj Mohammed^{1,2}

¹ *Laboratory of BioGeochemistry of Desert Environments. Faculty of Mathematics and matter sciences, Kasdi Merbah University, Ouargla 30000, Algeria.*

² *Kasdi Merbah University, Ouargla 30000, Algeria.*

*Corresponding author: bmebarka@yahoo.fr

The Asteraceae *Matricaria pubescens* (Desf.) is a spontaneous species from arid regions of Algeria. To get its volatile fraction, the aerial parts of this plant were subjected to a hydrodistillation using a cleavenger apparatus. A weak yield (0.1%) was obtained. The sample was analyzed by GC-MS on a non polar capillary column. Among identified components in this volatile composition were; spathuleol (10.59%), decanoic acid (5.03%), trans isoelemicin (4.70%), cadinol (1.59%).

Beside, the purpose of the present study was to continue the exploration of the biological effects of *M. pubescens*. Indeed, the prepared essential oil and its fragrant water have been the subject of an antibacterial investigation. However, most of the microorganisms tested were resistant to the tested samples.

Keywords: GC-MS, volatile composition, bioactivity, spontaneous plant, arid zones.

References:

1. R. Chemli, plantes medicinales et aromatiques de la flore de Tunisie, *CIHEAM -Options Méditerranéennes*, 119-126.
2. W. Tadrent, A. Kabouche, R. Touzani and Z. Kabouche, Chemotypes investigation of essential oils of "Guertoufa" herbs, *J. Mater. Environ. Sci.*, 5 (4) (2014) 1200-1205.
3. K. Maiza, A. Longeon, V. Hammiche, M. Guyot, F. Benabdesselam-Maiza, Biological Activities of Plants Collected in The Algerian Sahara, *Life Sciences Leaflets*, 2014, 52, 52-56.
4. R.P. Adams, Identification of Essential Oil Components by Gas Chromatography/Quadrupole Mass Spectroscopy. Academic Press, New York, Appendix I, II (1989).
5. http://www.flavornet.org/d_kovats_ov101.html, Accessed 27 / 02 / 2019.

ORAL PRESENTATION

IMPACT OF DIFFERENT CUTTING STAGES ON THE CONTENT, CHEMICAL COMPOSITION AND ANTIBACTERIAL ACTIVITE OF ESSENTIAL OILS FROM *SALVIA OFFICINALIS* L.

Belgin Coşge Şenkal¹, Tansu Uskutoğlu¹, Arzu Türker², Hülya Doğan³

¹ Yozgat Bozok University, Faculty of Agriculture, Field Crop Department, Yozgat, Turkey, belgin.senkal@bozok.edu.tr

² Bolu Abant İzzet Baysal University, Faculty of Arts and Sciences, Biology Department, Bolu, Turkey

³ Yozgat Bozok University, Vocational School of Technical Sciences, Department of Seed Science, Yozgat, Turkey

Objective: *Salvia officinalis* L. from Lamiaceae family is a worldwide cultivated aromatic herb [1]. Generally, this species has been used herbal tea for a wide range of disorders and illnesses [2]. Also, the essential oils of *S. officinalis* exhibit antioxidant, antimicrobial, antispasmodic, and stimulant properties [3]. The aim of the present study was to investigate the effect of different cutting stages on chemical composition and antibacterial activities of essential oils from *S. officinalis* L. cultivated in Turkey.

Material and Methods: Chemical composition of essential oils obtained by hydro-distillation from the dried aerial parts of *S. officinalis* cultivated in Bolu/Turkey ecological conditions harvested before flowering (BF), 50% of flowering (50%F) and full flowering (FF) stages were determined GC and GC/MS. The disc diffusion assay (Kirby-Bauer Method) was used to screen for antibiotic activity.

Results: The essential oil contents obtained were 0.30% in BF stage, 0.40% in 50%F, and 1.00% in FF stage. α -thujone (28.46%, 36.52% and 48.94%) were found to be the major component of the essential oils from BF, 50%F, and FF stages, respectively. The essential oils exhibited strong antibacterial activity against the microorganisms (*Salmonella typhimurium*, *Serratia marcescens*, *Proteus vulgaris*, *Enterobacter cloacae*, *Klebsiella pneumonia*, *Streptococcus pyogenes*, *Staphylococcus aureus*, *Staphylococcus epidermidis* and *Escherichia coli*) examined.

Conclusion/Discussion: The highest amount of essential oil and α -thujone content were obtained from the plants which were harvested in FF stages. The contents, chemical compositions and antimicrobial activities of essential oils from *S. officinalis* varied according to cutting stages.

Keywords: Disc diffusion, essential oil, GC/MS analysis, *Salvia officinalis* L.

References:

- [1] Khalil, R., & Li, Z-G. (2011). Antimicrobial activity of essential oil of *Salvia officinalis* L. collected in Syria. *African Journal of Biotechnology*, 10(42):8397-8402.
- [2] Dweck, A.C. (2000). The folklore and cosmetic use of various *Salvia* species. In: Kintzios, S.E. (ed) Medicinal and aromatic plants-industrial profiles, Sage, the genus *Salvia*. Harwood Academic Publishers, The Netherlands. pp.1-25.
- [3] Bernotienė, G., Nivinskienė, O., Butkienė, R., & Mockutė, D. (2007). Essential oil composition variability in sage (*Salvia officinalis* L.). *Chemija*, 18(4):38-43.

ORAL PRESENTATION

**MEDICINAL PLANTS AND PHYTOCHEMICALS: A SOURCE FOR GLUT4
STIMULATORS AND DIGESTIVE ENZYMES INHIBITORS**

Hilal Zaid^{1,2}, Sleman Kadan¹, Najlaa Bassalat² and Raghad Albzoor²

*Qasemi Research Center- Al-Qasemi Academic College, Baqa-El-Gharbia, Israel.
Faculty of Science, Arab American University/Palestine, P.O. Box 240, Jenin, Palestine.*

Diabetes is one of the world's most widespread diseases, which is characterized by persistent hyperglycemia and related to metabolic disorders. It affects over 327 million people and causes about 300,000 deaths annually [1]. Despite great advances in the prevention and therapy, existing treatments for this disorder have serious side effects. Plants used in traditional medicine represent valuable sources in the search for new medicinal compounds. In traditional system of medicine, at least 1200 species of medicinal plants are used for their antidiabetic attributes. However, only a small proportion of such plants have been scientifically validated [2].

The objective of this study was to assess the *in vitro* inhibitory activity of selected local anti-diabetic medicinal plants on carbohydrate and lipid digestive enzymes; namely, α -glucosidase, pancreatic α -amylase, and pancreatic lipase. The inhibitory activities of ethanol: water (50%:50%) herb extracts were evaluated using the enzymatic colorimetric assays. At a concentration of 0.2 mg/ml, among six herbal extracts tested, only *Allium sativum* (bulb) showed an inhibitory activity against the intestinal sucrase (α -glucosidase). *Cinnamomon cassia* (bark) showed an inhibitory activity against lipase. *C. cassia* (bark) and *Olea europaea* (leaves) inhibited pancreatic α -amylase with IC_{50} values of 24 ± 3.01 and 192.94 ± 6.4 μ g/mL, respectively. Tens of other local medicinal plants extracts were tested for GLUT4 activity and translocation to the plasma membrane (Fig. 1). These plants include but are not limited to *Abelmoschus esculentus* L., *Allium cepa* (bulb), *Allium sativum* (bulb), *Asparagus aphyllus* L., *Atriplex halimus*, *Cinnamomon cassia*, *Crataegus azarolus* L., *Gundelia tournefortii*, *Nigella sativa* (seeds), *Ocimum Basilicum*, *Olea Europea*, *Trigonella foenum-graecum*, *Teucrium polium* and *Urtica dioica*. GC/MS phytochemical analysis for most of these plants are applied and the potential active compounds are discussed (Fig. 2). The most effective plant extracts were selected and are being tested *in vivo* in diabetic mice model. The activity of these plants will be presented and discussed.

Keywords: Phytochemicals, diabetes, GLUT4, α -glucosidase, α -amylase, lipase, GC/MS.




Fig.1. Model demonstrating the detection of the membrane insulin-responsive glucose transporter 4 (stably expressing myc-tag) -GLUT4myc [3].




Fig.2. Some of the new detected phytochemicals

References:

1. **Zaid H.**, Tamrakar A.K., Razzaque M.S and Efferth T. (2018) Diabetes and Metabolism Disorders Medicinal Plants: A Glance at the Past and a Look to the Future. *Evidence-Based Complementary and Alternative Medicine (eCAM)*; In press.
2. Saad B., **Zaid H.**, Kadan S. & Shanak S. (2017). Anti-diabetes and Anti-obesity Medicinal Plants and Phytochemicals: Safety, Efficacy, and Action Mechanisms. Springer. Springer International Publishing AG 2017. ISBN 978-3-319-54101-3.
3. **Zaid H.**, Antonescu CN, Randhawa VK & Klip A. (2008). Insulin action on glucose transporters through molecular switches, tracks and tethers. *Biochemical Journal*, 413(2), 201-15.
4. Kadan S., Saad B., Sasson Y & **Zaid H.** (2016). *In vitro* evaluation of anti-diabetic activity and cytotoxicity of chemically analyzed *Ocimum basilicum* extracts. *Food Chemistry Journal*, 196, 1066-1074.

MESMAP-5 Draft Proceeding Book

ORAL PRESENTATION

MORPHOLOGY, YIELD AND QUALITY CHARACTERIZATIONS OF *NIGELLA SATIVA* L. AND *NIGELLA DAMASCENA* L.

Glsm Yaldiz, Mahmut amlica, Abdurrahman Baol, Halit Akin

Department of Field Crops, Faculty of Agriculture and Natural Sciences, Bolu Abant İzzet Baysal University, 14280 Bolu, Turkey. E-mail: g_yaldiz@hotmail.com

Objective/Purpose: *Nigella sativa* and *Nigella damascena* are two annual species of the family Ranunculaceae, used in traditional medicine for centuries for many diseases like cancer, cardiovascular complications, diabetes, asthma, kidney disease etc. The aim of this study is to determine the important agricultural and quality characters of 1 *N. sativa* cultivar (ameli) and 2 *N. damascena* populations (Ankara and Rize) in Bolu ecological conditions.

Material and Methods: One cultivar of *N. sativa* and two populations of *N. damascena* were used to conduct these studies. The field experiments were performed during two successive seasons (2017 and 2018) at the experimental farm (40°44'44'' N, 31°37'45'' E, 881), which is located at an elevation of approximately 881 m altitude, belonging to Bolu Abant İzzet Baysal University. (Bolu, Turkey).

Results: On average days to 50% seedling and flowering determined as 15.67-18.0 days in 2017, 14.33-16.0 days in 2018, 69.0-75.33 days in 2017, 65.67-75.33 days in 2018 respectively. Seed weights per capsule were 17.11-27.08 g in 2017 and 11.51-12.43 g in 2018, the 1000 seeds weights were between 2.18-2.84 g in 2017 and 2.37-2.77 g in 2018. Mean seed yield in the first year (38.02-60.18 kg/da) were higher than yield in the second year (31.88-37.20 kg/da). The essential oil content of the populations and cultivar was found to be very close to each other with 0.10-0.31%, and essential oil compositions were between 5.68-51.06 %. The mean fixed oil rate in populations and cultivar varied between 20 % and 29%, and fatty acid compositions were between 66.83-91.58%. The major inorganic matters were found as K⁺ (177.50-310.5 mg/g) and Ca⁺² (58.90-128.0 mg/g) and followed by Mg⁺² and Na⁺.

Conclusion: The plant height, the number of branches, the number of capsule and 1000-seed weight of populations varied according to years. Generally, in compared to populations and cultivar the best results were obtained in *N. sativa* cultivar.

Keywords: Morphology, *Nigella* sp., quality, yield

References:

[1] zel, A., & Koar, İ. (2018). Characterization on variety and populations of black cumin (*Nigella sativa* L.): I. agricultural properties. *Harran Tarım ve Gıda Bilimleri Dergisi*, 22(4), 533-543.

* This work was supported by Student Scientific Research Project Fund (2018.10.07.1281), Faculty of Agriculture and Natural Sciences, Bolu Abant İzzet Baysal University, Turkey.

ORAL PRESENTATION

**NATURAL INDUCED RESISTANCE IN CROP PLANTS BY PALESTINIAN
INDIGENOUS PLANT EXTRACTS AGAINST PESTS AND DISEASES**

Bushrah Karsou & Rana Samara

*Faculty of Agricultural Science and Technology, Palestine Technical University - Kadoorie, Tulkarm, Yafa
Street, Palestinian Territories*

The potential beneficial roles of indigenous plant extract have been emphasized in treatment of various plant diseases and insects. Plant and plant-derived products contain antioxidants, anti-inflammatory and repellents; that are thought to provide protection or treatment against many common pests and diseases. Extensive studies are now focusing on the positive role of plant extract derivatives as resistant elicitors thus insect pest and disease control. Negative cross-talk between the jasmonic and salicylic acid response pathways has been shown to increase the level of the plant's pathogen defenses at the expense of pest defenses. Plant activators are biological or chemical-derived stimulators of plant systemic acquired resistance (SAR) mediated through the salicylic acid (SA) pathway. Examples include plant pathogen *Pseudomonas syringae* and the chemical 1,2,3-benzothiadiazole-7-carbothioate (BTH or the commercial plant activator Actigard 50WG), both of which have been shown to prime the defenses in several crops. The objective is to determine under what circumstances pathogens might be negatively affected by plant SAR inducible defenses. A series of experiments with barley cultivars are compared according to relative growth rate when the inoculated artificially with loose smut pathogen on SAR induced or untreated crops leaves in combination with inhibitors of detoxification enzymes, antioxidants and multidrug resistance protein (MRP) transport.

The objectives of this study was to determine the effect of several indigenous plant extracts on the plant crops and evaluate their impact as elicitor for inducing plant resistance against disease.

References:

- Boughton, A.J., K. Hoover, and G.W. Felton. 2006. Impact of chemical elicitor applications on greenhouse tomato plants and population growth of the green peach aphid, *Myzus persicae*. *Entomologia Experimentalis et Applicata*. 120:175-188.
- Burketová, L., L. Trdá, P. Ott, and O. Valentová. 2015. Bio-based resistance inducers for sustainable plant protection against pathogens. *Biotechnol. Adv.* doi: 10.1016/j.biotechadv.2015.01.004.
- Jaradat, N., A. Eid, F. Abdelwahab, L. Isa, A. Abdulrahman, M. Abualhasan, and A.M.A. Hussein. 2015. "Phytochemical Analysis, Quantitative Estimations of Total Phenols and Free Radical Scavenging Activity of *Bupleurum subovatum* from Jerusalem. *Pharmaceutical Sciences*. 21.
- Kessler, A., I. T. Baldwin. 2001. Defensive Function of Herbivore-Induced Plant Volatile Emissions in Nature. *Science* 291: 2141-2144
- Mohammadi, M. and H. Kazemi. 2002. Changes in peroxidase and polyphenol oxidase activities in susceptible and resistant wheat heads inoculated with *Fusarium graminearum* and induced resistance. *Plant Science*. 162:491-498.
- Sabbobeh, R., H. Hejaz, A. Jahajha, S. Al-Akhras, H. Al-Jaas, and S. Abu-Lafi. 2016. "Antioxidant and Antimicrobial Activities of the Leaf Extract of *Salvia palaestina*. *Journal of Applied Pharmaceutical Science* Vol. 6:076-082.
- SAS Institute. 2008. *SAS Users Guide: Statistics*. Version 2. SAS Institute, Cary, NC, USA.
- Sighamony, S., Anees, I., Chandrakala, T. and Osmani, Z., 1986. Efficacy of certain indigenous plant products as grain protectants against *Sitophilus oryzae* (L.) and *Rhyzopertha dominica* (F.). *Journal of Stored Products Research*, 22(1): 21-23
- Spoel, S. H., and X. Dong. 201. How do plants achieve immunity? Defence without specialized immune cells. *Nature Reviews Immunology*, 12(2): 89-100.
- Tolman JH, McLeod DGR, Harris CR (2004) Cost of crop losses in processing tomato and cabbage in southwestern Ontario due to insects, weeds and/or diseases. *Canadian Journal of Plant Science*:915-921.
- Vincelli, P. 2016. Genetic Engineering and Sustainable Crop Disease Management: Opportunities for Case-by-Case Decision-Making. *Sustainability*. 8:495.

ORAL PRESENTATION

**INVESTIGATION OF ANTIMICROBIAL ACTIVITY OF SOME PLANTS AGAINST
E. COLI ISOLATED FROM BLOOD CULTURES**

Tuba Kuşman¹, Abdullah Dalar², Ömer Akgül³, Gülhan Bora³

¹ Department of Pharmacogony, Faculty of Pharmacy, Van Yuzuncu Yil University, 65080, Van, Turkey

² Department of Pharmaceutical Botany Faculty of Pharmacy, Van Yuzuncu Yil University, 65080, Van, Turkey

³ Department of Pharmaceutical Microbiology, Faculty of Pharmacy, Van Yuzuncu Yil University, 65080, Van, Turkey

Objective / Purpose: It is well known that plant-derived antimicrobials are often more efficient than antibiotics, even on resistant pathogens, likely owing to plant compounds synergistic activity [1,2].

Material and Methods: In this study, Plant samples were collected from Konalga village, Çatak/Van city, in the Eastern Anatolia on August 18th, 2018. the ground plant material was mixed with a 20-fold volume of acidified ethanol (80% ethanol, 19.9% H₂O and 0.1% trifluoroacetic acid, v/v/v), shaken for 2 h at room temperature (22°C) and centrifuged for 20 min at 4°C with the supernatant collected. The supernatants from the consecutive extractions were combined. Solvent was evaporated under reduced pressure at 40°C. The derived fraction was dissolved in purified water and freeze-dried under a vacuum at - 51°C to obtain a fine lyophilized powder used for analysis.

Thirty-seven *E. coli* were isolated from the patients in the intensive care unit. Disc diffusion and microdilution method were used to determine the sensitivity of *E. coli* isolates to plant extracts [3]. The plant extracts were dissolved in 12.5% dimethylsulfoxide and then sterile empty antibiotic discs (10 µL each) were absorbed.

Results: According to the disc diffusion test, it was observed that a low sensitivity diameter was formed against the extracts from *Helicrysum pallasii*, *Prangos meliocarpoides*, *Epilobium hirsutum*. In the *Verbascum cheiranthifolium* extract, it was determined that the sensitivity diameter did not occur. According to the microdilution method, *Helicrysum pallasii* showed sensitivity at 1000 to 250 µg/mL, *Prangos meliocarpoides* showed sensitivity at 1000 µg/mL, *Epilobium hirsutum* showed sensitivity at 1000 to 500 µg / mL and *Verbascum cheiranthifolium* showed no sensitivity.

Conclusion: The antimicrobial efficacy of the plant extracts on *E. coli* was found to be different and the antimicrobial effect was reduced in low concentrations.

Keywords: Antimicrobial, plant extracts, *E. coli*, disc diffusion test

References:

- [1] Pirvu, L., Nicorescu, V., Hlevca, C., Udeanu, D. I., & Nicorescu, I. (2015). Antimicrobial and synergistic activity of some whole and selective *Epilobium hirsutum* l.(great willowherb) extracts tested on standard and wild *Staphylococcus aureus* strains. *Farmácia*, 63(5), 690-695.
- [2] Bouarab Chibane, L., Degraeve, P., Ferhout, H., Bouajila, J., & Oulahal, N. (2018). Plant antimicrobial polyphenols as potential natural food preservatives. *Journal of the Science of Food and Agriculture*.
- [3] CLSI, C. (2012). Performance standards for antimicrobial susceptibility testing. Clinical and Laboratory Standards Institute (M100eS22), (s22nd Informational Supplement).

ORAL PRESENTATION

**THE ROLE OF MEDICINAL PLANTS USED IN TRADITIONAL MEDICINE FOR
DRUG DISCOVERY AND THEIR CONSERVATION IN PAKISTAN**

Muhammad Zafar¹

Department of Plant Sciences, Quaid-i-Azam University, Islamabad, Pakistan.
zafar@qau.edu.pk

Pakistan has a unique diversity of medicinal plants due to its geography and diverse environmental and ecological conditions. Medicinal plants are frequently used among the different communities in the world generally and in Pakistan predominantly for treating numerous diseases. Due to recent emerging trends in traditional medicine research and interest to use medicinal plants, a large portion of the medicinal plants has been exhausted, demonstrating a lack of traditional knowledge among communities using the plants. Therefore, an attempt is made in this study to concentrate on the role of medicinal plants used in traditional medicine for drug discovery. Questionnaire using Semi-structured interviews were conducted to list the medicinal uses of plants. Results were analyzed using different quantitative indices including Informant consensus factor (ICF), fidelity level (FL), use value (UV), frequency citation (FC) and relative frequency citation (RFC). In total of 101 plants species were reported for the medicinal purposes. Asteraceae was found to be most dominant family.

The leaves were noted as most frequently used parts (66%). Powder (54%) was the most commonly used preparation method. The highest ICF was reported for Glandular diseases (Diabetes Jaundice, diabetes, tonsils, hepatitis, liver inflammation). Fidelity level varies from 55 % to 97%. Species having high UV were *Cannabis sativa*, *Oxalis corniculata*, while the plant species with the highest RFC were *Ficus palmata*. High use value of medicinal plant should be screened for the pharmacological investigation for bio active phytochemical compounds that are essential for the synthesis of new drugs for numerous diseases. Their conservation is the need of the time for sustainable use and plant biodiversity.

Keywords: Diverse environment, Medicinal plants, Traditional medicine, Quantitative indices and plant biodiversity conservation.

ORAL PRESENTATION

**ESSENTIAL OIL COMPOSITION OF *CENTRANTHUS LONGIFLORUS* STEVEN
SUBSP. *LONGIFLORUS* FROM TURKEY**

Gülsüm Yıldız¹, Yavuz Bülent Köse², ¹Mine Kürkçüoğlu¹

¹ Department of Pharmacognosy, Anadolu University, Eskişehir, Turkey, gulsumyildiz@anadolu.edu.tr

² Department of Pharmaceutical Botany, Anadolu University, Eskişehir, Turkey

Objective / Purpose: *Centranthus* DC. (Caprifoliaceae) is represented by three species in the Flora of Turkey (1). *Centranthus longiflorus* Steven subsp. *longiflorus* widely distributes Northern, Southern and Central Anatolia and is used sedative and antispasmodic in the folk medicine and plant leaves are consumed as a spice in Turkey (2,3). The aim of this study is to determine the essential oil composition of aerial parts of *C. longiflorus* subsp. *longiflorus* collected from Tokat province of Turkey.

Material and Method: Essential oil of *C. longiflorus* subsp. *longiflorus* was hydrodistilled from aerial parts for 3 h using a Clevenger apparatus and analysed by gas chromatography (GC), and gas chromatography-mass spectrometry (GC-MS).

Results: The main constituents were identified as *neophytodiene* (19.5 %), *germacrene D* (13.8 %), *phytol* (8.9 %) and δ -*cadinene* (6.9 %) in the essential oil.

Conclusion / Discussion: The present work is the first contribution into the oil composition of *C. longiflorus* subsp. *longiflorus* from Turkey.

Keywords: *Centranthus longiflorus* subsp. *longiflorus*, Caprifoliaceae, essential oil, GC-FID, GC-MS

References:

- [1] Davis PH. (1972). In flora of Turkey and East Aegean Islands. Edinburgh University Press, Vol. 4 Edinburgh, UK, 558.
- [2] Suleyman, H., Guvenalp, Z., Kizilkaya, M., & Demirezer, L. O. (2007). Sedative effect of *Centranthus longiflorus* ssp. *longiflorus* in rats and the influence of adrenalectomy on its effect. *Yakugaku Zasshi*, 127(8), 1263-1265.
- [3] Kaval, İ., Behçet, L., & Çakılcıoğlu, U. (2015). Survey of wild food plants for human consumption in Geçitli (Hakkari, Turkey).

ORAL PRESENTATION

**INFLUENCE OF ANTIOXIDANT CAPACITY OF PLANT EXTRACTS ON
ENHANCEMENT OF ADAPTIVE CELL-MEDIATED RESPONSE IN
HERBIVORES**

Carmen Dana Șandru¹, Silvana Popescu², Mihaela Niculae¹, Elena Andruța Mureșan³, Brudașcă Gheorghe Florinel¹, Eموke Pall¹, Adrian Potarniche¹, Constantin Cerbu¹, Aurel Vasiu¹, Marina Spînu¹

¹Department of Clinical Sciences, Division of Infectious Diseases, University of Agricultural Sciences and Veterinary Medicine, Faculty of Veterinary Medicine, Cluj-Napoca, Romania

²Department of Animal production, Division of Hygiene and Welfare, University of Agricultural Sciences and Veterinary Medicine, Faculty of Veterinary Medicine, Cluj-Napoca, Romania

³Food Engineering Department, Faculty of Food Science and Technology, University of Agricultural Sciences and Veterinary Medicine Cluj-Napoca, Romania

Objective: This study monitored several alcoholic vegetal extracts from the antioxidant and immune cell stimulating adjuvant perspective in farmed herbivores, differing by their digestive physiology.

Material and Methods: Blood from randomly selected farmed ruminants (Romanian Spotted dairy cows, n=28 and Angora goats, n=19) and Romanian draft horses (n=27) was used to measure the *in vitro* effects on specific cell-mediated reactivity of alcoholic extracts of *Calendula officinalis*, *Echinacea angustifolia* and *E. purpurea*, by the *in vitro* whole blood blast transformation indices (BI%) after 60-72h of incubation. Cell growth was quantified by an orto-toluidine technique. All extracts were tested for their antioxidant capacity (free radical scavenging effect over DPPH (%)). Student's t- test was used to evaluate the significance of the differences.

Results: The antioxidant activity of plants rendered as follows: *E. purpurea*, *E. angustifolia* and *C. officinalis*. The *in vitro* growth indices were lowest in goats (*C. officinalis* - 58.52±10.02%, *E. angustifolia* - 50.06±11.67%, *E. purpurea* -50.79±10.98%) and higher in bovine (*C. officinalis* – 69.9±2.65%, *E. angustifolia* - 74.9±10.1%, p<0.05), and increased towards *E. angustifolia* versus *C. officinalis*. Similar BI% were obtained for the *Echinacea* species, which differed by their antioxidant capacity.

In vitro responses to *C. officinalis*, *E. angustifolia* and *E. purpurea* were the most pronounced similar for all extracts and independent on their antioxidant activity in horses.

Conclusion / Discussion: All the extracts showed immune-inhibiting effects in bovine and goats, but not in horses, supporting host-based differences. The immunological activity of the tested extracts was rather plant species than antioxidant activity dependent.

Keywords: Ruminants, equine, plant extracts, antioxidant, cell mediated immunity

References:

[1] Pellati F., Benvenuti S., Magro L., Melegari M., Soragni F. (2004) Analysis of phenolic compounds and radical scavenging activity of *Echinacea spp.* Journal of Pharmaceutical and Biomedical Analysis 35:289–301

ORAL PRESENTATION

BROMELAIN: A POSSIBLE PHYTOREMEDY TOWARDS BREAST CANCER

Amruta Kothare¹ Vinal Pardhi² Prajkta Chivte², Sakharam Muley¹ Tejashree Shanbhag*¹

1 Department of Life Sciences, K.C College, Churchgate, Mumbai- 400020, India

2 National Facility for Biopharmaceuticals, G. N. Khalsa College, Matunga, Mumbai-400019, India

Email - tvs310168@gmail.com

Objective: Bromelain is a cysteine protease present in various parts of pineapple (*Ananas comosus*) (L.Merr.). Bromelain has got clinical significance as anti-thrombotic, anti-inflammatory, agent etc. The current work aims at purification and comparative study of Bromelain activity in fruit as well as waste parts (core, crown, skin, and stem) of pineapple and their potential application as anti-cancer agent towards breast cancer.

Materials and methods: The enzyme was extracted and precipitated by ammonium sulphate (40%), dialysed and concentrated by TFF (tangential flow filtration) and further purified using IEX (ion exchange chromatography). The purified fractions were analysed by SDS-PAGE and MALDI ToF to confirm the molecular weight of the eluted protein. The activity of purified enzyme was confirmed by Casein Digestion activity assay as well as Zymography. The presence of a glycan moiety in the bromelain protease was detected using novel proteoglycan staining. The anticancer activity of potential fractions was determined by MTT assay on MCF-7 cell lines.

Results: The molecular weight of the purified enzyme from all parts fractions was 23kDa. The fold of purification for Bromelain proteases in the core, crown, fruit, skin and stem was 5.54, 3.04, 0.41, 12.58 and 18.49 respectively. Statistical analysis suggested specific activity optimum at 45°C temperature while the pH range to be 6.0-7.0. The results of SRB assay state that skin and crown bromelain extracts that are waste parts of pineapple, have GI50 value ≤ 10 making them potential cytotoxic agents towards breast cancer.

Conclusions: From the above study, bromelain protease one of the constituents of bromelain complex from pineapple fruit and its waste parts was purified and recovered with highest activity and purification fold. The potential cytotoxicity on breast cancer cell line MCF-7 was obtained from bromelain present in skin and crown of pineapple which are waste parts. This gives us an opportunity to explore their potential further and harness their efficiency in anticancer drug therapy or adjuvant/targeted drug delivery.

Keywords: Protease, Purification, Zymography, anticancer.

References:

- [1] V. Pardhi, P. Chivte, A. Kothare, T. Shanbhag, F. Biopharmaceuticals, G. N. K. College, L. Sciences, and K. C. College, "Research Article," vol. 7, 2016.
- [2] K. Babagana and M. Bala, "Comparative Study on The Extraction and Purification the Stem and Fruit Bromelain from Pineapple (*Ananas comosus*)," J. Nat. Sci. Res. www, vol. 5, no. 18, pp. 2225–921, 2015.

ORAL PRESENTATION

CHEMICAL COMPOSITION, ANTIOXIDANT ACTIVITY AND EFFECT ON HUMAN FIBROBLAST VISIBILITY OF EXTRACTS FROM OLEASTER (*ELAEAGNUS UMBELLATE*) FRUIT COLLECTED IN POLAND.

Zglinska Klara, Matusiewicz Magdalena, Daniluk Karolina, Balaban Jasmina, Niemiec Tomasz
Department of Animal Nutrition and Biotechnology, Faculty of Animal Science, Warsaw University of Life Sciences, Poland

Objectives: Oleaster is a plant come originally from Asia, where it is used in traditional medicine. Studies from Asian countries have shown that oleaster fruit extracts have a strong antioxidant activity. We wanted to investigate whether fruits growing in other climatic conditions will also exhibit these properties. The objective of this study was to determine the chemical composition of oleaster fruits grown in central Europe and the redox properties of berry extracts. In order to verify its potential use as medicine or cosmetic, cytotoxicity assays with fibroblast from human foreskin were made.

Methods: The fruit was collected from the farm located in Poland. Berry was analyzed for proximate contents according to AOAC International. Total phenols content was measured by the Folin–Ciocalteu method and mineral content by ICP-AES method.

Then, lyophilized samples were extracted with methanol. The solvent was evaporated and antioxidant activity was measured by DPPH and ferric reducing antioxidant power (FRAP) assays. Different concentrations of extract (1-6mg/ml) were used. As a reference, TROLOX standard curve (20-100 μ M) was made.

Furthermore, in vitro cell viability analysis was performed. Human normal fibroblast cell (HFFF-2) were incubated with oleaster extract (as 10% of medium). Cell Proliferation Kit II (XTT, Sigma-Aldrich) were made according to manufacturer protocol.

Results: Fresh berry contained 30,94% dried matter, 4,97% crude protein, 2,13% crude fat and 4,88% crude fiber. Proximate content of fruit collected in Poland was similar to fruit analyzed in Turkey [1]. Extracts showed the ability to reduce iron ions. The strength of the reaction was dose-dependent. Lyophilized extracts were an equivalent of 2,32 and 4,08 μ M Trolox for 3mg/ml and 6 mg/ml. DPPH assay show 86,91% inhibition at 6 mg/ml concentration. The XTT test showed no negative effect of extracts on human fibroblast cell viability. There was a trend towards increased proliferation in cells receiving oleaster extract (n/s).

Conclusions: The results of the study indicate that oleaster can be a valuable and safe raw material for cosmetics. The use of this fruit can be much wider because fibroblasts are the most common resident cells in ordinary connective tissue.

Keywords: Oleaster, autumn olive, *Elaeagnus umbellata*, antioxidant activity, fibroblast

References:

[1] Khattak K. (2012) Free radical scavenging activity, phytochemical composition, and nutrient analysis of *Elaeagnus umbellata* berry. J. Med. Plants Res. 6(39):5196-5203.

ORAL PRESENTATION

WILD PLANTS USED AS FOOD IN EDIRNE CENTRAL DISTRICT AND VILLAGES

Fatma Güneş

*Trakya University, Faculty of Pharmacy, Department of Pharmaceutical Botany, Edirne, Turkey.
drgunes@gmail.com*

Objective: The aim of this study is to identify the wild plants used as food by local people in Edirne province and villages, their local and Latin names, the parts used, forms of utilization.

Material and Methods: The study is executed in March-October in the years of 2013-2015 in Edirne and surrounding villages. Interviews were carried out face-to-face with the community. In this study, 18 villages of Edirne province were visited and interviews were performed with 26 persons in total. The identification of the plant species those determined to be in use was based on “Flora of Turkey and East Aegean Islands” (1,2). The plant samples identified are kept in Faculty of Pharmacy.

Results: As result of in this study, 41 taxa including in 24 families which are wild plants used as food, were recorded. In the table; the scientific names of the plants, families, local names, usable parts and utilization have been given alphabetically.

Conclusion: Previously, some ethnobotanical studies carried out in Trace (3-8). There is no study about wild plants used as food in Edirne province and villages.

Keywords: Edirne, ethnobotany, food, wild plants

Acknowledgement: I thanks to all the villagers of Edirne province, who have collaborated in the realization of this study. The study was carried out with the support of Trakya University (Project 2013/22).

References:

- [1] Davis, P.H., Miller, R.R., Tan, K. (1965-1985). *Flora of Turkey and the Aegean Islands*. Vol. 1-9, Edinburgh University Press, Edinburgh.
- [2] Güneş, A., Aslan, S., Ekim, T., Vura, I M., Babaç, M.T. (edlr) (2012). *List of Turkey Plants* (Vascular Plants). Nezahat Gökyiğit Botanik Garden and Flora Researchers Association Press, İstanbul.
- [3] Kültür, Ş., 2008. An ethnobotanical study of Kırklareli (Turkey), *Phytologia Balcanica* 14 (2), 279-289
- [4] Tuzlacı E, İşbilen DFA, Bulut G. (2010). Turkish folk medicinal plants, VIII: Lalapaşa (Edirne). *Marmara Pharmaceutical Journal*, 14:47–52.
- [5] Güneş, F., & Özhatay, N. (2011). An ethnobotanical study from Kars (Eastern) Turkey. *Biological Diversity and Conservation*, 4(1), 30-41.
- [6] Güneş, F. (2017). Food Plants Used in Meriç Town from Turkey. *Indian J of Pharmaceutical Education and Research*, 51(3).
- [7] Güneş, F. (2017). Medicinal plants used in the Uzunköprü district of Edirne, Turkey. *Acta Societatis Botanicorum Poloniae*, 86(4).
- [8] Güneş, F. (2018). An Ethnobotany Study in Enez Town from Edirne. *Current Perspectives on Medicinal and Aromatic Plants (CUPMAP)*, 1(1), 28-35.

ORAL PRESENTATION

CORRELATION BETWEEN CHEMICAL COMPOSITION AND ANTIOXIDANT ACTIVITY OF ESSENTIAL OILS FROM SIX AROMATIC MEDICINAL PLANTS GROWING IN ILLIZI AND GHARDAÏA (SOUTHERN ALGERIA)

Naima Hellali^{1,2}, Mebarka Bouziane², Mahfoud Hadj Mohammed²

¹ Faculty of Sciences and Technology, Department of process engineering, Ghardaïa University, BP 455, 47000 Ghardaïa, Algeria, hellali.naima@univ-ghardaia.dz

²Biogeochemistry Laboratory in Desert Environments, Faculty of Science, Kasdi Merbah University, BP 511, 30000 Ouargla, Algeria.

This study provides an assessment to the relationship between the chemical composition of the essential oils and the antioxidant activity of six aromatic medicinal plants growing in southern Algeria, namely *Cymbopogon schoenanthus*, *Cymbopogon Citratus*, *Artemisia judaica*, *Artemisia herba alba*, *Pituranthos chloranthus* and *Pituranthos scoparius*. The essential oils were obtained by hydrodistillation and analyzed by GC/MS. Their antioxidant activity was determined by using three methods of testing (DPPH, ABTS and iron reduction test (FRAP)), the results showed that the essential oil of *A. judaica* has the best antioxidant capacity in all tests. The statistical study allowed us to show a correlation between the chemical composition of the studied essential oils and the antioxidant activity and our results indicate that the high content of cyclic ether sesquiterpenes compounds (such as: Davana ether), the compounds have furan rings, or the existence of non-terpene alkenes/esters in the essential oil, increases its antioxidant effect.

Keywords: *C. schoenanthus*, *C. Citratus*, *A. judaica*, *A. herba alba*, *P. chloranthus*, *P. scoparius*, essential oil, antioxidant activity.

References:

- [1] Cimanga, K., Kambu, K., Tona .L., Apers, S., De Bruyne, T., Hermans, N., Totté, J., Pieters, L., Vlietinck, A.J., (2002), Correlation between chemical composition and antibacterial activity of essential oils of some aromatic medicinal plants growing in the Democratic Republic of Congo, *Journal of Ethnopharmacology*, 79(2), 213-220.
- [2] Budel, J.M., Wang, M., Raman, V., Zhao, J., Khan, S.I., Rehman, J.U., Techen, N., Tekwani, B., Monteiro, L. M., Heiden, G., Takeda, I. J. M., Farago, P.V., Khan, I. A., (2018), Essential Oils of Five Baccharis Species: Investigations on the Chemical Composition and Biological Activities, *Molecules*, 23, 2620; doi:10.3390/molecules 23102620.
- [3] Djeridane, A., Yousfi, M., Nadjemi, B., Boutassouna, D., Stocker, P., Vidal, N., (2006), Antioxidant activity of some algerian medicinal plants extracts containing phenolic compounds, *Food Chemistry*, 97(4), 654-660.
- [4] Hellali, N., Hadj Mohammed, M., (2008), Analyse par GC-MS de l'huile essentielle de la plante *Cymbopogon schoenanthus* de la région de Illizi, obtenue par hydrodistillation assistée par microondes. *PhytoChem. BioSub J.* 2(1):5.
- [5] Hellali, N., Hadj Mohammed, M., Ramdane, F., Talli, A., (2016), Antimicrobial and antioxidant activities of *Cymbopogon schoenanthus* (L.) spreng. essential oil, growing in Illizi – Algeria. *Journal of Medicinal Plants Research.* 10(14), 188-194.
- [6] Hellali, N., Hadj mahammed, M., Masrouk, H., (2017), Physico-chemical Study and Evaluation of Antimicrobial and Antioxidant Activities of *Artemisia judaica* L. Essential Oil, Growing in Illizi, Algeria. *Asian Journal of Chemistry*, 29(1), 181-186.
- [7] Elaïssi, A., Rouis, Z., Mabrouk, S., Bel Haj Salah, K., Aouni, M., Khouja, M. L., Farhat, F., Chemli, R., Harzallah-Skhiri, F., (2012), Correlation Between Chemical Composition and Antibacterial Activity of Essential Oils from Fifteen Eucalyptus Species Growing in the Korbous and Jbel Abderrahman Arboreta (North East Tunisia), *Molecules*, 17: 3044-3057.

ORAL PRESENTATION

THYMUS SIPYLEUS BOISS. SUBSP. ROSULANS (BORBAS) JALAS EXTRACTS AS ALTERNATIVE THERAPEUTIC AGENTS FOR THE TREATMENT OF SKIN INJURY

Oya Ustuner^{1a}, Ceren Anlas^{1b}, Tulay Bakirel^{1c}, Fulya Ustun Alkan^{1d}, Belgi Diren Sigirci^{2a}, Seyyal Ak^{2b}, Huseyin Askin Akpulat³

^{1a} Department of Pharmacology and Toxicology, Faculty of Veterinary Medicine, Istanbul University-Cerrahpasa, Avcilar 34320, Istanbul, Turkey, ovakeles@istanbul.edu.tr

^{1b} Department of Pharmacology and Toxicology, Faculty of Veterinary Medicine, Istanbul University-Cerrahpasa, Avcilar 34320, Istanbul, Turkey, cerenis@istanbul.edu.tr

^{1c} Department of Pharmacology and Toxicology, Faculty of Veterinary Medicine, Istanbul University-Cerrahpasa, Avcilar 34320, Istanbul, Turkey, tulbakir@istanbul.edu.tr

^{1d} Department of Pharmacology and Toxicology, Faculty of Veterinary Medicine, Istanbul University-Cerrahpasa, Avcilar 34320, Istanbul, Turkey, fustun@istanbul.edu.tr

^{2a} Department of Microbiology, Faculty of Veterinary Medicine, Istanbul University-Cerrahpasa, Avcilar 34320, Istanbul, Turkey, belgis@istanbul.edu.tr

^{2b} Department of Microbiology, Faculty of Veterinary Medicine, Istanbul University-Cerrahpasa, Avcilar 34320, Istanbul, Turkey, sak@istanbul.edu.tr

³ Department of Biology Education, Faculty of Education, Sivas Cumhuriyet University, 58140 Sivas, Turkey, hakpulat@cumhuriyet.edu.tr

Objective / Purpose: Wound can be described as a disruption in the structural and physiological continuity of skin, as a result of chemical, physical or immunological factors. In recent years, the demand for medicinal plants has increased globally in the management of wound healing, due to their less toxicity and fewer side effects. The extracts from *Thymus sipyleus* Boiss. subsp. *rosulans* (Borbab) J alas (TS) which is an endemic plant for the flora of Turkey, is used for the treatment of skin wounds in Turkish folk medicine. In this study, it was aimed to explain the mechanisms underlying the wound healing activity of this plant and to support the traditional use of TS extracts scientifically.

Material and Methods: The effects of different solvent (ethanol, n-hexane and water) extracts obtained from TS on wound healing were investigated on *in vitro* test systems, in relation to different phases of healing process. The effects of TS extracts on the inflammatory phase were determined by tumour necrosis factor alpha (TNF- α) assay and nitric oxide (NO) assay in RAW 264.7 mouse macrophages. The effects of extracts on proliferative phase were evaluated by MTT and Scratch assays on swiss 3T3 albino mouse fibroblast. Hydroxyproline assay was performed to elucidate the influence of TS extracts on collagen production. Antioxidant activities of extracts were determined by DPPH assay and Folin-Ciocalteu assay. As well as, the cytoprotective effects of extracts against H₂O₂-induced oxidative damage were evaluated in the co-treatment and pre-treatment protocols. Finally, antimicrobial activities of extracts were investigated against the most common bacteria found in skin wounds.

Results: We observed that TS extracts influenced the proliferative and inflammatory phases of healing process by increasing the proliferation and migration of fibroblasts and down-regulating the level of TNF- α and NO. The highest increase in the level of hydroxyproline was determined in cells treated with decoction extract (1.79 to 2.40 fold) at all applied concentrations. In addition, our results showed that TS extracts have antioxidant and antibacterial potential. The percentage of DPPH radical scavenging activities of extracts were ranged from 74.32% to 93.02%, at the highest concentration. Total phenolic compounds of extracts were calculated at the range of 78,15- 147,64 mg GAE/ g. DPPH radical scavenging activities and total phenolic content of n-hexane (non-polar) extracts were too low to be detected. All *Thymus* extracts exhibited complete protection against H₂O₂ induced oxidative damage on mouse fibroblasts.

Conclusion / Discussion: The extracts from TS accelerated the wound healing process by increasing fibroblast proliferation and migration as well as collagen synthesis and decreasing inflammation. We also concluded that TS extracts have potent effect on the protection of cells from oxidative damage, in relation to the antioxidant activity of plant. Our results indicated that TS can be an effective source to develop a therapeutic agent against skin wounds.

Keywords: Medicinal plants, *Thymus sipyleus* subsp. *sipyleus* var. *rosulans*., wound healing

ORAL PRESENTATION

ACUTE AND DERMAL TOXICITY STUDIES OF *ADANSONIA DIGITATA* L. SEEDS OIL ON WISTAR ALBINO RATS

Reem Hassan Ahmed¹ and El khatim Hassan Abdelgadir²

1: *Biology and Biotechnology Department, College of Applied and Industrial Sciences, University of Bahri, Sudan.*

2: *Pharmacology and Toxicology Department, Medicinal and Aromatic Plants and Traditional Medicine Research Institute, National Center for Research, Sudan*

Introduction: *Adansonia digitata* L. (Baobab tree) of family; Malvaceae is a multi-purpose tree species native to Africa. It is found abundantly in Western parts of Sudan and has been used extensively since ancient times in traditional medicine. Baobab seed oil has quality comparable with those of commercial oil and it has great nutritional and industrial potentials. Thus, research is recommended to validate its efficacy and safety for use.

Objectives: The present work is designed to evaluate the acute oral toxicity of *A. digitata* seeds oil and its observational dermal toxicity.

Methods: Three different doses (50, 250, 500 mg/kg) of the oil were administered to 3 groups of Wistar albino rats for 7 consecutive days. Hematology and biochemical parameters (AST, ALT, ALP, total protein, albumin, cholesterol and urea) values were investigated and compared to control un-treated rats. Dermal toxicity was assessed in another group of animals for 14 days' time.

Results: Oral administration of the oil caused a significant; dose dependent increases in AST, ALT and urea levels compared to the control. Gross pathological studies revealed congested heart and fatty changes in liver and kidney. While, dermal toxicity study revealed no observational signs of toxicity or irritation.

Conclusion: Oil extracted from *Adansonia digitata* L. seeds is safe to be used externally, however, it showed toxicity signs on oral administration.

Keywords: *Adansonia digitata*, Baobab oil, acute toxicity.

References:

- [1] Birnin U.A., Yauri and Garba S. (2011), Comparative Studies on Some Physicochemical Properties of Baobab, Vegetable, Peanut and Palm Oils. *Nigerian Journal of Basic and Applied Science*. 19(1):64-67
- [2] Emmy De Caluwé, Kateřina Halamová and Patrick Van Damme (2010) *Adansonia digitata* L. – A review of traditional uses, phytochemistry and pharmacology. *Afrika focus* — Volume 23, Nr. 1, pp. 11-5.
- [3] Environmental Protection Agency. Prevention, Pesticides and Toxic Substances (7101) EPA 712-C-98-192. August 1998. Health Effects Test Guidelines. OPPTS 870.1200. Acute Dermal Toxicity.

ORAL PRESENTATION

DETERMINATION OF HYBRIDIZATION IN *NEPETA* × *TMOLEA* BOISS. USING DIFFERENT APPROACHES

Taner Özcan

Department of Biology Education, Necatibey Education Faculty, Balıkesir University, Balıkesir, Turkey,
ozcant@balikesir.edu.tr

Objective / Purpose: The main purpose of this study is to examine morphological, micromorphological and molecular features of *Nepeta* × *tmolea* Boiss. hybrid specimens and its parents. For this purpose, trichome types and density are especially studied and one nuclear and two chloroplast DNA regions were sequenced, and after that distinct characters were revealed.

Material and Methods: Plant specimens were collected during the flowering stage (2016-2018) from natural habitats in Balıkesir, İzmir, and Muğla. *N.* × *tmolea* and *N. nuda* L. subsp. *nuda* were collected from Balıkesir and İzmir, and *N. viscida* Boiss. were picked up all of these locations. Stems, leaves, calyces, corollas, and nutlets of *N. viscida*, *N. nuda* subsp. *nuda* and their putative hybrid specimens *N.* × *tmolea* were investigated and photographed using a NeoScope JCM scanning electron microscopy. DNA isolations were performed using the DNeasy Plant Mini Kit, and the internal transcribed spacer (ITS) region of the nuclear ribosomal DNA (nrDNA) sequences and the rpl32-trnA(UAG) and trnA(Leu)-trnA(Phe) (trnL-F) regions of the chloroplast DNA (cpDNA) were used for molecular analysis. The PCR amplifications were performed using Shaw et al. (2007) protocol. Sequenced DNA was edited using Sequencer version 4.9.

Results: Field observations are very important to find out general characters. *N.* × *tmolea* has some intermediate characters between its parents. According to our findings, *N.* × *tmolea* specimens resembled *N. viscida* as general, they have a dense indumentum structure especially on their stems and leaves, but *N.* × *tmolea* samples were not viscid, and their stem, leaf and corolla colors were quite different from *N. nuda* subsp. *nuda* and *N. viscida*. 16 different individuals were sequenced to define their nrITS DNA data, and 32 different individuals for trnL-F and 15 different individuals for rpl32-trnA were sequenced to obtain these chloroplast DNA data. The most single nucleotide polymorphisms were obtained in nrITS data, but chloroplast DNA data has more ancestral information due to their insertions and deletions.

Conclusion / Discussion: Our data showed that the genetical methods are more compatible do define hybrid specimens and their parents. But, only one region could not be informative, different genome regions should be appropriate. *N.* × *tmolea* possibly has two DNA data from its parents.

Keywords: Hybridization, trichome, molecular, *N.* × *tmolea*.

References:

Shaw J, Lickey EB, Schilling EE, Small RL (2007). Comparison of whole chloroplast genome sequences to choose noncoding regions for phylogenetic studies in angiosperms: the tortoise and the hare III. *Am J Bot* 94: 275-288.

ORAL PRESENTATION

ANTI-FERTILITY AND ABORTIFACIENT EFFECT OF AQUEOUS SPROUTED SEEDS EXTRACT OF *TRIGONELLA FOENUM GRAECUM* (L.) IN MICE.

Oufkuir Sara¹, Ait Laaradia Mehdi¹, Laadrawi Jawad¹, Aboufatima Rachida², Chait Abederrahmane¹, Sokar Zahra¹.

¹: *Laboratory de Pharmacology, Neurobiology and Behavior, Cadi ayyad University, Semlalia faculty, Marrakesh, Morocco*

²: *Laboratory of Biological Engineering, Faculty of Science and Technology, Beni Mellal, Morocco*

Introduction: *Trigonella foenum graecum* or fenugreek is one of the oldest medicinal plants; the sprouts are used as spices and condiment to improve the flavor and the nutritive value of foods. Seeds have been revealed teratogenic and anti-fertilizing as it shown in our previous study carried out in Swiss albino mice (1), reason for why, we were interested in evaluating of sprouted seeds effect in mice reproduction.

Material and methods: Female mice have been treated orally with lyophilized aqueous extract of sprouted organic fenugreek seeds, from the first day of pregnancy until delivery, at doses of 0 (control), 200, 500, 800, 1000 mg/kg/day. Animals were observed for any intoxication symptoms all over the period of pregnancy. Additionally, Pregnancy abortion and deliverance indexes have been determined as follows:

- Pregnancy index: the percentage of pregnant females which have shown a vaginal plug.
- Abortion index: the percentage of the number of aborted offspring over pregnant animals.
- Deliverance index: the percentage of females delivering over pregnant animals.

The main phenolic compounds of lyophilized aqueous extract of germinated fenugreek seeds has been analyzed by high-performance liquid chromatography.

Results: all females survived until study termination, no abnormal clinical symptoms were observed in all groups, at any time during gestation. Oppositely, for the reproduction function, we have noticed a significant decrease of pregnancy index especially at 800 and 1000 mg/kg/day with a 25 % of fertility loss. The abortive effect starts at 500 mg/kg/day with 42% and increases to 66 % at 1000 mg/kg. The deliverance index shows a big decreasing at 800 and 1000 mg/kg where only 33 % of female mice end their gestation period. Spontaneous abortion took place between gestation days 16 and 18. HPLC analysis revealed the presence of gallic acid, caffeic acid, tyrosol, syringic acid and flavonoid rutin.

Discussion and conclusion: this result is important because it shows, for the first time, one of the negative aspects of use of germinated fenugreek seeds. Basing on this finding we could suggest that consumption of sprouted seeds of fenugreek should be avoided during pregnancy.

Keywords: Fenugreek, abortive, anti-fertilizing, sprouted seeds.

Reference

Khalki L, M'hamed SB, Bennis M, Chait A, Sokar Z, 2010, Evaluation of the developmental toxicity of the aqueous extract from *Trigonella foenum-graecum* (L.) in mice, *J Ethnopharmacol.*, 131(2), 321-5.

ORAL PRESENTATION

**FROM A TRADITIONAL REMEDY TO MODERN THERAPY; *IN VIVO*
ANTHEMORRHOIDAL STUDY OF *MALVA SYLVESTRIS* L.**

Ceylan Dönmez¹, Ufuk Koca Çalışkan¹, Petek Korkusuz², Özge Boyacıoğlu²

¹Department of Pharmacognosy, Faculty of Pharmacy, Gazi University, 06330, Ankara, Turkey
akaceylan@gmail.com

²Department of Histology and Embryology, Faculty of Medicine, Hacettepe University, 06100, Ankara, Turkey

Objective / Purpose: It is thought that about 50% of the world's population is suffering from hemorrhoids, an anorectal disease. Patients have tried to cope with the disease with both herbal and chemical products. *Malva sylvestris* belongs to Malvaceae family and its common English name is mallow, whereas Turkish name, is 'Ebegümeçi' in Turkey. It's leaves have potent anti-inflammatory, antioxidant, anti-complementary, anticancer and skin tissue integrity activity [1,2]. In this study, *in vivo* antihemorrhoidal and antiinflammatory activities of *M. sylvestris* aerial part extract was examined

Material and Methods: Hemorrhoid model was induced by croton oil in rats. Then the extract was implemented both externally (50%). A commercial herbal product was preferred as a positive control. After 7 days of the treatment, the animals were anesthetized, intracardiac blood samples were taken and divided into serum to determine tumor necrosis factor alpha (TNF- α) and vascular endothelial growth factor (VEGF). Sacrificed rats were weighed, following, 2 cm long anorectal tissues were cut and weighed to calculate anorectal coefficient (ARC). The tissues were fixed in formaldehyde (10%) for histopathological evaluations [3-5].

Results: According to ARC scores, biochemical and histopathological results, *M. sylvestris* leaf extract showed significant biological activity compared the controls. Value of ARC was found 1,86 \pm 0,09. Serum TNF- α and VEGF values were found 163,22 \pm 11,15 ng/L and 246,37 \pm 12,64 ng/L, respectively.

Conclusion / Discussion: The data demonstrated that the antihemoroidal activity is directly proportional to the effect on capillary permeability. Results will be discussed in detail in the oral presentation. The effectiveness of the traditional usage has been verified by scientific studies.

Keywords: hemorrhoid, *in vivo*, croton oil, mallow, *Malva sylvestris*

References:

- [1] Baytop, T. (1999). Türkiye'de Bitkiler ile Tedavi (Geçmişte ve Bugün)", 2. Baskı. Nobel Kitabevi.
- [2] Gasparetto, J. C., Martins, C. A. F., Hayashi, S. S., Otuky, M. F., & Pontarolo, R. (2012). Ethnobotanical and scientific aspects of *Malva sylvestris* L.: a millennial herbal medicine. *Journal of Pharmacy and Pharmacology*, 64(2), 172-189.
- [3] Azeemuddin, M., Viswanatha, G. L., Rafiq, M., Thippeswamy, A. H., Baig, M. R., Kavya, K. J., Shyam, R. (2014). An improved experimental model of hemorrhoids in rats: evaluation of antihemorrhoidal activity of an herbal formulation. *ISRN pharmacology*, 2014.
- [4] Gurel, E., Ustunova, S., Ergin, B., Tan, N., Caner, M., Tortum, O., Demirci-Tansel, C. (2013). Herbal haemorrhoidal cream for haemorrhoids. *Chinese Journal of Physiology*, 56(5), 253-262.
- [5] Dey, Y. N., Wanjari, M. M., Kumar, D., Lomash, V., Jadhav, A. D. (2016). Curative effect of *Amorphophallus paeoniifolius* tuber on experimental hemorrhoids in rats. *Journal of ethnopharmacology*, 192, 183-191.

ORAL PRESENTATION

**ANTICANCER, ANTIPROLIFERATIVE AND CELLULAR DNA
FRAGMENTATION ACTIVITIES OF *ASTRAGALUS ELONGATUS* SUBSP.
NUCLEIFERUS ON HUMAN CANCER CELLS**

Nazim Sekeroglu^a, Sevgi Gezici^b, Ceren Serin Tanrıöver^b, Fatih Yayla^c

^a Department of Food Engineering, Faculty of Engineering and Architecture, Kilis 7 Aralık University, 79000, Kilis, Turkey, E-mail: nsekeroglu@gmail.com, www.nazimsekeroglu.com

^b Department of Molecular Biology and Genetics, Faculty of Science and Literature; Advanced Technology Application and Research Center, Kilis 7 Aralık University, 79000 Kilis, Turkey

^c Department of Biology, Faculty of Science and Literature, Gaziantep University, Gaziantep, 27310, Turkey

Background and Objective: The genus *Astragalus* are of important medicinal plants that known as immunostimulants, hepatoprotective, antiviral and antidepressive agents. Their wide range of pharmacological effects led us to investigate biological activities of *Astragalus* species growing wild in Turkey. In our previous study, aerial parts and root extracts of *A. elongatus* subsp. *nucleiferus* were analyzed as their potential neuroprotective effects and antioxidant capacities along with polyphenolic contents. Taking our previous results, we aimed to evaluate potential anticancer and antiproliferative activities of ethanol and water extracts from both aerial parts and roots of that plant species in the presented research. **Material and Methods:** On this bases, 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) and trypan blue exclusion assays were performed to investigate cytotoxicity and cell proliferation potentials of the extracts against human lung carcinoma (A549), non-small lung cancer (H1299), glioma (C6) cancer cells, and non-tumorous human umbilical vein endothelial cells (HUVECs). All the conditions and chemicals were same as described in our previous research. The release of the enzyme lactate dehydrogenase (LDH) activity in the cultured cells was also analyzed according to the method of our previous research. Based on their cytotoxicity against A549, H1299 and C6 cancer cells, the concentrations of the extracts were selected for *in vitro* micronucleus (MN) assay. Additionally, apoptotic activity in the cells treated and untreated with the plant extracts were determined in terms of DNA fragmentation using the 'Apoptotic DNA-Ladder Kit'. All the experiments were carried out at least triplicate, and a linear regression analysis was performed to calculate IC₅₀ values for each cell line. **Results and Discussion:** The results showed that the roots of *A. elongatus* subsp. *nucleiferus* are able to induce growth inhibition and apoptosis in a concentration and time dependent manner. As a results of MTT and trypan blue assays, dose-dependent anticancer and antiproliferative effects were observed on human lung and brain carcinoma cells. The water extracts obtained from root was exerted higher anticancer activity, IC₅₀ values ranging from 5.81±0.46 to 18.24±0.12µg/mL (p<0.01) than the extracts obtained from the aerial parts of the plant. Besides increasing extract concentration lead to decrease the growth rate of the cancer cells, apoptosis was observed almost in all the cultured human malignant cells, which rapidly exhibited sign of apoptotic cell death as detected by DNA fragmentation. In contrary anticancer and cytotoxicity results, the ethanol extract of the root was also caused to highest apoptosis level and DNA fragmentation in a dose and time dependently. Regarding LDH activity, which gives information about the percentage of dead and necrotic cells, the results were found a similar activity profile as observed in MTT and trypan blue assays. Briefly, the extracts obtained from aerial parts and roots of the plant were demonstrated the highest LDH activity towards H1299 cells, which was followed by A549 cells. To summarize, anticancer and antiproliferative activity assays were resulted in the superiority of the water extracts obtained from the root of the plant against lung carcinoma cells. **Conclusion:** Overall, the results revealed that the root extracts of the plant could have remarkable anticancer and antiproliferative activities through enhancement of apoptosis and necrosis. The data obtained from our laboratory studies could be assumed as the first report for the literature, and the results should be validated by using further *in vitro* and *in vivo* techniques.

Keywords: Anticancer, apoptosis, *Astragalus* sp., DNA fragmentation, micronucleus, necrosis.

Acknowledgment: This study was financially supported by Scientific Research Project Unit of Kilis 7 Aralık University, Kilis-Turkey (Project code no: 1859LTP1)

References:

- [1] Gezici, S, Sekeroglu, N. 2019. Current perspectives in the application of medicinal plants against cancer: novel therapeutic agents. *Anti-Cancer Agents in Medicinal Chemistry*, 19, 1-11.
- [2] Gezici, S. 2018. Promising anticancer activity of lavender (*Lavandula angustifolia* Mill.) essential oil through induction of both apoptosis and necrosis. *Annals of Phytomedicine*, 7(2), 38-45.
- [3] Gezici S, Sekeroglu N, Kijjoa A. 2017. *In vitro* Anticancer Activity and Antioxidant Properties of Essential Oils from *Populus alba* L. and *Rosmarinus officinalis* L. from South Eastern Anatolia of Turkey. *Indian Journal of Pharmaceutical Education and Research*, 51(3):498-503.

ORAL PRESENTATION

ASSESSMENT OF CHOLINESTERASE AND TYROSINASE ENZYMES
INHIBITORY, PHYTOCHEMICAL COMPOSITION AND ANTIOXIDANT
PROPERTIES OF *MENTHA SPICATA* L.

Sevgi Gezici^{a,*}, Murat Tunçtürk^b, Lutfi Nohutcu^b, Nazim Sekeroglu^c

^a Department of Molecular Biology and Genetics, Faculty of Science and Literature; Advanced Technology Application and Research Center, Kilis 7 Aralık University, 79000 Kilis, Turkey,

E-mail: drsevgigezici@gmail.com, sevgigezici@kilis.edu.tr

^b Department of Field Crops, Faculty of Agricultural Engineering, Yuzuncu Yıl University, 65080 Van, Turkey,

E-mail: murattuncuturk@hotmail.com

^c Department of Food Engineering, Faculty of Engineering and Architecture; Advanced Technology Application and Research Center, Kilis 7 Aralık University, 79000 Kilis, Turkey,

E-mail: nsekeroglu@gmail.com, sekeroglu@kilis.edu.tr

Background and Objective: Because oxidative damage is one of the major factor contributing to neurodegeneration, combination of antioxidant activity and enzyme inhibition potential of the herbal extracts is closely related to combat neurodegenerative diseases and disorders efficiently. Spearmint (*Mentha spicata* L., Lamiaceae), an edible aromatic plant, are of important food sources used as flavoring, tea infusions and spicing in the world. Besides its importance in food, beverage, cosmetic and confectionary industries, there has been huge ethnobotanical information on the curative properties of *M. spicata* L. against common diseases. Even though biological properties of spearmint have been documented by previous scientific reports, one less known biological activity of spearmint is its neuroprotective activity. **Material and Methods:** In this concern, neuroprotective potentials, antioxidant capacities, mineral contents and total polyphenolic compositions of aqueous and methanol extracts of the leaves from *M. spicata* L. were aimed to examine in the current research. Neuroprotective potentials of the extracts were determined against inhibition of AChE, BChE, and tyrosinase enzymes, which are associated with pathogenesis of Alzheimer's disease. *In vitro* assays including DPPH, ABTS, FRAP, and CUPRAC were carried out to reveal antioxidant capacities of the extracts. The total phenolic and total flavonoid contents were analyzed spectrophotometrically. Mineral contents (K, Ca, Mg, Fe, Zn, Cu, Mn, Cd, and Pb) in spearmint leaves were also determined by Atomic Absorption Spectrometry. **Results:** A wide variation was observed in the mineral contents of the plant that Fe was found the major mineral with the values of 1471.41 ± 12.28 mg/kg, whilst Cu and Pb were determined as minor minerals. With regard to total polyphenolic components, the methanol extract possessed higher phenols and flavonoids contents (3.158 ± 0.52 mg/g extract as GAE and 1.31 ± 0.08 mg/g extract as QE, respectively). A significant correlation was found between total polyphenolic compositions and antioxidant capacities of the spearmint leaves. Correlation analysis indicated that the methanol extract, which possessed higher total phenols and flavonoids components than the water extract, was also found the more efficient in terms of antioxidant capacities and neuroprotective potentials. Both of the extracts demonstrated powerful inhibition on BChE, whereas they showed moderate inhibition on AChE and tyrosinase enzymes with the values ranging from $24.06 \pm 0.12\%$ to $57.68 \pm 0.18\%$. **Conclusions:** Consequently, our results indicate that *M. spicata* L., a natural antioxidant and neuroprotective agent, can provide considerable benefits against oxidative stress-related diseases.

Keywords: *Mentha spicata* L.; Alzheimer's Disease; neuroprotection; antioxidant; polyphenolic; mineral content

References:

- [1] Gezici, S., Sekeroglu, N., 2019. Neuroprotective potential and phytochemical composition of acorn fruits. *Industrial Crops and Products* 128, 13-17. <https://doi.org/10.1016/j.indcrop.2018.10.082>.
- [2] Senol, F. S., Sekeroglu, N., Gezici, S., Kilic, E., Orhan, İ. E. 2018. Neuroprotective potential of the fruit (acorn) from *Quercus coccifera* L. *Turkish Journal of Agriculture and Forestry*, 42(2), 82-87. <https://doi.org/10.3906/tar-1711-18>.
- [3] Gundogdu M, Tunçtürk M, Berk S, Sekeroglu, Gezici S. 2018. Antioxidant Capacity and Bioactive Contents of Mulberry Species from Eastern Anatolia Region of Turkey, 52(4), 96-101. <https://doi.org/10.5530/ijper.52.4s.82>.

ORAL PRESENTATION

NEUROPROTECTIVE EFFECTS OF ACTIVE PHYTOCONSTITUENTS AGAINST A β ₁₋₄₂-MEDIATED INJURY IN HIGH GLUCOSE-INDUCED PRIMARY HIPPOCAMPAL NEUROTOXICITY: *IN VITRO* HYPERGLYCEMIC ALZHEIMER'S DISEASE MODEL

Zubeyir Elmazoglu¹, Edgar Rangel-López², Abel Santamaria², Cimen Karasu¹

¹ Department of Medical Pharmacology, Faculty of Medicine, Gazi University, Ankara, Turkey,

² Laboratory of Excitatory Amino Acids, National Institute of Neurology and Neurosurgery Manuel Velasco Suárez, Mexico City, Mexico

Background: Alzheimer's Disease (AD) is the most common of dementia and characterized by progressive loss of memory and other cognitive functions among elderly people [1]. However, the underlying mechanisms of AD is still unclear. Hyperglycemia have an impact on oxidative stress pathways and neuroinflammatory signals in the brain, thereby connecting diabetes to neurodegeneration [2,3]. Four pathological free radical-related sources in the brain have been described which also feed the oxidative damage in diabetes: mitochondrial dysfunction, inflammation, AGEs/ALEs and high cytosolic ionic-calcium levels [4]. Thus, we aimed to investigate the neuromodulatory effects of well known antioxidant phytoconstituents, Oleuropein (OLE) and Rutin (RUT), in hyperglycemic (HG)-AD, *in vitro*.

Methods: AD model was designed by using embryonic primary hippocampal neurons. Cytotoxicity of OLE and RUT in various concentrations (1 nM-25 μ M) were tested by MTT and NRU assay. Long incubation periods (48 h) and high glucose (150 mM) were chosen to mimick chronic hyperglycemia. For the protection experiments, cells were pre-treated with OLE or RUT (10-500 nM), then co-treated with 150 mM glucose (24 h) and followed by incubation with 500 nM oligomeric A β ₁₋₄₂ (24 h) in high glucose conditions. MTT, NRU assays were held to evaluate toxicity and intracellular ROS generation levels were measured by DCFH-DA assay.

Results: Both OLE and RUT have cytotoxic effects at relatively high concentrations (> 10 μ M). While, OLE significantly reversed the high glucose toxicity in a wide range concentration (1-500 nM), RUT exerts only at 50-500 nM. Moreover, intracellular ROS levels were drastically increased in HG-AD model (p<0.01), leading to a significant cell death (p<0.01). RUT treated cells seemed to have higher resistance to ROS-mediated cell death (up to p<0.001), on the other hand OLE treated cells tended to have mild antioxidant effect in that HG-AD *in vitro* model.

Conclusion: Present results suggested that neuroprotection against HG and A β ₁₋₄₂ induced oxidative damage in primary hippocampal neurons may be attributable to redox modulatory effects of OLE or RUT.

Keywords: Hyperglycemia, Alzheimer Disease's, ROS, Oleuropein, Rutin, Neuroprotection

Acknowledgements: This study was financially supported by the Scientific and Technological Research Council of Turkey (Grant No: 31S088)

References:

- [1] Wijesekara N, et al., Impaired peripheral glucose homeostasis and Alzheimer's disease. *Neuropharmacology*. 2018 Jul 1;136(Pt B):172-181.
- [2] Zhao L, et al., Characteristic Metabolic Alterations Identified in Primary Neurons Under High Glucose Exposure. *Front Cell Neurosci*. 2018 Jul 17;12:207.
- [3] Markowicz-Piasecka M, et al., Metformin - a Future Therapy for Neurodegenerative Diseases: Theme: Drug Discovery, Development and Delivery in Alzheimer's Disease Guest Editor: Davide Brambilla. *Pharm Res*. 2017 Dec;34(12):2614-2627.
- [4] Uttara B, et al., Oxidative stress and neurodegenerative diseases: a review of upstream and downstream antioxidant therapeutic options. *Curr Neuropharmacol*. 2009 Mar;7(1):65-74.

ORAL PRESENTATION

SEA BUCKTHORN TREATMENT INFLUENCE ON THE MULTI-EFFECTOR IMMUNE RESPONSE IN SWINE EXPOSED TO POTENTIALLY PATHOGENIC MICROBIOME

Marina Spînu¹, Laura Andreea Rusu¹, Mihaela Niculae¹, Elena Andruța Mureșan², Jovan Bojkovski³, Aurel Vasîu¹, Eموke Pall¹, Diana Ioana Olah¹, Constantin Cerbu¹, Carmen Dana Șandru¹

¹Department of Clinical Sciences, Division of Infectious Diseases, University of Agricultural Sciences and Veterinary Medicine, Faculty of Veterinary Medicine, Cluj-Napoca, Romania

²Food Engineering Department, Faculty of Food Science and Technology, University of Agricultural Sciences and Veterinary Medicine Cluj-Napoca, Romania

³ Faculty of Veterinary Medicine, University of Belgrade, Belgrade, Serbia

Objective / Purpose: Free range systems expose swine to increased infectious pressure and further immune suppression, therefore alternative therapies, such as medicinal plant extracts could alleviate complex environmental stress.

Material and Methods: The research was performed on treated and untreated groups of extensively raised pigs, for identification of the carried microbiome and its sensitivity to antibiotics used in current therapy, subsequent to *in vivo* therapy with sea buckthorn (*Hippophae rhamnoides*) syrup. The antioxidant activity of the sea-buckthorn syrup was determined by assessing free radical scavenging effect over DPPH radical. Skin swabs were processed by classical microbiological methods, identification by RapIDTM and antibiotic sensitivity (Kirby-Bauer method, MAR) was evaluated against gentamicin, streptomycin, oxitetracycline, tylosin, amoxicillin-clavulanic acid, marbofloxacin, tulatromycin, cefotaxime and doxycycline. Immunological tests included *in vitro* blast transformation, serum MIC and MIB concentrations and N/L index.

Results: Strains from *Staphylococcus*, *Shigella*, *Kytococcus*, *Salmonella* and *Citrobacter* genera with MAR index of 0.22 - 0.33 were identified in the skin swabs. Although the antioxidant activity of the syrup was intermediate (54.65%) the N/L index was increased in syrup-treated piglets, similar to blast-transformation indices obtained from sea buckthorn treated piglets towards all extracts used *in vitro* indicating both an increased non-specific reactive capacity and specific cell-mediated immune potential. The sera from treated piglets showed a bacterial growth inhibiting capacity from dilution 1/4 to 1/64.

Conclusion / Discussion: The relationship between antioxidant capacity and antimicrobial protective response for sea-buckthorn syrup needs further investigation to improve the protective capacity against carried potentially pathogenic microbiome in swine.

Keywords: swine, sea-buckthorn, antioxidant, immune response, carried microbiome

References:

[1] Suryakumar, Geetha, Asheesh Gupta (2011). Medicinal and therapeutic potential of Sea buckthorn (*Hippophae rhamnoides* L.), Journal of Ethnopharmacology, 138(2):268-78

ORAL PRESENTATION

**IN VITRO ANTI-HELICOBACTER PYLORI AND ANTIMYCOBACTERIAL
ACTIVITY EVALUATION OF PLANTS FROM TURKEY**

Ayşe Esra Karadağ¹, Ayşegül Çaşkurlu¹, Fatma Tosun¹

¹ *Istanbul Medipol University, School of Pharmacy, Department of Pharmacognosy, 34810- Beykoz, Istanbul, Turkey, aysegulcaskurlu@gmail.com*

Objective: In this study, the extracts of *Ulmus minor* Mill. subsp. *minor* (Ulmaceae), *Lathyrus pratensis* L. (Leguminosae), *Glaucium leiocarpum* Boiss. (Papaveraceae) and *Echium vulgare* L. (Boraginaceae) were investigated for *in vitro* anti-*Helicobacter pylori* and anti-mycobacterial activities.

Material and Methods: The air-dried plant material was powdered and extracted with 70% ethanol by maceration. The extract was filtered and evaporated to dryness under reduced pressure in a rotary evaporator. The concentrated residue dissolved in a water-ethanol (90:10) mixture and then extracted with dichloromethane and ethyl acetate, respectively. Antibacterial activity was investigated by microdilution method against *Helicobacter pylori* ATCC 43504, *Mycobacterium smegmatis* ATCC 25291 and *Mycobacterium avium* clinical isolated¹⁻³.

Results: The ethyl acetate extract of *U. minor* subsp. *minor* showed MIC of 125 µg / mL activity against *H. pylori*. The other extracts showed weak or no inhibitory activity up to 2000 µg/mL concentrations against the tested microorganisms.

Conclusion: The ethyl acetate extract of *U. minor* subsp. *minor* will be further investigated to identify its active compounds that may provide a lead structure to develop a potential antibacterial drug substance for the treatment of *H. pylori* infections.

Keywords: anti-*Helicobacter pylori*, anti-mycobacterial, *Ulmus minor*, *Lathyrus pratensis*, *Glaucium leiocarpum*, *Echium vulgare*.

References:

- [1] Clinical and Laboratory Standards Institute M7-A7. Methods for Dilution Antimicrobial Susceptibility Tests for Bacteria That Grow Aerobically; Approved Standard-Seventh Edition, CLSI document A. Wayne, Pa. USA. 2006, 26(2).
- [2] Whitmire, J. M.; Merrell, D. S. Successful culture techniques for *Helicobacter* species: general culture techniques for *Helicobacter pylori*. *Methods Mol. Biol.* **2012**, 921.
- [3] Chung, G. A.; Aktar, Z.; Jackson, S.; Duncan, K. High-Throughput Screen for Detecting Antimycobacterial Agents. *Antimicrob. Agents Chemother.* **1995**, 39(10), 2235–2238.

ORAL PRESENTATION

NEMATICIDAL ACTIVITY OF ESSENTIAL OILS FROM *LAMIACEAE*

¹Argentieri Maria Pia, ²D'Addabbo Trifone, ¹Avato Pinarosa

¹Department of Pharmacy – Drug Sciences, University of Bari, Bari, Italy;

²Institute of Sustainable Plant Protection, National Council of Research, Bari, Italy

The *Lamiaceae* family contains important aromatic plants used in traditional and modern medicine and in food and pharmaceutical industries. Plants belonging to this family are mostly exploited for the extraction of essential oils (EOs). The genus *Monarda* L. includes about eighteen species of herbaceous, annual or perennial plants endemic of North America. Several biological activities have been reported for plant EOs and their main components, including a biocidal activity against plant soilborne pathogens and pests, which also includes plant-parasitic nematodes [1]. The EOs can play an important role in crop protection and have been proposed as an environmentally alternative to synthetic pesticides.

In this study, the EOs from two Italian species of *Monarda* (*M. didyma* and *M. fistulosa*) were evaluated for their *in vitro* activity on infective stages of plant-parasitic nematodes *Meloidogyne incognita*. EOs from both *Monarda* species were strongly active on *M. incognita* juveniles showing a LC₅₀ value of 1 µL mL⁻¹ after 24 h exposure. The main constituents of EOs, carvacrol, γ-terpinene, o-cymene and thymol were also evaluated for their *in vitro* activity. Among them, carvacrol showed higher nematocidal activity also at low concentrations and at short exposure time [2].

Keywords: Essential oils, carvacrol, *Meloidogyne incognita*, Nematicidal activity.

Reference:

[1] Bakkali F. et al.- *Food Chem Toxicol* (2008), 46: 446-475; [2] Laquale S. et al.- *Journal of Pest Science* (2018), 91(3): 1115-1125

ORAL PRESENTATION

NEW BIOACTIVE GYPSOGENIN COMPOUNDS

N. Gökçe Ulusoy¹, Safiye Emirdağ-Öztürk¹

¹ Ege University, Faculty of Science, Department of Chemistry, Bornova, Izmir, Turkey

Objective: The saponin groups have become quite remarkable on natural products. Many substances isolated from plant that has been reported pharmacological properties. Especially, Gypsogenin aglycone obtained from *Gypsophila arrostii* plant roots which is well known saponin. Gypsogenin aglycone has important biological activities such as antiviral, antitumor, anticarcinogenic, antioxidant and anticancer. Therefore, it is aimed to obtain new bioactive compounds that are functionally enriched in gypsogenin which show bioactive properties.

Material and Method: The starting material gypsogenin compound was obtained from *G. arrostii* plant roots water extract. Then, this gypsogenin aglycone and four different amines compounds were synthesized with ethanol at room temperature.

Conclusions: The study of the structures of the compounds (1-4) synthesized in our study which was determined by IR, UV, ¹H NMR, APT and LCMS analysis.


Conclusion / Discussion: In the last part of the study, biological activities of new compounds will be investigated.

Keywords: Gypsogenin, semi synthesis.

Acknowledgements: This project is supported by EGE UNIVERSITY BAP Project (17FEN064).

References:

- [1] Emirdag-Ozturk S.; Babahan I.; Ozmen A., (2014). Synthesis, characterization and in vitro anti-neoplastic activity of gypsogenin derivatives, *Bioorganic Chemistry*, vol.53, 15-23.
- [2] I. Arslan, A. Celik, M.F. Melzig, (2013). Nebulosides A–B, novel triterpene saponins from under-ground parts of *Gypsophila arrostii* Guss. var. *nebulosa*, *Bioorganic & Medicinal Chemistry*, 21 (5), 1279-1283.
- [3] Wang H., Fang B., Kong L., Li X., Feng Z., Wu Y., Uvdal K., Hu Z., (2018). A novel Schiff base derivative: Synthesis, two-photon absorption properties and application for bioimaging, *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy*, vol. 198, 304-308.



ORAL PRESENTATION

**LEVELS (PRODUCTION) AND BIOLOGICAL ACTIVITY OF SOLAMARGINE
AND SOLASONINE FROM *SOLANUM INCANUM***

Elsadig A. Eltayeb

*Department of Biology, College of Science, Sultan Qaboos University,
P.O. Box 36 PC 123, Muscat, Oman (eatayeb@squ.edu.om) +(968)99350060*

Solanum incanum is a species of nightshade native to the Middle East, the Sub-Sahara region and India. The plant, commonly known as thornapple and has diverse uses traditionally with reported biological activities in diabetes, parasitosis and cancer. These effects possibly arise from the steroidal alkaloid solasodine and its glycoalkaloids mainly solamargine and solasonine.

Optimal extraction of solasodine was performed using 2.5 N methanolic HCl followed by precipitation and extraction using chloroform. A fast and simple extraction method was developed and optimised to get maximum recovery of glycoalkaloids using a mixture of chloroform and methanol by sonication, in one-step. Similarly, a fast and accurate High Performance Thin Layer Chromatography (HPTLC) method for solasodine, solamargine and solasonine content was established and used to identify the developmental stages at which solasodine and its glycoalkaloids contents are maximal in the plant structures of *S. incanum* grown in Sultanate of Oman. The HPTLC method was validated for linearity, precision, recovery, LOD and LOQ.

The solasodine, solamargine and solasonine produced well separated compact bands at R_f values 0.30, 0.26 and 0.14, respectively on silica gel HPTLC plate using chloroform: methanol (9.25: 0.75 v/v) for solasodine and chloroform: methanol: 5% ammonia (7: 3: 0.5 v/v/v) for glycoalkaloids after visualization with anisaldehyde sulphuric acid reagent. The chromatograms were scanned at 530 nm wavelength in absorbance mode and simultaneous method was found linear ($r^2 = 0.99$) in concentration range of 40 – 2000 ng/ spot for solasodine and 50 – 2000 ng/ spot for both glycoalkaloids. The proposed method was also validated as per ICH guidelines and found precise, accurate, specific, robust and sensitive enough to carry out analysis of very minute quantity of solasodine, solamargine and solasonine in complex extracts. The method was applied for analysis solasodine and its glycoalkaloids content in small, young, immature and mature leaves with stem and root parts of *S. incanum* at intervals of 4 to 24 weeks for solasodine and up to 40 weeks for glycoalkaloids. The study revealed that the highest solasodine contents occur in the small leaves 12 – 20 weeks of plant growth and rich concentrations of glycoalkaloids with large variation at different stages of plant development.

In this study, solamargine displayed a superior cytotoxicity in human melanoma cancer cell lines at low concentration of 10 μ M, however it exhibited less cytotoxicity against breast cancer cell lines, bone metastasized and brain metastasized.

Solamargine appears to be selective agent targeting highly proliferating cancer cells; its effect was dependent on the dose of the compound and the type of the cancer cell line. Solamargine induces both necrosis and apoptosis in melanoma cancer cells and breast cancer cells and the mechanism was characterized. Solamargine up-regulated the expression of external death receptor, tumour necrosis factor receptor 1 (TNFR-1).

Keywords: *Solanum* glycoalkaloids, solasonine, solamargine

ORAL PRESENTATION

**KARYOMORPHOLOGY OF SOME MEDICINAL *LEPIDIUM* SPECIES
(BRASSICACEAE) FROM TURKEY**

Emrah Şirin

Selçuk University, Science Faculty, Department of Biology, Konya/Turkey
emrahsirin@selcuk.edu.tr

Objective / Purpose: The genus *Lepidium* L. is one of the largest genera of the Brassicaceae, consisting of 175 species. It is distributed worldwide, primarily in temperate and subtropical regions. *Lepidium* species has been used in folk medicine for curing some ailments such as diabetes, hypertension, asthma, bronchitis and rheumatic pains (Roughani and Miri, 2018). The aim of this study is to determine karyomorphological features of five species belonging to genus *Lepidium*. These species are also used as drugs.

Material and Methods: Many specimens were collected from different localities and populations belonging to genus *Lepidium* during the field studies in the year 2018. Mature seeds were selected and periodically germinated for chromosomal analyses. Chromosome counts were made on somatic metaphases in root tips using the squash technique. After germinating, the root tips were pretreated by 8-hydroxyquinoline and fixed by the Carnoy solution. Before staining, root tips were hydrolyzed with 5-N HCl, and stained with 1% aceto-orcein. At least 10 metaphases were examined per taxa; the best metaphase plate was photographed. Moreover, the coefficient of variation of the chromosome length (CV_{CL}), coefficient of variation of the centromeric index (CV_{CI}) and mean centromeric asymmetry (M_{CA}) were calculated. Idiograms and karyograms of these taxa were made by using the KAMERAM analysis system.

Results: The somatic chromosome number was counted as $2n=16$ for all studied species (*L. campestre* (L.) R.Br., *L. perfoliatum* L., *L. ruderale* L., *L. vesicarium* L. and *L. virginicum* L.). Each of species are diploid. AI, A1, A2, CV_{CL} , CV_{CI} and M_{CA} indexes were calculated for all taxa. The value of CV_{CL} ranges between 11.189–18.226 and the M_{CA} value varies between 12.69–21.53. According to the asymmetry indices it can be concluded that most of the taxa have symmetric karyotype. The most common karyotype formulas $14m + 2sm$ and $12m + 4sm$. Satellites were identified only in karyograms of most of *L. virginicum*.

Conclusion / Discussion: As a result of our study, we conclude that there are significant differences between chromosomal differences between chromosomal characteristics in *Lepidium* species and the reported notes about the intraspecific variation may be useful for taxonomy.

Acknowledgements: This study was financially supported by Selçuk University (BAP Project Number: 18401090).

Keywords: Asymmetry, Brassicaceae, Karyomorphology, *Lepidium*, Turkey

Reference:

Roughani, A., Miri, M.S., 2018. *Lepidium* species as antidiabetic herbal medicines. The First National Congress and International Fair of Medicinal Plants and Strategies of Persian Medicine that affects diabetes. MPSD 2018.

ORAL PRESENTATION

CHEMICAL AND BIOLOGICAL VALORISATION OF *CEDRUS ATLANTICA*

Bouamama H.¹, Nouaa S.¹ and Benharref A.²

¹Laboratory of Bioorganic and Macromolecular Chemistry, FST, UCA, PO BOX: 549, Av. Khattabi, 40000, Marrakech, Morocco

²Laboratory of Biomolecular Chemistry, Natural Products and Reactivity, Faculty of Sciences Semlalia, UCA, BP: 2390, Marrakech, Maroc Email: bouamamahafida@gmail.com



The present study concerns a medicinal and aromatic plant of the Moroccan High Atlas Mountains: *Cedrus atlantica* (Manetti ex Endl.) Carrière 1855. This work contributes mainly to the chemical and biological study of needles and wood of Atlas cedar (Oukaimeden region).

After extractions, chromatographic analysis, antioxidant and antibacterial activities of essential oils were investigated. The essential oils (EO) of needles and Cedar wood were obtained by steam distillation and then analyzed by gas chromatography (GC-FID). Which allowed detecting some major compounds: in the case of EO of fresh needles: 3 major compounds; EO needles after 2 weeks of drying: 2 compounds, for needle EO after 3 weeks of drying: 3 compounds and finally wood HE: 4 major compounds. This analysis allowed highlighting a effect of drying on the abundance of compounds. Open column chromatography and ¹H, ¹³C and DEPT ¹³⁵ NMR analysis were performed and Di-isooctyl phthalate and stearic acid have been identified.

In the other hand, quantitative estimation of total phenols by the method of Folin Ciocalteu revealed that the ethyl acetate extract of *Cedrus* needles contains a phenol content of 0.52 ± 0.049 mg EAG / g MS. The antioxidant activity was evaluated by DPPH free radical scavenging method (2,2- diphenyl-1-picrylhydrazil) and by FRAP iron reduction method. The quantitative antioxidant effect showed that AcOEt extract of *Cedrus* needles has an IC₅₀ value of 91.57 ± 1.05 µg / ml. This value approaches the BHT positive control: IC₅₀ = 77.18 µg / ml.

On the other hand, the essential oils of needles and wood of *Cedrus* have no effect on the reduction of DPPH. According to the FRAP method, those EOs show lower activity with higher IC₅₀% values (81.35 mg / ml and 2.5 ± 0.123 mg / ml, respectively). Antimicrobial activity was determined *in vitro* on two Gram-negative bacteria *Escherichia coli*, *Pseudomonas aeruginosa* and the Gram-positive strain *Staphylococcus aureus* by the disk diffusion method. The species tested were sensitive to the essential oil of *Cedrus* needles at concentrations of 0.5 to 20 mg / ml and for the wood at the concentrations of 0.83 to 83.3 mg / ml.

Keywords: *Cedrus atlantica*; Essential oil; GC / FID; Di-isooctyl phthalate; Stearic acid;

ORAL PRESENTATION

PHYTOCHEMICAL STUDY AND EVALUATION OF INHIBITORY POTENTIAL OF *ANGELICA PANCICII* VANDAS. ESSENTIAL OILS AND HEXANE EXTRACTS AGAINST ENZYMES INVOLVED IN ALZHEIMER DISEASE AND TYPE II DIABETES

Antoaneta Trendafilova¹, Suleyman Yur^{2,3}, Fatih Goger^{2,3}, Miroslav Rangelov¹, Milka Todorova¹, Ina Aneva⁴, Temel Ozek^{2,3}, Gulmira Ozek²

¹ Institute of Organic Chemistry with Centre of Phytochemistry, Bulgarian Academy of Sciences, 1113 Sofia, Bulgaria, trendaf@orgchm.bas.bg

² Department of Pharmacognosy, Faculty of Pharmacy, Anadolu University, 26470 Eskişehir, Turkey

³ Medicinal Plant, Drug and Scientific Research Center (AUBIBAM), Anadolu University, 26470 Eskişehir, Turkey

⁴ Institute of Biodiversity and Ecosystem Research, Bulgarian Academy of Sciences, 1113 Sofia, Bulgaria

Objective: In the present work, the essential oils and hexane extracts obtained from roots, leaf and fruits of Balkan endemic species *Angelica pancicii* Vandas. were investigated for their chemical composition, free radical (DPPH[•] and ABTS^{•+}) scavenging potential and inhibitory effects against acetylcholinesterase and porcine pancreatic α -amylase.

Material and Methods: *A. pancicii* was collected from Rila Mountain, Bulgaria in 2016. Air-dried leaves, fruits and roots were separately used for analysis. Essential oils were obtained with Clevenger apparatus and analyzed by GC-FID/MS. Hexane extracts were obtained by Soxhlet extraction. The amount of the individual coumarins in different plant organs was determined by HPLC-DAD method. Antioxidant effect was tested for their scavenging activity against DPPH and ABTS. Acetylcholinesterase inhibition was evaluated using Ellman's method. Antidiabetic potential was assessed *via* measurement of inhibition of α -amylase enzyme using KI/I₂ method.

Results: CG-FID/MS analyses of essential oils obtained from the different plant parts of *A. pancicii* resulted in the identification of 170 components. It has been found that essential oils differed significantly in their chemical composition. The fruit essential oil was rich in monoterpene hydrocarbons (84.3%), while sesquiterpenoids dominated in the leaf (64.0%) and the root (71.7%) oils. β -Phellandrene (69.1%), germacrene D (9.7%) and elemol (9.8%) were the main components in the fruit, leaf and root oils, respectively. Isoimperatorin, imperatorin, oxypeucedanin, oxypeucedanin hydrate, angeloylpangelin and umbelliprenin were isolated from hexane extract of the fruits and used for HPLC quantification. The main component in the fruits and roots was oxypeucedanin (326.7 \pm 1.3 and 261.9 \pm 1.5 mg/gDE, respectively). These extracts differed significantly in the content of oxypeucedanin hydrate, which was 15 times more in the roots than in the fruits. The leaves were poor in coumarins. The essential oils and extracts of *A. pancicii* demonstrated good anti-acetylcholinesterase activity (between 47.3 and 88.3 %). The leaf and fruit oils were the most active with values similar to that of the standard galantamin (88 %). The essential oils were poor scavengers of DPPH and ABTS radicals when compared with the hexane extracts. The highest TEAC values were obtained for leaf and root extracts (0.5 mM). The leaf extract exhibited also the best DPPH[•] scavenging activity (IC₅₀ 1.2 \pm 0.06 mg/ml). The studied samples demonstrated a weak inhibitory effect (Inh. % up to 31.9 \pm 1.1) towards α -amylase.

Conclusion: This is the first report on inhibitory potential of *A. pancicii* essential oils and hexane extracts against enzymes involved in Alzheimer disease and Type II diabetes. Our data indicated that the fruit and leaf essential oils from *A. pancicii* emerged as the sources of possible cholinesterase inhibitors and deserve future studies.

Acknowledgements: Authors thank to TUBITAK (Project No 116S021) and BAS (Bilateral project) for financial support.

Keywords: *Angelica pancicii* Vandas., essential oil, coumarins, DPPH and ABTS, α -amylase, acetylcholinesterase.

ORAL PRESENTATION

**IN VIVO ANTIDIABETIC ACTIVITY OF RHINACANTHINS ENRICHED EXTRACT
PREPARED BY MICROWAVE ASSISTED GREEN EXTRACTION PROCESS**

Muhammad Ajmal Shah^{a, b*}, Wantana Reanmongkol^{c, d}, Pharkphoom Panichayupakaranant^{b, d}

^aDepartment of Pharmacognosy, Faculty of Pharmaceutical Sciences, Government College University, Faisalabad 38000, Pakistan

^bDepartment of Pharmacognosy and Pharmaceutical Botany, Faculty of Pharmaceutical Sciences, Prince of Songkla University, Hat-Yai, Songkhla 90112, Thailand

^cDepartment of Clinical Pharmacy, Faculty of Pharmaceutical Sciences, Prince of Songkla University, Hat-Yai, Songkhla 90112, Thailand

^dPhytomedicine and Pharmaceutical Biotechnology Excellence Center, Faculty of Pharmaceutical Sciences, Prince of Songkla University, Hat-Yai, Songkhla 90112, Thailand

*E-mail address: ajmalshah@gcuf.edu.pk

Objective: *Rhinacanthus nasutus* is utilized as an herbal tea in China and a Thai traditional medicine for various ailments including diabetes. Rhinacanthins enriched extract (REE) is a semipurified *Rhinacanthus nasutus* leaf extract obtained by microwave assisted green extraction process. The present study was aimed to evaluate the antidiabetic potential of REE in comparison with its marker compound rhinacanthin-C (RC) (Fig 1).

Material and methods: REE was prepared simply by microwave assisted extraction process using ethanol and enriched by passing through ion exchange resin (Amberlite[®]) column. RC content in REE was analyzed by HPLC. In the present study, REE was tested for the antidiabetic potential in comparison with its marker compound RC and positive control glibenclamide employing streptozotocin induced diabetic rat model for 4 weeks. Numerous pathological parameters including fasting blood sugar, insulin, glycated hemoglobin, lipid profile, AST/ALT, creatinine and blood urea nitrogen levels were analyzed.

Results: HPLC analysis showed that REE contains 62 % w/w of RC. REE significantly normalized the blood sugar level, glycated hemoglobin, insulin level, lipid profile, AST/ALT, creatinine and blood urea nitrogen in diabetic rats without affecting the normal control.

Conclusion: The results of the *in vivo* study suggest that after safety assessment, REE could be employed as a standardized natural remedy for diabetes treatment.




Fig 1 Chemical structure of rhinacanthin-C

Keywords: Antidiabetic, *Rhinacanthus nasutus*, rhinacanthins enriched extract

References:

- [1] Shah, M.A., Khalil R., Haq Z.U., Panichyupakaranant, P. (2017). α -Glucosidase inhibitory effect of rhinacanthins-rich extract from *Rhinacanthus nasutus* leaf and synergistic effect in combination with acarbose. *Journal of Functional Foods*, 36(1), 325-331.
- [2] Shah, M.A., Muhammad, H., Mehmood, Y., Khalil, R., Ul-Haq, Z. and Panichayupakaranant, P. (2017). Superoxide scavenging and antiglycation activity of rhinacanthins-rich extract obtained from the leaves of *Rhinacanthus nasutus*. *Pharmacognosy Magazine*, 13(52), 652-658.
- [3] Shah, M.A., Jakkawanpitak, C., Sermwittayawong, D. and Panichayupakaranant, P., 2018. Rhinacanthins-rich extract enhances glucose uptake and inhibits adipogenesis in 3T3-L1 Adipocytes and L6 Myotubes. *Pharmacognosy Magazine*, 13(4), S817-821.

ORAL PRESENTATION

ENZYME INHIBITORY, ANTIOXIDANT ACTIVITIES AND PHYTOCHEMICAL STUDIES ON *Viscum album* SAMPLES COLLECTED FROM DIFFERENT HOST PLANTS

Sultan Pekacar, Hasya Nazlı Ekin, Nilüfer Orhan, Burçin Özüpek, Didem Deliorman Orhan

Department of Pharmacognosy, Faculty of Pharmacy, Gazi University, 06330, Hipodrom, Ankara, Turkey
sultanpkcr94@gmail.com

Objective/Purpose: Metabolic disorders such as diabetes and obesity are increasing day by day. In the treatment of these disorders, plants used in traditional folk medicine are also utilized. The main purpose of our study is to test these plants by *in vivo* and *in vitro* methods and isolation of active natural molecules from active plants. *Viscum album* L. (Loranthaceae) is a semi parasitic plant and traditionally used in the treatment of diabetes mellitus in Anatolia. In this study was planned to compare the *in vitro* antioxidant, antidiabetic (α -glucosidase and α -amylase enzyme inhibition), and antiobesity (pancreatic lipase enzyme inhibition) potentials of *V. album* samples collected from different host plants. The phytochemical contents of the extracts tested were examined by UV spectroscopy and HPLC techniques.

Material and Methods: *V. album* samples were collected from Beynam and Kızılcahamam (Ankara) in 2015 (March-May). α -amylase and α -glucosidase and pancreatic lipase inhibitory effects and antioxidant activities of the aqueous and ethanol extracts prepared from the samples were evaluated. Antioxidant activities of the extracts were determined by DPPH radical scavenging, ferric reducing, metal chelating and phosphomolybdenum assays. Additionally, total phenol and flavonoid contents of the extracts were investigated. Qualitative and quantitative analysis of phenolic acids in active extracts were performed by HPLC method.

Results: Ethanol extracts of *V. album* samples collected from *Salix*, *Abies* and *Pinus* species showed potent α -glucosidase enzyme inhibitor activity, especially at a concentration of 3 mg/ml. Also these extracts displayed strong ABTS scavenging (95.37% \rightarrow 100) and metal chelating activity (99.65-100.48%). Many of the extracts were ineffective on α -amylase and pancreatic lipase enzyme. Total phenol content calculated from gallic acid was 21.52-67.85 mg/g extract and total flavonoid content calculated from quercetin was determined as 10.31-13.11 mg/g extract.

Conclusion/Discussion: The findings obtained in the experiments demonstrated that *V. album* species possess potent antihyperglycaemic and antioxidant activity depending on host plant. Activity-guided isolation studies should be carried on *V. album* samples collected from *Salix*, *Abies* and *Pinus* species.

Keywords: α -glucosidase, α -amylase, pancreatic lipase, HPLC, *Viscum album* L.

Reference:

[1] D. Deliorman Orhan, M. Aslan, N. Şendoğdu, F. Ergun, E. Yeşilada. 2005. Journal of Ethnopharmacology, 98(95-102)

ORAL PRESENTATION

ENCAPSULATION OF *CAPSICUM FRUTESCENS* EXTRACTS WITH β – CYCLODEXTRIN AND 2-HYDROXYPROPYL – β – CYCLODEXTRIN

Enkelejda Goci^{1,2}, Entela Haloci², Jona Berberi¹

¹Pharmacotherapeutic Research Center, Faculty of Medical Sciences, Aldent University, Tirana, Albania, enkelejda.goci@ual.edu.al

²Department of Pharmacy, Faculty of Medicine, University of Medicine, Tirana, Albania

Purpose: *Capsicum frutescens* extracts and capsaicin topical formulation are used in the treatment of such diseases like: rheumatoid arthritis, diabetic neuropathy, osteoarthritis, muscular pain, etc. The aim of this study is the evaluation of recovery percent of *Capsicum frutescens* alcoholic extracts during encapsulation in β – cyclodextrin and 2 – hydroxypropyl – β – cyclodextrin forms.

Materials and Methods: The *Capsicum frutescens* fruits were collected from region of Korca at the southeast of Albania. The alcoholic (ethanol 95°C) extracts were obtained by the methods of maceration process. The extracts contain the capsaicin alkaloid, which is a liposoluble molecule. For improvement of the bioavailability and solubility of this molecule is proposed the encapsulation process in β – cyclodextrin and 2 – hydroxypropyl – β – cyclodextrin. Cyclodextrins are cyclic oligosaccharides with a hydrophilic outer surface and a somewhat lipophilic central cavity. They are able to form water-soluble inclusion complexes with many lipophilic water-insoluble drugs.

Inclusion complexes between the *Capsicum* extracts and β -cyclodextrin and 2 – hydroxypropyl – β – cyclodextrin were prepared by co-precipitation method with ratios: 10:90, 20:80 and 30:70 (w/w) for β – cyclodextrin and ratios: 1:1 and 1:9 for 2 – hydroxypropyl – β – cyclodextrin. For all the encapsulation complexes were evaluate the percent of recovery.

Results: From the study it was concluded that capsaicin can capsule successfully in β -cyclodextrin and 2 – hydroxypropyl – β – cyclodextrin with a best powder recovery 92.4% in the ratio of 30:70 for the complex with β – cyclodextrin and a 90.1% recovery for the complex capsaicin: 2 – hydroxypropyl – β – cyclodextrin at the ratio 1:1.

Keywords: Capsicum extract, β -cyclodextrin, 2 – hydroxypropyl – β – cyclodextrin, recovery.

References:

- [1] Zhu G., Xiao Z., Zhu G., Rujunzhou, Niu Y. (2016). Encapsulation of 1-menthol in hydroxypropyl- β -cyclodextrin and release characteristics of the inclusion complex. Pol. J. Chem. Tech., Vol. 18, No. 3.
- [2] I.L. Domínguez-Cañedo & C.I. Beristain-Guevara. (2011). Microencapsulation of habanero chilli (*capsicum chinense*) oleoresin in β -cyclodextrin and antioxidant activity during storage. European Drying Conference – EuroDrying 2011, Spain.
- [3] Yingying Zhao, Chaonan Sun, Feng Shi, Caleb Kesse Firempong, Jiangnan Yu, Ximing Xu & Weiming Zhang (2016) Preparation, characterization, and pharmacokinetics study of capsaicin via hydroxypropyl-beta-cyclodextrin encapsulation, *Pharmaceutical Biology*, 54:1, 130-138.
- [4] Haloci E., Goci E. et al. Screening the antibacterial activity of satureja Montana essential oil before and after inclusion in β -cyclodextrin.

ORAL PRESENTATION

INVESTIGATION OF THE ANTIMICROBIAL ACTIVITY OF EXTRACTS
OBTAINED FROM TESBI (*STYRAX OFFICINALIS* L.)

Tansu Uskutoğlu¹, Belgin Coşge Şenkal¹, Cüneyt Cesur², Hülya Doğan³

¹ Department of Field Crops, Faculty of Agriculture, Yozgat Bozok University, Yozgat, Turkey,
tansuuskutoglu@bozok.edu.tr

² Department of Energy Systems Engineering, Faculty of Engineering, Karaman Mehmetbey University,
Karaman, Turkey

³ Department of Seed Science, Vocational School of Technical Sciences, Yozgat Bozok University, Yozgat,
Turkey

Objective / Purpose: Tesbi (*Styrax officinalis* L.) is a shrub grows in forestland. The resin of the species *S. officinalis*, named by the past as ‘storax’, was used in traditional medicine in the Mediterranean basin, notably for antiseptic purposes and against respiratory diseases [1]. This resin was also used by Romans, Egyptians, Phenicians and Ionions to treat a variety of ailments and as an incense [2]. To our knowledge, there is no work on leaf extract of this species. The aim of this study was to determine the antibacterial activity of leaf extract.

Material and Methods: 4 g of each tesbi leaf (finely grounded with labrotary blender) was extracted with five different solvents (ethanol / methanol / acetone / ethyl acetate / chloroform) and the extracts were dried at 40 °C by rotary evaporator. The disc diffusion technique (Kirby-Bauer method) is used for the evaluation of anti-microbial activity [3]. 7 Gram (-) bacteria (*Escherichia coli*, *Pseudomonas aeruginosa*, *Salmonella typhimurium*, *Serratia marcescens*, *Proteus vulgaris*, *Enterobacter cloacae* and *Klebsiella pneumoniae*) and 2 Gram (+) bacteria, *Staphylococcus aureus*, *Staphylococcus epidermidis*) were used in the tests.

Results: As a result of the experiment, it was understood that tesbi leaf extracts have moderate effect on *S. aureus*, *P. vulgaris*, *S. epidermidis* and no effect on other bacteria.

Conclusion / Discussion: In this preliminary study tesbi leaf extract shows antibacterial activities, but more investigation is needed, especially leaf extracts should be investigated in HPLC and which component shows antibacterial activity should be determined. Also this results, should be verify with agar dilution and broth dilution methods and minimum inhibitory concentration will be determine.

Keywords: Antibacterial activity, *P. vulgaris*, *S. aureus*, *S. epidermidis*, *Styrax officinalis* L.

References:

- [1] Viano, J., Ruzzier, M., Rabier, J., Tayoub, G., Masotti, V., Girard, G. (2006). Essential oil composition of leaf, flower and stem of *Styrax (Styrax officinalis* L.) from south-eastern France. *Flavour Fragr. J.*, 21: 809–912.
[2] Akgul, Y.Y., Anil, H. (2003). Benzofuran from seeds of *Styrax officinalis*. *Fitoterapia* 74: 743–745.
[3] Prescott, L.M., Harley, J.P., Klein, D.A. (1990). Antimicrobial chemotherapy. In: Microbiology, Wm. C. Brown Publishers, Dubuque, IA, p. 253-255.

ORAL PRESENTATION

ETHNOMEDICINAL STUDY OF *CORNUS MAS L.* FROM THE FOLK AND TRADITIONAL MEDICINE OF REPUBLIC OF MACEDONIA

Biljana Bauer

Department of Pharmacognosy, Faculty of Pharmacy, University of Ss Cyril and Methodius, Majka Tereza No 47, 1000 Skopje Republic of Macedonia

biba@ff.ukim.edu.mk

Objective/Purpose: This study was carried out concerning ethnomedicine on preparations obtained from *Cornus mas* L. from the folk and traditional medicine of Republic of Macedonia, an area so far less frequently studied from the perspective of folk traditional medicinal, local food and handicraft and other uses.

Results: European cornel acts as astringent and antidiarrhoeic. Tea, juice, fresh and dry fruits because of its antimicrobial, anti-inflammatory action are used at diarrhea, colitis and other intestine disorders. Dry fruits are used directly as a medicine against diarrhea. It acts chemostatic and protective on capillary. Yörüks in Republic of Macedonia collected cornels primarily for medicinal herbal infusions for internal cleansing.

Cornels as mountainous fruits were used by Macedonians as a food long time ago. Fruits are used dry as a food. Dry fruits are used for preparing compote or directly in nutrition in winter and spring period. From the fruit jam, marmalade, compote and juice are prepared at home and in the food industry. Dry fruits are kept in rakiya and sugar for obtaining liqueur. Alcohol beverage *rakiya* - *dryankoitsa* was prepared from cornels. Manufacture of hammer in Ohrid or stick in Mariovo (*koledashki*) for Christmas Eve (*kolede*) is made from a cornel branch. Handmade handle for Macedonian traditional instrument *gajdarka*, in Kumanovo is manufactured from cornel because it is smooth and not cracked. Cornel is a symbol of durability, solidity, and with its hardness influences the health.

Conclusion/Discussion: The district, from the ethnobotanical point of view, shows traces of the influences of the neighbouring regions. Some medicinal uses are linked to beliefs or residual forms of magic prescriptions.

Keywords: European cornel, medicinal, food, handicraft, use

Reference:

[1] Nedelcheva, A., Pieroni, A., Dogan, Y. (2017), Folk food and medicinal botanical knowledge among the last remaining Yörüks of the Balkans, *Acta Soc Bot Pol*, 86(2), 3522.

ORAL PRESENTATION

**CYTOTOXIC FLAVONOIDS FROM THE ETHYL ACETATE EXTRACT OF THE
AERIAL ROOTS OF *FICUS BENGHALENSIS* LINN**

Saloni Jain and Tabassum Khan

*Dept. of Pharmaceutical Chemistry & QA, SVKM's Dr. Bhanuben Nanavati College of Pharmacy, Mumbai,
India*

Email address: tabassum.khan@bncp.ac.in

Objective/Purpose: The aerial roots of *Ficus benghalensis* are used as an immunomodulatory medicine by traditional healers in rural areas of Maharashtra, India. Our lab has demonstrated the antioxidant and immunomodulatory activity of the roots thereby validating their traditional use. The antioxidant activity of natural phytochemicals/extracts is related to other bioactivities like antiproliferative activity hence we thought it prudent to evaluate the aerial roots for potential anticancer activity. Our earlier studies reported the presence of flavonoids in the ethyl acetate extract and literature reports indicate potent anticancer properties of flavonoids by inducing apoptosis and G1 cell cycle arrest via different pathways. The objective was to evaluate the ethyl acetate extract for potential anticancer activity, isolate and characterize some anticancer compounds from the ethyl acetate extract of the aerial roots of *Ficus benghalensis*.

Materials and Methods: Preliminary *in vitro* studies on the successive solvent extracts of the aerial roots indicated good antioxidant and anticancer activity for the ethyl acetate extract. Bioactivity guided fractionation of the ethyl acetate extract using preparative TLC resulted in four fractions. The fractions were subjected to LC-UV and LC-MS analysis to identify the flavonoids present. Cytotoxicity studies were conducted by the MTT assay using doxorubicin as the standard drug on A549 cell line (lung cancer cell line), MDA-MB-231 cell line (breast cancer cell line), Hela cell line (cervical cancer cell line).

Results: Seven compounds (kaempferol 3-O-acetyl-glucoside, quercetin 3-arabinoside-7-rhamnoside, quercetin 3-O-(6"-malonyl-glucoside) 7-O-glucoside, naringenin-feruloyl hexose, pelargonidin 3-O-rutinoside, epicatechin gallate, cyanidin 3-O-rutinoside) were identified from this extract and are reported for the first time from this plant. Cytotoxicity studies using the MTT assay indicated a significant anticancer activity on A549 cell line (lung cancer cell line), MDA-MB-231 cell line (breast cancer cell line), Hela cell line (cervical cancer cell line) with IC₅₀ values comparable to doxorubicin standard for the extract and its fractions.

Conclusion/Discussion: The bioactive ethyl acetate rich extract is a potential source of anticancer flavonoids and their glycosides which could serve as templates for anticancer drug design and discovery. Quercetin and related flavonoid analogues identified are reported to induce apoptosis on different cancer cell models by affecting the intrinsic apoptotic proteins via the Erk pathway. The functional groups present in the identified flavonoids such as 2, 3 double bond, 3-OH group and 4-oxo on the C ring are reported to affect the function of mitochondria membrane by decreasing the fluidity via inhibiting the respiratory chain of mitochondria or causing uncoupling, which might partially be attributed to apoptosis.

Keywords: *Ficus benghalensis*, aerial roots, cytotoxicity, A549 cell line, MDA-MB-231 cell line, Hela cell line kaempferol 3-O-acetyl-glucoside, quercetin 3-arabinoside-7-rhamnoside, quercetin 3-O-(6"-malonyl-glucoside)- 7-O-glucoside,

ORAL PRESENTATION

SEASONAL EDIBLE PLANTS OF MAHA RASHTRA(INDIA) - A POTENTIAL
MEDICINAL TREASURY

*Sagarika Damle¹, Archana Thite²

1. Department of Life Sciences, Kishinchand Chellaram College, Mumbai, India

2. Department of Life Sciences, Kishinchand Chellaram College, Mumbai, India

*Author for correspondence: sagarika.damle@kccollege.edu.in

Objective/Purpose: India's rich biodiversity of flora offers a unique opportunity to study different plant species important to human health. Most of these are looked upon as a rich source of bioactive molecules, useful in developing therapeutics. In today's times, the urban population is often neglecting inclusion of nutraceutically important indigenous food sources by opting for exotic ones. Leafy seasonal vegetables, is one such class of plants, with nutritional as well as therapeutic value. These leafy plants, from plant families such as Amaranthaceae, Liliaceae, Caesalpinae etc, are used for treating ailments by tribal population. They are either used as juice or eaten in cooked form by common population. This research project aimed at re-discovering some of these seasonal plants that are grown in rural areas adjoining metro city of Mumbai, India and exploring their nutritional as well as antibiotic/antioxidant properties, that would make them, a potential target for nutraceutics or medicinal formulations.

Material and Methods: Different leafy vegetables such as Math, Shevra, Bharangi, Ghol, Mayalu, Takla, Phodshi available in monsoon season in Mumbai, were procured from local vendors. The dried leaf powders were used as sample for estimating proteins, micro, macronutrients and heavy metals by ICP-AES. For antimicrobial analysis, samples were extracted using three solvents viz, ethanol, acetone and chloroform and were tested against three laboratory strains of microorganisms, *Escherichia coli*, *Staphylococcus aureus*, *Klebsella pneumoniae*, using Agar cup diffusion method. Antioxidant activity was checked for Phodshi with a comparison between edible and non-edible plant part and for Bharangi, the comparison was done for the cooked and uncooked leaves for their scavenging capacity by DPPH assay.

Results: In all the plants under study, Math and Bharangi showed a high % of protein which is uncommon in leaves. In elemental analysis, Shevra was found to be rich in Magnesium, Potassium, Iron and Zinc, where as Takla, Ambadi and Alu were good source of Calcium and Copper. For the study of antibacterial effect, maximum inhibition was observed by the plant extract of Mayalu while least inhibition was found by Takla. Between the three solvents used, inhibitory action was Chloroform > Ethanol > Acetone. Comparing microorganisms, maximum inhibition was seen against *S. aureus* followed by *E. coli* and least against *K. pneumoniae*. The inedible or waste part of Phodshi, was effective in controlling *Corynebacterium* and the scavenging role of its edible portion was found to be as high as 99% in 10 µg/ml of sample. In Bharangi plant, cooked sample showed increased antioxidant activity than the fresh or dried samples.

Conclusion/Discussion: All plants under study showed promising results for nutrient composition, antimicrobial efficacy and antioxidant effects, which can be considered further for identifying bioactive components of therapeutic use. Since these plants are seasonal, there is a need for creating awareness regarding their dietary and health benefits. Most of these indigenous leafy vegetables are less known to the urban population thus growers are faced with low demand. If this downtrend continues, these seasonal plants with a therapeutic potential, would soon disappear from map of available edible vegetables. It is, therefore, imperative to investigate their potential for contributing active biomolecules for medicinal purpose.

Keywords: Seasonal plants, Nutraceutics, ICP-AES, Antimicrobial, Indigenous

References:

- [1] Almeida M.R. (2003) –Flora of Maharashtra, Orient Press India, I & IV. A
- [2] Indian Council of Medical research for National Institute of Nutrition(2009)- Nutrient requirements and recommended dietary allowances for Indians, Mineral requirements and Trace mineral requirements , Chapter 8-11, pp 156-245
- [3] Trivedi P.C. (2006), Medicinal Plants, Traditional Knowledge, I.K. International Pvt.Ltd. India
- [4] Krishna A.R.Sing AJaleel V.A., Sreena R, Karthikeyan S.Gothandam K.M.(2013) –Morphological, Phytochemical and antibacterial properties of wild and indigenous plant (*Amorphophallus commutatus*) , Academic Journals, 7(13),744-748
- [5] Philip Mauleux, Songklanakarinn- The use of stable free radical diphenylpicryl hydrazyl for estimating antioxidant activity, J.Sc.Technol.2004,26(2):211-219.
- [6] Anjali Soni and Sheetal Sosa(2013) – Phytochemical Analysis and Free Radical scavenging Potential of Herbal and Medicinal Plant Extracts , Journal of Pharmacognocny and Phytochemistry, 2(4):22-29.
- [7] Manal I.A.,Musa H.A.,Yousif K.H. and Sabahelkhiar M.K.(2013)- Investigation of Antibacterial activity extraction from tow Medicinal plants available in Sudan, Research Journal of Recent Sciences,2(10), 6-9.
- [8] Bujbalrao R, Dr. Damle S, Dr. Thite A, Krishnamurthi B. (2015)- 'Elemental analysis of some seasonal plants of Maharashtra', published in International Research Journal of Biological Sciences (Online), ISSN: 2278-3202, Vol 4 (4) pp 1-15, April 2015.
- [9] www.wisegeek.com/what-are-the-medicinal-uses-of-Portulaca-oleracea.htm(2014)

ORAL PRESENTATION

**ERBERIS PLANTS WITH ORANGE, RED AND PURPLE FRUITS:
ANTIOXIDANT AND ANTI-HYPERTENSIVE ACTIVITIES**

Eda Şensu^a, Kadriye Nur Kasapoğlu^a, Mine Gültekin-Özgülven^a, Evren Demircan^a, Sevcan Erşan^c, Ayla Arslaner^d, Beraat Özçelik^{a,b}

^a*Istanbul Technical University, Faculty of Chemistry&Metallurgical Engineering, Department of Food Engineering, Istanbul, Turkey*

^b*BIOACTIVE Research. & Innovation Ltd., ITU ARI-3 Teknokent, Maslak, İstanbul, Turkey*

^c*Bayburt University, Faculty of Engineering, Department of Food Engineering, Bayburt, Turkey*

^d*Institute of Food Science and Biotechnology, University of Hohenheim, Garbenstrasse 25, 70599 Stuttgart, Germany*

The genus *Berberis* belonging to the Berberidaceae family is widely grown in Europe, North Africa, Asia, Iran and in the Central Anatolia region of Turkey (1). The parts (fruit, root, leaf, branch) of this plant exhibit various health effects such as anti-hypertensive, anti-diabetic, anti-microbial, anti-cancer, and antioxidant activity (3) since it is a rich source of polyphenolics, as well as carotenoids (2). Therefore, the present study was performed to investigate phenolic and carotenoid content and some biological properties such as antihypertensive and antioxidant activity of *Berberis* plant with orange, red and purple fruits obtained from Bayburt Province in Turkey. Individual phytochemical profile was analyzed by HPLC-DAD/ESI-MS. Antioxidant activity was determined by CUPRAC and DPPH assays, antihypertensive activity was detected using Angiotensin converting enzyme (ACE) inhibitory activity. According to the results, purple *Berberis* fruit had the highest phenolic content (81.42 mgGAE/gDW), anthocyanin content (455.91 mg cyn-3-gly/100gDW) and antioxidant activity (112 mgTE/g and 290 mgTE/gDW, DPPH and CUPRAC assays, respectively). All kinds of *Berberis* leaves had phenolics in the range of 24.81-42.62 mgGAE/gDW while all *Berberis* branches contained between 15.01 and 21.36 mgGAE/gDW. There is no significant difference among all kinds of branches and leaves in terms of antioxidant activities. Chlorogenic acid was the major phenolic compound found in all fruits and leaves. However, epigallocatechingallate was the major phenolic compound in all branches. Delphinidin-3-glycoside, cyanidin-3-glycoside and peonidine-3-glycoside were the main anthocyanins found in purple *Berberis* fruit. Surprisingly, orange, red and purple fruits contained similar amount of carotenoids (~4mg/100gDW). The highest ACE inhibition activity was found in red *Berberis* fruit (73.84%). However, orange and purple fruits did not show ACE inhibition activity. *Berberis* branches and leaves had ACE inhibition activity over 50%. The order of ACE inhibition activity in *Berberis* plants was red fruit>orange branch>purple leaf~red branch>purple branch>orange leaf~red leaf (p<0.05). No correlation was detected between phenolics and ACE inhibition activity while a high correlation was observed between phenolics and antioxidant activity. The purple fruit exhibited the highest antioxidant activity due to anthocyanins and other phenolics while red fruit exhibited the highest anti-hypertensive activity. Therefore, *Berberis* which is unknown in Turkey but it is produced and consumed in Iran is used for dietary supplement and functional food production.

Keywords: Anthocyanins, Angiotensin converting enzyme inhibitor, Antioxidant, *Berberis*, Carotenoids, Phenolic compounds.

References:

- [1] Akbulut, M., Çalışır, S., Marakoğlu, T., & Coklar, H. (2009). Some Physicochemical and nutritional properties of barberry (*Berberis vulgaris* L.) fruits. *Journal of food process engineering*, 32(4), 497- 511.
- [2] Končić, M. Z., Kremer, D., Karlović, K., & Kosalec, I. (2010). Evaluation of antioxidant activities and phenolic content of *Berberis vulgaris* L. and *Berberis croatica* Horvat. *Food and chemical toxicology*, 48(8-9), 2176-2180.
- [3] Imenshahidi, M., & Hosseinzadeh, H. (2016). *Berberis vulgaris* and berberine: an update review. *Phytotherapy research*, 30(11), 1745-1764.

ORAL PRESENTATION

**EVALUATION OF THE RELIABILITY OF PLANTS USAGE IN FOODS: A
REVIEW**

Ezgi Şenol¹, Beraat Özçelik*^{1,2}

^{1*} *Department of Food Engineering, Faculty of Chemical and Metallurgical Engineering, Istanbul Technical University, Istanbul, Turkey, ozcelik@itu.edu.tr; ezgisenol23@hotmail.com*

² *BIOACTIVE Research and Innovation Food Manufacturing and Chemical and Metallurgical Engineering, Istanbul, Turkey*

Functional food is a concept that have been put forward in recent years. A food given an additional function (often one related to health-promotion or disease prevention) by adding new ingredients or more of existing ingredients is known as functional. Functional foods are products that provide significant benefits on the human physiology and metabolic functions beyond the body's need for essential nutrients, thus enabling to protect from diseases and achieve a healthier life. Although the concept of functional foods is new, its application is very old. As the side effects of the drugs have been learned and health problems have increased, the orientations to functional foods have increased and the products are diversified.

Functional foods have been even more than normal food (Ozcelik, 2016). The functional food markets, whose trade volume has exceeded \$ 100 billion around the worldwide, has increased in terms of quantity and diversity in developed countries such as the USA, the European Union and Japan. In Turkey, the functional products market for 2015, it is stated that the ratio reached to 600 million Turkish Liras (Anonym,2014). The usage of plants for food industry to obtain functional food products has become a notable issue in recent years. A rising interest for medicinal and aromatic plants and the active substances obtained from them has led to studies to determine the effects of these plants. The purpose of his review is to determine and evaluate the reliability of plants in terms of food and drugs effects.

Keywords: Functional food, plant, reliability.

ORAL PRESENTATION

IN VITRO CYTOTOXICITY AND IN SILICO MUTAGENICITY ASSESSMENTS OF THE ISOLATED COMPOUNDS FROM *SIDERITIS GERMANICOPOLITANA* BORNM. SUBSP. *GERMANICOPOLITANA*

Büşra Karpuz¹, Fatma Ayaz², Esra Emerce³, Bilge Şener¹, İhsan Çalış⁴, Hayri Duman⁵, M. Iqbal Choudhary⁶, Nurgün Küçükboyacı¹

¹ Department of Pharmacognosy, Faculty of Pharmacy, Gazi University, Ankara, Turkey

² Department of Pharmacognosy, Faculty of Pharmacy, Selçuk University, Konya, Turkey

³ Department of Pharmaceutical Toxicology, Faculty of Pharmacy, Gazi University, Ankara, Turkey

⁴ Department of Pharmacognosy, Faculty of Pharmacy, Near East University, Nicosia, Turkish Republic of Northern Cyprus

⁵ Department of Biology, Faculty of Science, Gazi University, Ankara, Turkey

⁶ International Center for Chemical and Biological Sciences, University of Karachi, Karachi, Pakistan

Objective / Purpose: The genus *Sideritis* L. belongs to the Lamiaceae family and consists of more than 150 species which are mainly distributed in Mediterranean region. In Turkey, *Sideritis* sp. are commonly known as “Dağ çayı, Yayla çayı, Ada çayı” and represented by 46 species with high endemism ratio [1]. *S. germanicopolitana* Bornm. subsp. *germanicopolitana* is a perennial herb which is endemic to Turkey, has been used as herbal tea in Anatolia [2]. The aim of the present study was to investigate the chemical constituents of *S. germanicopolitana* subsp. *germanicopolitana* and evaluate some of the isolated compounds for cytotoxic and mutagenic activities.

Material and Methods: Aerial parts of *S. germanicopolitana* subsp. *germanicopolitana* were collected from Söğüt, Bilecik, Turkey in July 2017, at the flowering period. The dried and powdered plant was extracted with methanol at room temperature. The methanol extract was subsequently fractionated by *n*-hexane, chloroform and *n*-butanol. The *n*-hexane and chloroform fractions were subjected to chromatographic methods for the isolation and purification of the compounds. The structures of the compounds were elucidated by means of spectroscopic analysis. The cytotoxic activity of some of the isolated compounds was evaluated on MCF-7, A549, HeLa and BEAS-2B cell lines by MTT assay. In addition, their *in silico* mutagenicity assessments were performed by combining of knowledge-based and statistical-based approaches.

Results: We isolated a number of diterpenoids, flavonoids and one fatty acid derivative. Among them, we have identified a diterpenoid isomer [major: 7-*epi*-candiciandiol; minor: sideridiol (**1**)] from *n*-hexane fraction, and also a diterpenoid isomer [major: foliol; minor: isofoliol (**2**)] and two flavones [penduletin 4'-methyl ether (**3**) and penduletin (**4**)] from chloroform fraction. Compounds **1-4** showed inhibitory effects on cell proliferation, especially compound **1** was found to have remarkable cytotoxic activity against MCF-7 (IC₅₀=6.46 µg/mL), A549 (IC₅₀=2.26 µg/mL), and HeLa cells (IC₅₀=4.48 µg/mL). All compounds were found to be non-mutagenic in *in silico* analyses. Other isolated compounds are under research.

Conclusion / Discussion: All identified compounds were isolated from *S. germanicopolitana* subsp. *germanicopolitana* for the first time. In conclusion, *S. germanicopolitana* subsp. *germanicopolitana* was displayed to be a potential natural source for developing cytotoxic agents and furthermore, no mutagenic activity was observed in the isolated compounds.

Keywords: *Sideritis*, *S. germanicopolitana* subsp. *germanicopolitana*, diterpenoid, flavonoid, cytotoxicity, mutagenicity.

Acknowledgements: This study was supported by Gazi University Research Foundation [grant number 02/2018-09]

References:

- [1] Güvenç, A., ve Duman, H. (2010). Morphological and anatomical studies of annual taxa of *Sideritis* L. (Lamiaceae), with notes on chorology in Turkey. *Turkish Journal of Botany*, 34, 83-104.
[2] Huber-Morath. (1982). *Sideritis* L. In: Davis PH, editor. Flora of Turkey and the east Aegean Islands. Volume Seven. Edinburgh: University Press; p.178-199.

ORAL PRESENTATION

EVALUATION OF THE *IN VITRO* WOUND HEALING ACTIVITY OF ISOLATED CONSTITUENTS FROM *CHRYSOPHTHALMUM MONTANUM* (DC.) BOISS.

Fatma Ayaz¹, Mert İlhan², Esra Küpeli Akkol², Nezhun Gören³, İhsan Çalış⁴, Muhammad Iqbal Choudhary⁵, Nurgün Küçükboyacı²

¹ Department of Pharmacognosy, Faculty of Pharmacy, Selçuk University, Konya, Turkey,

² Department of Pharmacognosy, Faculty of Pharmacy, Gazi University, Ankara, Turkey,

³ Department of Pharmacognosy, Faculty of Pharmacy, Yeni Yüzyıl University, İstanbul, Turkey,

⁴ Department of Pharmacognosy, Faculty of Pharmacy, Near East University, Nicosia, Turkish Republic of Northern Cyprus,

⁵ H.E.J. Research Institute of Chemistry, International Center for Chemical and Biological Sciences, University of Karachi, Karachi, Pakistan

Objective: *Crysophthalmum montanum* (DC.) Boiss. is commonly distributed in the eastern and southeastern parts of Turkey, as well as the North of Iraq and Syria. In Turkey, the aerial parts of *C. montanum* is traditionally used to treat common cold by sniffing into nose as powder, and applied with cotton on injured body for wound healing by preparing decoction method [1]. In our study, we aimed to investigate the inhibitory effects of the isolated constituents from *C. montanum* on metalloproteinase enzymes using *in vitro* assays to explain the potential wound healing activity mechanism.

Material and Methods: As a part of our ongoing researches on *C. montanum*, we isolated secondary metabolites after the multistep chromatographic purifications on the chloroform and *n*-hexane subextracts of the aerial parts of *C. montanum*. We investigated the potential wound healing activity of the isolates by using inhibitory activity assays on hyaluronidase, collagenase and elastase enzymes, all of which have essential role in the wound healing process.

Results: Our phytochemical studies on *C. montanum* led to the isolation of five sesquiterpene lactones (**1-5**), namely 6 α -acetoxy-4 α -hydroxy-1 β H-guaia-9.11(13)-dien-12.8 α -olide (**1**), 6 α -acetoxy-4 α -hydroxy-9 β .10 β -epoxy-1 β H-guaia-11(13)-en-12.8 α -olide (**2**), 4 α ,6 α -dihydroxy-1 β ,5 α ,7 α H-guaia-9(10),11(13)-dien-12,8 α -olide (**3**), (4 α ,5 α ,8 β ,10 β)-4,10-dihydroxy-1,11(13)-guaidien-12,8-olide (**4**), and (4 α ,5 α ,8 β ,10 α)-4,10-dihydroxy-1,11(13)-guaiadien-12,8-olide (**5**), as well as a steroidal glycoside mixture (**6**) and a flavonoid, chrysosplenol C (**7**) from the chloroform subextract, together with a mixture of triterpenes, taraxasterol acetate and Ψ -taraxasterol acetate (**8**) from the *n*-hexane subextract. Compounds **7** and **8** displayed significant inhibitory effects with the value of 28.53 and 24.41 % for collagenase, 32.28 and 29.31 % for elastase enzymes, respectively. However, none of the compounds showed any inhibitory activities on hyaluronidase enzyme.

Conclusion: Our results provide the phytochemical and biological evidence for the usage of *C. montanum* in Turkish folk medicine for wound healing. It was revealed that compounds **7** and **8** can promote wound healing process by inhibiting the collagenase and elastase enzymes.

Keywords: *Crysophthalmum montanum*, collagenase, elastase, hyaluronidase, wound healing

Reference:

[1] Arasan, S., Kaya, I. (2015). Some important plants belonging to Asteraceae family used in folkloric medicine in Savur (Mardin/Turkey) area and their application areas, *Journal of Food and Nutrition Research*, 3(5), 337-340.

ORAL PRESENTATION

**ANTI-*HELICOBACTER PYLORI* ACTIVITY OF TWO MEDICINAL PLANTS;
FENUGREEK (*TRIGONELLA FOENUM-GRAECUM* L.) AND CUMIN (*CUMINUM
CYMINUM* L.)**

Bouhenni Hasna¹, Doukani Koula¹, Tabak Souhila¹

*Laboratory of Agrobiotechnology and Nutrition in Semi Arid Zones
Faculty of Nature and Life Sciences, University of Ibn Khaldoun, Tiaret (14000), Algeria
E-mail : bouhennihasna@gmail.com*

The bacterial pathogen *Helicobacter pylori* colonizes in over half of the world's population. The infection with this bacterium is an important cause of peptic ulcer disease and eradication of this organism greatly reduces the recurrence rate of ulcers. *H. pylori* also can cause chronic gastritis and gastric cancer inducing inflammatory responses and pathological changes in the gastric microenvironment. The growing problem of antibiotic resistance by the organism demands the search for novel candidates from plant-based sources. In the present study, we evaluated the *in vitro* anti-*H. pylori* activity of two medicinal plants (*Cuminum cyminum* L. and *Trigonella foenum-graecum* L.) on clinical isolates of *H. pylori*. Gastric biopsy samples were obtained from Algerian patients presenting with gastroduodenal complications. *Helicobacter pylori* was isolated from the specimens following standard microbiology procedures, characteristics of colonies were identified by Gram's stain, urease, oxydase and catalase tests. The disc diffusion method was used to determine the susceptibility of *H. pylori* isolates to methanol extracts of cumin and fenugreek at different concentrations. The results showed that methanolic extract of these plants demonstrated significant *in vitro* anti-*H. pylori* activity against the clinical isolates of *H. pylori*.

Keywords: *Helicobacter pylori*, Fenugreek, Cumin, Antibacterial effect

ORAL PRESENTATION

**ETHNOMEDICINAL AND AROMATIC SALT TOLERANT PLANTS USE VALUE
IN KHEWRA SALT MINE, PUNJAB, PAKISTAN**

Muhammad Zafar & Moona Nazish

¹*Department of Plant Sciences, Quaid-i-Azam University, Islamabad, Pakistan*

**Corresponding author e-mail: zafar@qau.edu.pk*

It is the principal quantitative investigation of halophytic vegetation with their customary therapeutic and fragrant uses in Khewra salt mine, Punjab, Pakistan. In this investigation, the reliance of nearby individuals on halophytes gives pragmatic proof about conventional social insurance frameworks. An aggregate of 151 respondents including 13 customary healers were talked with utilizing quick evaluation strategy. The reported information was mapped and examined quantitatively and in GIS too. The mineral portrayal of the dirt was completed utilizing X-beam diffraction (XRD) strategy. Ethno-restorative information was gathered for 38 salt tolerant plant species having a place with 23 families. Results showed the Fabaceae as much of the time referred to plant family (5 species) and *Acacia nilotica* (L.) Delile is the most referred to plant species. Herbs (half) were the prevailing living thing and leaf (37%) was the generally utilized plant section (24 reports). The general significance of the halophytes and homogeneity of the respondents' learning was dictated by utilizing source agreement factor (ICF) and use esteem (UV). The most noteworthy ICF (0.62) was found for gastrointestinal illness class. Relative recurrence of reference and UV ranges from 0.08 to 0.3 and 0.22 to 0.48 separately. The XRD examination demonstrated the copper sulfide nature of soil which is the pointer of halophytes dispersion in the investigation territory. Lion's shares of the halophytes have been seen to have solid pharmacological and sweet-smelling proof. It is the main recorded provide details regarding halophytes utilized for therapeutic purposes in this locale. The discoveries of this study will be valuable for the safeguarding of Ethnomedicinal and sweet-smelling legacy and for future pharmacological, sweet-smelling and phytochemical screening of salt tolerant plants to discover their potential for normal medication improvement.

Keywords: Halophytes, Medicinal, aromatic, GIS, X-ray diffraction, Khewra Salt Mines, Phytochemicals, Pakistan.

Acknowledgements: Authors acknowledge Pakistan Science Foundation (PSF) Pakistan under letter No. PSF/P&D/TG-ND (384)19, by providing travel grant for oral presentation and to attend Symposium (MESMAP-2019).

ORAL PRESENTATION

**CHEMICAL CHARACTERIZATION AND BIOLOGICAL STUDY OF THE
SPECIES *SENECIO CINERARIA***

Zine El Abidine Ababsa^{1,2}, Wahiba Kara ALI³, Kamel Medjrubi², Salah Akkal², Nacira Abidli⁴

¹ *Département des Sciences de la matière, Faculté des sciences exactes et de la nature et vie, Université Larbi BEN M'HIDI, Oum El Bouaghi- Algérie (ababsapharm@gmail.com)*

² *Unité de Valorisation des ressources naturelles, Molécules Bioactives et Analyses Physico-Chimiques et biologiques, Faculté des Sciences Exactes, Département de Chimie, Université Mentouri de Constantine 1- Algérie*

³ *Laboratoire de Biologie et de l'Environnement, Faculté des Sciences de la Nature et Vie, Université Mentouri - Constantine 1- Algérie.*

⁴ *Département de Biologie, école nationale supérieure des enseignants ENS, Kouba, Alger- Algérie*

Senecio cineraria is a perennial shrub of Mediterranean origin, belongs to the family of *Asteraceae* (Dupont et Guignard, 2015)¹, it is used in pharmaceutical preparations and in homeopathy. This study is a scientific contribution to the determination of certain phytochemicals, as well as the study of some in vitro biological activities of the methanolic extract of the plant. The analysis of the extract by colorimetric tests was revealed the presence of flavonoids, alkaloids and tannins. Qualitatively, the TLC analysis of the extract showed the presence of a multitude of flavonoid varieties. The flavonoid assay showed a significant content on the order of 60.16 mg EQ / g E. The study of the antioxidant power by the DPPH method has shown that the concentration which traps 50% of the DPPH ° (IC50) radical is 0.35 mg / ml. EMSC revealed a significant anti-hemolytic effect compared to the positive control. This is proportional to the concentration of the extract used during the test. The antibacterial potential of the extract was confirmed on strains: *Staphylococcus aureus*, *Listeria monocytogenes*, *Klebsiella oxytoca*, with MICs of 10 mg / ml, 20 mg / ml and 2.5 mg / ml respectively. Chronometric coagulation tests (TCK and TQ) have shown that the extract has significant anticoagulant activity.

¹ F. Dupont et J- L. Guignard. Abrégé Botanique. Elsevier /Masson. 16 ème édition. 2015.

ORAL PRESENTATION

PHENOLIC CONTENTS AND ANTI-OXIDANT ACTIVITIES OF *ABIES NUMIDICA* LEAVES EXTRACTS

Djamila Benouchenne, Inès Bellil, Douadi Khelifi

Laboratory of Genetic Biochemistry and Plant Biotechnologies, Faculty of Natural Sciences and Life, University of Mentouri Brothers Constantine 1, 25000 Constantine, Algeria. dbenouchenne1992@gmail.com, dkhelifi@yahoo.fr bines07@yahoo.fr

Objective / Purpose: Recently, polyphenols have attracted a great deal of attention due to their roles in the plant and on human health. There is a strong need to find new effective anti-oxidants from natural sources to prevent human diseases. Algerian fir (*Abies numidica*) is an endemic tree, belongs to *Pinaceae* family, used in traditional medicine as anti-inflammatory, for cataplasm and for the respiratory system. The objective of this study is to extract secondary metabolites from Algerian fir leaves (AFL) and evaluate the anti-oxidant activities of its extracts.

Material and Methods: the air-dried powdered leaves were extracted with 80% methanol and fractionated by dichloromethane (CH₂Cl₂), ethyl acetate, and n-butanol. The total phenolic (TPC) and total flavonoid (TFC) contents of ethyl-acetate and n-butanol fractions were estimated. Furthermore, the antioxidant activities of ethyl-acetate, n-butanol extracts were evaluated by DPPH, CUPRAC assays.

Results: the results showed that ethyl-acetate of AFL is rich in total phenol (490,76±6,65mg gallic acid equivalent/ g extract) and total flavonoids (143,19±5,59 mg quercetin equivalent/g extract) and showed a high anti-oxidant activities in DPPH and CUPRAC assays.

Conclusion / Discussion: it can be concluded that extracts of Algerian fir can be considered as strong anti-oxidant agents, and it is very important to evaluate more assays to determine the biological activities in order to exploit it in human requires.

Keywords: phenolic contents, anti-oxidant activity, leaves extracts, *Abies numidica*.

ORAL PRESENTATION

PHYSICOCHEMICAL AND PHYTOCHEMICAL CHARACTERIZATION AND ANTIBACTERIAL ACTIVITY OF SPINELESS CACTUS (*OPUNTIA FICUS INDICA* F. *INERMIS*) CLADODES IN THE REGION OF TIARET (ALGERIA)

Koula Doukani^{*1}, Brahim Kamel Louacini¹

¹Laboratory of Agro-Biotechnology and Nutrition in Semi Arid Zones, Faculty of Nature and Life Sciences, Ibn Khaldoun University, Po. Box n° 78 Zaaroura –Tiaret (Algeria)

* Corresponding author: kouladoukani@gmail.com

Opuntia ficus indica commonly known as barbary fig has many medicinal properties that have recently been approved. The objective of this study was to promote the therapeutic potential of *Opuntia ficus indica* f. *inermis* cladodes in the region of Tiaret by their physicochemical and phytochemical characterization and antibacterial activity evaluation. Physicochemical analyzes have shown that young and old *Opuntia ficus indica* f. *inermis* L. cladodes are rich in water, minerals (potassium and calcium) and fiber but low in crude protein.

The total phenolic compounds and flavonoids contents were 36.4 mg GAE / g DW and 16.6 mg QE / g DW for the young cladodes respectively and 17.6 GAE / g DW and 15.35 mg QE / g DW for the old cladodes respectively. Phytochemical screening revealed the presence of tannins, steroids and reducing sugars in both types of cladodes.

The results of antibacterial effect of young cladode methanolic extract carried out by the diffusion method of disks against three pathogenic strains: *E. coli* ATCC 25922, *S. aureus* ATCC 25923 and *P. aeruginosa* ATCC 27853 showed the sensitivity of *S.aureus* and *E. coli* to this extract. The cladodes of *Opuntia ficus indica inermis* L. may be a potential source of biological molecules for medicine, nutrition and biotechnology.

Keywords : Cladode, *Opuntia ficus indica* f. *inermis*, physicochemical, phytochemical, antibacterial.

ORAL PRESENTATION

DEVELOPMENT OF MICROEMULSION CONTAINING *PLANTAGO MAJOR* EXTRACTS: FORMULATION AND EVALUATION OF TOPICAL ANTI-INFLAMMATORY ACTIVITIES

Asih Triastuti^{1*}, Oktavia Indrati², Farida Hayati³

¹ Laboratory of Pharmaceutical Biology, Department of Pharmacy UII, 55584 Yogyakarta, Indonesia
asih.triastuti@uii.ac.id

² Laboratory of Pharmaceutical Technology, Department of Pharmacy UII, 55584 Yogyakarta, Indonesia

³ Laboratory of Pharmacology, Department of Pharmacy UII, 55584 Yogyakarta, Indonesia

Objective / Purpose: *Plantago major* L. (Plantaginaceae) is widely used to treat various diseases. It is reported to have anti-microbial, anti-diabetic, anti-spasm, anti-inflammatory activity and wound healing properties. *P. major* contains biologically active compounds such as polysaccharides, flavonoids, iridoid glycosides, terpenoids, and alkaloids¹. Despite several publications supporting health benefits of this plant, not much work has been done to develop an anti-inflammatory formulation containing extracts of *P. major*. Therefore, the aim of this work was to formulate extract of *P. major* into microemulsion to increase economical value of this plant and to support the society with affordable dosage form.

Material and Methods: Microemulsions are clear, optically isotropic and thermodynamically stable systems generally composed of a blend of oil, water and surfactant(s)². Three formulas were developed with variation of surfactants tween 80: lecithin, Formula 1 = 10:1, Formula 2 = 11:1 and Formula 3 = 12:1 respectively, along with cosurfactant (propylene glycol) and extract of *P. major* (0.3%). Microemulsions were stored at 4°C, 25°C, and 40°C for 4 weeks, and were characterized regarding their organoleptic, homogeneity, pH, and viscosity. Particle size was determined on day 0 and week 4 using Horiba®SZ-100 nano particle analyzer. Formula 3 was investigated for its anti-inflammatory activity using croton-oil induced ear edema in mice³. Percentage of inflammatory inhibition was calculated and confirmed by histological section of ears

Results: During 4 week of storage, all formula showed homogeneity and stability and met the particle size requirements for microemulsions. Topical application of microemulsions reduced ear edema and pro-inflammatory cells in the tissue and its activity were similar to hydrocortisone 1% (p > 0.05).

Conclusion / Discussion: Our data indicated that *P. major* extracts can be developed into modern formulation with potent anti-inflammatory activity.

Keywords: *Plantago major*, microemulsion, topical anti-inflammation, croton oil-induced ear edema

References:

- [1] Samuelsen A.B. The traditional uses, chemical constituents and biological activities of *Plantago major* L. A review *Journal of Ethnopharmacology*. 2000;71:1–21.
- [2] Fanun M. Microemulsions as delivery systems. *Current Opinion in Colloid and Interface Science*. 2012;17(5):306-313. doi:10.1016/j.cocis.2012.06.001
- [3] Gábor M. The mouse ear as a model for cutaneous irritation. *Journal of Toxicology - Cutaneous and Ocular Toxicology*. 2002;21(3):191-202. doi:10.1081/CUS-120013039

ORAL PRESENTATION

UNTARGET METABOLOMICS ANALYSES OF THE NEUROPROTECTIVE POTENTIAL OF AYAHUASCA ON HUMAN NEUROBLASTOMA CELLS

Albert Katchborian Neto¹, Wanderleya Toledo dos Santos², Ana Cláudia Chagas de Paula², Michael Murgu³, Marisi Gomes Soares¹, Daniela Aparecida Chagas-Paula¹

¹Chemistry Institute, Federal University of Alfenas, Minas Gerais, Brazil

²Department of Pharmaceutical Sciences, Federal University of Juiz de Fora, Minas Gerais, Brazil

³Waters Corporation, Alameda Tocantins 125, 27th floor, Alphaville, 06455020, São Paulo, Brazil

Objective/Purpose: The available treatments of Parkinson's Disease (PD) target mainly the pathology symptoms and are responsible for severe adverse effects on the host. The aim of this research was to analyse the neuroprotective role of Ayahuasca and its plant matrices: *Banisteriopsis caapi* and *Psychotria viridis* on the viability of human neuroblastoma cells line (SH-SY5Y) by using chemical metabolomic approaches.

Material and Methods: The Ayahuasca beverage and the vegetal material were obtained from a syncretic religion in Brazil 'O Santo Daime'. The *B. caapi* vines and *P. viridis* leaves were harvested in the city of Piedade-SP, Brazil in 2017. The extracts, fractions and isolated alkaloids were characterized by HPLC-UV-DAD. The neuroprotective effects of the samples were screened for 48 h and 72 h assays on the viability of SH-SY5Y cells through the MTT and Calcein-AM. Whereas the cell death was evaluated by propidium iodide staining. The 6-hydroxydopamine (6-OHDA) was used to assess neurodegeneration in PD. The UPLC-HRMS metabolome data was handled using Waters MS software MassLynxTM V4.1 and MarkerLynx XS 3.0.1, which the multivariate statistical analyses were performed. The molecular formula determination was based on the WATERS scientific i-FIT norm. The metabolites were dereplicated using Dictionary of Natural Products (DNP) and online databases.

Results: Ayahuasca as well as the crude extracts of *B. caapi* and *P. viridis*, and the hexanic fractions exhibited neuroprotective effects at the non-toxic concentrations tested. In addition, lesser neuroprotective effects were also obtained for the *B. caapi* hydroethanolic fraction and for the harmine alkaloid. The PCA-X provided an initial look at the dataset, while the cross-correlation statistical analysis of the MS data performed by OPLS-DA regression has revealed key metabolites for the activity. Thus, we identified besides the harmine others 26 potential biomarkers, indicated by S-plot dispersion and Variables Important for Projection (VIP) that have contributed most to the difference in the metabolite profiles of the active samples.

Conclusion / Discussion: The data demonstrated that the evaluated crude extracts have noteworthy neuroprotective effects. Thus, Ayahuasca based plants definitely could be considerate promising for the treatment of PD. The potential of untargeted metabolomic analysis has been underlined as it has generated chemically relevant data for the research. Therefore, the highlighted metabolites and other plant extracts containing them are likely to be in further consideration for the PD.

Keywords: Metabolomics, Ayahuasca, Neuroprotection, Neurodegenerative diseases

References:

[1] YULIANA, N. D. et al. Metabolomics for bioactivity assessment of natural products. *Phytotherapy Research*, v. 25, n. 2, p. 157–169, 2011.

ORAL PRESENTATION

ANTIOXIDANT ACTIVITY OF THE FRUIT EXTRACTS OF *WITHANIA SOMNIFERA* AND *SOLANUM NIGRUM*

Iram Nizam¹, M.Mushfiq²

1 AMU Girls High School, Aligarh Muslim University, Aligarh, U.P., India,

2 Gulistan Apartments, Anoopshahar Road, Aligarh, U.P., India

iramjaipur@rediffmail.com

Objective and Purpose: Antioxidants are known for the prevention and treatment of oxidative stress related disorders, such as diabetes and cardiovascular diseases. Synthetic antioxidants have considerable amount of toxicity, therefore there is an interest in developing herbal drugs. Medicinal herbs possess many phytochemicals which are efficient scavengers of free radicals. The present study was conducted to assess the antioxidant activity of two medicinal herbs, *Withania Somnifera* and *Solanum nigrum* from the family Solanaceae.

Material and Methods: The dried berries (10g each) were extracted in alcohol, separately. Total phenolic content was determined by Folin- Ciocalteu Method and antioxidant activity was measured by DPPH radical scavenging activity.

Results: The total phenolic content and antioxidant activity of *Solanum nigrum* was considerably higher than *Withania somnifera*. However, both herbal extracts showed good amount of total phenolic content and antioxidant activity. The alcoholic extract of *Withania somnifera* shows more than 50% DPPH radical scavenging activity at 200µg/ml while the ethanol extract of *Solanum nigrum* showed above 80% DPPH radical scavenging activity at the same concentration.

Keywords: Antioxidants, total phenolic content, medicinal herbs, *Withania somnifera*, *Solanum nigrum*

References:

- [1] Hsieh, C.C., Fang, H.L., Lina, W.C. (2008). Inhibitory effect of *Solanum nigrum* on thioacetamide-induced liver fibrosis in mice. *Journal of Ethnopharmacology*, 119, 117-121.
- [2] Dhar, R.S., Verma, V., Suri, K.A., Sangwan, R.S., Satti, N.K., Kumar, A., Tuli, R., Qazi, G.N. (2006). Phytochemical and genetic analysis in selected chemotype of *Withania somnifera*. *Phytochemistry*, 67, 2269-2276.

ORAL PRESENTATION

**EVALUATION OF SOME BIOLOGICAL ACTIVITIES OF THE BULBS OF
CORYDALIS SOLIDA (L.) CLAIRV. SSP. *INCISA* LIÉDEN SPECIES GROWING IN
TURKEY**

Mehtap Kılıç¹, Erdal Kaya², Bilge Şener¹

¹Department of Pharmacognosy, Faculty of Pharmacy, Gazi University, 06330 Ankara, Turkey,
klcmehtap89@gmail.com

²Department of Ornamental Plant Breeding and Agronomy, Atatürk Horticultural Central Research
Institute, 77102 Yalova

Objective: The continuing research for the determination of bioactive secondary metabolites from Turkish geophytes as therapeutic agents for demantia is mainly based on the need for drug candidates affected to brain areas. This presentation describes for the first time the *in vitro* anticholinesterase properties and cytotoxic activities of the bulb extracts and alkaloidal fractions of *Corydalis solida* (L.) Clairv. ssp. *incisa* Lidén. Furthermore, the content of the active alkaloid fractions of the bulbs was determined by LC-Q-TOF-MS/MS.

Material and Methods: The bulbs of *Corydalis solida* (L.) Clairv. ssp. *incisa* Lidén were collected from Edirne province. The plant species were also preserved as *ex-situ* in Yalova. The ethanolic and alkaloidal extracts were prepared from the bulbs. The anticholinesterase activities of the extracts and fractions were tested by modification of the Ellman's method. The cytotoxic activities of the extract were determined using the MTT assay against 3T3 and MCF-7 cell lines. The optimization of LC-MS conditions was used in ESI in the positive ion mode.

Results: The alkaloidal extract of the bulbs exhibited a highest activity against AChE and BChE with IC₅₀ values of 0,0167 ± 0,0006 mg/ml and 0,1759 ± 0,0039 mg/ml (galanthamine 0,0068 ± 0,0005 mg/ml and 0,3444 ± 0,0082 mg/ml as positive control), respectively. Among the fractions obtained from the alkaloidal extract, protoberberine-type alkaloids exerted the most promising activity against both cholinesterases, with IC₅₀ values of 0,0224 ± 0,0002 mg/ml and 0,1831 ± 0,0053 mg/ml for AChE and BChE, respectively. The ethanolic extract did not show any cytotoxicity against 3T3 normal fibroblast cell line, however it was found to be active against MCF-7 cell line at 50 µg/ml concentration as 57.38 % inhibition.

Conclusion: The present study showed for the first time the *in vitro* activity of *Corydalis solida* ssp. *incisa* alkaloidal extract and fractions from bulbs as neuroprotective potential and their metabolite profile by LC-Q-TOF-MS/MS. Besides, the anticholinesterase assays on alkaloidal extract and its fractions showed that protoberberine-type alkaloids were the most potent inhibitor of both AChE and BChE.

Keywords: *Corydalis*, isoquinoline alkaloids, anticholinesterases, cytotoxicity

Acknowledgement: This study was funded by the grant of Gazi University Scientific Research Projects Unit (Grant No: 02-2017-13).

References:

[1] Kaya, E.(ed.), (2014), Geophytes of Turkey, Vol. 3, Atatürk Central Horticultural Research Institute, Edition No: 96, Yalova.

ORAL PRESENTATION

**MEDICINAL PLANTS USED IN THE TREATMENT OF SKIN DISORDERS IN
TURKISH VETERINARY MEDICINE**

Ceren Anlas¹

¹ *Department of Pharmacology and Toxicology, Faculty of Veterinary Medicine, Istanbul University-Cerrahpasa, Avcilar 34320, Istanbul, Turkey, cerenis@istanbul.edu.tr*

Over centuries, people have developed their own system of keeping animals healthy and productive using age-old home remedies, surgical and manipulative techniques, husbandry strategies and associated magicoreligious practices. In this context, ethnoveterinary medicines (EVM) are used extensively and quite effectively in many parts of the world for primary health care treatment to make domestic animals productive and healthy. These medicals, considered often safer, cheaper than conventional drugs, can provide useful alternatives to modern animal health care systems.

Medicinal plants are regarded as an integral component of EVM and have been successfully used for treatment of many diseases in traditional veterinary medicine. Turkey has very rich flora with more than 11.000 species of plants, of which around 34.5% are endemic, as a result of its ecological features and geographical location. Especially in recent years, the use of phytotherapeutic remedies has gained popularity in veterinary practice as well as human medicine, in Turkey. In this context, skin diseases and wounds (18.7%) have been reported as the most treated disorders with medicinal plants in Turkish veterinary medicine, following gastrointestinal diseases (26%).

Medicinal plants used in the treatment of skin wounds in animals can regulate the wound healing by affecting key steps for the healing process such as proliferation, migration, collagen deposition or angiogenesis. In addition, many medicinal plants can accelerate the healing process of common skin diseases in animals such as mange, ringworm or dermatitis, due to their multiple pharmacological properties such as antiparasitic, antifungal, antioxidant or antimicrobial activity.

In this review about the medicinal plants used for the treatment of skin disorders in Turkish veterinary medicine, it was aimed to present current knowledges on traditional uses of selected plants from a scientific view point.

Keywords: Medicinal plants, skin disorders, veterinary medicine

ORAL PRESENTATION

LEAVES ANTIMICROBIAL ACTIVITY OF SOME *GLYCYRRHIZA* L. SPECIES

Faruk Karahan¹, Cumhuri Avşar²

¹ Department of Biology, Faculty of Science and Art, Hatay Mustafa Kemal University, Hatay, , Turkey

² Department of Biology, Faculty of Science and Art, Sinop University, Sinop, Turkey

Objective / Purpose: The genus *Glycyrrhiza* includes well-known traditional medicinal plants growing in various parts of the World. Antimicrobial effect studies on leaves are limited although there are many previous studies on antibacterial activity of root extract from licorice. In the study is aimed to determine in-vitro antibacterial effect of leaves extract of *G. glabra* L., *G. flavescens* Boiss. and *G. echinata* L. species.

Material and Methods: Plant samples of these *Glycyrrhiza* species were collected from 17 different localities of the East Mediterranean part of Turkey in 2016. The taxonomic identification was carried out based on related literature. The plant leaves were shade-dried at room temperature and then ground into powder. The leaves metanolic extracts were evaluated for antimicrobial activities against 7 bacterial and 2 yeast strains using discdiffusion and minimum inhibitory concentration methods.

Results: Our results indicated that the leave extracts were more effective against Gram-positive bacteria than against Gram-negative ones. In addition, the extracts had higher antimicrobial effect against *Candida* species than against bacteria. Max. Inhibition zones values were determined as 20 mm in *G. flavescens*, 16 mm in *G. glabra* and 15 mm in *G. echinata*.

Conclusion / Discussion: Results indicated that different populations of same species had got different level in terms of biological activity. This study also supported the traditional use of licorice and as well as suggested that it may also be its beneficial role in the treatment of other infections. The obtained results indicated that different environmental conditions in each habitat might affect the contents of chemical compounds and the biological activity in the natural licorice populations.

Keywords: *Glycyrrhiza*, licorice, antimicrobial activity, Turkey

MESMAP-

ORAL PRESENTATION

QUORUM SENSING INHIBITION ACTIVITY OF VOLATILE OILS APPLIED IN AROMATHERAPY

Ufuk Koca-Çalışkan^{1*}, Elif Burcu Bali², Demet Erdönmez³, Müjgan Özfenerci¹

¹ Department of Pharmacognosy and Phytotherapy Programme, Faculty of Pharmacy, Gazi University, Ankara, Turkey, ukoca@gazi.edu.tr

² Department of Medical Services and Techniques, Programme of Medical Laboratory Techniques, Vocational School of Health Services, Gazi University, Ankara, Turkey

³ Department of Biology, Faculty of Science and Letters, Aksaray University, Aksaray, Turkey

Objective/Purpose: Quorum sensing (QS), the cell-to-cell signalling system in bacteria, is a molecular mechanism of gene regulation, in which bacteria adapt their behavior according to cell density and the surrounding environment. Multidrug-resistance bacteria commonly use this system for developing the resistance; therefore, quorum sensing inhibition (QSI) could be an effective approach for the prevention of bacterial infections, and the discovery of new anti-QS agents. In this study, it is aimed to investigate QSI activity of selected volatile oils used in 'Aromatherapy' on *Chromobacterium violaceum* biosensor system.

Material and Methods: Volatile oils of *Citrus limon*, *Mentha piperita*, *Eucalyptus globulus*, *Citrus sinensis*, (linalol), *Citrus aurantium bergamia*, *Melaleuca alternifolia*, and their blends, were tested for their ability to inhibit QS. The bioreporter strain *Chromobacterium violaceum* CV026 the signaling molecule N-(β -ketocaproyl)-L-homoserine lactone (3-oxo-C6-HSL) was used with violacein pigment inhibition and diffusion assays.

Results: All of the aromatherapeutic volatile oils inhibited the violacein pigment of CV026 (11.32%-90.07%) and exhibited violacein inhibition zone varying from 7.00 to 32.33 mm. The most powerful volatile oils against CV026 were *C. aurantium bergamia*, *M. piperita* and *M. Alternifolia* respectively.

Conclusion/Discussion: This study indicates that the volatile oils may be the potential to be used as natural QS inhibitors for the treatment of bacterial infections in aromatherapy.

Keywords: Aromatherapeutic volatile oils, *Chromobacterium violaceum* CV026, quorum sensing, aromatherapy

References

- [1] July Fong, Chaodong Zhang, Renliang Yang, Zhao Zhi Boo, Soon Keat Tan, Thomas E. Nielsen, Michael Givskov, Xue-Wei Liu, Wu Bin, Haibin Su&Liang Yang (2018). Combination Therapy Strategy of Quorum Quenching Enzyme and Quorum Sensing Inhibitor in Suppressing Multiple Quorum Sensing Pathways of *P. Aeruginosa*. *Scientific Reports*, 8:1155.
- [2] Nicholas A Boire, Stefan Riedel and Nicole M Parrish (2013) Essential Oils and Future Antibiotics: New Weapons against Emerging 'Superbugs' *J Anc Dis Prev Rem*, Vol 1(2): 105.

ORAL PRESENTATION

ANTIOXIDANT, ANTIGLYCATION AND ANTIDIABETIC POTENTIAL OF A NATURAL SWEETENER, *STEVIA REBAUDIANA*

Pamela Jha¹, Renitta Jobby², Sonit Kumari³, Rasika Ghodmare⁴, Nitin Desai⁵, Ahmad Ali⁶

^{1,2,3,4,5}Amity Institute of Biotechnology, Amity University, Mumbai-Pune Expressway, Bhatan, Panvel, Mumbai-410206, pamelajha@gmail.com

⁶Department of Life Sciences, University of Mumbai, Santacruz East, Mumbai-400098

Objective / Purpose: The antidiabetic potential of leaf aqueous *Stevia* extract, a natural, non-caloric sugar substitute, is known. The mechanism is still not clear. This study proposes one of the underlying mechanisms of controlling diabetes by evaluating the extract's antioxidant and antiglycating potential in an in vitro glycation system.

Material and Methods: The plants of *Stevia rebaudiana* Morita I variety, was obtained from organic Innovation, Guwahati, Assam. The aqueous extract, prepared from leaves of stevia were used for antioxidant assays like FRAP and DPPH. The same extracts were used for antidiabetic assays (α -glycosidase inhibitory assays and α -amylase inhibitory assays) and antiglycating assays (measurement of browning and early glycation end products).

Results The IC₅₀ value for aqueous extract of *Stevia* was found to be 300 μ g/ml, by both DPPH inhibition and ABTS radical scavenging assays. The same concentration of *Stevia* showed 20.35 \pm 1.67% inhibition of α -amylase and 8.30 \pm 0.95% of α -glycosidase inhibition. It was observed that there was significant reduction in fructosamine formation and browning of *in vitro* glycation system with *Stevia* extract as compared to the control.

Conclusion/Discussion This study proved that the leaf extract from *Stevia rebaudianana* can be used as the source of antiglycating agents in alleviating the symptoms of diabetes.

Keywords: Antidiabetes, antioxidants, antiglycation, stevia, sweetener.

References:

- 1) Jha P., Momin AR, Kumar D, Ali A, 2018. Reversal of glycoxidative damage of DNA and protein by antioxidants, *Annals of Phytomedicine* 7(1): 1-5.
- 2) Jha P., Ali A., Bio-nanoparticles in the treatment of glycation-induced secondary complications of diabetes, Chapter 12, *Nanomaterials and Plant potential*, Springer (Accepted- May 2018)

ORAL PRESENTATION

**CHANGES IN THE WEED SPECIES IN FIELDS OF MILK THISTLE,
CORIANDER, CHICKPEA AND FORAGE PEA WHICH ARE SOWN ON
DAMAGED BY FROST AREAS OF WINTER OILSEED CANOLA**

Grozi Delchev

Department of Plant Production, Faculty of Agriculture, Trakia University, Stara Zagora, Bulgaria

delchevgd@dir.bg

Objective / Purpose: The aim of this experiment is to investigate the changes in the weed species in fields of 4 crops - milk thistle, coriander, chickpea and forage pea which are sown on damaged by frost crops of winter oilseed canola.

Material and Methods: During 2016 - 2018 was conducted a field experiment. On areas with damaged by frost winter oilseed canola, at the spring were sowed and investigated 4 spring crops: 1 milk thistle cultivar - Silmar (*Silybum marianum* Gaertn.); 1 coriander cultivar - Lozen (*Coriandrum sativum* L.); 1 chickpea cultivar - Kabule (*Cicer arietinum* L.); 1 forage pea cultivar - Mir (*Pisum sativum* L.). The same variants were planted on areas under conventional soil cultivation for each of these crops. Weed control was carried out by different herbicide combinations and herbicide tank mixtures.

Conclusion / Discussion: Coriander and milk thistle are suitable crops for sowing on areas after damaged by frost winter oilseed canola, when using herbicide combination Stomp aqua + Stratos ultra or herbicide tank mixture Zencor + Shadow. After plowing of canola crops, it is more appropriate to sow chickpea in which weed control is carried out by soil treatment with herbicide Merlin flex, followed by foliar treatment with herbicide tank mixture Challenge + Shadow. Herbicide combination Dual gold + Listego should not be used in chickpea. On plowed areas with damaged by frost winter canola hybrids without any problems can be sown forage pea, in which weed control is carried out by herbicide combinations Stomp aqua + Korum or Dual gold + Listego.

Keywords: milk thistle, coriander, chickpea, forage pea, weeds, herbicides.

ORAL PRESENTATION

INVESTIGATION OF VOLATILE COMPONENTS AND VARIOUS *IN VITRO* BIOLOGICAL ACTIVITIES of *Tilia tomentosa* (SILVER LINDEN) MOENCH FLOWERS

Nihal Karakaş¹, Mehmet Evren Okur², Ayşe Esra Karadağ³, Ayşegül Çaşkurlu³,
Şükran Kültür⁴, Betül Demirci⁵

¹ *Istanbul Medipol University, School of Medicine, Beykoz, Istanbul, Turkey*

² *University of Health Sciences, Faculty of Pharmacy, Department of Pharmacology, Uskudar, Istanbul, Turkey*

³ *Istanbul Medipol University, School of Pharmacy, Department of Pharmacognosy, 34810- Beykoz, Istanbul, Turkey, ayseesraguler@gmail.com*

⁴ *Istanbul University, Faculty of Pharmacy, Department of Pharmaceutical Botany, Beyazıt, Istanbul, Turkey*

⁵ *Anadolu University, Faculty of Pharmacy, Department of Pharmacognosy, Eskişehir, Turkey*

Objective: *Tilia tomentosa* Moench. (Malvaceae) flower *n*-hexane extract was investigated for its *in vitro* biological activities and volatile constituents by Gas Chromatography / Mass Spectroscopy (GC / MS).

Material and Methods: The dried plant material was powdered and extracted with *n*-hexane by maceration. The antioxidant capacity was tested *in vitro* DPPH and ABTS radical scavenging methods. Antimicrobial activity was investigated by microdilution method against microorganisms of *Pseudomonas aeruginosa* ATCC 10145, *Escherichia coli* NRLL B-3008, *Staphylococcus aureus* ATCC 6538, *Helicobacter pylori* ATCC 43504 and *Mycobacterium smegmatis* ATCC 25291. The effect on cell viability *in vitro* was determined by measuring the metabolic activity of the cells.

Results: *T. tomentosa* *n*-hexane extract shows that it has moderate levels of antioxidant activity. In addition, the increased activity of the extract on *in vitro* cell viability was determined.

Conclusion: *T. tomentosa* *n*-hexane extract has a positive effect on cell viability, it is rich in camphor, it affects *S. aureus* at a dose of 125 mg / mL and it has a certain level of antioxidant activity. For these reasons, the extract may be used as a cell regenerator.

Keywords: Cell culture, *Tilia tomentosa*, Antimicrobial, Cell regenerator, GC / MS.

ORAL PRESENTATION

ANTIDIABETIC AND ANTIGLYCATING PROPERTIES OF *SPIRULINA PLATENSIS*

Ahmad Ali*, Additiya Paramanya

University Department of Life Sciences, University of Mumbai, Vidyanagari, Santacruz (E), Mumbai, Maharashtra, INDIA

*Corresponding author (Email: - ahmadali@mu.ac.in)

Objectives/Purpose: Diabetes is a major health concern for the world population. Natural resources are being explored for the antidiabetic and antiglycating properties to combat Diabetes. This study was carried out to evaluate both these properties in *Spirulina platensis*, well-known blue - green alga.

Material and Methods: The Antidiabetic property was checked using the extent of inhibition of alpha-amylase enzyme. The extract of *Spirulina platensis* was also incubated with the *in vitro* glycation system consisting of sugar and protein. The amount of glycation products in the presence/absence of *Spirulina* extracts were measured by established methods like NBT assay, carbonyl content, Thioflavin T assays and total AGEs. The DNA damage induced by glycation was also checked by agarose gel electrophoresis.

Results: The results obtained with alpha-amylase inhibition indicate that the *Spirulina* extracts have significant antidiabetic potential. The significant decrease in the amount of glycation in the presence of *Spirulina* extract also suggests the antiglycating potential of phytochemicals present in this organism.

Conclusion/Discussion: These results indicate the significance antidiabetic and antiglycating potential of the extract of *Spirulina platensis*. Further characterization will help in the identification of active constituent from *Spirulina* which can be used for the prevention of Diabetes.

Keywords: Antiglycation, Carbonyl content, DNA damage, Glycation, Protein aggregation, *Spirulina*

References:

- 1) Jha P, Momin AR, Kumar D, Ali A, 2018. Reversal of glycoxidative damage of DNA and protein by antioxidants, Annals of Phytomedicine. 7(1): 1-5.
- 2) Pandey R, Kumar D, Ali, A. 2018. Nigella sativa Seed Extracts Prevent the Glycation of Protein and DNA. CUPMAP. 1:1-7.

ORAL PRESENTATION

PHYTOCHEMICAL PROFILES AND BIOLOGICAL ACTIVITIES OF ENDEMIC MEDICINAL PLANTS FROM KAZAKHSTAN

Aizhamal Baiseitova³, Zhamilya Abilova², Janar Jenis^{1, 2*}

¹ The research center for Medicinal plants, al-Farabi Kazakh National University, Almaty, Kazakhstan

² Shanghai Institute of Materia Medica, Chinese Academy of Science, Shanghai, China

³ Division of Applied Life Science (BK21 plus), IALS, Gyeongsang National University, Jinju, 52828, Republic of Korea

Objective / Purpose: For thousands of years, medicine and natural products have been closely linked through the use of traditional medicines. In Kazakhstan grow over six thousand kinds of plants and these natural resources have been efficiently used in the treatments for a variety of diseases in Kazakh traditional medicine. We focused our attention on study of the bioactive chemical constituents of endemic medicinal plants such as *Ikonnikovia kaufmaniana*, *Artemisia heptapotamica*, *Artemisia albicerata* and *Limonium michelsonii* Lincz.

Material and Methods: The flowering plants of *I. kaufmaniana* and *A. heptapotamica* Poljak were collected in July, 2016, from Almaty region of Kazakhstan. Phytochemical profiles from ethyl acetate extract of *I. kaufmanniana* were quantified by HPLC-ESI MS/MS. The antioxidant activities of all compounds were screened using three different radical sources (DPPH, ORAC, and hydroxyl radicals). As well as, the bioactive constituents from *A. heptapotamica* obtained by using different chromatographic methods including MCI, Sephadex LH-20, silica gel, ODS, preparative HPLC and semi – preparative HPLC. The structures of all isolated compounds mainly achieved by extensive analysis of MS, 1D and 2D NMR spectroscopic data, and ECD spectrum.

Results: Isolation and structural identification processes of *I. kaufmaniana* revealed phenolic compounds having dihydroflavanonol, flavonol, isoflavone and flavanol skeletons and most compounds had significant antioxidant activity against three radical sources, and their efficacies were found to differ by their functionality and skeleton. The phytochemical investigation of *A. heptapotamica* resulted new sesquiterpene lactone dimer, together with 10 known compounds including three dimers, four guaianolides and three *seco*-guaianolides.

Conclusion / Discussion: Secondary metabolites of endemic medicinal plants were investigated for the first time, and related biological activities were evaluated.

Keywords: phytochemical profile, *Artemisia heptapotamica*, *Ikonnikovia kaufmaniana*, biological activity.

References:

- [1] María José Abad Martínez, Luis Miguel Bedoya Del Olmo, Luis Apaza Ticona. (2012). The *Artemisia* L. Genus: A Review of Bioactive Sesquiterpene Lactones. *Studies in Natural Products Chemistry* 37:43-65.
- [2] Akhmetova A, Mukhitdinov N, Ydyrys A. (2014). Anatomical structure of virginal plants of *Ikonnikovia Kaufmanniana* (Regel) lincz. *Mod Phytomorphology*. 6:205–208.

ORAL PRESENTATION

**INVESTIGATION OF THE EFFECT OF SEVEN PLANT SPECIES ON ENZYMES
THAT ASSOCIATED WITH DIABETES AND OBESITY**

Tuğba Günbatan, Hasya Nazlı Ekin, Burçin Özüpek, Sultan Pekacar,
İlhan Gürbüz, Didem Deliorman Orhan

Department of Pharmacognosy, Faculty of Pharmacy, Gazi University, Etiler 06330, Ankara, Turkey

Objective: In this study, measuring the inhibitory effect of seven plant species on enzymes that associated with carbohydrate and lipid metabolism was aimed. Their antioxidant activities and total phenol and flavonoid contents were also evaluated.

Material and Methods: Methanol and water extract were prepared from aerial parts of *Alcea biennis* Witerl, *Anchusa leptophylla* Roem. & Schult., *Alhagi pseudalhagi* Desv., *Berberis crataegina* DC., *Ceratonia siliqua* L., *Scabiosa rotata* M.Bieb. and inflorescence of *Tilia argentea* DC. and their in vitro α -amylase, α -glucosidase and pancreatic lipase inhibitory activities were evaluated. On the other hand their total phenol and flavonoid contents, metal chelating, super oxide dismutase, reducing power and antioxidant activities were also determined [1-11].

Results: Methanol and waters extracts of *A. pseudalhagi* (45.21 and 358.7 $\mu\text{g/mL}$ IC₅₀) and *T. argentea* (81.86 and 411.6 $\mu\text{g/mL}$ IC₅₀) attracted attention with their high α -glucosidase and pancreatic lipase inhibitory activities. Studied extracts were determined not to have effect against α -amylase, except methanol extract of *S. rotata*. Generally, they also showed high antioxidant activity in CUPRAC, ABTS, SOD, Metal Chelating and DPPH methods. The highest total antioxidant activity was observed in *B. crataegina* methanol extract.

Conclusion: Extracts from *A. pseudalhagi* and *T. argentea* came into prominence with their high α -glucosidase and pancreatic lipase inhibitory activities. More detailed investigations on these extracts should be done.

Keywords: α -glucosidase, α -amylase, pancreatic lipase.

References:

- [1] Ali, H., Houghton, P., and Soumyanath, A. (2006). α -Amylase inhibitory activity of some Malaysian plants used to treat diabetes; with particular reference to *Phyllanthus amarus*. *Journal of Ethnopharmacology*, 107(3), 449-455.
- [2] Lam, S. H., Chen, J. M., Kang, C. J., Chen, C. H., and Lee, S. S. (2008). α -Glucosidase inhibitors from the seeds of *Syagrus romanzoffiana*. *Phytochemistry*, 69(5), 1173- 1178.
- [3] Kim, Y. S., Lee, Y. M., Kim, H., Kim, J., Jang, D. S., Kim, J. H., and Kim, J. S. (2010). Anti-obesity effect of *Morus bombycis* root extract: anti-lipase activity and lipolytic effect. *Journal of Ethnopharmacology*, 130(3), 621-624.
- [4] Matthaas, B. (2002). Antioxidant activity of extracts obtained from residues of different oilseeds. *Journal of Agricultural and Food Chemistry*, 50, 3444-3452.
- [5] Apak, R., Güçlü, K., Özyürek, M. ve Karademir, S. E. (2004). A novel total antioxidant capacity index for dietary polyphenols, vitamin C and E, using their cupric ion reducing capability in the presence of neocuproine: CUPRAC method. *Journal of Agricultural and Food Chemistry*, 52, 7970-7981.
- [6] Gülçin, İ. (2007). Comparison of in vitro antioxidant and antiradical activities of L-tyrosine and L-Dopa. *Amino Acids*, 32(3), 431-438.
- [7] Dinis, T. C., Maderia, V. M., & Almeida, L. M. (1994). Action of phenolic derivatives (acetaminophen, salicylate, and 5-aminosalicylate) as inhibitors of membrane lipid peroxidation and as peroxyl radical scavengers. *Archives of Biochemistry and Biophysics*, 315(1), 161-169.
- [8] Kosalec, I., Bakmaz, M., Pepeljnjak, S., & Vladimir-Knezevic, S. (2004). Quantitative analysis of the flavonoids in raw propolis from northern Croatia. *Acta Pharmaceutica (Zagreb, Croatia)*, 54(1), 65-72.
- [9] Orhan, N., Deliorman Orhan, D., Gokbulut, A., Aslan, M., & Ergun, F. (2017). Comparative analysis of chemical profile, antioxidant, in-vitro and in-vivo antidiabetic activities of *Juniperus foetidissima* Willd. and *Juniperus sabina* L. *Iranian Journal of Pharmaceutical Research*, 16(Suppl), 64-74.
- [10] Oyaizu, M. (1986). Studies on products of browning reaction. *The Japanese Journal of Nutrition and Dietetics*, 44(6), 307-315.
- [11] Zongo, C., Savadogo, A., Ouattara, L., Bassole, I. H. N., Ouattara, C. A. T., Ouattara, A. S., . . . Traore, A. S. (2010). Polyphenols content, antioxidant and antimicrobial activities of *Ampelocissus grantii*(Baker) Planch.(Vitaceae): A medicinal plant from Burkina Faso. *International Journal of Pharmacology*, 6(6), 880-887.

ORAL PRESENTATION

**THE EFFECTS OF NITROGEN AND FARM MANURE APPLICATIONS ON
ESSENTIAL OIL CONTENT OF DILL (*Anethum graveolens* L.)**

Ferit Özen¹, Gülsüm Yaldiz², Mahmut Çamlica²

¹*Medicinal and Aromatic Plants Department, Mudurnu Süreyya Astarçı Vocational School, Mudurnu Bolu, Turkey*

²*Department of Field Crops, Faculty of Agriculture and Natural Sciences, Bolu Abant İzzet Baysal University, 14280 Bolu, Turkey. E-mail: g_yaldiz@hotmail.com*

Objective/Purpose: *Anethum graveolens* L. or dill, belonging to Apiaceae (Umbelliferae) family, is an annual aromatic herb known for culinary and medicinal use since ancient times. It has been used in ayurvedic medicines since ancient times and it is a popular herb widely used as a spice and also yields essential oil. The present study aimed to determine the effects of different doses of chemical (ammonium nitrate) and organic fertilizer (farm manure) applications on the essential oil content of dill (*Anethum graveolens* L.).

Material and Methods: In this study, different doses of farm manure (750, 1000, 1250 and 1500 kg/da) and ammonium nitrate (3, 6, 9, 12 kg/da) with a control (no manure) were applied by sowing. The experiments were arranged in the Completely Randomized Blocks Design with three replicates. Each experimental plot consisted of five rows, with a distance of 0.3 m between rows and 0.2 m between plants, and a plot size of 3.3 m². The essential oil percentage was subjected to hydro-distillation for 3 hours using a modified Clevenger apparatus.

Results: The hydro-distilled essential oil content ranged from 0.28% to 6.16%. The highest essential oil content were obtained in the treatments with farm manure doses of 750 kg/da and control application, which were significantly different compared to the other treatments. Among all of the essential oils included in the study, dill seeds showed the highest essential oil yield, followed by dill leaves and dill herbs. As suggested by the present study, a suitable ratio of farm manure in the soil can increase the essential oil content.

Conclusion: This organic manure could thus be considered as suitable substitutes for chemical fertilizers when growing medicinal plants that are gaining both an increased importance and demand.

Keywords: Dill, essential oil, farm manure.

Reference:

[1] Kazemi, M., & Abdossi, V. (2015). Chemical composition of the essential oils of *Anethum Graveolens* L. Bangladesh Journal of Botany, 44(1), 159-161

ORAL PRESENTATION

ELEMENTAL ANALYSIS OF LEAFY VEGETABLES AS NUTRITIONAL & MEDICINAL BENEFACTORS

Archana Thite* & Sagarika Damle

Department of Life Sciences, Kishinchand Chellaram College, Mumbai, India

**Email - archana.thite@kccollege.edu.in*

Objective/Purpose: A geographical biodiversity so rich that it's considered a blessing & a source of natural wealth. This has been the outlook of Indians for generations, who have created traditional cuisines which are abundant in nutritional & medicinal benefits. Leafy vegetables form a part of meals which contain a variety of minerals. Near Mumbai are some regions of the Western ghats, a hot spot of rich biodiversity, where plants from families such as Amaranthaceae, Liliaceae, Fabaceae are often consumed as leafy vegetables. Yet there are some leafy vegetables which are lesser known to urban people. They are indigenous, occur in natural wild habitats and are available only during monsoon. Their benefits are known to the local tribes who consume them to ward off monsoon ailments. Ca^{+2} , Na^+ , K^+ , Cu^{+2} , Zn^{+2} , Fe^{+2} etc. are elements which are important for normal metabolism in human being. Most of the elements are quantified along with their effects. Our aim was to analyse some common and uncommon leafy vegetables for the mineral content. On understanding their mineral content, the dietary benefits of uncommon leafy vegetables can be popularised among urban population.

Materials and Methods: Different leafy vegetables; all Season/common plants: *Portulaca oleracea* (Ghol), *Basella rubra* (Mayalu), *Vigna unguiculate*(Chawli), *Colocassia esculenta*(Alu) and *Cassia tora* (Takla) and seasonal/uncommon plants (Monsoon): *Amorphophallus commutatus* (Shevra), *Amaranthus viridis* (Pandhara math), *Clerodendron indicum* (Bharangi), *Ipomoea aquatica* (Nalichi bhaji), *Smithia conferta*(Kauli), *Hibiscus cannabinus*(Ambadi) were procured from Mumbai local and rural vendors. The leaves were dried at a constant weight and powdered. The sample powders were acid digested and then subjected to ICP-AES analysis.

Results: The macro-elements K^+ , Ca^{+2} , Na^+ were present in adequate quantities in both the plant groups. Minor-elements were checked and Cu^{+2} was also present in adequate amounts but the uncommon plants had more content of Copper as compared to common ones. *Amorphophallus commutatus* (Shevra) had Zn^{+2} present in good quantity, where as all other plants showed its absence.

Discussion: The macro-elements are all essential for the normal metabolism of human body. Potassium maintains Na^+/K^+ - ATPase pump, muscular activities and electrolyte balance in the body. Its deficiency results in formation of kidney stones and cardiac diseases whereas in excess it causes accumulation of calcium in bones. Calcium acts as neurotransmitter, maintains muscular functioning, helps in proper bone formation, promotes absorption of Ca^{+2} from intestines by calcitonin, also helps Na^+/K^+ pump. Na^+ helps to maintain homeostasis. The minor-elements are necessary for enzyme activation.

Key words: Minerals, leafy vegetables, ICP-AES.

References:

- [1] Almeida M.R. (2003) –Flora of Maharashtra, Orient Press India, I & IV. A
- [2] Indian Council of Medical research for National Institute of Nutrition(2009)- Nutrient requirements and recommended dietary allowances for Indians, Mineral requirements and Trace mineral requirements , Chapter 8-11,pp 156-245
- [3] Trivedi P.C. (2006), Medicinal Plants, Traditional Knowledge, I.K.International Pvt.Ltd. India
- [4] Carmen Cristina Elekes et al., The appreciation of mineral element accumulation level in some herbaceous plants species by ICP–AES method, Environmental Science and Pollution Research, 17(6), 1230-1236 (2010)
- [5] Arun Kumar et al., Analysis of nutritional elements in Indian medicinal herbs used to cure general weakness, Natural Science 4(4), 211-215 (2012) doi:10.4236/ns.2012.44032 8. Pavia D.L., Lampman
- [6] Anjali Soni and Sheetal Sosa(2013) – Phytochemical Analysis and Free Radical scavenging Potential of Herbal and Medicinal Plant Extracts , Journal of Pharmacognocny and Phytochemistry, 2(4):22-29.
- [7] G.M., Kriz G.S. and Engel R.G., Introduction to Organic Laboratory Techniques: A Microscale Approach 3rd Edition Saunders College Publishing: New York, NY, (1999) 9. Kos V. et al., Journal of Analytical Chemistry, 354(5-6), 648-652 (1996)
- [8] Bujbalrao R, Dr. Damle S, Dr. Thite A, Krishnamurthi B. (2015)- 'Elemental analysis of some seasonal plants of Maharashtra', published in International Research Journal of Biological Sciences (Online), ISSN: 2278-3202, Vol 4 (4) pp 1-15, April 2015.
- [9] www.wisegeek.com/what-are-the-medicinal-uses-of-Portulaca-oleracea.htm(2014)
- [10] http://www.ntgb.org/plants/plant_details.php(2015)

ORAL PRESENTATION

CHEMICAL ANALYSIS OF WILD PLANT *INULA HELENIUM* SUBSP. *PSEUDOHELENIUM* GRIERSON GROWING IN VAN LAKE AROUND.

Rüveyde Tunçtürk^a, Lütfi Nohutçu^a, Murat Tunçtürk^a, Tamer Eryiğit^d

^a*Yuzuncu Yil University, Agricultural Faculty, Field Crops Department, Van-Turkey*

^b*Tamer Eryigit, Y.Y.U. Gevas Vocational School, Department of Plant and Animal Production, Van-Turkey*

The genus *Inula* belongs to the Compositae family and comprises 27 species growing wild in Turkey. *Inula helenium* L. subsp. is of the commonly consumed wild edible plants as food and medicinal purposes inhabitants in Eastern Anatolia. It has been used in traditional Anatolia medicine for abdominal distension and pain, acute enteritis and bacillary dysentery. Wild plants gathered from nature are cheaper food and important for human health. Thus, in the present study nutritional value and mineral composition of used parts of selected *Inula helenium subsp. pseudohelenium* Grierson was investigated.

In laboratory analysis, dry matter, total ash, % N, crude protein, crude fiber and pH were examined as nutritional value. Useful minerals (Ca, Cu, Fe, K, Mg, Mn, Na, P, S and Zn) and heavy metals (Cd, Co, Cr and Pb) that hazardous elements for livings were also determined. Results in this research showed that *Inula helenium subsp. pseudohelenium* Grierson contains high value of macro elements such as sodium (1.07 mg/g), potassium (19.16 mg/g), magnesium (2.12 mg/g) and calcium (14.17 mg/g). Also, it has maximum micro-elements such as iron (246.24 mg/kg), manganese (52.98 mg/kg), copper (40.27 mg/kg) and zinc (26.84 mg/g). Therefore, we can conclude that It has high nutritional values from the viewpoint of mineral elements such as potassium, calcium, iron, manganese and zinc.

Keywords: Mineral content, Wild plant, *Inula helenium* L, East Anatolia

ORAL PRESENTATION

NEUROPROTECTIVE AND COGNITIVE ENHANCEMENT POTENTIALS OF XANTHONE-ENRICHED FRACTION OF *GARCINIA MANGOSTANA* AND α -MANGOSTIN

Vikneswaran Murugaiyah *¹, Ning Tiang, Zurina Hassan

¹Discipline of Pharmacology, School of Pharmaceutical Sciences, ²Centre for Drug Research, Universiti Sains Malaysia, 11800 Penang, Malaysia

Objective: Mangosteen (*Garcinia mangostana*), also known as “The Queen of Fruit” has been used traditionally in Southeast Asia to treat against inflammation, skin infection, wound and diarrhea. The present study aimed to investigate the neuroprotective and cognitive enhancement potentials of xanthone enriched fraction (XEF) from *G. mangostana* and its major constituent, α -mangostin.

Material and Methods: The neuroprotective effects of XEF and α -mangostin were studied in various stressors-induced neurotoxicity models in neuroblastoma cell lines and rat primary cells. Their effect on cognition was investigated in chronic cerebral hypoperfusion (CCH) rats, prepared by permanent bilateral common carotid arteries occlusion (PBCCAO). Two weeks after surgery, CCH rats were orally administered (single and 14 days repeated dose) with XEF and α -mangostin prior to locomotor activity and Morris water maze, long term potentiation (LTP) evaluation.

Results: The stressors-induced neurotoxicity caused reduction in cell viability of 25 to 45%. At lower concentration range of 0.25-1 μ g/mL, XEF and α -mangostin showed significant and concentration dependent neuroprotection in all test models. Among them, α -mangostin showed the most promising neuroprotective effect, especially in the glutamate-induced neurotoxicity. The *in vivo* studies showed no effect on the rat's locomotor activity. However, α -mangostin (50mg/kg) and XEFGM (100mg/kg) significantly reversed the cognitive impairment induced by PBCCAO in the spatial learning test. In addition, α -mangostin also showed significant improvement in the reference memory. LTP results revealed that α -mangostin improved the basal synaptic transmission but has no improvement on the inhibition of LTP observed in CCH rats. Likewise, no changes were observed in the protein expressions of BDNF and CAMKII α in hippocampus of the treated rats.

Conclusion: The present study suggests that XEF and α -mangostin are potential protective agents against oxidative stress and excitotoxicity-induced neurodegeneration and ameliorated learning and memory deficits in CCH rats, worthy of further investigation.

Keywords: *Garcinia mangostana*; α -mangostin; Xanthone enriched fraction; Neuroprotective; Cognitive enhancement; Chronic cerebral hypoperfusion.

ORAL PRESENTATION

MEDICINAL PLANTS AS IMMUNOSTIMULANTS IN FISH FARMING

Ataman Bilge Sari

İstanbul University – Cerrahpaşa, Faculty of Veterinary Medicine Department of Veterinary Pharmacology and Toxicology, Istanbul, Turkey

For a long time, disinfectants and antibiotics have been used effectively for the treatment of fish diseases. It has been revealed that overuse of chemicals carry potential risks such as environmental problems, residual accumulation in tissue, resistance to antibiotics, and reduction of the efficiency. These problems have led some researchers and fish farmers to consider alternative natural sources from plants that would minimize environmental harm while allowing increased aquaculture production. Main problems of aquaculture production are high stock density and consequently the decrease in water quality, the increase of stress and infection risk. After the occurrence of an infection, the control and treatment are more challenging, and it leads to mass deaths and economic losses. Because of the difficulties in treatment, prophylactic methods like vaccination or immunostimulants have gained importance in the protection of fish health.

It has been shown that the use of immunostimulants can be an effective tool for increasing the resistance of fish against disease. Immunostimulant medicinal plants have been reported to stimulate non-specific immune mechanisms also have additional effects like stimulation appetite or weight gain and also antibacterial and antiparasitic properties due to alkaloids, flavonoids, terpenoids and saponins. The advantage of stimulating nonspecific immune response is that it acts against all pathogens since it is not selective for against only specific agents and it can be used in different kinds of aquaculture systems conveniently.

The aim of this presentation is to give information about the medicinal plants, plant extracts, application methods and the mode of actions on the nonspecific immune system of fish.

Keywords: Immunostimulant, Medicinal Plants, Aquaculture.

ORAL PRESENTATION

**STUDIES ON MEDICINAL AND AROMATIC PLANTS UNDER THE FARMERS
CONDITIONS IN THE PROVINCE OF KAYSERİ**

Ahmet Yenikalayci^{1*}, Kemal Gül², Mahmut Güneş³

⁽¹⁾: *Muş Alparslan University, Faculty of Applied Sciences, Herbal Production and Technologies
Department*

⁽²⁾ *Directorate of Provincial Agriculture and Forestry, Herbal Production and Plant Health Branch
Directorate/KAYSERİ*

⁽³⁾ *County Agriculture and Forestry Directorate /KAYSERİ*

*: *Responsible writer e-mail: a.yenikalayci@alparslan.edu.tr*

This study comprises the results of demonstrative researches on some Medicinal and Aromatic Plants carried out by Kayseri Directorate of Provincial Agriculture and Forestry under the farmers' situations in 2015-2017 within the scope of the Project of Improving the Medicinal-Aromatic and Dye Plants Cultivations sustained by BÜGEM (General Directorate of Improving Plant Production).

In this Project, 5 different plants consisting of Lavandin, Lemon Balm, Orchis, Black Seed and Oregano were studied in 11 counties for 3 years. Favorable results were obtained from Lavandin, Lemon Balm, Black Seed and Oregano plants in terms of cultivation under the Kayseri ecological situations. For Lavandin (*Lavandula x intermedia*) species, dry flower yield was found to be 30-127 kg/da, essential oil rate in dry flowers was found to be % 2.72-9.00, for Lemon Balm (*Melissa officinalis* L.) species, dry herb yield was found to be 525-610 kg/da, essential oil rate in dry leaf was found to be % 0.07-0.24, and for Oregano (*origanum onites* L.) species, essential oil rate was found to be % 4.64.

In the cultivation of Black Seed, although there is no any trouble in terms of the cultivation, weed problem seemed as an important problem in front of us. Since there aren't any herbicides authorised by the Ministry of Agriculture and Forestry, sufficient yield wasn't obtained. Although applying different planting times and mulching methods in orchis plants, favorable results couldn't be obtained in terms of cultivation because of the difficult winter conditions and frost damage.

Keywords: Farmer Conditions, Medicinal Plants, Farming.

ORAL PRESENTATION

**SOLVENT DEPENDING IMMUNOLOGICAL ACTIVITY OF NETTLE PLANT
EXTRACTS IN SHEEP WITH CONTAGIOUS ECTHYMA**

Aurel VasIU¹, Mihaela Niculae¹, Elena Andruța Mureșan³, Brudașcă Gheorghe Florinel¹, Eموke Pall¹,
Constantin Cerbu¹, Marina Spînu¹, Silvana Popescu², Carmen Dana Șandru¹,

¹*Department of Clinical Sciences, Division of Infectious Diseases, University of Agricultural Sciences and
Veterinary Medicine, Faculty of Veterinary Medicine, Cluj-Napoca, Romania*

²*Department of Animal production, Division of Hygiene and Welfare, University of Agricultural Sciences and
Veterinary Medicine, Faculty of Veterinary Medicine, Cluj-Napoca, Romania*

³*Food Engineering Department, Faculty of Food Science and Technology, University of Agricultural Sciences
and Veterinary Medicine Cluj-Napoca, Romania*

Objective / Purpose: Contagious ecthyma is a highly economically impacting disease of sheep. The purpose of this study was to evaluate the immunological effects of nettle extracts in different solvents, also depending of the age category, on the adaptive cell-mediated response in sheep with clinical disease.

Material and Methods: Twelve adult ewes and their lambs, from a 360 animal flock, during a spontaneous outbreak of orf were subject to treatment which started on day 0, while blood was sampled in parallel, from diseased lambs and their mothers and also clinically healthy controls on days 0 and 10. The antioxidant capacity (scavenging effect over DPPH,%) of the nettle extract was established. The *in vitro* blast transformation test was performed to evaluate the effects of two nettle extracts (DMSO and ethylic alcohol). The significance of the differences was estimated by Excel program.

Results: The scavenging effect in the nettle extract was of 68.5±2.03. Both the alcoholic and DMSO nettle extracts significantly (p<0.05-p<0.01) increased the *in vitro* blast transformation indices after the treatment in diseased adult sheep (4.8±1.03 and 7.06±0.92%) and lambs (13.39± 2.45 and 19.68±5.42%), when compared to initial values and those obtained in healthy animals.

Conclusion / Discussion: The *in vitro* blast transformation indices were influenced by the health status of the animals, the age and the solvent used to nettle extraction, allowing the best choice of extraction to achieve stronger adaptive immune responses.

Keywords: *Urtica dioica* L., orf, sheep, adaptive immunity, blast transformation.

Reference:

[1] Gülçin I. Küfrevioğlu Ö. I., Oktay M., Büyükkuroğlu M. E. (2004) Antioxidant, antimicrobial, antiulcer and analgesic activities of nettle (*Urtica dioica* L.) Journal of Ethnopharmacology 90: 205–215

ORAL PRESENTATION

HIV-2 VPX/VPR REGULATE THEIR STABILITY USING TWO MOTIFS

Kazunori Shimagaki¹, Ryoko Koga¹, Haruna Fujino¹, Halil Ibrahim Ciftci¹, Minami Yamamoto¹, Hiroshi Tateishi¹, Masami Otsuka¹, Yoshihiro Yamaguchi², Mikako Fujita³

¹ Department of Bioorganic Medicinal Chemistry, Faculty of Life Sciences, Kumamoto University, Japan

² Environmental Safety Center, Kumamoto University, Japan

³ Research Institute for Drug Discovery, School of Pharmacy, Kumamoto University, Japan

Objective / Purpose: HIV-2 and some SIVs have similar two proteins Vpx and Vpr, and expression level of Vpx is much higher than that of Vpr. The Vpx has unique two motifs, a zinc-binding site HHCC, and a polyproline motif (PPM) which consists of seven consecutive prolines from the P103 residue. In contrast, the Vpr does not have these two regions. We already showed that zinc-binding site of HIV-2 Vpx withstands the instability of the protein (1), and introduction of the zinc-binding residue to HIV-2 Vpr increases its expression level (2) in a cell. About the PPM, we reported that this motif promotes translation, multimerization, and infectivity in macrophages (3, 4).

Materials and Methods: Expression vector for Vpx/Vpr or its mutants was transfected into 293T cells, and the cells were lysed. Immunoblotting of the lysates were then conducted. E.coli was transformed with expression vector for Vpx or its mutants, and lysed. The lysates were centrifuged, and immunoblotting was performed using the supernatants and precipitates.

Results and Discussion: The expression level of seven point PPM mutants of Vpx was low in a mammalian cell. In the presence of proteasome inhibitor or with co-expression of Gag, the expression level was recovered. This recovery was not seen in the zinc-binding site mutants, demonstrating that zinc-binding site and PPM prevent instability of the Vpx differently in a cell. Next, the ratio of H39L mutant (one of the zinc binding sites H39 was changed to L) and P109A mutant (one of PPM amino acid P109 was changed to A) of Vpx in soluble and insoluble fractions after lysis of transformed E. coli was examined, and compared to that of WT. The result showed that more amounts of proteins are insoluble in H39L and P109A, indicating instability of protein structure. Finally, PPM was introduced into HIV-2 Vpr mutant having zinc-binding site HHCC. This Vpr chimeric protein with both zinc-binding site and PPM has further higher expression level than the mutant having only zinc-binding site.

Conclusion: HIV-2 Vpx oppose the instability by carrying both zinc-binding site and PPM, in contrast, HIV-2 Vpr keeps low expression level by not having these two motifs. The motifs are dual tools to regulate their stability in Vpx/Vpr.

Keywords: HIV-2, Vpx, Vpr, zinc-binding site, poly-proline motif

References:

- Yamamoto *et al.* J Gen Virol, 98, 275 (2017)
Koga *et al.* FEBS Open Bio, 8, 146 (2017)
Miyake *et al.* J Gen Virol, 95, 179 (2014)
Ciftci *et al.* FEBS Lett, 589, 1505 (2015)

ORAL PRESENTATION

ANALYSIS OF CORRELATION BETWEEN HEAVY AND ESSENTIAL METALS IN *SALIX ALBA* L. (FAM. SALICACEAE) IN HIGHLY POLLUTED AREA IN KOSOVO

Demush Bajraktari¹, Biljana Bauer², Aneta Dimitrovska², Zoran Kavrakovski², LulzimZeneli³

¹ Kosovo Medicines Agency, Prishtina, Republic of Kosovo, demushbajraktari@yahoo.com

² Faculty of Pharmacy, University of Ss Cyril and Methodius, Majka Tereza No 47, 1000 Skopje, Republic of Macedonia

³ Faculty of Education, University of "Fehmi Agani", Str. "Ismail Qemali", n.n. 50 000 Gjakovë, Republic of Kosovo

Objective/Purpose: Plants, like all other living things, need essential elements for their growth and development. It is determined that different plant cultures, grown in contaminated area have indicated difference in the proportion of the metal absorption, accumulation, distribution and also showed significant difference in concentration in different parts of plant. The selected useful macroelements (Ca, Mg) play a major role in building a plant structure, while microelements (Fe, Cu, Zn, Mn), are principally involved in enzymatic process. The main aim of this study is to investigate the concentration of selected elements in willow bark samples *Salix alba* L. (Fam. *Salicaceae*) in highly polluted area in Kosovo and the correlation between heavy toxic metals in soil and selected elements in plant.

Material and Methods: In highly polluted area near Kosovo Thermo Power Plants (KTPP) in Kosovo, 30 samples of willow bark were taken from the trees in the height 1.5 - 2 m, from the ground and 30 samples of soil were taken in around area 1 - 2 m from the willow tree (30 - 40 cm depth). All samples were collected in distance 30 km along the river stream of Sitnica which passes near the KTPP. Soil samples were air dried, while bark samples initially were thoroughly washed with deionized water to remove any particles or surface impurities attached to the samples, then were dried in thermostat for 24 hours on 65°C. Metal contents were measured by inductively coupled plasma optical emission spectrometry using Perkin-Elmer Optima 2100DV.

Results: Our results indicated accumulation of toxic heavy metals and useful elements in soil and plant. The average values of useful elements in willow bark samples ranged for Ca: 5260.00 – 22280.00 mg/kg; Mg: 840.00 - 1680.00 mg/kg; Fe: 66.79 - 910.75 mg/kg, Cu: 5.09 - 28.66 mg/kg, Zn: 56.39 - 140.94 mg/kg, Mn: 19.68 - 392.75 mg/kg and Al: 67.79 - 894.77 mg/kg.

Conclusion/Discussion: Our data indicated that the correlation between heavy metals in soil and useful elements in plant (Ni/Zn $r = 0.558$; Cr/Cu, $r = 0.343$ and Cr/Mn, $r = 0.385$) and also correlation between heavy toxic metals with useful ones in plant samples Ni/Al ($r = 0.883$), Ni/Fe ($r = 0.790$) Ni/Zn ($r = 0.327$), Cd/Fe ($r = 0.393$) and Cr/Fe ($r = 0.793$)

Keywords: Heavy metals, essential metals, correlation, willow

References:

[1] Jordanoska, B., Stafilov, T., Pelivanoska, V. (2018), Accumulation and availability of trace elements from soil into oriental tobacco grown in Macedonia. *Environmental Engineering and Management Journal*, 17 (6), 1491-1500.

ORAL PRESENTATION

**SYNTHESIS AND CHARACTERIZATION OF SILVER NANOPARTICLES USING
VARIOUS PLANT EXTRACTS AND THEIR VARIOUS ENVIRONMENTAL
APPLICATIONS**

Priya Sundarajan*¹ & Sangeetha Shetty²

^{1, 2} Department of Life Science and Biochemistry, St. Xavier's College (Autonomous), Mumbai, India.
Presenting and corresponding author. Email: priya.s@xaviers.edu

Objective/ Purpose: Silver nanoparticles are being used in numerous technologies and incorporated into a wide array of consumer products that take advantage of their desirable optical, conductive, and antibacterial properties. Considering the advantages of green synthesis of silver nanoparticles, the present study aims at synthesis of silver nanoparticles (AgNPs) using various plant extracts and their characterization.

Materials and methods: The AgNPs were synthesized using about 24 different plant extracts; Turmeric, Henna, Onion, *Moringa*, to name a few. The effect of varying pH and temperature of these extracts on the synthesis of AgNPs was studied. Characterization of nanoparticles was carried out by UV –Vis spectroscopy, Scanning Electron Microscopy (SEM) and XRD in order to determine the size and purity of the samples. They were screened for their antibacterial property against various bacterial cultures using agar diffusion method and zone of inhibition was measured. Minimum inhibitory concentration of silver nanoparticles against *E. coli* and *S. aureus* was determined using broth dilution method. The AgNPs were used for remediation of toxic dyes from water samples and also screened for their anticancer properties against lung cancer cell lines.

Results: The AgNPs that were synthesized using the various plant extracts turned from colourless to brown after 24hrs indicating that ionic silver (Ag⁺) has reduced to metallic silver (Ag⁰). Those SNPs which had a characteristic absorption maximum between 400nm to 480nm indicating the formation of nanoparticles, were selected for further studies namely SNPs made from extracts of Arjuna, Turmeric, Betel-leaf and Henna. Effect of varying pH and temperature on the synthesis of SNPs were determined, indicating that they could be best synthesized within 24hrs at 45°C and are stable at pH 4 (Arjuna), 6 (Turmeric) and having maximum absorbance between 400nm to 480nm. The synthesized extracts were concentrated by lyophilization and characterized by SEM and XRD. The size of the SNPs was confirmed by SEM to be between 70 to 80 nm.

The AgNPs showed good antimicrobial property against the organisms used, and was found to be in the range of 20-100 µg/ml. This antimicrobial effect of the synthesized AgNPs was to disinfect water contaminated with *E. coli* and was found to be effective in eliminating *E.coli* from water samples and hence could be used for water treatment. The nanoparticles were also found to be efficient in the remediation of toxic dyes from waste water samples. The AgNPs were screened for their anticancer properties and were found to be inhibitory to lung cancer cell lines.

Conclusion: From our studies it is clear that the AgNPs synthesized using the various plant extracts namely, by the green synthesis method can be used for various environmental applications.

Keywords: Green synthesis, Silver nanoparticles (AgNPs), Antimicrobial activity, SEM, Water treatment, dye remediation, anticancer effects.

References:

1. Anuj, S.A. and Ishnava, K.B. (2013) Plant Mediated Synthesis of Silver Nanoparticles Using Dried Stem Powder of *Tinospora cordifolia*, Its Antibacterial Activity and Its Comparison with Antibiotics. *International Journal of Pharmacy and Biological Sciences*, 4, 849-86.
2. Gonsalves C, Salanke P, Shetty S and Sundarajan P. (2016); Green Synthesis and Characterization of Silver Nanoparticles. *Int. J. of Adv. Res.* 4(8). 1563-1568] (ISSN 2320-5407).
3. Edison, T.J.I. and Sethuraman, M.G. (2012) Instant Green Synthesis of Silver Nanoparticles Using *Terminalia chebula* Fruit Extract and Evaluation of Their Catalytic Activity on Reduction of Methylene Blue. *Process Biochemistry*, 47, 1351-1357. <http://dx.doi.org/10.1016/j.procbio.2012.04.025>
4. Patil, R.S., Kokate, M.R. and Kolekar, S.S. (2012) Bioinspired Synthesis of Highly Stabilized Silver Nanoparticles Using *Ocimum tenuiflorum* Leaf Extract and Their Antibacterial Activity. *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy*, 91, 234-238. <http://dx.doi.org/10.1016/j.saa.2012.02.009>.

ORAL PRESENTATION

**MICROWAVE HYDRO-DIFFUSION AND GRAVITY OF ESSENTIAL OIL FROM
ZINGIBER OFFICINALE ROSC.**

Lalita Piarasd¹, Panomkorn Kwakhong¹, Rachadaporn Benchawattananon², Kanyarat Holasut^{1*}

¹ *Department of Chemical Engineering, Faculty of Engineering, Khon Kaen University, Khon Kaen 40002, Thailand.*

² *Integrated Science, Forensic Science, Faculty of Science, Khon Kaen University, Khon Kaen 40002, Thailand.*

**Corresponding Author: Holasut, K.; E-mail address: kanyarat@kku.ac.th*

Objective / Purpose: In this study, the optimal conditions of microwave hydro-diffusion and gravity (MHG) to extract the essential oil from Zingiber officinale Rosc (Ginger) were investigated. MHG has been compared with hydrodistillation (HD) in terms of extraction time, chemical composition, cost and energy of the operation.

Material and Methods: Fresh rhizomes of Zingiber officinale Rosc. (Age about 12 months) extracted by MHG and HD methods. Analysis chemical composition of essential oil by gas chromatography – mass spectrometry (GC-MS)

Results: The optimized conditions of MHG was microwave power of 720 W and extraction time of 35 min, identified 45 compounds by GC-MS. On the other hand, HD methods takes 180 min to extraction and only identified 4 compounds.

Conclusion / Discussion: Results of this study showed that MHG better than HD: shorter extraction time, better chemical composition, reduce cost and less than energy consuming.

Keywords: Green extraction, Hydro-diffusion, Solvent-free microwave extraction, Essential oil, Ginger.

References:

- (1) Vatanya L., Natta L., Pornthip D., & Kesra V. (2010). Chemical compositions and antioxidant activity of Zingiber officinale Roscoe Essential oils. *Journal of Science and Technology*. 37(3). 297-312. (In Thai)
- (2) Asma F., Anne-Sylvia F.T., Franco V., Mehrez R., & Farid C. (2010). A surprising method for green extraction of essential oil from dry spices: Microwave dry-diffusion and gravity. *Journal of Chromatography A*. 1217. 7345-7350.
- (3) Asma F., Hasnia B., Rafik B., Zima N., Walid E., Metrez R., & Jallour B. (2017). Efficiency of the optimized microwave extraction assisted extraction on the yield, chemical composition and biological activities of Tunisian Rosmarinus officinalis L. essential oil. 105. 224-233.

ORAL PRESENTATION

THERAPEUTIC POTENTIAL OF OLIVE LEAF POLYPHENOLS (ZeyEX®) IN HUMAN OSTEOARTHRITIC CHONDROCYTES *IN VITRO*

Zehra Aydin Bek¹, Zubeyir Elmazoglu¹, Berna Goker², Berivan Bitik³, Cem Nuri Aktekin⁴,
Cimen Karasu¹

1. Department of Medical Pharmacology, Faculty of Medicine, Gazi University, Ankara Turkey

2. Department of Rheumatology, Faculty of Medicine, Gazi University, Ankara Turkey

3. Department of Rheumatology, Faculty of Medicine, Baskent University, Ankara Turkey

4. Department of Orthopaedic Surgery, Faculty of Medicine, Yildirim Beyazıt University, Ankara Turkey

Objective: Osteoarthritis (OA) is characterized by degeneration of articular cartilage. It is among the most common diseases all over the world and leading cause of chronic morbidity in older population. OA patients have limited treatment options. Inflammation and oxidative stress plays an important role in (OA pathogenesis) the progressive loss of articular cartilage¹. Olive leaf polyphenols like oleuropein and hydroxytyrosol have antioxidant and antiinflammatory effects². Chondroprotective effects of olive leaf extracts has been shown to in animal models³, but the cellular mechanism(s) are unclear. Thus, we aim to investigate the cellular action mechanism of olive leaf polyphenolic extract mixture (ZeyEX®) in human osteoarthritis.

Methods: For primary chondrocyte culture, joint tissues were obtained from OA patients undergoing total knee arthroplasty⁴. Isolated primary chondrocyte cells were treated with different concentrations of ZeyEX®. Cell viability (MTT assay), proliferation (RTA- Icelligence) and ROS production (DCFH-DA assay) were analyzed. Anti-inflammatory effects were revealed with inflammatory markers, IL-1B, IL-6, TNF-a (LUMINEX) and caspase-1 levels (ELISA). Lipid peroxidation, 4-HNE-, AGE-adduct and glutathione peroxidase (GPx) were also measured by ELISA kits.

Results: Olive leaf polyphenols (ZeyEX®) increase cell viability at nanomolar concentrations and reduce intracellular ROS production. ZeyEX® also ameliorates chondrocyte proliferation via regulation of inflammatory markers, lipid peroxidation/protein oxidation (4-HNE, AGE-adducts) and antioxidant enzyme, GPx.

Conclusion: This study, which implies that olive leaf polyphenols (ZeyEX®) is a potential therapeutic and redox modulatory agent for OA *in vitro*, is leading the way for future research.

Keywords: Olive leaf, ZeyEX®, Osteoarthritis, Primary human chondrocytes

Acknowledgements: This study was financially supported by the Scientific and Technological Research Council of Turkey (Grant No: 31S088)

References:

- (1) Marchev AS, Dimitrova PA, Burns AJ, Kostov RV, Dinkova-Kostova AT, Georgiev MI. (2017) Oxidative stress and chronic inflammation in osteoarthritis: can NRF2 counteract these partners in crime? *Ann N Y Acad Sci.* 1401:114-135.
- (2) ALHaithloul HAS, Alotaibi MF, Bin-Jumah M, Elgebaly H, Mahmoud AM. (2019) Olea europaea leaf extract up regulates Nrf2/ARE/HO-1 signaling and attenuates cyclophosphamide-induced oxidative stress, inflammation and apoptosis in rat kidney. *Biomed Pharmacother.* 111:676-685.
- (3) Takuma M, Haruka K, Mutsuto W, Toshiki M, Kenshiro M, Akane T, Hiroshi M, Yoshihiro N. (2018) Olive leaf extract prevents cartilage degeneration in osteoarthritis of STR/ort mice. *Biosci Biotechnol Biochem.* 82:1101-1106.
- (4) Isolation and Cell Culture of Primary Human Chondrocytes Technical Reference Guide (Lonza).

ORAL PRESENTATION

EFFICACY AND SAFETY STUDY OF THE NEW FORMULATION OF HERBAL COFFEE ON WEIGHT CONTROL

Nuttinee Teerakulkittipong^{1*}, Yanisa Chanthophat¹, Papawarin Han-gla¹, Phatcharat Lueangworachartkul¹, Suthabodee Muongmee¹, Monsiri Chetkhtkran¹ and Suchada Sukontanon¹

¹Department of Biopharmaceutical science, Faculty of Pharmaceutical Sciences, Burapha University, 169 Saensuk Subdistrict, Muang District, Chonburi, Thailand, nuttinee@go.buu.ac.th*, s.sukontanon@gmail.com,

Objective / Purpose: Overweight and obesity are the majorly underlying factors in public health leading to clinical complications. A new coffee formula with natural extracts was developed to increase metabolic rate and would possibly be used as an alternative for weight control. We aimed to formulate and study on the efficacy, safety, and satisfaction of the coffee formula containing five herbal extracts compared with placebo.

Materials and Methods: Herbal coffee have been made from the mixture of five herbal extracts from Guarana (*Paullinia cupana*), Lotus leaves (*Nelumbo nucifera*), Turmeric (*Curcuma loga*), Ginger (*Zingiber officinale*) and Black ginger (*Kaempferia parviflora*) with coffee in vary proportions to obtain twelve different formulations, each herbal coffee bag contained approximately 14 g of the product and have been selected based on the stability test and favor. 24 overweight participants were included and randomized into 2 groups, 12 participants for each in which one group received coffee only while another one received the herbal coffee. Once daily consumption for 30 days was assigned in both groups as well as the body composition that was examined on day 0, 14, and 30.

Results: The results indicated that the formulation number 5 showed the promising result in stability test and taste preferences. Study results showed the group receiving herbal coffee were statistically different in body fat percentage and serum LDL-cholesterol decrease compared within group (P=0.073, 0.036 respectively at 90% and 95% CI). The coffee were found to be safe in both group of participants. The satisfactory level was moderate.

Conclusion / Discussion: In summary, this new herbal coffee tended to be safe with the promising results for weight control in a short term study. For statistically significant difference of outcomes, further experiment on efficacy and safety in longer period should be developed.

Keywords: Overweight, obesity, herbal coffee

References:

1. Islam D, Huque A, Mohanta LC, Lipy EP, Rahman MN, Sultana A, Sheuly UK. Studies on the Hypoglycemic and Hypolipidemic Effects of Nelumbo nucifera Leaf in Long-Evans Rats. Journal of Diabetes Mellitus. 2017 Jul 11; 7(03):55
2. Promthep K., Eungpinichpong W., Sripanidkulchai B., Chatchawan U. Effect of Kaempferia parviflora extract on physical fitness of soccer players: A randomized double-blind placebo-controlled trial. Med. Sci. Monit. Basic Res. 2015; 21:100–108.
3. Han LK, Gong XJ, Kawano S, Saito M, Kimura Y, Okuda H. Antiobesity actions of Zingiber officinale Roscoe. Yakugaku zasshi: Journal of the Pharmaceutical Society of Japan. 2005 Feb;125(2):213-7.
4. Choi BK, Park SB, Lee DR, Lee HJ, Jin YY, Yang SH, Suh JW. Green coffee bean extract improves obesity by decreasing body fat in high-fat diet-induced obese mice. Asian Pacific journal of tropical medicine. 2016 Jul 31;9(7):635-43.
5. Dietpill-reviews.co.uk. (2018). Guarana for Weight Loss. [online] Available at: <http://dietpill-reviews.co.uk/guarana-for-weight-loss/> [Accessed 12 Feb. 2018].)

ORAL PRESENTATION

PHYTOCHEMICAL PROFILE AND ANTIMICROBIAL POTENTIAL OF EXTRACTS OBTAINED FROM *THYMUS MARSCHALLIANUS* WILD

Mihaela Niculae¹, Daniela Hanganu², Iliora Oniga², Daniela Benedec², Radu Giupana¹,
Carmen Dana Sandru¹, Marina Spinu¹

¹ Department of Infectious Diseases, Faculty of Veterinary Medicine, University of Agricultural Sciences and Veterinary Medicine Cluj-Napoca, Romania, niculaemihaela1@gmail.com

² Department of Pharmacognosy, Faculty of Pharmacy, University of Medicine and Pharmacy "Iuliu Hațieganu" Cluj-Napoca, Romania, danahcluj@gmail.com

Objective / Purpose: Previous studies highlighted for several medicinal plants belonging to the Lamiaceae family multiple valuable biological activities and therapeutic applications [1]. This study was aimed to evaluate the phytochemical profile and antimicrobial potential of two extracts obtained from *Thymus marschallianus* Willd., an Eurasian species of Lamiaceae family.

Material and Methods: Two extracts were prepared from *Thymus marschallianus* Willd. harvested from spontaneous Flora and culture, respectively. Extract preparation was made by maceration with ethanol 70%, 10 days, at room temperature, using blooming aerial part of the plants. Qualitative and quantitative characterization of the extracts (polyphenolic, flavonoid and phenylpropane derivatives profiles) was performed by chromatographic (HPLC- MS/ESI+ analysis) and spectrophotometric methods. The disk diffusion method was used for the *in vitro* antimicrobial screening, while minimum inhibitory concentrations and minimum bactericidal concentrations were established based on the broth microdilution assay.

Results: The chromatographic analysis allowed the identification of 6 flavonoids, 2 aglycons (quercetol, luteolin), 4 flavonoids (derivatives of quercetol, luteolin and apigenin) and rosmarinic acid and its derivative, methyl-rosmarinate. Spontaneous flora extract presented a higher content in flavonoid aglycon, but less rosmarinic acid. The spectrophotometric results showed similar values for total polyphenols, total flavonoids, and total caffeic acid derivatives. Also, both extracts displayed promising antimicrobial potential against Gram positive and Gram negative bacteria, with the highest efficacy expressed towards the *Bacillus cereus* > *Staphylococcus aureus* > *Escherichia coli* reference strains. The MBC/MIC ratio indicated bactericidal activity against Gram positive bacteria.

Conclusion / Discussion: Our data suggested that the extracts from *Thymus marschallianus* may represent sources of antimicrobials and further studies are intended to investigate the mechanism of action and also the potential cytotoxicity.

Keywords: *Thymus marschallianus* Willd., phytochemical profile, *in vitro* antimicrobial efficacy

References:

[1] Khoury, M., Stien, D., Eparvier, V., Ouaini, N., El Beyrouthy, M. (2016). Report on the Medicinal Use of Eleven Lamiaceae Species in Lebanon and Rationalization of Their Antimicrobial Potential by Examination of the Chemical Composition and Antimicrobial Activity of Their Essential Oils. Evidence-based complementary and alternative medicine: eCAM 2016, 2547169. doi:10.1155/2016/2547169

ORAL PRESENTATION

**HEADSPACE SOLID PHASE MICROEXTRACTION (HS-SPME) ANALYSIS OF
SIDERITIS OZTURKII AYTAC & AKSOY**

Tuna Uysal, Ela Nur Şimşek Sezer, Meryem Bozkurt

*Department of Biology, Faculty of Art and Science, University of Selçuk, Konya,
Turkey, tuysal@selcuk.edu.tr*

Objective / Purpose: The genus *Sideritis* (Lamiaceae) is represented by more than 150 bushy species and annual or perennial plants. Many studies have been conducted to elucidate the chemical composition and pharmacological activities of plants of the genus. *Sideritis ozturkii* Aytac & Aksoy is an endemic plant to Turkey and used as herbal tea and folk medicine in central Anatolia. The purpose of this study was the detection of the volatile oil components of *S. ozturkii* by solid phase micro-extraction/gas chromatography/mass spectrometry (SPME-GC-MS) technique.

Material and Methods: The plant material was collected from Konya-Derebucak province in 2016. The collected plant material washed and dried without sunlight. Dried and powdered samples were used directly. Two SPME fibers, 50/30 µm DVB/CAR/PDMS and 85 µm Carboxen/PDMS were preferred for analysis. SPME procedure were analysed on a Shimadzu QP2010 ULTRA FID GC-MS system. The identification of volatile components was carried out by comparing three libraries with W9N11, SWGDR4G4, and SWGDR4G5. Analyses were performed triplicate.

Results: The volatile oil compositions were identified in *Sideritis ozturkii* samples by SPME-GC/MS analysis using two different fibers. The major constituents of dried leaves detected with 50/30 µm DVB/CAR/PDMS fiber were octanoic acid (11.70 %) and nonanoic acid methyl ester (16.68 %) for 85 µm Carboxen/PDMS fiber, respectively. On the other hand, the major constituents of flowers were detected with both fibers were hexanoic acid (24.32 %- 26.34 %) and hexanal (18.67 % -16.68 %) respectively.

Conclusion: The comparative volatile oil composition of *S. ozturkii* leaves and flowers were revealed via SPME method for the first time with this study.

Keywords: Paşa çayı, Solid phase microextraction, Turkey.

ORAL PRESENTATION

ANTIBACTERIAL ACTIVITY AND COMPOSITION OF ESSENTIAL OILS OF LEAVES AND BASIL FLOWERS OF THREE DIFFERENT GROWING SEASONS UNDER ARID CLIMATE

Nabila Slougui^{1,2}, Ahlem Tlili², Hadjer Bentayeb², Mahfoud Hadj Mahammed²

¹Department of the preparatory class National Polytechnic School of Constantine., Algeria,
slougui.nabila@gmail.com

² Biogeochemistry Laboratory of the desert Environments, Kasdi Merbah University, Ouargla, Algeria

Objective / Purpose: The evaluation of the antibacterial activity of essential oils of basil grown under arid climate has been studied by many researchers [1]. We wanted to study this property in relation to the growing seasons of the plant as the first parameter of study, and what part of plant that will present a better antibacterial activity as the second parameter of study. The relationship between this activity and the majority composition of the essential oil has been developed.

Material and Methods: The antibacterial activity of our 6 essential oils was evaluated on three two Gram-negative strains: *Escherichia coli* ATCC 25923 and *Pseudomonas aeruginosa* ATCC 27853 and one gram positive: *Staphylococcus aureus* ATCC 25922. The culture media used are nutrient agar (GN) for the maintenance of bacterial strains and Mueller Hinton agar (MH) for the study of bacterial susceptibility to HE. The diffusion method in Agar medium (Aromatogramme) is used for the study of antibacterial activity which allows to determine the sensitivity of the different bacterial species vis-à-vis our essential oils. The majority composition of essential oils was made by a Shimadzu GC-MS – QP2020.

Results: The results obtained showed that the essential oil of the different parts of the *Ocimum basilicum* L. Had no antibacterial activity marked against the strain of *Pseudomonas aeruginosa* (ATCC 27853). According to our results, the essential oil of basil Autumn flowers inhibits bacterial growth with inhibition diameters of 21.5 and 25 mm against *Escherichia coli* and *Staphylococcus aureus* respectively. This has a direct relationship with its majority composition: Linalool (34.99%), methylcinnamate (21.09%) and Methylchavicol (14.53%).

Conclusion / Discussion: The essential oils of the flowers showed antibacterial activities better than that of the leaves and this in the three study seasons.

Keywords: Activité antibactérienne, *Ocimum basilicum* L, huile essentielle

References:

[1] Mith, H., Yayi-Ladékan, E., Kpoviessi, S, D, S., Bokossa, I, Y., Moudachirou, M., Daube, G. & Clinquart, A., (2016) Chemical Composition and Antimicrobial Activity of Essential Oils of *Ocimum basilicum*, *Ocimum canum* and *Ocimum gratissimum* in Function of Harvesting Time, *Journal of Essential Oil Bearing Plants*, 19 (6), 1413-1425.

ORAL PRESENTATION

THE ESSENTIAL OIL YIELD AND COMPOSITIONS of TARRAGON (*Artemisia dracunculus* L.) CULTIVATED IN BAYBURT

S.Ayşe Çelik¹ İrem Ayrın¹ Sefa Akbulut² Rabia Serpil Günhan³ Asuman Kan³ Yüksel Kan¹

¹Selçuk University, Agriculture Faculty, Department of Medicinal Plants, Konya/Turkey

²Karadeniz Technical University, Faculty of Forest, Department of Forest Engineering,
Trabzon/Turkey

³Konya Technical University, Technical Sciences Vocational School of Higher Education, Department of Food
Processing, Konya/Turkey

E-mail: sadiyeayse.sayar@gmail.com phone + 90-539 799 33 63

Objective / Purpose: In the study, the essential oil yield and compositions obtained from herb of tarragon (*Artemisia dracunculus* L.) cultivated in Bayburt, Turkey ecological conditions were investigated.

Material and Methods: The air-dried herb of tarragon was subjected to hydrodistillation for 3h using a Clevenger-type apparatus to produce essential oil. The GC-MS analysis was carried out with Agilent 7890 GC-MS system. The relative percentages of the separated compounds were calculated from total ion chromatograms. The identification of the oil components was based on the Wiley and NIST mass spectral library.

Results: The essential oil yield of tarragon herb was determined as 0.3%. The most important essential oil component of metil kavikol is 65.3%. The major essential oil components of tarragon were determined as (*Z*)-beta-Osimen (7.9%), (*E*)-beta-Osimen (5.7%), Timol (3.4%), respectively.

Conclusion / Discussion: Tarragon (*Artemisia dracunculus* L.) was cultivated commercially in Bayburt province. According to the results of this study, it was determined that essential oil yield and components of tarragon were determined. Tarragon, a medical and aromatic plant, is known to be rich in content in terms of seconder metabolits. This exotic plant for Turkey has been used traditionally as spices herb, cooking, and salads.

Keywords: Tarragon, *Artemisia dracunculus* L., essential oil, composition

ORAL PRESENTATION

**THE EFFECTIVENESS OF THAI TRADITIONAL HERBAL DECOCTION
CONSUMED IN THE NORTHEAST OF THAILAND**

Rachadaporn Benchawattananon^{1*} and Kanyarat Holasut^{2*}

*1*Integrated Science, Forensic Science, Faculty of Science, Khon Kaen University, Khon Kaen 40002, Thailand.*

*2*Chemistry, Faculty of Engineer, Khon Kaen University, Khon Kaen 40002, Thailand.*

Corresponding Author: Benchawattananon, R.; E-mail address: rachadaporn@kku.ac.th

Objective / Purpose: The study is studied the effectiveness of Thai traditional herbal decoction consumed in Khon Kaen and Chaiyaphum province located in the northeast of Thailand. The information on the use of decoction in the treatment of fatigue are flu-like syndrome and diabetes.

Material and Methods: The research was a gather from 10 traditional herbal healers 3 monks and the other village through 80 interviews. All of the conversations with the information were conducted in the Thai language. The data were recorded as documents, questionnaires, sound recording and photographs.

Results: The finding were 1) 86% commonly used cause of get well sooner than consumed tablets and capsule pattern. 2) 90% specific to personal up to weight, age, allergy and symptoms. 3)92% Safe without chemicals effect to their health 4) 79% Following the herbal ways by believe in ancestors 5) 84% Believe in social media.

Conclusion / Discussion: The Thai traditional herbal decoction consumed in Khon Kaen and Chaiyaphum province located in the northeast of Thailand. The information on the use of decoction in the treatment of fatigue, flu-like syndrome and diabetes. People still consumed Thai traditional herbal decoction in their daily lives. The promoting Thai traditional herbal knowledge should be transferred for younger generations.

Keywords: Thai traditional herbal decoction, Thailand, flu-like syndrome

References:

- (1) Somtanuek C., Promchit S., Thaya J., Saowapa P., & Sompop P.(2012).Medicinal plants used with Thai traditional Medicine in modern healthcare service: A case study in Kabchoeng Hospital, Surin Province,Thailand.Journal of Ethnopharmacology. 141(2012),193-203.
- (2) Somasundaram A.& Kenichi W.(2017).Japanese Kampo Medicine for the Treatment of Common Disease: Foguson Inflammation.143-164.

ORAL PRESENTATION

STABILITY STUDY OF OSELTAMIVIR EXTEMPORANEOUS SUSPENSION

Thatsanand Xayavong¹, Theera Rittirod²

¹Graduate student, ²Assoc. Professor,
Faculty of Pharmaceutical Sciences, Khon Kaen University, Khon Kaen 40002, Thailand

*Corresponding author : E-mail: Theerarit@gmail.com

Objective / Purpose: The stability of Oseltamivir Extemporaneous Suspension (OSEL ES) used in Queen Sirikit National Institute of Child Health (QSNICH), Bangkok Thailand was investigated.

Material and Methods: Solution of 10 mg/ml OSEL ES were compounded and stored in three different temperature conditions; room temperature, refrigerated temperature (7 ± 3 °C) and accelerated temperature (45 ± 5 °C). Triplicate samples were periodically taken for 12 weeks and assayed by high-performance liquid chromatography.

Results: All of the samples stored under room temperature and refrigerated conditions showed more than 90% of their initial concentrations of OSEL remained throughout the course of the study. For the accelerated temperature 62.91% of the initial concentration of OSEL remained at 12th week, which was significantly lower than the other conditions ($p < 0.05$). In addition, samples stored at 45 ± 5 °C showed changes in color and reductions in pH over the course of the experiment.

Conclusion / Discussion: In conclusion temperature above room temperature should be included for stability testing of extemporaneously prepared suspensions in tropical climates.

Keywords: oseltamivir phosphate, extemporaneous suspension, physicochemical tests, stability test

References:

Albert K., & Bockshorn J. Chemical stability of Oseltamivir in oral solutions. *Pharmazie*.2007;62:678-682.
Thatsanand Xayavong and Theera rittirod, Stability of Oseltamivir Extemporaneous Suspension, *Asia Pacific J. Sci and Tech.* 24 (1) Jan-March 2019 (in Press)

ORAL PRESENTATION

EFFECTS OF OLIVE LEAF (*OLEA EUROPEA L.*) EXTRACT ON THE GROWTH PERFORMANCE AND SOME BLOOD PARAMETERS OF COMMON CARP (*CYPRINUS CARPIO*)

Ebru Yilmaz¹, Deniz Çoban¹, Murat Er²

¹Adnan Menderes University, Faculty of Agriculture, Department of Aquaculture, Aydın, Turkey, ebruyilmaz@adu.edu.tr

²Adnan Menderes University, Faculty of Veterinary Medicine, Department of Animal Nutrition and Nutritional Diseases, Aydın, Turkey

Objective / Purpose: Herbs or spices have been reported to promote various functions like growth, appetite stimulation, antistress, immune functions, skin coloration, egg hatching rates, hematological and biochemical status and also increase disease resistance in fish culture due to different active components. Olive leaf is the most important source of oleuropein known in nature. Several researchers revealed the fact that oleuropein has many pharmacological qualities including antioxidant, antimicrobial, antiaterogenic and antiviral activities. The olive leaf extract that we used in study include oleuropein at the rate of 2.3%. In this study, the effects of olive leaf extract on growth performance and some blood parameters of common carp were evaluated. For this purpose, 3 different experimental diets containing rate of 0% (OLE0), 0.25% (OLE025) and 0.5% (OLE05) olive leaf extract were prepared. This study consisted of 3 repetitions for each group with 71.58±0.66 g and 200 L aquarium as 25 fish per aquarium for 60 days. The objective of this study was to determine the effects of olive leaf extract on the growth performance and some blood parameters in the common carp.

Material and Methods: *Cyprinus carpio* fishes were obtained from a local fish farm in İzmir, Turkey. Olive leaf extract was obtained from Talya Herbal product. It was incorporated to a laboratory-prepared feed rate of 0% (control group), 0.25% and 0.50% for diets designed OLE0, OLE025 and OLE05, respectively. A control group was prepared without any supplements. Proximate analyses of the diets were performed using standard methods. Haematological parameters (RBC count, Hct ration, Hgb concentration) were determined by using an automatic analyser (Mindray BC-3000 Plus, China). Creatinine, total protein, glucose, albumin, globulin, uric acid, urea, glutamic oxaloacetic transaminase, glutamic pyruvic transaminase and cholesterol were assessed by the automatic analyzer apparatus (Reflotron plus, Roche, Germany) using commercially available Reflotron kits (Roche Diagnostics, Germany). The analyses of the data obtained in the experiment were done by using the SPSS 17 statistics program. One way variance analysis (ANOVA) was applied to the data which were then subjected to Tukey multiple comparison test. The differences between groups were evaluated to be $p < 0.05$.

Results: The three diets were equally accepted by the fish and there was no mortality or disease in any treatment. There were no significant differences ($P > 0.05$) in average weight gain, feed conversion ratio (FCR) and specific growth rate (SGR). The RBC count, Hgb concentration, Hct ratio, Mean corpuscular hemoglobin concentration (MCHC), Mean corpuscular volume (MCV) and Mean corpuscular hemoglobin (MCH) in the treatment groups were statistically similar to the values recorded for the control group ($p > 0.05$). In serum biochemical findings of fish, olive leaf extract was found to be a significant difference among the groups ($p < 0.05$).

Conclusion / Discussion: In conclusion, findings of present study indicate that feeding *Cyprinus carpio* with diet containing 0.5% olive leaf extract over a period of 60 days might be adequate to improve fish immune parameters.

Keywords: *Cyprinus carpio*, Olive leaf (*Olea europea L.*) extract, Blood parameter, Growth performance

References:

- [1] Adams, S.M., & Greeley, M.S. (2000). Ecotoxicological Indicators of Water Quality: Using Multi-Response Indicators to Assess the Health of Aquatic Ecosystems. *Water, Air, and Soil Pollution*, 123(1), 103-115.
- [2] AOAC. (2000). Official methods of analysis. Association of Official Analytical Chemists. (17th ed.) Washington, USA.
- [3] Awad, E.S. (2010). Studies on Plantbased Dietary Supplements for Control of *Aeromonas hydrophila* Infections in Rainbow Trout (*Oncorhynchus mykiss* Walbaum) (PhD Thesis). Heriot Watt University, UK.
- [4] Baba, E., Acar, Ü., Yilmaz, S., Zemheri, F., & Ergün, S. (2018). Dietary olive leaf (*Olea europea L.*) extract alters some immune gene expression levels and disease resistance to *Yersinia ruckeri* infection in rainbow trout *Oncorhynchus mykiss*. *Fish Shellfish Immunology*, 79, 28-33.
- [5] Bain, B.J., Lewis, S.M., & Bates, I. (2006). Dacie and Lewis Practical Haematology, Philadelphia, Elsevier Press., 600 pp.
- [6] Bircan, C. (2006). Determination of aflatoxin contamination in olives by immunoaffinity column using high-performance liquid-chromatography. *Journal of Food Quality*, 29(2), 126-138.

ORAL PRESENTATION

LEAF ANATOMY OF THE GENUS *ORIGANUM* L. (LABIATAE) IN TURKEY

Narin Sadıkoğlu¹, Mehmet Efe²

¹ Department of Pharmacognosy, Faculty of Pharmacy, İnönü University, Malatya, Turkey,
organum@hotmail.com

² Department of Nursing, School of Health, Muş Alparslan University, Muş, Turkey

Objective / Purpose: The plants of the genus *Origanum* L, which is used as oregano in the Labiatae family, are mainly used as traditional folk medicine and spices. In 2018 with addition of new taxa to the flora, the genus *Origanum* is represented by 67 natural taxa in the world, 34 of them occurs in Turkey [1]. These are mainly found in the Mediterranean region. In particular, there are difficulties in distinguishing closely allied taxa. A detailed study has been carried out in order to distinguishing these taxa according to anatomical characteristics. The study is carried out in order to contribute to the identification of taxa that cannot be diagnosed morphologically if it contains missing diagnostic character.

Material and Methods: Leaf samples are taken from a collection of the specimens collected from the nature between 2001-2012 by Narin Sadıkoğlu, located in the Herbarium of İnönü University, Faculty of Pharmacy. Cross-sections are taken from the center of the leaves at 15 & 25 microns thickness with a microtome.

Results: The leaves are isolateral, covered with a layer of cuticle. Epidermis consists of single-layered rectangular cells. Cover and glandular hairs, Labiatae type glandular hairs on the epidermis can be seen. The stomata are 2 (diacytic)-3, rarely 4 (anisocytic) of Labiatae type and they are hygromorphic or mesomorphic. Palisade parenchyma cells are single layered and have much chloroplasts. Sponge parenchyma cells are in 1-5 layered and intercellular spaces are evident.

Conclusion / Discussion: *Origanum* taxa differs in the characters such as cover hair, glandular hair, shape of upper and lower epidermis, number and size of cells, stomata type, crystals, cuticle, width of the vascular bundle. According to the data obtained as a result of microscopic analysis of the taxa, the distinctive basic characters were determined.

Keywords: *Origanum*, leaf anatomy, Labiatae, Turkey

Acknowledgements: The authors would like to thank to Bio. İsmet Gürhan for photographing and preparing the materials, Fedli Ekici, the Director of Muş State Hospital for allowing the use of the microtome and Laboratory Assistant Murat Çelik for his assistance in cross-section. This study was supported by Research Fund of the İnönü University. Project Number: 2011/155.

References:

[1] Celep, F., Dirmenci, T. (2017). Systematic and Biogeographic overview of Lamiaceae in Turkey. *Nat. Volatiles & Essent. Oils*, 4(4), 14-27.

ORAL PRESENTATION

**CHARACTERIZATION OF VOLATILE COMPOUNDS OF *MENTHA PULEGIUM*
VIA HS-SPME**

Ela Nur Şimşek Sezer, Meryem Bozkurt, Nihal İlhan, Tuna Uysal

*Department of Biology, Faculty of Art and Science, University of Selçuk, Konya,
Turkey, elasimsek@selcuk.edu.tr*

Objective / Purpose: *Mentha pulegium* L. is a herbaceous perennial plant that belonging to the Lamiaceae family. The plant grows up in moist areas of the mountains and plains. In therapeutic applications, this plant and/or its active constituents have been used traditionally for several disorders. Their diuretic, antispasmodic, digestive, diaphoretic, sedative, antiseptic, stimulant and expectorant effects have been reported. The aim of this study is the determination of the volatile and semi-volatile oil content of *Mentha pulegium*.

Material and Methods: The *Mentha pulegium* species was collected from its natural habitat. The powdered plant samples were analyzed directly on a Shimadzu QP2010 ULTRA FID GC-MS system. Volatile compounds were detected by solid-phase microextraction/gas chromatography/mass spectrometry (SPME-GC-MS) method. Two SPME fibers preferred for analyses; 85 µm Carboxen/PDMS and 85 µm polyacrylate. The components were identified by comparing three libraries with W9N11, SWGDR4G4, and SWGDR4G5. Analyses were performed three times and the mean values were taken into consideration.

Results: The volatile oil compositions were identified in *Mentha pulegium* samples by SPME-GC/MS analysis. According to the analyses, significant differences were observed between the two SPME fibers. The major volatile component for 85 µm Carboxen/PDMS fiber were 2-Isopropyl-5-methylhex-2-enal (30.37%) and for 85 µm polyacrylate fiber 2(3H)-Benzofuranane, 3a,4,7,7a-tetrahydro-3a-methyl (48.59%), respectively. In addition, the most common and shared components to both fibers were, p-Menthone (14.15% - 8.82 %), phenol,5-methyl-2-(1-methylethyl)(8.68 % - 7.67 %).

Conclusion: *Mentha pulegium* is a plant that is rich in essential oils and this study is the first study about the volatile content of plant with SPME method.

Keywords: Yarpuz, SPME-GC-MS, Turkey.

ORAL PRESENTATION

**CYTOTOXIC AND GENOTOXIC EFFECTS OF WATER EXTRACTS OF
ZYGOPHYLLUM CORNUTUM AND *ATRACYLIS GUMMIFERA* USING *ALLIUM
CEPA* ASSAY**

Awatif Boumaza¹, Messaouda Khallaf¹, Mouna Cherairia¹

¹Département de Biologie, Faculté des Sciences de la Nature et de la Vie et Sciences de la Terre et de l'Univers.
Université 8 Mai 1945 Guelma

Objective / Purpose: The interest in herbal products in the world is fueled by the rising costs of drugs. The safety of herbal products depend on the safety of its sources, and this should be established before these are developed as herbal medicinal products. The potential genotoxic effects of aqueous extracts of *Zygophyllum cornutum* and *Atractylis gummifera*, which are commonly used in traditional medicine to treat a variety of disease conditions, was investigated using *Allium cepa* assay.

Material and Methods: Both extracts of 2, 4, 6, 8, 10 and 12 mg/mL were tested on root meristems of *A. cepa* to determine the half maximal effective concentration (EC50) as a parameter of cytotoxicity. Synthetic water was used as negative control [1]. Based on EC50 corresponding values, three concentrations were used for each plant to evaluate the genotoxic effects on *A. cepa* [1][2].

Results: The results showed a dose-dependent decrease in roots growth with both plants, but it was more remarkable with *A. gummifera*. According to the obtained EC50 values, *Z. cornutum* (9.38 mg/ml) is considered less toxic than *A. gummifera* (5.38 mg/ml). The results of genotoxic test showed that mitotic index decreased as the concentrations of the extract increased. For *Z. cornutum*, the three tested concentrations 6, 10 and 12 mg/ml had respectively 83.56, 77.49 and 65.12 % mitotic indexes, while with *A. gummifera*, the three tested concentrations 2, 6 and 12 mg/ml had respectively 67.92, 58.19 and 22.38 % mitotic indexes.

A dose-dependent increase of total chromosome aberrations (TCA) was also observed. For *Z. cornutum* (concentration mg/ml; TCA%): (6 ; 3.85), (10 ; 5.25) and (12 ; 6.87). For *A. gummifera*: (2 ; 13.81), (6 ; 32.35) and (12 ; 48.67).

Conclusion / Discussion: The results of this study suggested that the aqueous extracts of *Zygophyllum cornutum* and *Atractylis gummifera* exerted significant genotoxic and mitodepressive effects at the tested concentrations.

Key words: *Allium cepa*, *Atractylis gummifera*, Chromosome aberrations, cytotoxicity, genotoxicity, Mitotic index, *Zygophyllum cornutum*

References:

- [1] Rank, J., (2003). The method of *Allium* anaphase–telophase chromosome aberration assay. *Ekologija*, 38–42
[2] Fiskesjo, G., (1985). The *Allium* test as a standard in environmental monitoring, *Hereditas* 102 99–112.

ORAL PRESENTATION

**ADVERSE EFFECTS OF CONVENTIONAL DRUG-MEDICINAL PLANT
INTERACTIONS IN VETERINARY MEDICINE**

Yigit Gunes^{1a}, Tulay Bakirel^{1b}, Oya Ustuner^{1c}

^{1a}*Department of Pharmacology and Toxicology, Faculty of Veterinary Medicine, Istanbul University-Cerrahpasa, Avcilar 34320, Istanbul, Turkey, yigit.gunes@istanbul.edu.tr*

^{1b}*Department of Pharmacology and Toxicology, Faculty of Veterinary Medicine, Istanbul University-Cerrahpasa, Avcilar 34320, Istanbul, Turkey, tulbakir@istanbul.edu.tr*

^{1c}*Department of Pharmacology and Toxicology, Faculty of Veterinary Medicine, Istanbul University-Cerrahpasa, Avcilar 34320, Istanbul, Turkey, oyakeles@istanbul.edu.tr*

The use of herbal remedies for the prevention and the treatment of a variety of illnesses in small animals have been increasing nowadays. Because of the increased interest in natural products in veterinary medicine, more information is needed regarding their use and potential hazards. Although extensive literatures are available on the effectiveness of herbal extracts and components, there is no profound information on the safety assessments of the combined use of herbal medicines and conventional drugs in veterinary medicine.

Particularly, the interaction of the biological active constituents contained in herbal remedies with conventional drugs is a major concern. Drug-herb interactions refer to the possibility that an herbal constituent may alter the pharmacologic effects of a conventional drug. Possible pharmacokinetic interactions include those that alter the absorption, metabolism, distribution, or elimination of a drug or herbal constituent, resulting in an increase or decrease in the concentration of active agent at the site of action. In addition, the interaction of the herbal medicines and conventional drugs may lead to potential toxic effects. Some interactions are documented in clinical trials, some are inferred from in vitro experiments, and others are suspected only on theoretical grounds. Numerous case reports in literature serious and potentially life-threatening adverse effects following human and animal exposure to herbal preparation are seen.

We are of the opinion that the review not only highlights the most updated research results available in the evaluation of the possible risks of herb-drug interactions, but also provides some essential skills for the healthcare researchers and veterinary clinicians to solve some relevant issues they may encounter in this field.

Keywords: Medicinal plants, conventional drugs, interaction, veterinary therapy

ORAL PRESENTATION

**DETERMINATION OF VOLATILE AND SEMI-VOLATILE OIL CONTENT OF
ISATIS TINCTORIA L. VIA SPME-GC-MS**

Meryem Bozkurt, Ela Nur Şimşek Sezer, Nihal İlhan, Tuna Uysal

Department of Biology, Faculty of Art and Science, University of Selçuk, Konya,
Turkey, mbozkurt@selcuk.edu.tr

Objective / Purpose: *Isatis tinctoria* L. is a species of the genus Brassicaceae and generally distributed in Central Asia. The plant is also used for medical purposes as well as being used as a colorant. *Isatis* is one of the most effective herbs used in herbal medicine, antiseptic and antiviral drugs in traditional Chinese medicine. The aim of this study is to determine and characterize the volatile oil components found in the leaves and flowers of *Isatis tinctoria* via the SPME-GC-MS method.

Material and Methods: The plant material was collected from the Ardıçlı Village (Konya Province) and preserved under suitable conditions. 65 µm PDMS/DVB and 85 µm polyacrylate fibers were used to determine the volatile and semi volatile oil composition of the samples. Analyzes were performed using the Shimadzu QP2010 ULTRA FID GC-MS system. The components were identified by comparing three libraries with W9N11, SWGDR4G4 and SWGDR4G5.

Results: According to the data obtained from the analysis of *Isatis tinctoria*, in leaf content 120 and 78 compounds and in flower content 123 and 87 compounds were detected respectively. According to the analysis, with 65 µm PDMS/DVB fiber, the most amount of components in the leaf benzeneetanol (4.67 %) and undecane,5-methyl (3.93 %), the most common components in the flower, morpholine, 4-octadecyl- (18.68 %), 4-Amino-furazan-3-carboxylic acid (3-morpholin-4-yl-propyl)-amide (11.91 %). With 85 µm polyacrylate fiber, the most amount of components in the leaf, benzene, 2,4-diisocyanato-1-methyl- (33.41 %) and pyrrole-3-carbonitrile, 5-formyl-2,4-dimethyl (14.47 %), in the flower content 2,2,3,3,4,4,4-Heptafluoro-N-(2-morpholin-4-yl-ethyl)-butyramide (19.67 %) and morpholine, 4-octadecyl- (18.07 %), respectively.

Conclusion: This study is the first study in terms of characterization and comparison of volatile and semi volatile oil components by SPME-GC-MS analysis from leaves and flowers of *Isatis tinctoria* L. species from Turkey.

Keywords: Çivitotu, solid phase micro-extraction, Turkey.

ORAL PRESENTATION

**THE CHEMICAL COMPOSITION OF *DELPHINIUM SZOWITSIANUM* BOISS.
SPECIES SPREADING IN VAN REGION**

Murat Tuncturk¹, Tamer Eryigit², Ruveyde Tuncturk¹, Lütü Nohutçu^a

¹Van Yuzuncu Yil University, Agricultural Faculty, Field Crops Department, Van-Turkey
Y.Y.U. Gevas Vocational School, Department of Plant and Animal Production, Van-Turkey
E-mail: ruveydetuncturk@yyu.edu.tr

The family Ranunculaceae, which includes the Delphinium species, has 59 genera and 2500 species worldwide. In Turkey, it has 18 genus and 216 species and its endemism rate (23.5%) is quite a high family. *Delphinium szowitsianum* Boiss. is a perennial herbaceous species. It has been spread in the Eastern Anatolia Region of our country and it is known as "zarif hezaren".

In the study, it is aimed to determine some nutrients and mineral substance contents of *Delphinium szowitsianum* Boiss which is naturally grown around the Van lake in the Eastern Anatolia Region.. In plant samples, some nutrients and mineral contents such as total ash, crude protein, pH, crude cellulose and N, Na, Mg, K, Ca, P, S, Mn, Fe, Cu, Zn, Cd, Co have been investigated. As a result of the research; it has been determined as crude protein ratio 4.60%, pH 5.99, total nitrogen content 0.74 %, crude ash content 6.31% and crude cellulose content 28.20%. In addition, while some mineral substance contents were determined as K 5.31 g/kg, Ca 16.37 g/kg, P 1.07 g/kg, Mn 35.41 mg/kg, Fe 459.69 mg/kg and Zn 32.23 mg/kg, heavy metal contents was determined as Cr 0.49 mg/kg, Cd 0.11 mg/kg, Co 1.35 mg/kg and Pb 0.44 mg/kg.

Key Words: *Delphinium szowitsianum* Boiss, Eastern Anatolia, Heavy Metal, Medicinal Plants

1.

ORAL PRESENTATION

**THE ESSENTIAL OIL YIELD AND COMPOSITIONS OF SUMMER SAVORY
(*SATUREJA HORTENSIS* L.) COLLECTED IN TURKEY**

Emine Saliha Toprak¹, Hasan Hüseyin Yiğitbaşı¹, Himmet Özcan¹, İrem Ayran¹, Sadiye Ayşe Çelik¹
Hayal Akyıldırım Beğen²

¹*Selçuk University, Agriculture Faculty, Department of Medicinal Plants, Konya/Turkey*

²*Artvin Coruh University, Health Services Vocational School, Artvin/Turkey*

e-mail: salihaemine@gmail.com, phone + 90-541 712 80 95

Objective / Purpose: In this study, it was investigated essential oil yield and compositions of Summer Savory (*Satureja hortensis*) collected from Artvin, Turkey. Common name of *Satureja hortensis* is Summer Savory.

Material and Methods: The air-dried herb parts of *Satureja hortensis* were subjected to hydrodistillation for 3 h using a Clevenger-type apparatus to produce essential oil. The GC-MS analysis was carried out with Agilent 7890 GC-MS system. The relative percentages of the separated compounds were calculated from total ion chromatograms. The identification of the oil components was based on the Wiley and NIST mass spectral library.

Results: The essential oil yield of Summer Savory was determined as 3.0 % . Thymol amount from essential oil composition of *Satureja hortensis* was found 42.7 % . Other main components were carvacrol (3.2 %), gamma-terpinene (25.0 %) and p-cymene (13.7 %).

Conclusion / Discussion: In this study, thymol was determined main compound from Summer savory essential oil. But in other studies, carvacrol was determined as the main compound. According to the results of this study, the observed differences in the constituents of summer savory essential oil collected from Artvin may be due to different climatic, soil factors and the cultivation conditions.

Keywords: Summer Savory, *Satureja hortensis*, Essential oil, Thymol.

ORAL PRESENTATION

**THE ESSENTIAL OIL YIELD AND COMPOSITIONS OF *NIGELLA SATIVA* L. AND
NIGELLA DAMASCENA L. CULTIVATED IN KONYA ECOLOGICAL
CONDITIONS**

İrem Ayran¹, S.Ayşe Çelik¹, Asuman Kan², Yüksel Kan¹

¹*Selçuk University, Agriculture Faculty, Department of Medicinal Plants, Konya/Turkey*

²*Konya Technical University, Technical Sciences Vocational School of Higher Education, Department of Food Processing, Konya/Turkey*

E-mail: irem.ayran@gmail.com, phone + 90-505 939 31 35

Objective / Purpose: In this study, it was investigated essential oil yield and compositions of seeds of *Nigella sativa* L. and *Nigella damascena* L. cultivated in Konya ecological conditions, Turkey.

Material and Methods: The air-dried seeds of black cumins species were subjected to hydrodistillation for 3h using a Clevenger-type apparatus to produce essential oil. The GC-MS analysis was carried out with Agilent 7890 GC-MS system. The relative percentages of the separated compounds were calculated from total ion chromatograms. The identification of the oil components was based on the Wiley and NIST mass spectral library.

Results: Essential oil yield of black cumin seeds was determined as 0.1% in *Nigella sativa* L. and 2.5% in *Nigella damascena* L.. The most important essential oil component of *Nigella sativa* L., timokinon content is 15.4 % and the major essential oil components of *Nigella damascena* L.. were determined as beta-Element(45.9%) . Timokinin, is the important essential oil component, was not found in *Nigella damascena* L.

Conclusion / Discussion: According to the results of this study, information about essential oil yield and components of every both species was been evaluated. The black cumin seeds have a very important place due to fixed oil, volatile oil and nutrients constant. The oil obtained from the seeds of the plant is used in the treatment of many diseases with medical purposes as well as being used as a spice and flavor in food products.

Keywords: Çörekotu, *Nigella sativa* L., *Nigella damascena* L., essential oil, composition, timokinon

ORAL PRESENTATION

**THE ROLE AND USE OF OLIVE OIL AND OLIVE LEAVES AS
A FUNCTIONAL FOOD**

Rukiye Özkaya¹, Hakan Çetinkaya², İbrahim Samet Gökçen², Mehmet Koç²

1 The Institute for Graduate Studies in Science and Engineering, Kilis 7 Aralık University, Kilis, Turkey

2 Department of Horticulture, Faculty of Agriculture, Kilis 7 Aralık University, Kilis, Turkey

*Corresponding author: hcetinkaya67@gmail.com

Objective / Purpose: The main feature of the Mediterranean diet is the pharmaceutical and nutraceutical effects of plant-derived foodstuffs and beverages include mainly fruits, vegetables, herbs and spices used in human nutrition. These phenylpropanoid-rich foods and beverages are important as pharmanutrients due to their antioxidant, anti-inflammatory, antimutagenic, cardioprotective, antihypersensitive and antitumoral activities. Olive (*Olea europaea* L.) is a characteristic plant of the Mediterranean region.

The olive plant has been an important source of nutrition and medicine for centuries. Due to the recognition of its high dietetic nutritional value, olive oil and olive leaves have been used for ancient times. The leaves, fruits and fruit oil of the olive tree contain significant amounts of bioactive and phenolic compounds. Oleuropein, hydroxytyrosol, catechin, chlorogenic acids, p-coumaric acid, caffeic acid, 3-hydroxycinnamic acid are some of the important compounds. The amount of these substances varies according to factors such as tree age, fruit load, cultivar, harvest time, biotic and abiotic stress conditions, processing, and storage status of the crop. Nutritional and health benefits of olive and olive products stand out it as a functional food.

Conclusion / Discussion: Olives, olive products and olive oil used in the Mediterranean diet are becoming increasingly important in both the nutrition and the pharmaceutical industry due to the bioactive substances it contains.

Keywords: *Olea europaea* L., Olive oil, olive leaves, polyphenol, functional food

ORAL PRESENTATION

AS A GOOD SOURCE OF MELATONIN: GRAPE AND GRAPE PRODUCTS

İbrahim Samet Gökçen^{1*}, Hakan Çetinkaya¹, Birhan Kunter², Nurhan Keskin³

¹ Department of Horticulture, Faculty of Agriculture, Kilis 7 Aralık University, Kilis, Turkey

² Department of Horticulture, Faculty of Agriculture, Ankara University, Ankara, Turkey

³ Department of Horticulture, Faculty of Agriculture, Van Yüzüncü Yıl University, Van, Turkey

*Corresponding author: ibrahimsamett@gmail.com

Abstract: Melatonin is known to be naturally found in a wide range of mammals and birds mainly responsible for the circadian rhythm of the body. It was then identified in plants. Studies look for melatonin in higher plants have been focused on the physiological roles in plants and also beneficial health effects for humans as a bioactive plant based compound. Recent studies have been indicated that melatonin in the edible plants has health-promoting effects on human health due to its anti-inflammatory, antioxidant and anticancer features as well as the therapeutic effect on cardiovascular diseases. Melatonin has been identified and quantified in different plants such as grapes, walnuts, sweet cherries, tomatoes and cucumbers. In this context grape and grape products provide a wide range of natural products as pharmanutritional effects. However, melatonin content may alterable according to cultivars, ecological conditions and some agrochemical treatments. Grape and grape products are represented to be good melatonin sources. In this review, we focus on melatonin content of grape and grape products and its effects on human health.

Keywords: *Vitis vinifera* L., circadian rhythm, foodstuffs, phytochemicals

POSTER PRESENTATIONS

MESMAP-5 Draft Proceedings Book

POSTER PRESENTATION

**THE EFFICACY OF NEW BIOAPIFIT[□] ANTI-HEMORRHOIDAL OINTMENT
VERSUS STANDARD TREATMENT PROTOCOL**

Višnja Oreščanin^{1*}, Štefica Findri Guštek², Josipa Rodić³

¹OREŠČANIN Ltd., Laboratory for herbal drugs development, A. Jakšića 30, Zagreb, Croatia,
vorescan@gmail.com

²Findri-Guštek Centre for healthcare, Ninska 5, Sesvete, Croatia, stefica.findri.gustek@gmail.com

³Family medicine clinic, Vilima Korajca 19, Zagreb, Croatia, josipa.rodic@dedominis.com

Objective / Purpose: Hemorrhoids are defined as varicosities of the veins of the hemorrhoidal plexus, often complicated by inflammation, pain, thrombosis, and bleeding. The disease correlate positively with age and more than half of the population of both sexes aged 50 years and older will develop hemorrhoid symptoms during their lifetime. The primary objective of this study was assessment of the efficacy of new multi-herbal ointment consisted of honey, *Cera flava*, glycerin, the oil macerates of *Achilea millefolium* L., *Plantago major* L., *Quercus robur* L., *Salvia officinalis* L., *Olea europaea* L., *Polygonum aviculare* L., *Calendula officinalis* L., *Matricaria chamomilla* L., essential oils of *Melaleuca alternifolia*, *Timus vulgaris* ct. Thymol and *Origanum vulgare* for the treatment of hemorrhoids of grade 1 to 3 while the second goal was the comparison of its treatment potential with standard approach applied under the same condition.

Materials and methods: The experimental group consisted of 66 participants ranging from 27 to 69 years was treated with new Bioapifit[□] anti hemorrhoidal ointment while the control group consisted of 40 participants ranging from 25 to 89 years was treated with Faktu[□] ointment (50 mg/g of policresulene and 10 mg/g of cinchocaine). Either Bioapifit[□] or Faktu[□] ointment was applied externally three times a day for ten days onto clean perianal area and rectally once a day (before bedtime) using appropriate applicator. The evaluation of the patients before and following the therapy was done in terms of pain (0-10), defecation discomfort (0-10), bleeding severity (0-4), anal itching severity (0-4) and overall subjective discomfort (0-10). For statistical evaluation Statistica 11.0 software package was employed.

Results: Before the therapy the mean values and standard deviations in the experimental group for pain, defecation discomfort, bleeding, itching and overall subjective discomfort were 6.6±0.9, 6.9±1.0, 2.2±0.7, 2.6±0.6 and 7.0±0.8, respectively. A significant decrease of all five parameters was obtained after only three days of the treatment with mean values and standard deviations of 2.2±0.8, 2.5±1.0, 0.6±0.5, 1.3±0.7 and 3.0±0.8 for pain, defecation discomfort, bleeding, itching and overall subjective discomfort, respectively. After five days of the therapy, there was no bleeding while the mean value and standard deviation were reduced to 0.5±0.6, 0.9±0.7, 0.6±0.6 and 1.4±1.1 for pain, defecation discomfort, itching and overall subjective discomfort, respectively. At 10th day of the treatment all five parameters were graded 0 by all 66 patients. The control patients treated with Faktu[□] ointment also showed a significant decrease of all five parameters after only three days of the therapy while in the end of the treatment overall subjective symptoms decreased from 3.8±1.0 to 0.9±0.9. Bioapifit[□] anti-hemorrhoidal ointment showed better results for all treatment period compared to Faktu[□] ointment.

Conclusion / Discussion: Presented results revealed that the preparation combining herbal ingredients with the honeybee's products could be used safely and effectively for the treatment of the symptoms of hemorrhoidal disease. The combination of the ingredients with soothing, calming, anti-inflammatory, coating, pH adjusting and lubricating effect with those with proven astringent, vasoconstrictor and haemostatic activity resulted in alleviation followed by complete disappearance of the symptoms like pain, bleeding, itching and defecation discomfort.

Keywords: hemorrhoidal disease, herbal preparations, honeybee's products, ointment

POSTER PRESENTATION

ANTIOXIDANT ACTIVITY AND FREE AGLYCONE COMPOUNDS IN AERIAL PARTS OF THREE SPECIES OF *PISTACIA* GROWING IN ALGERIA

N.Hamlat^{1,2} A. Hassani¹ S.Boubekeur³

¹Laboratoire de Recherche sur les Produits Bioactifs et Valorisation de la Biomasse, Ecole Normale Supérieure BP 92 Kouba, Alger, Algérie.

² Faculté de médecine Ziania, université Alger1, Alger, Algérie

³Laboratoire de chimie analytique Saidal Alger.

Pistacia a genus of flowering plants from the family anacardiaceae, contains about twenty species. In Algeria among them three are more popular including *P.Atlantica* Desf, *P.Lentiscus* L and *P.Therbinthus* L. Different parts of these species have been used in traditional medicine for various purpose like antiseptic, gastrointestinal and respiratory tract disorder. Also they have pharmacological activities such as: antioxidant and antimicrobial.

The aim of the present study was to identify an important class of phenolic compounds: flavonoïds and specially free aglycones, contained in the leaves and seeds of the quoted three species of *Pistacia* growing in Algeria, and to determine the antioxidant activity using the 1,1-diphenyl-1-picrylhydrazyl (DPPH).

The results allowed as to identify several compounds such as: quercitine myrecitine, luteoline and kaempferol. Antioxydant capacity (%) [IC₅₀ (0.0123 for leaves and 0.0121 for fruits of *Pistacia terebenthus*), (IC₅₀ (2.546 for leaves and 99.996 for fruits of *Pistacia atlantica*) and (IC₅₀ (11.748 for leaves and 52.292 for fruits of *Pistacia lentiscus*)).

According to the results it may be concluded that the extracts of the three species of *Pistacia* revealed considerable antioxidant activity.

Key words: *Pistacia*, flavonoïds, antioxidant activity, HPLC.

References:

- [1]:Z.Memariami, M.Mobli, M.H.S.Surmaghi, M.R.S.Ardekani, R.Rahimi, 2013, Scientific World Journal
- [2]:P.Lebreton; M.Jay; B.Vorin; M.P.Boucher, Analyse qualitative et quantitative des flavonoïdes, chimie analytique (1967) 49,375,385.
- [3]:J.B.Harborne, Phytochemical methods a guide to modern techniques of plant analysis, Chapman and Hall, Ed,London (1973) 277.
- [4]:A.Asghar Hatamnia, N.Abbaspour, R.Darvishzadeh, antioxidant activity and phenolic profile of different parts of Bene (*Pistacia atlantica* Subsp Kurdica) fruits, Food Chemistry (2014), volume 145,306-311.
- [5]:R.Hemma, S.Belhadj, C.Ouahchia, F.Saidi, antioxidant activity of *Pistacia lentiscus* methanolic extracts, Revue agrobiologia(2018),8 (1), 845-852.
- [6]:H.Hosseinzadeh, S.A.S.Tabassi, N.M.Moghadam, M.Rashedinia, S.Mehri, antioxidant activity of *Pistacia Vera* fruits, leaves and gum extracts, Iranian Journal of Pharmaceutical research (2012), 11 (3), 879-887.

POSTER PRESENTATION

**CHEMICAL PROFILE AND ANTIOXIDANT ACTIVITY OF NEW SELECTIVE
ZINNIA ELEGANS FRACTIONS**

Ana Flavia Burlec¹, Łukasz Pecio², Mariusz Kowalczyk², Cornelia Mircea³, Oana Cioancă⁴,
Andrei Corciovă¹, Alina Nicolescu⁵, Cătălina Stan⁶, Wiesław Oleszek², Monica Hăncianu⁴

¹Department of Drug Analysis, Faculty of Pharmacy, "Grigore T. Popa" University of Medicine and Pharmacy,
Iasi, Romania, flavia_burlec@hotmail.com

²Department of Biochemistry and Crop Quality, Institute of Soil Science and Plant Cultivation, State Research
Institute, Pulawy, Poland

³Department of Pharmaceutical Biochemistry and Clinical Laboratory, Faculty of Pharmacy, "Grigore T.
Popa" University of Medicine and Pharmacy, Iasi, Romania

⁴Department of Pharmacognosy, Faculty of Pharmacy, "Grigore T. Popa" University of Medicine and
Pharmacy, Iasi, Romania

⁵"Petru Poni" Institute of Macromolecular Chemistry, Iași, Romania

⁶Department of Pharmaceutical and Biotechnological Drug Industry, Faculty of Pharmacy, "Grigore T. Popa"
University of Medicine and Pharmacy, Iasi, Romania

Objective / Purpose: *Zinnia elegans* (syn. *Zinnia violacea*) is a very popular ornamental plant of the *Asteraceae* family, widely cultivated for the impressive range of flower colors and persistent bloom [1,2]. Given its easy growth and high adaptability to improper landscape conditions [1], we decided to investigate its potential use as a resource of valuable secondary metabolites with biochemical actions.

Material and Methods: Preliminary detection of classes of compounds found in a methanolic extract obtained from *Zinnia elegans* inflorescences was achieved using high-resolution LC-MS techniques. The extract has been further subjected to Solid Phase Extraction and separation using Sephadex LH-20 column chromatography, which allowed for different fractions to be obtained. The LH-20 fractions have been investigated regarding their antioxidant properties through lipoxygenase inhibition and metal chelating activity assays. Moreover, one representative fraction was further purified through preparative HPLC, and NMR spectroscopy analyses using a Bruker Avance III HD 500 MHz spectrometer were carried out for establishing the exact structure.

Results: The obtained results indicate that the investigated fractions contain several polyphenolic compounds such as chlorogenic acids and apigenin, kaempferol and quercetin glycosides, as well as nitrogen containing compounds. Antioxidant tests showed that some of the analyzed fractions had similar EC₅₀ values as the positive control. The NMR analysis revealed the presence of a new kaempferol glycoside.

Conclusion / Discussion: Given the promising antioxidant results, fractions obtained from *Zinnia elegans* inflorescences justify further purifications and structural analysis of compounds, while the investigation of other biological actions could prove of significant applicability.

Keywords: *Zinnia elegans*, HR-MS, lipoxygenase, metal chelation

Acknowledgement: This work was supported by a mobility grant of the Romanian Ministry of Research and Innovation, CNCS-UEFISCDI, project number PN-III-P1-1.1-MC-2017-0956, within PNCDI III.

References:

1. Bajaj, Y. (1997). *High-tech and micropropagation VI*. Berlin: Springer.
2. Odenwald, N., & Turner, J. (2006). *Identification, selection, and use of southern plants for landscape design*. Baton Rouge: Claitor's Pub. Division.

POSTER PRESENTATION

***IN VIVO* ANTI-ULCERATIVE AND ANTIDIARRHEAL ACTIVITIES OF THE
TETRACLINIS ARTICULATA SPECIES OF THE CUPRESSACEAE FAMILY**

Zine El Abidine Ababsa^{1,2}, Mohamed Tahar Derouiche³, Jean Pierre Mahy⁴,
Kamel Medjroubi², Salah Akkal²

¹ Département des Sciences de la matière, Faculté des sciences exactes et de la nature et vie, Université Larbi BEN M'HIDI, Oum El Bouaghi- Algérie (ababsapharm@gmail.com)

² Unité de Valorisation des ressources naturelles, Molécules Bioactives et Analyses Physico-Chimiques et biologiques, Faculté des Sciences Exactes, Département de Chimie, Université Mentouri de Constantine 1- Algérie

³ Faculté de médecine, département de pharmacie, Université Salah BOUBNIDER de Constantine 03. Algérie.

⁴Institut de chimie moléculaire et des matériaux d'Orsay (ICMMO), Université paris sud 11. France.

Tetraclinis articulata (Thuya of Barbary), endemic to North Africa, is used as a traditional medicine for the treatment of many diseases. We studied the antidiarrheal properties of *Tetraclinis articulata* n-butanol extract in male and female Swiss albino mice to support its traditional use. The antidiarrheal activity of the plant extract was evaluated in a model of castor oil-induced diarrhea in mice and compared to loperamide (a reference inhibitor of diarrhea). The effect of n-butanol extract on gastrointestinal motility was determined by the oral administration of charcoal and Castrol oil-induced intestinal fluid accumulation (enteropooling).

The n-butanolic extract of the species *tetraclinis articulate* showed antidiarrheal activity significantly inhibited gastrointestinal motility and castor oil induced enteropoolysis, similar to the inhibition achieved in loperamide treated mice. Ulcer is the most common gastrointestinal disturbance resulting from an inadequate gastric mucosal defense. Several drugs are available in the market to address the disease; however, these drugs are associated with unnecessary side effects.

Previous research have confirmed the efficacy of plant extracts for possible treatment of the disease. This research aims to evaluate the anti-ulcer properties of medicinal plants. N-butanol extract from the leaves of *Tetraclinis articulata* is evaluated for his anti-ulcer activity. The extract showed inhibitory activity with exhibiting more than 70% inhibition. We conclude that *Tetraclinis articulata* extracts are potential sources of new anti-ulcer agents and antidiarrheal drugs.

POSTER PRESENTATION

**IN VIVO ANTI-INFLAMMATORY EFFECTS OF SOME EXTRACTS FROM
*PELARGONIUM SP.***

Cristina Iancu¹, Cornelia Mircea^{1*}, Oana Cioanca², Catalina Stan³, Nina Filip⁴, Cristinel Stan⁵,
Gabriela Tatarina⁶, Monica Hăncianu²

¹Department of Pharmaceutical Biochemistry and Clinical Laboratory, Faculty of Pharmacy, "Grigore T. Popa" University of Medicine and Pharmacy, Iași, Romania

²Department of Pharmacognosy, Faculty of Pharmacy, "Grigore T. Popa" University of Medicine and Pharmacy, Iași, Romania

³Department of Drugs Industry and Pharmaceutical Biotechnology, Faculty of Pharmacy, "Grigore T. Popa" University of Medicine and Pharmacy, Iași, Romania

⁴Department of Biochemistry, Faculty of Medicine, "Grigore T. Popa" University of Medicine and Pharmacy, Iași, Romania

⁵Department of Anatomy, Faculty of Medicine, "Grigore T. Popa" University of Medicine and Pharmacy, Iași, Romania

⁶Department of Organic Chemistry, Faculty of Pharmacy, "Grigore T. Popa" University of Medicine and Pharmacy, Iași, Romania

*corresponding author: corneliamircea@yahoo.com

Objective: Various *in vivo* tests are used to evaluate the anti-inflammatory properties of chemical compounds [1]. The most common test is the TPA-induced oedema because it produces a strong reaction in the dermic inflammation. TPA activates the protein kinase-C, but also the nuclear factor KB and the inflammatory mediators such as TNF- α , the TL-1 β and the MIP2 macrophages [2]. In our study we evaluated the anti-inflammatory effects of some methanolic extracts obtained from *Pelargonium hispidum*, *Pelargonium radens*, *Pelargonium peltatum*, and *Pelargonium zonale*.

Material and Methods: The anti-oedematous/anti-inflammatory effect of the investigated samples was demonstrated on a TPA-induced animal model, when, after the treatment, there was a reduction of the typical inflammatory parameters (ear base size, length and mass) [3].

Results: Generally, the values obtained were determined after a single topical application of TPA and are consistent with the literature data [4]. The maximum expression of oedema was observed 24 hours after the administration of TPA with important differences for the groups treated with the investigated *Pelargonium* extracts. For the same dose (5 mg/mL) applied topically, the most intense oedema reduction effect was present in the group 4 treated with methanol extract of *P. zonale*, in which inflammation compared to the negative control (TPA group) decreased by about 70%, and, compared to the positive control (treated with indomethacin), decreased by about 30%. The other groups treated with the *P. radens* extract (group 3) and *P. grandiflorum* extract (group 2) showed a reduction of the inflammatory process by about 50% compared to the negative control. *P. hispidum* methanol extract (group 1) showed an anti-edematous effect of 45% compared to the negative control.

Conclusion / Discussion: Inflammation is a pathological process whose evolution involves the presence of enzymes, proinflammatory chemical mediators, extravasation of fluids, cell migration, tissue damage, but also tissue recovery processes. Thus, our data indicated that the methanol extract of *Pelargonium zonale* has a promising potential in reducing the inflammatory processes.

Keywords: anti-inflammatory effect, *Pelargonium species*, TPA-induced assay, methanol extract.

References:

- [1] Hvid, H., Teige, I., Kvist, P.H., Svensson, L., Kemp, K. (2008). TPA induction leads to a Th17-like response in transgenic K14/VEGF mice: a novel *in vivo* screening model of psoriasis. *Int Immunol*, 20, 1097-1106.
- [2] Williams, L.A.D., Connar, A.O., Latore, D.O., Ringer, S., Whittaker, J.A., Conard, J., Vogler, B., Rosner, H., Kraus W. (2008). The *in vitro* anti-denaturation effects induced by natural products and non-steroidal compounds in heat treated (immunogenic) bovine serum albumin is proposed as a screening assay for the detection of anti-inflammatory compounds, without the use of animals, in the early stages of the drug discovery process. *West Indian Med J*, 57, 327-331.
- [3] Passos, G.F., Medeiros, R., Marcon, R., Nascimento, A.F.Z., Calixto, J.B., Pianowski, L.F. (2013). The role of PKC/ERK1/2 signaling in the anti-inflammatory effect of tetracyclic triterpene euphol on TPA-induced skin inflammation in mice. *Euro J of Pharmacol*, 698, 413-420.
- [4] Dehelean, C.A., Șoica, C., Ledeți, I., Aluș, M., Zupko, I., Gălușcan, A., Cinta-Pinzaru, S., Munteanu, M. (2012). Study of the betulin enriched birch bark extracts effects on human carcinoma cells and ear inflammation. *Chem Central J*, 6, 137-145.

POSTER PRESENTATION

NEW GYPSOGENIN-AMINE COMPOUNDS

Ece Sözer¹, Safiye Emirdağ-Öztürk¹, N. Gökçe Ulusoy¹, Nurettin Yaylı², Halil Ciftci^{3,4}, Mohamed O. Radwan^{3,4,5}, Mikako Fujita⁶, Masami Otsuka⁴

¹ Department of Chemistry, Faculty of Science, Ege University, Bornova, Izmir, Turkey

² Faculty of Pharmacy, Karadeniz Technical University, Trabzon, Turkey

³ Department of Drug Discovery, Science Farm Ltd., Kumamoto, Japan

⁴ Department of Bioorg. Medicinal Chemistry, School of Pharmacy, Kumamoto University, Kumamoto, Japan

⁵ Chemistry of Natural Compounds Department, National Research Centre, Cairo, Egypt

⁶ Research Institute for Drug Discovery, School of Pharmacy, Kumamoto University, Kumamoto, Japan

Background / aims: Gypsogenin compound, a natural saponin of pentacyclic triterpene, belongs to the carnations family (Caryophyllaceae). The Gypsogenin compound is found in Gypsophila plant along with sugar chains that contribute to various biological properties such as antiviral, antitumor, antioxidant and anti-cancer. We have targeted some of the semi-synthesis reactions in order to make the higher activity more active.

Material and Method: We synthesized new gypsogenin derivatives (**1-3**). The progress of the reactions was followed by sodium triacetoxy borohydride in DCE. Purification was carried out using chromatographic methods.


Results: Up to now, in our continuous research, the elucidation of the synthesized compounds (**1-3**) was determined by IR, UV, ¹H NMR, APT and LCMS analysis.

Conclusion / Discussion: In the last part of the study, biological activities of new compounds will be investigated.

Keywords: Gypsogenin, semi synthesis

References:

- [1] Emirdag-Ozturk S.; Babahan I.; Ozmen A., (2014). Synthesis, characterization and in vitro anti-neoplastic activity of gypsogenin derivatives, *Bioorganic Chemistry*, vol.53, 15-23.
[2] Karaca E.Ö. (2018). Synthesis of new Schiff base compounds and identification of their structures, *Journal of Polytechnic*, 21(1), 245-249.



This project is supported by TUBITAK (117R034).

POSTER PRESENTATION

CHEMICAL CONSTITUENTS OF *ARTEMISIA ALBICERATA*

A. Amantay¹, M. Dyusebaeva¹, J. Jenis^{1, 2*}

¹ Faculty of Chemistry and Chemical Technology, al-Farabi Kazakh National University, Almaty, Kazakhstan
E-mail: alzhan.zh@mail.ru

² Shanghai Institute of Materia Medica, Chinese Academy of Science, Shanghai, China

Objective / Purpose: *Artemisia albicerata* is of great interest and occupy an important place among the vast variety of medicinal plants in Kazakhstan due to its medical and pharmacological properties and has a rich chemical composition consisting of amino acids, fatty acids, flavonoids and other low-molecular substances. The aim of the present research was the investigation of chemical composition and developing a technology for producing a biologically active complex based on *A. albicerata*.

Material and Methods: The aerial part of *A. albicerata* utilized in the present survey was collected in Almaty region, Kazakhstan at the end of August in 2018. The elemental constituent in the ash was investigated by applying the method of multi-element atomic emission spectral analysis. Gas-liquid chromatography was used for determination of amino acids and fatty acids. Conditions for obtaining a biologically active complex from *A. albicerata* were: extractant - 70% ethyl alcohol, the ratio of raw material and extractant - 1:8, time of triple extraction - 3 days, temperature - 10 ° C.

Results: Moisture content (6.62 %), extractives (12.71 %) and total ash (7.96 %) were identified. Eleven macro-micro elements were obtained from the ash of plant, main of them were Ca (485.110 µg/ml), K (481.660 µg/ml), Mg (97.950 µg/ml), Na (46.820 µg/ml), Fe (30.2939 µg/ml). In addition, twenty amino and eight fatty acids were identified from *A. albicerata*. The major amino acid contents were glutamate (2615 mg/100g), aspartate (1296 mg/100g) and alanine (890 mg/100g), while the composition of fatty acids mainly was oleic (19.6 %) and linoleic (68.4 %) acids.

Conclusion / Discussion: The results demonstrated constituent of important macro-micro elements, amino and fatty acids, main classes of biologically active substances in the aerial part of *A. albicerata*. Consequently, the results indicate a more thorough study is needed.

Key words: *Artemisia albicerata*, macro-micro elements, amino-, fatty acids.

References:

[1] Alukina L.S., Ryakhovskaya T.V. - 16 (2). - Alma-Ata: Academy of Sciences of the Kazakh SSR, 1980. - 187-192 p.

POSTER PRESENTATION

**LIPOSOLUBLE CONSTITUENTS FROM THE AERIAL PART OF
*LIGULARIA NARYNENSIS***

Nurlybekova A.K.¹, Amantay A.¹, Ye Yang³, Abilov Zh.A.¹, Jenis J.^{1,2,3*}

¹Faculty of Chemistry and Chemical Technology, Al-Farabi Kazakh National University, Almaty, Kazakhstan

²Research Center for Medicinal Plants of Al-Farabi Kazakh National University, Almaty, Kazakhstan

³Shanghai Institute of Materia Medica, Chinese Academy of Sciences, Shanghai, China

*E-mail: janarjenis@mail.ru

Objective: *Ligularia* is a medicinally important genus of the family Compositae that comprises about 180 Eurasian species, 17 species growing in mountains of Kazakhstan [1]. Phytochemical investigations of various *Ligularia* species showed the presence of monoterpenes, sesquiterpenes, diterpenes, triterpenes, alkaloids, steroids, flavonoids, lignans, and other skeleton type compounds, as well as some of these compounds showed various biological activities such as anticancer, antibacterial, antihepatotoxicity, antioxidant, and antithrombus [2]. In this work, fifty liposoluble constituents in chloroform extract from the aerial part of *Ligularia narynensis* from Kazakhstan have been identified by GC-MS method for the first time.

Material and Methods: To determine the liposoluble constituents' composition was made erenow of the raw material used GC/MS device. The chloroform extract from the aerial part of *L. narynensis* was analyzed by Electron Impact Ionization (EI) method on Agilent 7890A-5975C GC-MS (Gas Chromatograph coupled to Mass Spectrometer) fused silica capillary column (30m x 0.25mm; 0.25 µm film thickness), coated with HP-5MS were utilized. The carrier gas was helium (99.999 %). The column temperature was programmed from 50°C (held for 10 min), with 10°C/min rate program to increase temperature to 300°C. The latter temperature maintained for 40 min (Acquisition parameters full scan; scan range 30-1000 amu). The injector temperature was 310°C. Injection: with a 1 µl. Detector ion source (EI-70eV). Samples were injected by splitting with the split ratio 5:1.

Results: The liposoluble constituent in aerial part of *L. narynensis* was analyzed by GC-MS method. Total fifty compounds were separated and their relative contents were determined by area normalization in which the major constituents were Olean-12-ene, 3-methoxy-, (3.beta.)-(18.24 %), 12-Oleanen-3-yl acetate, (3.alpha.)-(9.70 %), Linoleic acid ethyl ester (8.62 %), Lupeol (6.29 %), and A'-Neogammacer-22(29)-en-3-ol, acetate, (3.beta.,21.beta.)-(5.71 %).

Conclusion: Presence of these bioactive constituents indicated that the plant extract possesses anti-inflammatory, anticoronary and anticancer activities. Further and comprehensive investigation is scheduled to be implemented in the next research stage.

Key words: *Ligularia narynensis*, chloroform extract, liposoluble constituents, GC-MS.

Acknowledgement: The work was supported by grants from Ministry of Education and Science of the Republic of Kazakhstan (0118PK00458).

References:

- [1] Baitenov, M.S. (2001). Flora of Kazakhstan [Flora Kazahstana] // Gylym, Kazakhstan. P.206-207.
[2] Wang, Y.M., Zhao, J.Q., Yang, J.L., Tao, Y.D., Mei, L.J., Shi, Y.P. (2017). *Biochemical Systematics and Ecology* 72, 8-11. <https://doi.org/10.1016/j.bse.2017.03.007>

POSTER PRESENTATION

ANTIBACTERIAL ACTIVITY OF CONIFERS

Izabela Korona-Glowniak^{1,2}, Diana Ivanova^{2a}, Anna Malm¹, Alexander Tashev³, Paraskev Nedialkov⁴, Galina Kalotova², George Angelov²

¹ Department of Pharmaceutical Microbiology, Medical University of Lublin, 20-093 Lublin, Poland, anna.malm@umlub.pl

² Institute of Chemical Engineering, Bulgarian Academy of Sciences, 1113 Sofia, Bulgaria

³ Department of Dendrology, University of Forestry, 1756 Sofia, Bulgaria, altashev@abv.bg, altashev@mail.ru

⁴ Faculty of Pharmacy, Medical University, 1000 Sofia, Bulgaria

Objective / Purpose: The discovery of new antimicrobial agents is necessary in view of the appearance of multidrug resistant bacterial strains, allergies and other side effects of conventional antibiotic treatment, rendering difficult the therapy of many infectious diseases. This study is aimed at identification of conifers with efficient antibacterial activities against Gram-positive (GP) and Gram-negative (GN) bacteria.

Material and Methods: *Juniperus sabina* var. *balkanensis* was collected from the Rhodope Mountains.³ *Sequoiadendron giganteum* was from the Botanic garden area of the Bulgarian Academy of Sciences. *Chamaecyparis pisifera*, *Chamaecyparis lawsoniana*, *Cupressus arizonica* and *Taxus baccata* were from the Arboretum of the University of Forestry, Sofia. The extracts, obtained as it was described,⁴ were screened for antimicrobial activity towards reference strains (ATCC). Minimal inhibitory concentrations (MIC) were tested by two-fold microdilution method at concentrations ranging from 0.039 to 10 mg/ml.

Results: Antibacterial activity of various conifer extracts was studied on harmful pathogens or commensal microorganisms that can become pathogenic in immunocompromised patients. In general, GP bacteria were more sensitive than GN strains to the tested extracts that may result from different structure of their cell wall. In the group of studied conifer extracts, the best activity was determined for the leaves extracts of *Juniperus sabina* var. *balkanensis* R.P. Adams & A.N. Tashev against GP bacteria: *Micrococcus luteus*, *Bacillus subtilis* and *Bacillus cereus* (MIC=0.078 mg/ml); *Staphylococcus epidermidis* (MIC=0.156 mg/ml); *Staphylococcus aureus*, *Streptococcus pyogenes* and *Streptococcus pneumoniae* (MIC=0.313 mg/ml); *Enterococcus faecalis* and *Streptococcus mutans* (MIC=0.625 mg/ml). *J. sabina* var. *balkanensis* showed growth-inhibitory effects also against some GN bacteria, such as *Klebsiella pneumoniae* (MIC=2.5 mg/ml) etc. *Chamaecyparis pisifera* (Siebold & Zucc.) Endl., *Sequoiadendron giganteum* (Lindl.) J. Buchh., *Cupressus arizonica* Greene also exhibited efficient activities against the studied GP bacteria, compared to less active *Chamaecyparis lawsoniana* (A. Murray) Parl. and *Taxus baccata* L. leaves extracts.

Conclusion / Discussion: In the group of studied conifers, the leaves extract of *Juniperus sabina* var. *balkanensis* possessed superior antibacterial activity against a panel of GP and GN bacteria. *J. sabina* varieties are easy for cultivation and therefore they are feasible for industrial exploitation. Comparatively good activities against the studied GP bacteria exhibited also the leaves extracts of *Chamaecyparis pisifera*, *Sequoiadendron giganteum*, *Cupressus arizonica*, whereas *Chamaecyparis lawsoniana* and *Taxus baccata* leaves extracts showed less activity. Identification of antibacterial conifer metabolites is in progress.

Keywords: antimicrobial activity, Coniferae.

Acknowledgements: The work is performed within the Bulgarian Science Fund contract DN 07/25 (2016); 'Erasmus plus' contract (2017-2020) between the Medical University of Lublin and the Bulgarian Academy of Sciences; project "Development of green eco-technologies" of the Bulgarian Academy of Sciences.

² Both are considered first authors.

³ Adams, R.P., Schwarzbach, A.E., Tashev, A.N. (2016). *Phytologia*, 98(2), 100-111.

⁴ Ivanova, D.I., Tashev, A.N., Nedialkov, P.N. et al. (2018). *Bulg. Chem. Commun.* 50(C), 144-150

POSTER PRESENTATION

MEDICINAL PLANTS OF BULGARIA

Alexander TASHEV & Violeta DIMITROVA

University of Forestry, Faculty of Forestry, Department of Dendrology, 10 Kliment Ohridski Blvd., 1756 Sofia, Bulgaria. E-mail: vilydi@abv.bg; altashev@abv.bg

Objective/Purpose: Bulgaria is a small country in Southeast Europe, but its geographic location, varied terrain and climate contribute to great floristic diversity. The medicinal plants are part of it. They represent 23.2% of the country's flora. The aim of the present paper is to present information about the full composition of the medicinal plants of Bulgarian flora as well as characteristics of the systematical structure and classification according to biological type, life form, floristic geoelements, period of flowering, and ecological groups.

Materials and methods: The complete list of medicinal plants in Bulgaria was compiled on the basis of the Medicinal Plants Act [1], extensive literature analysis and own researches [2, 3]. The life forms were identified according to Raunkiaer [4] and for floral elements the classification of Walter [5] was used. The ecological forms were identified according to Flora of Republic Bulgaria, vol. I-XI.

Results: As a result of the conducted surveys it was found that there are 950 medicinal species of vascular plants in the country belonging to 468 genera and 126 families. The richest family containing medicinal plants was Rosaceae (98 specie). The most medicinal plants existed among perennial herbaceous plants (539 species). Geoelements with an European component were predominant (408 species). The most plants bloom from June to August (121 species). The ecological structure of medicinal plants shows that mesophytes (435 species); heliophytes (545 species) and thermophilous plants (607 species) were dominated. The results of the study include also a detailed information about the national and international conservation importance of the analyzed species. 65 species were included in the Red Book of Bulgaria. There were 81 species protected by the Biodiversity Law. Among the medicinal plants are 16 Balkan and 4 Bulgarian endemic species. The international significance of these plants was defined by the fact that 33 of the species were under protection by different international conventions.

Conclusion/Discussion: The ratio between the biological types, the predominant geoelements and the life forms shows that it is typical for the temperate continental climate zone and the flora of our country. The total percentage of protected species (7%) is not high, which is positive because most species are subject to use by the local population, the pharmaceutical and perfumery industry.

Key words: medicinal plants, flora, systematic structure, geoelements, biological types, conservation significance, Bulgaria

References:

- [1] Medicinal Plants Act. State Gazette (SG), issue 29/7.04.2000, amended SG No. 96/2017. (in Bulgarian).
- [2] Tashev, A. 2017. Medicinal Plants of the Bulgarian Flora. In: Tichonchuk P., Senchik A., Subbekin B., Timchenko N. (eds.) Protection of the ecological environment and rational use of small-fruits resources: *Proceedings of the 9th China-Russia Scientific Forum of Forest-Ecological Construction with International Participation*, Heihe City, China. pp. 387-392 (in Russian).
- [3] Tashev A., Tsavkov E. 2008. Medicinal plants of the Bulgarian dendroflora. – *Phytologia Balcanica*, 14(2): 269-278.
- [4] Raunkiaer, S. 1934. The Life Form of Plants and Statistical Plant Geography. Clarendon Press, Oxford. 632 p.
- [5] Assyov B., Petrova A. 2012. Conspectus of the Bulgarian vascular flora. *Horology and floristic elements*. Bulgarians Biodiversity Foundation, 492 p. (in Bulgarian).

POSTER PRESENTATION

POLYPHENOLS FROM *ATHAMANTA SICULA* L.

Kabouche A¹, Bouratoua A¹, Kabouche Z¹, Al Abdulmagid A²,
Voutquenne-Nazabadioko L², Morjani H³

¹Université des frères Mentouri-Constantine, Département de chimie, Laboratoire d'Obtention des Substances Thérapeutiques (LOST), Campus Chaabet-Ersas, 25000 Constantine, Algeria

²ICMR-UMR CNRS 7312, Groupe Isolement et Structure, Campus Sciences, Bât. 18, BP 1039, 51687 Reims, France

³MEDyC UMR CNRS 7369, URCA, Faculté de Pharmacie, SFR CAP Santé, 1, rue du Maréchal-Juin, 51096 Reims, France

Introduction: The genus *Athamanta* (Apiaceae) consists of about nine species found in the South Eastern Europe and North Africa. Previous phytochemical investigation on this genus reported the isolation of the flavonoids and propenphenols. In the present work, we described the isolation of polyphenols from the dichloromethane, ethylacetate and *n*-butanol fractions of *A. sicula* L. The present work aims to determine the polyphenols of *Athamanta sicula*, collected in North-eastern Algeria

Material and methods: Air dried and powdered aerial parts (2.5 kg) of *A. sicula* were macerated with 70% MeOH then concentrated under reduced pressure. The residue was dissolved in water and sequentially partitioned with light petroleum, CH₂Cl₂, EtOAc and *n*-Butanol. The CH₂Cl₂ fraction (6g) was subjected to silica gel column chromatography eluting with gradient of CHCl₃-Toluene (0-100% CHCl₃) and CHCl₃- EtOAc (0-100% EtOAc). Subfractions underwent further fractionations led to a phytosterol and three propenphenols (**1-4**). The, EtOAc fraction was submitted to silica gel CC using a step gradient solvent system *n*-hexane-acetone (0-100% Acetone) to give seven fractions. The fraction 3 was also subjected to silica gel CC but eluted with an isocratic system EtOAc:MeOH:H₂O (10:1:0.5) to yield three flavonoids (**5-7**). The *n*-butanol fraction (15g) was fractionated by polyamide SC6 column chromatography followed by TLC purification leading to the isolation of one flavone glycoside (**8**). Compounds **3-5** were tested for their cytotoxicity activity against fibrosarcoma HT 1080 cells by means of the MTT method

Results. Eight known compounds were isolated *Athamanta sicula*, their structure elucidation was based on analyses of spectroscopic data 1D-2D NMR and HRESIMS techniques and by comparing data with those reported in literature . They exhibited a moderate cytotoxicity.

Discussion. Stigmasterol (**1**), anthricinol (**2**), myristicin (**3**), apiol (**4**), apigenin (**5**), luteolin (**6**), diosmetin (**7**), and luteolin-7-O-β-D-glucoside (**8**). Compounds **2-4** are characteristic of Apiaceae family, anthricinol (**2**) is new in *Athamanta* genus. Only myristicin exhibited a moderate cytotoxic activity with IC₅₀ = 45 μM. As a conclusion: *Athamanta sicula* is a good source of polyphenols

Keywords: *Athamanta sicula* ; flavonoids ; propenphenyls ; cytotoxicity.

POSTER PRESENTATION

ANTIBACTERIAL ACTIVITY OF *PEGANUM HARMALA* SEED EXTRACTS

Elif Güzel^{1*}, Hüseyin Servi², Cansu Vatanserver³

^{1*}*School of Pharmacy, Altınbaş University, Istanbul, Turkey, elifguzel96@hotmail.com*

²*Department of Pharmaceutical Botany, Faculty of Pharmacy, Altınbaş University, Istanbul, Turkey*

³*Department of Pharmaceutical Microbiology, Faculty of Pharmacy, Altınbaş University, Istanbul, Turkey*

Objective / Purpose: *Peganum harmala* L. is a member of Nitrariaceae family which has traditional usages as anthelmintic, lactagogue, antispasmodic, antipyretic effects and stimulation of the motor tracts of the cerebrum in folk medicine [1,2]. Previous studies illustrate that methanolic extracts of *Peganum harmala* seed had good antibacterial activity against *S. aureus*, *P. aeruginosa*, *K. pneumoniae*, *P. vulgaris*, and *E. coli*, but it was more susceptible in *S. aureus* [2]. By contrast, *n*-hexane extract from the seed of *P. harmala* showed no antimicrobial activity against both Gram (+) and Gram (-) bacteria [3]. Furthermore, depending on the former experiments, methanolic extracts of the seeds also demonstrated antibacterial activity against *B. cereus* bacteria [4]. In addition to all these, methanolic extracts also indicated antimicrobial activity against *Branhamella catarrhalis* [5]. The objective of this study is to determine the antibacterial activity of *Peganum harmala* seed extracts.

Material and Methods: The hexane, ethyl acetate and methanol extracts from the seed of *Peganum harmala* were obtained by maceration method. Antibacterial activity of seed extracts was observed against *S. aureus* ATCC 29213, *B. cereus* ATCC 14579, *P. aeruginosa* ATCC 27853 and *E. coli* ATCC 25922 by using a broth microdilution and disc diffusion assays. The experiments were started 8 mg/mL for broth microdilution assay and 10 mg/disc extracts were used for disc diffusion assay. Chloramphenicol was used as a positive control for both assays. All the experiments were performed in duplicate.

Results: Hexane and ethyl acetate extracts did not show activity against any of the tested bacteria. However, methanol extract showed inhibitory activity against the tested microorganisms. The highest inhibitory activity was observed against *B. cereus*, the MIC value was 1 mg/mL and the disc diffusion results were 24 mm at 10 mg/disc.

Conclusion / Discussion: The methanol extract of seed showed antibacterial activity against Gram (+) and Gram (-) bacteria. But the extract was more susceptible to *B. cereus* than other tested bacteria.

Keywords: *Peganum harmala*, extract, antibacterial activity

References:

- [1] Hassan, I. (1967). Some folk uses of *Peganum harmala* in India and Pakistan. *Economic Botany*, 21(3), 284-284.
- [2] Prashanth, D., & John, S. (1999). Antibacterial activity of *Peganum harmala*. *Fitoterapia*, 70(4), 438-439.
- [3] Shahverdi, A. R., Monsef-Esfahani, H. R., Nickavar, B., Bitarafan, L., Khodae, S., & Khoshakhlagh, N. (2005). Antimicrobial activity and main chemical composition of two smoke condensates from *Peganum harmala* seeds. *Zeitschrift für Naturforschung C*, 60(9-10), 707-710.
- [4] Darabpour, E., Bavi, A. P., Motamedi, H., & Nejad, S. M. S. (2011). Antibacterial activity of different parts of *Peganum harmala* L. growing in Iran against multi-drug resistant bacteria. *EXCLI journal*, 10, 252.
- [5] Sokmen, A., Jones, B. M., & Erturk, M. (1999). The in vitro antibacterial activity of Turkish medicinal plants. *Journal of ethnopharmacology*, 67(1), 79-86.

POSTER PRESENTATION

**DETERMINATION OF ANTIBACTERIAL ACTIVITY FROM SEED EXTRACTS
OF *PISTACIA TEREBINTHUS***

Elif Güzel^{1*}, Cansu Vatansever², Hüseyin Servi³,

^{1*}School of Pharmacy, Altınbaş University, Istanbul, Turkey, elifguzel96@hotmail.com

²Department of Pharmaceutical Microbiology, Faculty of Pharmacy, Altınbaş University, Istanbul, Turkey

³Department of Pharmaceutical Botany, Faculty of Pharmacy, Altınbaş University, Istanbul, Turkey

Objective / Purpose: *Pistacia terebinthus* L. is a member of Anacardiaceae family that has been used as a traditional natural remedy for different purposes all around the world. The plant has been used as an aphrodisiac and expectorant in Greece, as a diuretic and laxative in Jordan, as a hypotensive and antiseptic in Spain, antidiarrhea and astringent properties in Iran, and finally stomach ache, asthma and as an antidiabetic in Turkey [1]. Previous studies demonstrate that ethanolic extracts of *Pistacia terebinthus* showed antimicrobial activity against *Staphylococcus aureus*, *Proteus vulgaris*, *Bacillus cereus*, and *Mycobacterium smegmatis* depending on the disc diffusion tests [2]. Additionally, ethanolic extracts of the plant also showed antimicrobial activity against *S. aureus* in microdilution experiments, whereas there is no indication of antimicrobial property on *E. coli* [3]. The objective of this study is to determine the antibacterial activity of *Pistacia terebinthus* seed extracts.

Material and Methods: The plant material (350 g) was macerated with hexane, ethyl acetate, and methanol, respectively. Antibacterial activity of seed extracts was observed against *S. aureus* ATCC 29213, *B. cereus* ATCC 14579, *P. aeruginosa* ATCC 27853 and *E. coli* ATCC 25922 by using a broth microdilution and disc diffusion assays. The experiments were started 8 mg/mL for broth microdilution assay and 10 mg/disc extracts were used for disc diffusion assay. Chloramphenicol was used as a positive control for both assays. All the experiments were performed in duplicate.

Results: The inhibitory activity was observed against the tested microorganisms for methanol extract. The methanol extract showed the highest inhibitory activity against *S. aureus* and *B. cereus*. The inhibition zone diameters were 14 mm at 10 mg/disc for *S. aureus* and *B. cereus*. The minimum inhibition concentration (MIC) values were 0.5 mg/mL for *S. aureus* and 1 mg/mL for *B. cereus*. Ethyl acetate extract showed inhibitory activity against *E. coli* and *P. aeruginosa*. The MIC values were 8 mg/mL and the inhibition zone diameters were 8 mm at 10 mg/disc for both bacteria. However, the same extract did not show activity against *S. aureus* and *B. cereus*. Hexane extract showed inhibitory activity against *P. aeruginosa*, the MIC value was 8 mg/mL and the inhibition zone diameter was 9 mm at 10 mg/disc.

Conclusion / Discussion: *Pistacia terebinthus* seed methanol extract has antibacterial activity against both Gram-positive and Gram-negative bacteria. The methanol extract was susceptible against *S. aureus* and *B. cereus*. The ethyl acetate extract has inhibitory activity against Gram-negative bacteria. All the seed extracts were susceptible against *P. aeruginosa*.

Keywords: *Pistacia terebinthus*, extract, antibacterial activity

References:

- [1] Bozorgi, M., Memariani, Z., Mobli, M., Salehi Surmaghi, M. H., Shams-Ardekani, M. R., & Rahimi, R. (2013). Five *Pistacia* species (*P. vera*, *P. atlantica*, *P. terebinthus*, *P. khinjuk*, and *P. lentiscus*): a review of their traditional uses, phytochemistry, and pharmacology. *The Scientific World Journal*, 2013.
- [2] Dulger, B., & Gonuz, A. (2004). Antimicrobial activity of certain plants used in Turkish traditional medicine. *Asian J. Plant Sci*, 3(1), 104-107.
- [3] Kavak, D. D., Altok, E., Bayraktar, O., & Ülkü, S. (2010). *Pistacia terebinthus* extract: As a potential antioxidant, antimicrobial and possible β -glucuronidase inhibitor. *Journal of Molecular Catalysis B: Enzymatic*, 64(3-4), 167-171.
- [4] Hacibekiroğlu, I. I., Yılmaz, P. K., Haimi, N., Kılınç, E., Tolun, V., & Kolak, U. (2015). In vitro biological activities and fatty acid profiles of *Pistacia terebinthus* fruits and *Pistacia khinjuk* seeds. *Natural product research*, 29(5), 444-446.

POSTER PRESENTATION

ANTIBACTERIAL ACTIVITY OF *SAMBUCUS NIGRA* SEED EXTRACTS

İrem Şengür¹, Cansu Vatansever², Hüseyin Servi³

¹Faculty of Pharmacy, Altınbas University, Istanbul, Turkey

²Department of Pharmaceutical Microbiology, Faculty of Pharmacy, Altınbas University, Istanbul, Turkey

³Department of Pharmaceutical Botany, Faculty of Pharmacy, Altınbas University, Istanbul, Turkey

Objective / Purpose: *Sambucus nigra* L., is 3-4 meters in length, a small tree with unified leaves and fragrant [1]. It belongs to the Adoxaceae family. The plant is located in the Northern Turkey and Iraq, Western Syria and Iran. Infusion or decoction from flowers, leaves and stem barks of *Sambucus nigra* used as diuretic, diaphoretic and laxative effects [2]. Previously, the antimicrobial activity of ethanol extract from *Sambucus nigra* was studied. The ethanol extract showed inhibitory activity against *Bacillus cereus*, *Escherichia coli* ve *Pseudomonas aeruginosa* [3]. The objective of this study is to determine the antibacterial activity of *Sambucus nigra* L. seed extracts against *Staphylococcus aureus* ATCC 29213, *B. cereus* ATCC 14579, *P. aeruginosa* ATCC 27853 ve *E. coli* ATCC 25922 by using a broth microdilution and disc diffusion assays.

Material and Methods: The hexane, ethyl acetate and methanol extracts from the seed of *Sambucus nigra* L. were obtained by maceration method. The experiments were started 8 mg/mL for broth microdilution assay and 10 mg/disc extracts were used for disc diffusion assay. Chloramphenicol was used as a positive control for both assays. All the experiments were performed in duplicate.

Results: Methanol extract showed inhibitory activity against tested microorganisms. The minimum inhibition concentration (MIC) value for methanol was 8 mg/mL and the disc diffusion result was 12 mm for all bacteria. Ethyl acetate extract showed inhibitory activity against *B. cereus*. The MIC values of ethyl acetate extract were 8 mg/mL for *B. cereus* and the disc diffusion result was 8 mm. Hexane extract did not show inhibitory activity against any of the tested bacteria.

Conclusion / Discussion: The methanol extract of seed showed antibacterial activity against Gram-positive and Gram-negative bacteria. Hexane extract is not susceptible to any bacteria. Ethyl acetate extract is not susceptible to tested bacteria except that *B. cereus*.

Keywords: *Sambucus nigra* L., extract, antibacterial activity

References:

- [1] Özmen, M . (2010). Hatay-Erzin Başlamış Köyünden Derlenen Bitki Adları. Erdem, (58), 255-268. Retrieved from <http://dergipark.gov.tr/erdem/issue/43857/539427>
- [2] Dündar, S., (2010). “*Sambucus* L. türleri üzerinde fitoterapötik çalışmalar.” Gazi Üniversitesi Sağlık Bilimleri Enstitüsü, Farmakognozi-Fitoterapi Yüksek Lisans Programı.
- [3] Hearst, C., McCollum, G., Nelson, D., Ballard, L. M., Millar, B. C., Goldsmith, C. E., & Rao, J. R. (2010). Antibacterial activity of elder (*Sambucus nigra* L.) flower or berry against hospital pathogens. *Journal of Medicinal Plants Research*, 4(17), 1805-1809.

POSTER PRESENTATION

DETERMINATION OF ANTIBACTERIAL ACTIVITY OF *LINUM USITATISSIMUM* SEED EXTRACTS

İrem Şengür¹, Hüseyin Servi², Cansu Vatansever³

¹Faculty of Pharmacy, Altınbas University, Istanbul, Turkey

²Department of Pharmaceutical Botany, Faculty of Pharmacy, Altınbas University, Istanbul, Turkey

³Department of Pharmaceutical Microbiology, Faculty of Pharmacy, Altınbas University, Istanbul, Turkey

Objective / Purpose : Flaxseed or linseed (*Linum usitatissimum* L.) is an annual herb belongs to the Linaceae family. Consumption of flaxseed has been demonstrated to have a multitude of positive health benefits including decreasing rate of tumor growth, reducing serum cholesterol level and decreasing the incidence of breast, prostate, and colon cancers [1]. Previously, the antimicrobial activity of *n*-hexane, ethyl acetate, butanol and water extracts from *L. usitatissimum* was studied by disc diffusion assay. The ethyl acetate and butanol extracts showed inhibitory activity against *S. aureus*, *B. cereus* and *E. carotovora* [2]. The objective of this study is to determine the antibacterial activity of *Linum usitatissimum* L. seed extracts against *S. aureus* ATCC 29213, *B. cereus* ATCC 14579, *P. aeruginosa* ATCC 27853 and *E. coli* ATCC 25922 by using a broth microdilution and disc diffusion assays.

Material and Methods: The hexane, ethyl acetate and methanol extracts from the seed of *Linum usitatissimum* L. were obtained by maceration method. The experiments were started 10 mg/mL for disc diffusion assay and 8 mg/mL for broth microdilution assay. Chloramphenicol was used as a positive control for both assays. All the experiments were performed in duplicate.

Results: Hexane extract showed inhibitory activity against *E. coli*, *P. aeruginosa* and *S. aureus*. The MIC value was 8 mg/mL and the disc diffusion result was 10 mm at 10 mg/mL. Ethyl acetate extract showed inhibitory activity against *S. aureus* and *P. aeruginosa*. The MIC values of ethyl extract were 0.5 mg/mL for *S. aureus* and 8 mg/mL for *P. aeruginosa*. The diffusion results were 8 mm for *P. aeruginosa* and 12 mm for *S. aureus* at 10 mg/mL. The methanol extract showed inhibitory activity against *S. aureus*. The MIC value was 8 mg/mL and the disc diffusion result was 8 mm at 10 mg/mL.

Conclusion / Discussion: Seed extracts of the plant did not show activity against *B. cereus*. These extracts are not susceptible to *B. cereus*. But the extracts showed activity against *S. aureus* at different concentrations. Ethyl acetate extract was more susceptible against *S. aureus* than other extracts.

Keywords: *Linum usitatissimum* L., extract, antibacterial activity

References:

- [1] Anwar, F., & Przybylski, R. (2012). Effect of solvents extraction on total phenolics and antioxidant activity of extracts from flaxseed (*Linum usitatissimum* L.). *Acta scientiarum polonorum. Technologia alimentaria*, 11(3), 293-301.
- [2] Bakht, J., Ali, H., Khan, M. A., Khan, A., Saeed, M., Shafi, M., ... & Tayyab, M. (2011). Antimicrobial activities of different solvents extracted samples of *Linum usitatissimum* by disc diffusion method. *African Journal of Biotechnology*, 10(85), 19825-19835.

POSTER PRESENTATION

THE ANTIBACTERIAL ACTIVITY OF *ELAEAGNUS ANGUSTIFOLIA* L. FRUIT AND LEAF EXTRACTS

Irmak Alev¹, Cansu Vatansever², Hüseyin Servi³

¹Faculty of Pharmacy, Altınbaş University, Istanbul, Turkey

²Department of Pharmaceutical Microbiology, Faculty of Pharmacy, Altınbaş University, Istanbul, Turkey

³Department of Pharmaceutical Botany, Faculty of Pharmacy, Altınbaş University, Istanbul, Turkey

Objective: *Elaeagnus angustifolia* L. (Russian olive, oleaster) belongs to Elaeagnaceae family. The plant has been used as traditional folk medicine for many years due to it is anti-inflammatory, muscle relaxant, anti-ulcerogenic, antimicrobial, antinociceptive, antitumor, and antioxidant effects [1]. Previously, the antibacterial activity of crude methanol extract, *n*-hexane, ethyl acetate, chloroform and water fractions of *E. angustifolia* L. was determined with well diffusion assay. The crude extract, hexane, ethyl acetate, and water fractions showed activity with different inhibition zone diameters against *E. coli* and *S. aureus*. And only the hexane and ethyl acetate fractions showed inhibitory activity against *P. aeruginosa*. But the chloroform fraction had no antibacterial activity against tested microorganisms [2]. The purpose of this study is to determine the antibacterial activity of *Elaeagnus angustifolia* L. fruit and leaf extracts against *E. coli* ATCC 25922, *S. aureus* ATCC 29213, *P. aeruginosa* ATCC 27853 and *B. cereus* ATCC 14579 by using broth microdilution and disc diffusion methods.

Material and Methods: Maceration method was used to obtain the hexane, ethyl acetate and methanol extracts from the fruit and leaf of *Elaeagnus angustifolia* L. The experiments were began with 8 mg/mL for broth microdilution assay and 10 mg/disc extracts were used for disc diffusion assay. Chloramphenicol was used as a control antibiotic for both methods. All the experiments were performed in duplicate.

Results: The ethyl acetate extract of fruit was more effective than the other extracts for tested microorganisms. The minimum inhibition concentration (MIC) values of fruit ethyl acetate extract were 1 mg/mL for *B. cereus*, 2 mg/mL for *E. coli* and *S. aureus* and 8 mg/mL for *P. aeruginosa*. The inhibition zone diameters of fruit ethyl extract were 15 mm for *E. coli*, 14 mm for *S. aureus*, 12 mm for *P. aeruginosa* and 10 mm for *B. cereus*. The hexane and ethyl acetate extracts from leaves showed inhibitory activity against tested microorganisms at 8 mg/mL. But the fruit hexane extract did not show activity against *E. coli*. The leaf methanol extract did not show activity *E. coli* and *S. aureus*. The fruit methanol extract did not show antibacterial activity against *B. cereus*.

Discussion / Conclusion:

In the current study reveals that the fruit and leaf extracts of *E. angustifolia* had inhibitory activity against different bacteria at different concentrations. And the antibacterial activity may change according to the plant part and extract type which is used. As an example, the leaf methanol extract was not susceptible against *E. coli* and *S. aureus* but fruit methanol extract was susceptible against these two bacteria.

Keywords: *Elaeagnus angustifolia* L., extract, antibacterial activity

References:

- [1] Afshar, F. H. & Asnaashari, S. & Torbati, M. 2016. "Essential oil from flowers and leaves of *Elaeagnus angustifolia* (Elaeagnaceae): Composition, radical scavenging and general toxicity activities". *Advanced Pharmaceutical Bulletin*. 6(2): 163–169.
- [2] Khan, S. & Khan, A. & Shah, A.A & Shah, S. M. & Hussain, S. & Ayaz, M. & Aya, S. 2016. Heavy metals content, phytochemical composition, antimicrobial and insecticidal evaluation of *Elaeagnus angustifolia*. *Toxicology and Industrial Health*, 32(1), 154-161.

POSTER PRESENTATION

**MUTAGENIC *IN VITRO* ACTIVITY AND GENOTOXIC EFFECT OF
ZYGOPHYLLUM CORNUTUM METHANOLIC EXTRACT**

Awatif Boumaza¹, Messaouda khallaf¹, Mouna Cherairia¹

¹*Département de Biologie, Faculté des Sciences de la Nature et de la Vie et Sciences de la Terre et de l'Univers.
Université 8 Mai 1945 Guelma*

Objective/Purpose: *Zygophyllum cornutum* *coss* is an Algerian medicinal plant widely used in the treatment of different diseases mainly diabetes. The purpose of our study was to determine possible mutagenic/genotoxic effect of this plant.

Materials and Methods: Three concentrations (5, 15 and 50 mg/ml) of *Zygophyllum cornutum* methanolic extract were screened to the presence of mutagenic activity by Ames test using TA100 and TA98 strains of *Salmonella tiphymurium* with and without metabolic activation (S9 microsomal fraction) according to the method of K. Mortelmans and E. Zeiger, 1983 [1]. The micronucleus assay on human lymphocytes was used to evaluate genotoxic effect according to the method described by Fenech et morley, 1985 [2]. The frequency of the micronucleus (MN) and the proliferation index (PI) were calculated according to Fenech et morley, 1985.

Results: The Ames test showed negative results without microsomal activation while mutagenic activity was observed in the presence of microsomal activation. The number of the revertant colonies observed at 5, 15 and 50 mg/ml was respectively 49, 77 and 89 with TA98 and 88, 95 and 55 with TA100. In the micronucleus test, a non significant increase of the micronucleus frequency was observed at 15 mg/ml 8.66 with PI of 1.70 . At 50 mg/ml, the frequency of MN increased (24.76) and PI decreased (1.55) significantly.

Conclusion/Discussion: Even that further studies must be carried out, the mutagenic activity and the genotoxic effect of *Zygophyllum cornutum* should be taken in consideration when used as therapeutic plant.

Key words: Ames test, genotoxicity, micronucleus test, *Zygophyllum cornutum*.

References:

- [1] Mortelmans, K., Zeiger, E. (1983). The Ames *Salmonella*/microsome mutagenicity assay. *Mutation Research* 455 (2000) 29–60
[2] Fenech, M., Morley, A. (1985). Measurement of micronuclei in lymphocytes. *Mutat. Res.*, 147: 29-36

POSTER PRESENTATION

BIOLOGICAL ACTIVITIES OF *THYMUS FONTANESII* POLYPHENOLIC EXTRACT: *IN VITRO* AND *IN VIVO* EFFECT ON BACTERIAL ADHESION AND COLONISATION

Boubakeur B^{1,2}, Khadem H^{1,2}, Drabo M. S³, Tir Touil Meddah A².

Faculty of nature and life sciences, IbnKhaldoun University, Tiaret, Algeria.

Laboratory of Bioconversion, Health Safety and Microbiological Engineering, University of Mascara, Algeria.

Laboratory of Applied Biochemistry and Immunology - CRSBAN, UFR-SVT 03 B.P. 7021 Ouagadougou 03,

University of Ouagadougou, Burkina Faso.

Boubakeurbadra82@yahoo.fr

Objective / Purpose: Polyphenols are plants secondary metabolites that are not absorbed in the small intestine but can be degraded by colonic microbiota to simpler and more easily absorbable bioactive compounds with potential physiological effects. They can act as potent antioxidants and have diverse effects on microbes. They appear to be effective antimicrobial agents against pathogens and to improve probiotic growth and metabolisms. A polyphenol-rich diet may then improve the intestinal status and possibly attenuate oxidative stress and suppress inflammations. These attributes express the potential antioxidant and microbial effects of those molecules. Furthermore, these compounds have technological aspects as they can enhance metabolism of LAB in dairy industry. The aim of this study was therefore to assess the *in vitro* and *in vivo* influence of *Thymus fontanesii* polyphenolic extract on the bacterial adhesion and colonization.

Material and Methods: The polyphenolic content of *Thymus fontanesii* was extracted in methanol by sonication. For the assessment of *in vitro* polyphenolic effect on bacterial adhesion and biofilm formation, bacterial cultures, at fixed density ($\sim 10^7$ germs/ml), are supplemented by the polyphenolic extract of *T. fontanesii* in increasing final concentrations: 0 μ g/ml, 50 μ g/ml, 100 μ g/ml and 200 μ g/ml. For the *in vivo* data, against negative control group (no treatment has been achieved), test group was force-fed 4 weeks with 100 μ l of polyphenolic extract.

Results: Cynamic, caffeic, carboxylic, furilic, M anisic and vanilic acids, berberine, catechine, resorcinol and rutine were the important phenolic acids and flavonoids identified by HPLC in the extract. Considerable anti-adhesive activity of extract has been recorded; the adhesion of the pathogenic strains tested was reduced in presence of the polyphenolic extract of *Thymus fontanesii* at lethal concentrations ranging from 100 μ g / ml to 200 μ g / ml, the strains *Bacillus cerius* 11798, *Bacillus subtilis* 3633, and *Escherichia coli* 10536 proved to be more sensitive, they completely lost their ability to form a biofilm. At sublethal concentrations (from 50 to 100 μ g/ml), *Staphylococcus aureus* 25922 and *Pseudomonas aeruginosa* 27853 develops a very high propensity to form a biofilm protector against the toxic effect of extract. The *in vivo* tests were shown that the polyphenolic extract shaped Wistar Rat gut microbiota in favour of beneficial bacteria and in detriment of pathogenic bacteria, and prevented bacterial translocation.

Conclusion / Discussion: Therefore, these natural molecules from medicinal plant can be used as alternative or additive agents to prevent the oxidative stress and modulate the bacterial activity and their adhesion, colonization and translocation capacity.

Keywords: Microbiota; bacterial translocation; polyphenols; bacterial adhesion; *Thymus fontanesii*.

References:

- [1] Boubakeur B., Tirtouil A., Khadem H., Meddah B., Ahcen S. (2016). An assessment of the effect of aqueous extract from *Thymus fontanesii* on growth, aggregation and biofilm formation of pathogenic and probiotic bacteria. *J Appl Environ Bio Sci.* 6 (7):51-60.
- [2] Khalil, S.K and A. Rowayd 2010. Influence of gallic acid and catechine polyphenols on probiotic properties of *Streptococcus thermophilus* CHCC 3534 strain. *Word journal of Microbiology and Biotechnology.*, 11: 2069-2079.
- [3] Kos, B., J. Suskovic., S. Vukovic., M. Simprag., J. Frece and S. Matosic, 2003. Adhesion and aggregation ability of probiotic strain *Lactobacillus acidophilus* M92. *Journal of Applied Microbiology.* 94: 981-987.
- [4] Parkar, S. G., Stevenson, D. E., Skinner, M, A. 2008. The potential influence of fruit polyphenols on colonic microflora and human gut health. *International Journal of Food Microbiology.* 124, 295-298.
- [5] Quideau, S., Deffieux, D., Douat-Casassus, C., Pouysegou, L. 2011. Plant polyphenols: chemical properties, biological activities, and synthesis. *Chem Int Ed.* 50, 586-621.
- [6] Verhelst, A.C. M., B.C. Schroyen., B.C.T. Buys., A, C. Niewold, 2010. The effects of plant polyphenols on enterotoxigenic *Escherichia coli* adhesion and toxin binding. *Livestock Science.* 133: 101-103.

POSTER PRESENTATION

**IN VITRO TOXICOLOGICAL EVALUATION OF SOME MEDICINAL PLANTS
USED BY WOMEN IN NORTHERN MAPUTALAND, SOUTH AFRICA**

Samukelisiwe Clerance Ngubane¹, Helene de Wet¹, Sandy van Vuuren²

¹Department of Botany, Faculty of Science and Agriculture, University of Zululand, Private Bag 1001,
KwaDlangezwa 3886, South Africa.
deweth@unizulu.ac.za

²Department of Pharmacy and Pharmacology, Faculty of Health Science, University of Witwatersrand, 7 York
Road, Parktown 2193, South Africa

Purpose: Medicinal plants are still used as a primary source of healthcare in northern Maputaland because of the strong cultural beliefs in medicinal plants. This increases the potential exposure to plant toxicity. Women in this area use also medicinal plants to maintain healthy pregnancy/delivery and to treat gynaecology problems [1]. There are very few scientific reports on the toxicity of medicinal plants in this area. The lack of toxicological information on these plants raises concerns about the safety for the unborn child. The aim of this study was to evaluate the safety of 33 medicinal plants used by women in northern Maputaland to treat gynaecology and obstetrics complaints.

Material and Method: An Ethnobotanical survey was conducted using a structured questionnaire to investigate medicinal plants used to treat gynaecology and obstetrics complaints. All plant material was collected on site, dried and ground into powder. Methanol-dichloromethane (1:1) and aqueous extracts were prepared. An acute cytotoxicity study was conducted using an *in-vitro* assay called the Brine Shrimp Lethality Assay (BSLA). Cytotoxicity was determined by the death of *Artemia franciscana* after the exposure to the plant extract. In cases where roots were being used by the people, leaves (17) were simultaneously collected for comparison. The results were recorded after 24 and 48 hours.

Results: The ethnobotanical survey resulted in the documentation of 33 medicinal plant species. The people in northern Maputaland use mostly water to prepare their herbal medicine. Eleven aqueous extracts were cytotoxic, and the same number was also observed with the ethanol-dichloromethane extracts. However, in some cases, only the aqueous extracts were toxic or only the organic extracts. Aqueous extract of *Euphorbia tirucalli* stem was considered the most cytotoxic by killing 100% of the brine shrimps within 24 hours in all replicates. However, this plant was only used by one interviewee. The most mentioned plants were *Bridelia cathartica* (18 times to treat dysmenorrhoea, amenorrhoea, menorrhagia, oligomenorrhoea infertility and to prevent miscarriage) and *Ranunculus multifidus* (13 times for genital warts). The aqueous root extract of *B. cathartica* was non-cytotoxic but the organic extract and the leaves were cytotoxic. Both the aqueous and organic extracts of *R. multifidus* indicated signs of cytotoxic activities. Overall leaf extracts were highly cytotoxic when compared to their respective root extracts, only three of the leaf extracts were non-cytotoxic in this experiment.

Conclusion: Literature showed that the majority of medicinal plants reported to have medicinal use in northern Maputaland have not been assessed for their safety. The number of cytotoxic extracts encountered indicate the importance of evaluating safety of traditional medicine. Subsequently, the communities need to be aware of such adversities.

Keywords: Cytotoxicity, Medicinal plants, Women problems, northern Maputaland, South Africa

References:

[1] De Wet, H., & Ngubane, S. (2014). Traditional herbal remedies used by women in a rural community in northern Maputaland (South Africa) for the treatment of gynaecology and obstetric complaints.

POSTER PRESENTATION

IN-VITRO SUSCEPTIBILITY TESTING OF TEA TREE OIL ACNE CREAM IN COMBINATION WITH SALICYLIC ACID AND NICOTINAMIDE

Saiqa Ishtiaq, Rabia Arshad, Sairah Hafeez Kamran

Punjab University College of Pharmacy, University of the Punjab, Allama Iqbal Campus, Lahore, Pakistan, saiqa.pharmacy@pu.edu.pk

The tea tree oil acne cream was prepared in Negiban dispensing pharmacy according to SOP by levigating the specific amount of ingredients; tea tree oil (2%), nicotinamide(3%) and salicylic acid(3%). The formulation is then tested for its susceptibility by microwell dilution method on 15 samples of *Staphylococcus aureus* (*S. aureus*) and five samples of *Propionibacterium acnes* (*P. acnes*) collected from the out-patient department of Skin, Mayo Hospital, Lahore. After finding the susceptibility, minimum inhibitory concentration of the cream was estimated to establish its effectiveness, by 96 well plate method. The results of the samples (1 to 15) against *S. aureus*, majority of the samples have showed maximum zone of inhibitions at dilution 1:5. For *P. acnes*, samples (1 to 5) majority no of the samples depicted maximum zone of inhibition at dilution ratio 1:10. The results suggests that the tea tree oil acne cream is effective in a narrow dilution range and shows potent anti-acne effect. While, the results of the MIC of the *S. aureus* and *P. acnes*, the cream showed 0.074, 0.069 at concentration 12.5 mg/ml and 0.023 at concentration 12.5 mg/ml, 0.007 at concentration 6.25 mg/ml, 0.073 at concentration 3.125 mg/ml and 0.072 at concentration 0.02 mg/ml, respectively. The results confirm the susceptibility and effectiveness of the formulation against the bacterial strains responsible for acne.

Key words: Tea Tree Oil, *Staphylococcus aureus*, *Propionibacterium acnes*

References:

- Abbasi, AM., Khan, MA., Ahmad, M., Zafar, M., Khan, H., Muhammad, N., & Sultana, S. (2009). Medicinal plants used for the treatment of jaundice and hepatitis based on socio-economic documentation. *African Journal of Biotechnology*, 8(8).
- Amsterdam, D. (1996). Susceptibility testing of antimicrobials in liquid media. *Antibiotics in laboratory medicine*, 4, 61-143.
- Bakkali, F., Averbeck, S., Averbeck, D., & Idaomar, M. (2008). Biological effects of essential oils—a review. *Food and chemical toxicology*, 46(2), 446-475
- Bassett, IB., Pannowitz, DL., & Barnettson, RS. (1990). A comparative study of tea-tree oil versus benzoylperoxide in the treatment of acne. *The Medical Journal of Australia*, 153(8), 455-458.
- Bassolé, IHN., & Juliani, HR. (2012). Essential oils in combination and their antimicrobial properties. *Molecules*, 17(4), 3989-4006.
- Burt, S. (2004). Essential oils: their antibacterial properties and potential applications in foods a review. *International journal of food microbiology*, 94(3), 223-253.
- Burt, SA., & Reinders, RD. (2003). Antibacterial activity of selected plant essential oils against *Escherichia coli* O157: H7. *Letters in applied microbiology*, 36(3), 162-167.

POSTER PRESENTATION

IMPACT OF SOME HERBICIDES AND HERBICIDE COMBINATIONS ON SOWING CHARACTERISTICS OF CORIANDER (*CORIANDRUM SATIVUM* L.)

Grozi Delchev

Department of Plant Production, Faculty of Agriculture, Trakia University, Stara Zagora, Bulgaria
delchevgd@dir.bg

Objective / Purpose: The aim of this investigation was to establish the influence of some soil-applied herbicides, foliar-applied herbicides and herbicide combinations on sowing characteristics of the coriander seeds and the quantity of waste grain.

Material and Methods: The research was conducted during 2013 - 2015 on pellic vertisol soil type. Under investigation was Bulgarian coriander cultivar Lozen 1 (*Coriandrum sativum* L.). Factor A included no treated control, 6 soil-applied herbicides – Tender EC (S-metolachlor) - 1.5 l/ha, Silba SC (metolachlor + terbuthylazine) - 3.5 l/ha, Sharpen 33 EC (pendimethalin) - 5 l/ha, Merlin flex 480 SC (isoxaflutole) - 420 g/ha, Smerch 24 EC (oxyfluorfen) - 1 l/ha, Raft 400 SC (oxidiargil) - 1 l/ha and 5 foliar-applied herbicides – Kalin flo (linuron) - 2 l/ha, Eclipse 70 DWG (metribuzine) - 500 g/ha, Sultan 500 SC (metazachlor) - 2 l/ha, Corrida 75 DWG (tribenuron-methyl) - 20 g/ha, Lontrel 300 EC (clopyralid) - 500 ml/ha. Factor B included no treated control and 1 antigraminaceous herbicide – Tiger platinum 5 EC (quizalofop-P-ethyl) - 2.5 l/ha. Soil-applied herbicides were treated during the period after sowing before emergence. Foliar-applied herbicides were treated during rosette stage of the coriander.

Conclusion / Discussion: Herbicides Merlin flex and Lontrel and herbicide combinations Merlin flex + Tiger platinum and Lontrel + Tiger platinum proven decrease germination energy of the coriander seeds. Laboratory seed germination and lengths of primary germ and primary root are decreased by herbicide Merlin flex and herbicide combination Merlin flex + Tiger platinum only. Herbicide Merlin flex and herbicide combination Merlin flex + Tiger platinum do not proven decrease waste grain quantities. High yields of coriander seeds are obtained by foliar treatment with antigraminaceous herbicide Tiger platinum after soil-applied herbicides Raft, Smerch, Sharpen, Silba and Tender. Tank mixtures of Tiger platinum with foliar herbicides Kalin flo, Eclipse, Sultan, Corrida and Lontrel also lead to obtaining of high seed yields. The use of the soil-applied herbicide Merlin flex does not increase the seed yield, due to its higher phytotoxicity against coriander. Alone use of soil-applied or foliar-applied herbicides leads to lower yields due to they must to combine for full control of weeds in coriander crops.

Keywords: coriander, herbicides, herbicide combinations, seed yield, sowing characteristics

POSTER PRESENTATION

PROFILING OF PHENOLIC COMPOUNDS BY HPLC–DAD, ANTIOXIDANT AND ENZYME INHIBITORY ACTIVITIES OF *MARRUBIUM VULGARE* L. AND *MARRUBIUM PARVIFLORUM* FISCH. & MEY. SUBSP. *PARVIFLORUM*

Fatma Ayaz¹, Nuraniye Eruygur¹, Yavuz Bağcı², H. Filiz Ayyıldız³

¹ Department of Pharmacognosy, Faculty of Pharmacy, Selcuk University, Konya, Turkey,

² Department of Pharmaceutical Botany, Faculty of Pharmacy, Selcuk University, Konya, Turkey,

³ Department of Basic Pharmaceutical Sciences, Faculty of Pharmacy, Selcuk University, Konya, Turkey

Objective: The genus *Marrubium* L., belonging Lamiaceae, is represented by 20 species in Turkey, twelve of which are endemic. *Marrubium* species is traditionally used in the treatment of various diseases such as asthma, pulmonary inflammation and hypoglycemi; for analgesic, sedative and cholagogue effects. According to the literature, it was found that *Marrubium* species have various therapeutic effects such as hypoglycemic, antihypertensive, antispasmodic, antiproliferative, antioxidant, hepatoprotective, hypolipidemic, gastroprotective, antibacterial and vasorelaxan as well as bioactive secondary metabolites, such as diterpenes, sterols, phenylpropanoids and flavonoids. We aimed to investigate the antioxidant and enzyme inhibitory activities of the methanol and water extracts of *M. vulgare* L. (MV) and *M. parviflorum* Fisch. & Mey. subsp. *parviflorum* (MP) growing in Turkey.

Material and Methods: We conducted *in vitro* preliminary bioassays on the plant extracts for their evaluation of antioxidant activities (DPPH, ABTS, iron chelating, β -carotene/linoleic acid system, TPC and TFC) and enzyme inhibition effects (acetylcholinesterase, butyrylcholinesterase, and tyrosinase). We also analysed the plant extracts for the qualitative and quantitative determination of phenolic compounds by HPLC-DAD using various mobile phases having different elution gradients and run times.

Results: According to our results, the methanol extract of MP was found the highest antioxidant properties for DPPH and ABTS assays while the water extract of MV having the most iron chelating and lipid peroxidation capacity. In addition, while the methanol extracts of MP and MV showed more inhibition effects on the cholinesterase enzymes, it was found that water extracts of MP and MV had more antityrosinase activities. When the plant extracts are compared in terms of phenolic substance content of MP and MV, we observed that trans-ferulic acid, elagic acid, caffeic acid, and 1,2-dihydroxy benzene were determined as the main components in the extracts.

Conclusion: As a conclusion, our study can provide scientific results for further investigation on the developing novel antioxidant, anticholinesterase, and antityrosinase agents from natural sources.

Keywords: Antioxidant activity, enzyme inhibition, HPLC, *Marrubium vulgare*, *Marrubium parviflorum*,

Acknowledgment: This work was supported by Selcuk University-Scientific Research Project No: 18401140.

POSTER PRESENTATION

**PRELIMINARY PHYTOCHEMICAL SCREENING OF FOUR SAHARAN PLANTS USED
IN SOUTH WEST OF ALGERIA**

¹*Mahmoudi K., ¹Cheriti A et ²Didi Oueld El Hadj M

¹*Phytochimie and Organic Synthesis Laboratory, University of TAHRI Mohamed of Bechar, Algeria.*

²*Ecosystem Protection in Arid and Semi-arid Area Laboratory, University of KASDI Merbah of
Ouargla, Algeria.*

**E-mail: Kheira.mahmoudi05@yahoo.fr*

Phytochemical surveys are now acted as the first step towards the discovery of useful drugs. The present investigation was carried out to assess the qualitative phytochemical analysis of different parts of four Saharan plants belonging to Solanaceae and Asclepiadaceae family namely: *Daturastramonium*, *Hyoscyamusmuticus*, *Pergulariatomentosa* and *Calotropisprocera*.

Qualitative phytochemical analysis of this plant confirms the presence of various phytochemicals like saponins, flavonoids, steroids, unsaturated sterols and terpenes and alkaloids in both the plants extracts, where as unsaturated sterols, cardenolides and tannins were found to be absent in some extracts of parts of plants tested.

Keyword: qualitative phytochemical analysis, *Calotropis procera*, *Pelgularia tomentosa*, *Datura staramonium* and *Hyoscyamus muticus*.

POSTER PRESENTATION

**PHYTOCHEMISTRY AND CYTOTOXIC POTENTIAL OF
LITSEA DECCANENSIS BARK.**

Tabassum Khan^{1*} and Anju Kashyap²

¹Department of Pharmaceutical Chemistry and Quality Assurance, SVKM's Dr. Bhanuben Nanavati College of Pharmacy, Mumbai, Maharashtra, India.

²Department of Quality Assurance, SVKM's Dr. Bhanuben Nanavati College of Pharmacy, Mumbai, Maharashtra, India.

Objective: The genus *Litsea*, family Lauraceae, contains approximately 200 plant species mainly distributed in the tropical and subtropical regions of the world. *Litsea* species have been reported to possess a wide variety of pharmacological activities such as anticancer, anti-inflammatory, antimicrobial, antioxidant, antidiabetic, anti-HIV, insecticidal to name a few. There are very few phytoconstituents reported from the bark of *Litsea deccanensis* Gamble, viz. caryophyllene oxide, p-sitosterol, actinodaphnine and litsomentol. Given the varied pharmacological activity of the leaves and stem of this plant we thought it prudent to investigate the bark for secondary metabolites with potential biological activity. These are the first reports of biological activity of the bark of *Litsea deccanensis* and identification of new phytoconstituents.

Material and methods: The bark was collected from Amboli Ghats, Maharashtra state, dried and powdered. Successive soxhlet extraction was carried out using petroleum ether 60-80°C, toluene, dichloromethane, ethyl acetate, methanol and water. Alkaloid rich fraction of the bark powder was prepared using reported method. The successive solvent extracts and the alkaloid rich fraction were screened for cytotoxic activity using Adriamycin as standard. LC-MS analysis of the solvent extracts and the alkaloid rich fraction were conducted. The total alkaloid content in the bark was determined using aluminium chloride assay.

Results: The successive solvent extracts exhibited significant cytotoxic activity on the selected cell lines. The alkaloid rich fraction showed cytotoxic activity on Human Breast Cancer cell line MCF-7, Human Breast Cancer cell line MDA-MB-231, Human Cervical Cancer cell line HeLa, Human Leukemia cell line U-937. The alkaloid rich fraction exhibited the most potent activity against all the selected cell lines. The LC-MS data of alkaloid rich fraction indicated the presence of alkaloids such as 3,4-methylene dioxy methamphetamine (MDMA), arecoline, salsolidine, boldine, hydrocodone, nedocromil, phytosphingosine and colchicine. The total alkaloid content was found to be 0.13 µg of atropine equivalent/100 mg of extract.

Conclusion: The successive solvent extracts and the alkaloid rich fraction were found to exhibit cytotoxic activity on breast, cervical and leukemia cell lines with the alkaloid rich fraction being the best and potentially responsible for the observed cytotoxic effect. This paper presents the first report on new phytoconstituents identified in *Litsea deccanensis* bark. It also presents for the first time the cytotoxic potential of the bark opening new avenues for drug design and development.

Keywords: Cytotoxicity, *Litsea deccanensis*, alkaloid rich fraction, alkaloids, MCF-7, HeLa, U-937.

References:

1. Kong, D.G., Zhao, Y., Li, G.H., Chen, B.J., Wang, X.N., Zhou, H.L., Lou, H.X., Ren, D.M. and Shen, T., 2015. The genus *Litsea* in traditional Chinese medicine: an ethnomedical, phytochemical and pharmacological review. *Journal of ethnopharmacology*, 164, pp.257,263.
2. T. R. Govindachari, N. Viswanathan and P. A. Mohamed. Structure of litsomentol, a new tetracyclic triterpene. CIBA Research Centre.
3. MUKHERJEE, P. K. *QUALITY CONTROL AND EVALUATION OF HERBAL DRUGS: Approaches for Evaluating Natural Products and ... Traditional Medicine.*; ELSEVIER, 2018.
4. John, B.; Sulaiman, C. T.; George, S.; Reddy, V. R. K. Spectrophotometric Estimation of Total Alkaloids in Selected *Justicia* Species. *Int. J. Pharm. Pharm. Sci.* **2014**, 6 (5), 647–648.
5. Salamah, N.; Ningsih, D. S. Total Alkaloid Content in Various Fractions of *Tabernaemontana Sphaerocarpa* Bl. (Jembirit) Leaves. *IOP Conf. Ser. Mater. Sci. Eng.* **2017**, 259 (1). <https://doi.org/10.1088/1757-899X/259/1/012017>.
6. Skehan, P.; Storeng, R.; Scudiero, D.; Monks, A.; McMahon, J.; Vistica, D.; Warren, J. T.; Bokesch, H.; Kenney, S.; Boyd, M. R. New Colorimetric Cytotoxicity Assay for Anticancer-Drug Screening. *J. Natl. Cancer Inst.* **1990**, 82 (13), 1107–1112.
7. Vichai, V.; Kirtikara, K. Sulforhodamine B Colorimetric Assay for Cytotoxicity Screening. *Nat. Protoc.* **2006**, 1 (3), 1112–1116. <https://doi.org/10.1038/nprot.2006.179>.
8. Khandelwal, K. R. *Practical Pharmacognosy: Techniques and Experiments*; Nirali Prakashan, 2006.

POSTER PRESENTATION

IN VITRO ANTIOXIDANT PROSPECTION OF ONION OIL FROM ROMANIAN SOURCE

Gabriela Tataringa¹, Ana Maria Zbancioc¹, Catalina Stan², Cornelia Mircea³, Monica Hancianu⁴

¹Department of Organic Chemistry, Faculty of Pharmacy, “Grigore T. Popa” University of Medicine and Pharmacy, Iași, Romania

²Department of Pharmaceutical and Biotechnological Drug Industry, Faculty of Pharmacy, “Grigore T. Popa” University of Medicine and Pharmacy, Iași, Romania

³Department of Biochemistry, Faculty of Pharmacy, “Grigore T. Popa” University of Medicine and Pharmacy, Iași, Romania

⁴Department of Pharmacognosy, Faculty of Pharmacy, “Grigore T. Popa” University of Medicine and Pharmacy, Iași, Romania

Objective / Purpose: Onion is botanically known as *Allium cepa* and it belongs to the Liliaceae family. It has been used as a medicinal remedy from time immemorial, that's probably because its chemical composition. The main organic compounds are flavonoids and organosulfur compounds. This study was designed to evaluate the antioxidant activities of Romanian onion oil.

Material and Methods: The essential oil of *Allium cepa* was obtained by hydrodistillation and then it was analyzed by GC-MS. In vitro antioxidant activity was done using DPPH scavenging activity, reducing power and hydrogen peroxide scavenging assay

Results: Onion oil was able to reduce DPPH radicals, with IC₅₀ value of 0.781 mg/mL; the results indicate that onion oil exhibited a significant DPPH radical scavenging activity, about 8 folds more active than ascorbic acid. For all investigated concentrations, the absorbance of the mixture was high, thus the reducing power was significant. Hydrogen peroxide scavenging activity showed that the antioxidant components of onion oil (probably sulfur compounds) are good electron donors, IC₅₀ being 0.36 mg/mL.

Conclusion: Taking into account the obtained results, we can state that *Allium cepa* oil can play a major role in alleviating the number of oxidative stress by reducing the oxidative damage of cellular component caused by ROS.

Keywords: onion oil, antioxidant activity

References:

- [1] Helen, A., Krishnakumar, K., Vijayammal, P.L., Augusti K.T. (2000). Antioxidant effect of onion oil (*Allium cepa* Linn) on the damages induced by nicotine in rats as compared to alpha-tocopherol. *Toxicol Lett*, 116(1-2), 61-8.
- [2] Ye, C.L., Dai, D.H., Hu, W.L. (2013). Antimicrobial and antioxidant activities of the essential oil from onion (*Allium cepa* L.). *Food Control*, 30(1), 48-53.
- [3] Vazquez-Armenta, F., Cruz-Valenzuela, M., Ayala-Zavala, J. (2016). Onion (*Allium cepa*) Essential Oils. In: Preedy, V.R. (Ed.), *Essential Oils in Food Preservation, Flavor and Safety*. Academic Press, 617-623.
- [4] Zohri, A.N., Abdel-Gawad, K., Saber, S. (1995). Antibacterial, antidermatophytic and antitoxigenic activities of onion (*Allium cepa* L.) oil. *Microbiological Research*, 2(150), 167-172.

POSTER PRESENTATION

PROFILING ENZYME INHIBITORY POTENTIAL OF CYCLOARTANE TYPE OF TRITERPENE GLYCOSIDES FROM THE ROOTS OF THREE *ASTRAGALUS* SPECIES

F. Sezer Şenol Deniz¹, Ilkay Erdogan Orhan¹, İhsan Çalış²

¹ Department of Pharmacognosy, Faculty of Pharmacy, Gazi University, Ankara, Turkey

² Department of Pharmacognosy, Faculty of Pharmacy, Near East University, Lefkosa, TRNC

Objective / Purpose: *Astragalus* L., the largest genus in the family Fabaceae, is represented by 64 sections and about 475 taxa with 202 endemic species in the flora of Turkey. The roots of various *Astragalus* species represent very old and well-known drugs in traditional medicine, used for antiperspirant, diuretic, and tonic purposes as well as for the treatment of nephritis, diabetes, leukemia, and uterine cancer. In the present study, the isolated compounds from the roots of different *Astragalus* species growing in Turkey were subjected to high-throughput screening against acetylcholinesterase (AChE), butyrylcholinesterase (BChE), and tyrosinase (TYR), the fundamental enzymes associated with the pathology of Alzheimer's and Parkinson's diseases, also collagenase and elastase enzymes, related to inflammation and cosmetic purposes.

Material and Methods: Eleven cycloartane triterpene glycosides were isolated from the roots of *Astragalus melanophrurius* (astrasieversianin II, astrasieversianin X, astragaloside I, astragaloside IV, astragaloside VI, cyclocanthoside E, and cyclocanthoside G), *A. oleifolius* (macrophyllsaponin A, macrophyllsaponin B, macrophyllsaponin C, and macrophyllsaponin D), and *A. cephalotes* var. *brevicalyx* (cyclocanthoside E). Structures of the isolated compounds were elucidated by means of following spectroscopic methods; UV, 1D NMR (¹H NMR, ¹³C NMR, and DEPT-135) and 2D NMR (COSY, HSQC, and HMBC). ChE, TYR, collagenase, and elastase inhibitory activity of the compounds were screened using ELISA microtiter assays at 1 mg/mL stock concentration.

Results: Our results indicated that the isolated compounds had a low ChE inhibitory activity, whereas they were inactive against TYR and elastase. The most active compound against collagenase was macrophyllsaponin B (35.41 ± 3.96 %), while the other compounds had a modest level of collagenase inhibitory activity.

Conclusion / Discussion: Cycloartane type of triterpene glycosides isolated from the roots of three *Astragalus* species were herein shown to have only no to low or moderate level of enzyme inhibitory effect against the mentioned enzymes, which represent the first study on them in this regard.

Keywords: Cholinesterase, tyrosinase, elastase, collagenase, enzyme inhibition, cycloartane glycosides, *Astragalus*

POSTER PRESENTATION

ANTIOXIDANT ACTIVITY AND PHENOLIC CONTENT OF *PEGANUM HARMALA* USED IN ALGERIAN FOLK MEDICINE

Habiba Daroui-Mokaddem^{1,2}, Assia Zeghib^{2,3}, Fouad Zeghib⁴, Linda Achouri³, Sara Abbas³, Khouloud Fradj³, Amel Moussaoui³, Soraya Hioun⁵, Zahia Kabouche² and Belgacem Djabri⁶

¹Université Badji-Mokhtar, Annaba, Département de Médecine, Faculté de Médecine, 23000. ANNABA, ALGERIE. E-mail : habiba.mokaddem@yahoo.com

²Université des frères Mentouri-Constantine, Département de chimie, Laboratoire d'Obtention de Substances Thérapeutiques (LOST), 25000, CONSTANTINE, ALGERIE.

³Université Larbi Tébessi-Tébessa, Département de Biologie Appliquée, Faculté des Sciences Exactes et des Sciences de la Nature et de la Vie, 12000, TEBESSA, ALGERIE.

⁴Université Larbi Tébessi-Tébessa, Laboratoire Eau et Environnement (LEE), 12000, TEBESSA, ALGERIE.

⁵Université Larbi Tébessi-Tébessa, Département des êtres vivants, Faculté des Sciences Exactes et des Sciences de la Nature et de la Vie, 12000, TEBESSA, ALGERIE.

⁶Université Larbi Tébessi-Tébessa, Département de Biologie Appliquée, Laboratoire des molécules bioactives et applications (LMBAA), 12000, TEBESSA, ALGERIE.

The aim of this study is to determine the total phenolic content and the antioxidant potential of *Peganum harmala* (Zygophyllaceae), medicinal plant widely used in Algerian folk medicine.

Air-dried and powdered aerial parts of *Peganum harmala*, harvested from the area of Tébessa (North-Eastern Algeria), were extracted by percolation using solvents with increasing polarity, successively: petroleum ether (PE), dichloromethane (DM), ethyl acetate (EA) and methanol (ME), to yield dry extracts. The capacity of the obtained extracts to inhibit the free radical 1, 1-diphenyl-2-picrylhydrazyl (DPPH) was measured according to the method of Loo et al. (2008). Total phenolic content was estimated as gallic acid equivalents per milligram of dried plant extract, according to the Folin-Ciocalteu phenol reagent method (Li et al., 2007).

The ME extract showed, on the one hand, the best DPPH inhibition percentage at test-concentration, and, on the other hand, the highest yield which was 4.3, 5.6 and 15.8-fold higher than that of PE, DM and EA extracts, respectively. Its total phenolic content was 2-fold higher than that of PE, DM and EA extracts.

Peganum harmala may be suggested as a new potential source of natural antioxidant via its ME extract justifying the Algerian folk medicine use of this plant. Further investigations are necessary in order to refine its antioxidant potential and to determine its phytochemical composition.

Keywords: *Peganum harmala*, percolation, extracts, antioxidant, phenolic content.

References

- Li, H.B.; Cheng, K.W.; Wong, C.C.; Fan, K.W.; Chen, F.; Jiang, Y. Evaluation of antioxidant capacity and total phenolic content of different fractions of selected microalgae. *Food Chem.* 2007, 102, 771-776.
Loo, A.Y.; Jain, K.; Darah, I. Antioxidant activity of compounds isolated from the pyrolygneous acid, *Rhizopora apiculata*. *Food chem.* 2008, 107, 1151-1160.

POSTER PRESENTATION

**PHYTOCHEMICAL SCREENING AND ANTIBACTERIAL ACTIVITY OF
PITURANTHOS SCOPARIUS AERIAL PARTS HARVESTED FROM THE AREA OF
TÉBESSA (ALGERIA)**

Assia Zeghib^{1,2}, Habiba Daroui-Mokaddem^{2,3}, Fouad Zeghib⁴, Soraya Hioun⁵, Khouloud Fradj¹, Zahia Kabouche² and Belgacem Djabri⁶

¹ Université Larbi Tébessi-Tébessa, Département de Biologie Appliquée, Faculté des Sciences Exactes et des Sciences de la Nature et de la Vie, 12000, TEBESSA, ALGERIE.

² Université des frères Mentouri-Constantine, Département de chimie, Laboratoire d'Obtention de Substances Thérapeutiques (LOST), 25000, CONSTANTINE, ALGERIE.

³ Université Badji-Mokhtar, Annaba, Département de Médecine, Faculté de Médecine, 23000. ANNABA, ALGERIE.

E-mail : habiba.mokaddem@yahoo.com

⁴ Université Larbi Tébessi-Tébessa, Laboratoire Eau et Environnement (LEE), 12000, TEBESSA, ALGERIE.

⁵ Université Larbi Tébessi-Tébessa, Département des êtres vivants, Faculté des Sciences Exactes et des Sciences de la Nature et de la Vie, 12000, TEBESSA, ALGERIE.

⁶ Université Larbi Tébessi-Tébessa, Département de Biologie Appliquée, Laboratoire des molécules bioactives et applications (LMBAA), 12000, TEBESSA, ALGERIE.

The aim of this study is to determine the phytochemical screening and the antibacterial potential of aerial parts from *Pituranthos scoparius* (Coss. & amp; Dur) Benth. & amp; Hook. (Apiaceae), endemic species widely used in Algerian folk medicine.

Air-dried and powdered aerial parts of *Pituranthos scoparius*, harvested from the area of Tébessa (North-Eastern Algeria), were extracted by percolation using solvents with increasing polarity, successively: petroleum ether (PE), dichloromethane (DM), ethyl acetate (EA) and methanol (ME), to yield dry extracts. The plant aerial parts were screened for the presence of key families of phytochemicals according to the standardized methods (Dohou *et al.*, 2003; Rizk, 1982; Razafindrambao, 1973 and Bouquet, 1972). The antibacterial activity of the obtained dried plant extracts was evaluated against selected pathogenic bacteria, using the well agar diffusion method.

The phytochemical screening of the plant aerial parts highlights a variety of secondary metabolites mainly represented by flavonoïds, saponins and alcaloïds. The ME extract, on the one hand, showed against the tested pathogenic bacteria, the best antibacterial effect which was either similar or better than that of some antibiotics-controls and, on the other hand, it exhibited the highest yield which was nearly 2-fold higher than that of PE, DM and EA extracts all together. The good antibacterial effect of PE extract was relativized at view of its extraction lower-yield.

The *Pituranthos scoparius* ME extract may be suggested as a new potential source of natural antibacterial, justifying the Algerian folk medicine use of this endemic species. Further investigations are necessary in order to refine its antibacterial potential and to determine its phytochemical composition.

Keywords: *Pituranthos scoparius*, percolation, extracts, antibacterial activity, phytochemical screening.

References:

- Bouquet A. (1972). Plantes Médicinales du Congo Brazzaville. Ed: O.R.S.T.O.M.
Dohoo N. Yamin K. Tahrouch S. Idrissi Hassani L. M. Badoc A. et Gmira N. (2003). Screening phytochimique d'une endémique ibéro marocaine *Thymelaea lythroides*. Bull. Soc pharm. Bordeaux. 142, pp : 61-78.
Rizk A.M., (1982). Constituents of plants growing in Qatar, *Fitoterapia*, 52 (2) : 35-42.
Razafindrambao R.S. (1973). Etude d'une Plante Médicinale Malgache *Buxus madagascariensis* Baill et ses variétés. Ed: O.R.S.T.O.M

POSTER PRESENTATION

STUDY OF THE VARIETAL BEHAVIOR OF 6 BASIL VARIETIES (*OCIMUM BASILICUM* L) GROWN FOR THE FIRST TIME IN ALGERIA UNDER ARID CLIMATE

Nabila Slougui^{1,2}, Naima Draoui³, Cherifa Khaddar³, Mahfoud Hadj Mahammed²

¹ Department of the preparatory class National Polytechnic School of Constantine., Algeria,
slougui.nabila@gmail.com

² Biogeochemistry Laboratory of the desert Environments, Kasdi Merbah University, Ouargla, Algeria

³ Department of Agronomy, Kasdi Merbah University, Ouargla, Algeria

Objective / Purpose: In Algeria the production of aromatic and medicinal plants is very small and difficult to evaluate and almost all of our needs is imported. The objective of our trial is to study the behavior of six (06) Varieties of basil (*Ocimum basilicum* L.) (O.B.) in the edapho-climatic conditions of the Ouargla region (southeast Algeria).

Material and Methods: Our experimentation took place at the station of the Technical Institute for Development of Saharan Agriculture (I.T.D.A. S) of Hassi Ben Abdallah, which is at an altitude of 157 m, latitude of 32 °. 52 ' north and longitude of 5 °. 26 ' East. The plant material consist 6 varieties of basil: *Ocimum basilicum* V1, *O. B. purperescens* V2, *O. B. Minimum* V3, *O. B. Marceillais* V4, *O. B. cinnamon* V5 and *O. B. citriodora* V6. The experimental device adopted [1] is in complete random blocks, consisting of 6 treatments and 4 repetitions. Each block contains 6 elementary plots of 2 m in length on 1.5 wide. To achieve the plotted objectives we studied a number of growth parameters (stem height, branch width and leaf area), flowering (flowering date), yield (number of leaves per plant, total biomass,...), The cycle time and the mortality rate.

Results: The mortality rate is very low and sometimes nil in some varieties, purple (V2), Dwarf Compact (V3) and cinnamon (V5) showed a lower mortality rate of 3.12% against the highest mortality rate is marked by OM (V4) with a E average of 40.62% this can be explained by the fact that Marseille basil (V4) is less suited to the climatic conditions of arid zones.

Conclusion / Discussion: The statistical study of our agronomic results gave us significant values to highly significant, which confirms the success of the culture of the six varieties of basil of European origin under arid climate.

Keywords: Varietal behavior, *Ocimum basilicum* L, growth parameters, yield parameters, arid climate.

References:

[1] Carron, C, A., Rey, C, H. and Brutin, (2017). Essai de variétés de basilic en montagne. *Revue suisse vitic. Arboric. Hortic*, 38(1), 51-55.

POSTER PRESENTATION

**NEUROPROTECTIVE EFFECT OF *HYPERICUM THYMOPSIS* AGAINST
CHRONIC EXPOSURE TO ALUMINUM CHLORIDE AND ALZHEIMER'S
DISEASE**

Salima Douichene*, Wahiba Rached, Nessrine Ammari, Nouredine Djebli

*Department of Biology, laboratory of pharmacognosy and phytotherapy, faculty of life and Natural
Sciences, Mostaganem university Algeria*

The effect of Aluminum chloride was investigated to describe the associated behavioral and brain modifications. We don't know enough about the biological chemistry of chronic and sub-acute exposure to Aluminium to be able to predict its impact on human health. Although the hypothesis of a link between Aluminium and Alzheimer's disease (AD) has been supported by several epidemiological studies. Extract of *Hypericum thymopsis*(HTE), a well known medicinal plant, is used for the treatment of depression, has been explored in the present study for its protective role against Aluminum neurotoxicity. This data suggests that HTE may be a candidate for application in neurodegenerative diseases such as Alzheimer's disease. Results of this study demonstrate that Aluminiumneurotoxicity play an important role in the development of anxiety disorders, depression and memory deficit in mice, these alterations of behavioral activities can be a cause of development of Alzheimer's disease. HTE possesses significant antioxidant activity and renders neuroprotection which was more pronounced at the dose of 200mg/ kg against Al induced neurotoxicity. Chronic administration of *Hypericum thymopsis* significantly improved retention in both tasks, attenuated oxidative damage, Aluminum concentration in Aluminum treated mice (p 0,05). *Hypericum* have neuroprotective effects against Aluminum-induced cognitive dysfunction and oxidative damage.

Keywords: Aluminum, Alzheimer's Disease, *Hypericum Thymopsis* Extract (HTE) Neuroprotection, Mice

POSTER PRESENTATION

IN VITRO ENZYME INHIBITORY ACTIVITY OF THREE *FERULAGO* SPECIES GROWING IN TURKEY

Tuğba Günbatan¹, Mert Ilhan¹, Esra Küpeli Akkol¹, İlhan Gürbüz¹, Hayri Duman², Ceyda Sibel Kiliç³

¹Department of Pharmacognosy, Faculty of Pharmacy, Gazi University, Etiler 06330, Ankara, Turkey

²Department of Biology, Faculty of Science, Gazi University, 06330, Etiler, Ankara, Turkey

³Department of Pharmaceutical Botany, Faculty of Pharmacy, Ankara University, 06100, Tandogan, Ankara, Turkey

Ferulago species are named as “Çakşır” or “Çağşır” in Turkey and represented with 35 taxa (18 of them are endemic). In this study, hyaluronidase, collagenase and elastase inhibitory potential relevant to wound healing activities of the ethanol extracts of the roots of three *Ferulago* species were investigated.

Material and Methods: 80% ethanol extracts were prepared from the roots of 3 *Ferulago* species [*F. cassia* Boiss., *F. humilis* Boiss., *F. macrosciadia* Boiss. & Balansa,] and their *in vitro* hyaluronidase, collagenase and elastase inhibitory activities at 100 µg/ml concentration were evaluated [1-5].

Results: Ethanol extract of *F. macrosciadia* was the most active one against collagenase and elastase with 34.83% and 41.20% inhibition values, respectively. None of the studied extracts exerted significant inhibitory activity on hyaluronidase.

Conclusion: To the best of our knowledge, we herein report wound healing activity of three *Ferulago* species for the first time. According to the data we obtained, the most effective species in respect to collagenase and elastase inhibition tests was found to be *F. macrosciadia*.

Keywords: Apiaceae, Collagenase, Elastase, *Ferulago*, Hyaluronidase

References:

- [1] Lee, K.K., Choi, J.D. (1999). The effects of *Areca catechu* L. extracts on anti ageing. *International Journal of Cosmetic Sciences*, 21, 285-294.
- [2] Sahasrabudhe, A., Deodhar, M. (2010). Anti-hyaluronidase, anti-elastase activity of *Garcinia indica*. *International Journal of Botany*, 6, 299-303.
- [3] Barrantes, E., Guinea, M. (2003). Inhibition of collagenase and metalloproteinases by aloins and aloe gel. *Life Sciences*, 72, 843-50.
- [4] Melzig M.F., Loser, B., Ciesielski, S. (2001). Inhibition of neutrophil elastase activity by phenolic compounds from plants. *Pharmazie*, 56, 967-70.
- [5] Tumen, I., Guragac, F.T., Keles, H., Reunanen, M., Kupeli Akkol, E. (2017). Characterization and Wound Repair Potential of Essential Oil *Eucalyptus globulus* Labill., *Fresenius Environmental Bulletin*, 26 (11), 6390-6399.

POSTER PRESENTATION

**COMPARATIVE ANALYSIS OF RARE AND ENDANGERED PLANTS OF HIRKAN
DENDROFLORA**

T.S.Mammadov, A.G.Rasulova, S.B.Bagirova

*Institute of Dendrology of ANAS Azerbaijan, Baku, Mardakan settle., S.Yesenin str.89, Az 1044
dendrary@mail.az*

In the research work has been comparatively analyzed the Hirkan dendroflora and have been collected herbariums of rare and endangered trees and shrubs. There are identified 26 families of 40 genuses 60 species of rare and endangered trees and shrubs naturally spreading areas, distribution of individuals in populations, have been performed dendrochronical analyzes on old samples.

As a result of the observations have been determined, that in the Hirkan flora the majority of species in the areas have been diminished and decreased compared to previous years, many species are in danger of being disappeared.

From this point of view by the monitorings of the area were studied threat criterias, the reasons of decline in plant varieties, those plants have been identified based on version of IUCN 3.1.

In the research work have been processed out the protection measures, the biological features of species, the causes of change in natural resources.

Keywords: Hirkan National Park, 3rd relict, endem plants, forest, flora

References:

1. Azerbaijan USSR "Red Book". Baku: İşiq, 1989, 544 p.
2. Azerbaijan rare trees and shrubs. II Volume. Baku: Azerb.USSR Academ.Science. Publishing, 1964, 220 p.
3. Azerbaijan trees and shrubs. I cild.Baku: Elm, 1970, 321p.
4. Hajiyev V., Musayev S. Recommended plants and plant formations for Azerbaijan "Red and Green Books". Baku: Elm, 1996, 40 p.
5. Prilipko L.I. Azerbaijan trees and shrubs. I Volume. Baku: Azerb.USSR Academ.Science. Publishing, 1961, 319 p.
6. Mammadov T. S. And others. "Dendroflora of Azerbaijan". I-II-III Volumes. Baku: Elm, 2011-2017, 388 p/
7. "Flora of Azerbaijan".I – VIII Volumes.Baku:Academ.Science.of Azerbaijan USSR, 1950-1961, 566 p.
8. T.S.Mammadov, E. O. Isgender , T.H.Talıbov. "Azerbaijan rare trees and shrubs", Baku 2016, 378 p.
9. Azerbaijan's "Red Book". II Volume, Baku, 2013, 668 p.

POSTER PRESENTATION

ROS SCAVENGING AND REDUCING CAPACITY OF AROMATIC HERBS

Blagica Jovanova¹, Svetlana Kulevanova², Tatjana Kadifkova Panovska¹

¹ Department of Applied Biochemistry, Faculty of Pharmacy, University of Sts. Cyrill and Methodius, Skopje, Macedonia

² Department of Pharmacognosy, Faculty of Pharmacy, University of Sts. Cyrill and Methodius, Skopje, Macedonia

Objective / Purpose: The regular use of many aromatic herbs has been reported for prevention and treatment of numerous chronic conditions^{1,2}. Despite their widespread utilization, their antioxidant properties have not been studied in detail yet. Moreover, their regular use in prevention from diseases associated with excess of free reactive oxygen species (ROS) lacks scientific background. The purpose of the current study is to examine the antioxidant potency of frequently used aromatic herbs in diet and traditional medicine: tuber of *Allium ascalonicum* L. (AAR), *Allium cepa* L. (ACR), *Allium sativum* L. (ASR) and *Orchis mascula* (L.) L. (OMT), herba of *Melissa officinalis* L. (MOH), seeds of *Nigella sativa* L. (NSS) and *Trigonella foenum-graecum* L. (TFGS).

Material and Methods: Ethanol extracts were evaluated for their antioxidant activity by the following *in vitro* assays: DPPH (2,2'-diphenyl-1-picrylhydrazyl), FRAP (ferric reducing antioxidant power), NSSOH (non-site-specific-) and SSOH (site-specific hydroxyl radical-2-deoxy-d-ribose degradation) assays. Additionally, contents of total polyphenols and flavonoids were measured by Folin-Ciocalteu and AlCl₃ method.

Results: Obtained results showed prominent antioxidant activity *in vitro* and significant amount of polyphenols in the extracts. MOH was identified as extract with highest radical scavenging (IC₅₀ 1.54 mg/mL) and ferric reducing ability (28.3%). The ferric reducing power of extracts is decreasing as follows: MOH > ASR > NSS > TFGS > OMT > AAR > ACR. Additionally, MOH and ASR also manifested metal-chelating ability. Moreover, high correlation was established between total polyphenols and the antioxidant assays showing the evident role of polyphenols in the antioxidant defense.

Conclusion / Discussion: Overall, the data obtained for ethanol extracts of *Melissa officinalis* and *Allium sativum* in the current study demonstrate they could be considered as potential source of highly bioactive compounds against oxidative stress in biological systems. Additionally, they could provide protective role from chronic diseases associated with excessive amount of ROS.

Keywords: antioxidant activity, plant extracts, traditional medicine, aromatic herbs

References:

- [1] Booker, A., Johnston, D. and Heinrich, M. (2012). Value chains of herbal medicines—Research needs and key challenges in the context of ethnopharmacology. *Journal of Ethnopharmacology*, 140(3), 624-633.
[2] Heinrich, M. and Jäger, K.A. (2015). *Ethnopharmacology*. John Wiley & Sons – UK

POSTER PRESENTATION

**ANTIOXIDANT ACTIVITY OF FOUR NITROGEN ORGANIC COMPOUNDS:
CORRELATION BETWEEN THEORETICAL AND EXPERIMENTAL DATA**

Rafik Bensegueni^{1,2}, Mounia Guergouri², Chawki Bensouici³ & Mustapha Bencharif ².

¹*Université Mohamed Cherif Messaadia, 41000 Souk Ahras, Algérie.*

²*Laboratoire de Chimie des Matériaux Constantine, Université Frères Mentouri Constantine1, 25000 Constantine, Algérie.*

³*Centre de Recherche en Biotechnologie, Ali Mendjeli Nouvelle ville UV03, Constantine, Algérie.*

Objective / Purpose: The antioxidant compounds searching, continues to attract scientists, pharmaceutical and industrial actors. In the present study, we evaluated both theoretically and experimentally, the antioxidant activity of a set of four nitrogen organic compounds, one semicarbazone, one oxime and two phenylhydrazones, noted SEM, OXN, SIP and NIP, respectively.

Material and Methods: First, our four compounds were synthesized and characterized. Then, DPPH, CUPRAC and FRAP tests were used to evaluate their antioxidant activity. Furthermore, a DFT study (geometry optimization and frequency calculations) was carried out on the four molecules permitting the calculation of some molecular descriptors (BDE, IP and PA). All computations were performed using Gaussian 09 software.

Results: The tests of antioxidant activity were very fulfilling showing a well-established antioxidant power for our compounds. The SIP compound exhibits the highest DPPH scavenging activity which was even greater than that of the used standard antioxidants butylatedhydroxyanisole (BHA) and butylatedhydroxytoluene (BHT), with an IC₅₀ of 3.81 ± 0.26 against 6.14 ± 0.41 and 12.99 ± 0.41 µg/ml, respectively. On the other hand, the resulting BDE values are in a good agreement with the experimental results.

Conclusion / Discussion: The obtained results show an excellent correlation between the theoretical and the experimental data, which may allow these compounds to be used either in pharmaceutical or food industries. Farther tests are recommended in order to reinforce these findings.

Keywords: Antioxidant activity, DFT, molecular descriptors, nitrogen organic compounds.

References:

- [1] Blois, M.S. (1958) Antioxidant determinations by the use of a stable free radical. *Nature*. 81, 1199-1200.
- [2] Gaussian 09, Revision E.02. Frisch MJ, Trucks GW, Schlegel HB, Scuseria GE, Robb MA, Cheeseman JR et al. Gaussian, Inc., Wallingford CT, 2016.
- [3] Leopoldini, M., Russo, N., Toscano, M. (2011). The molecular basis of working mechanism of natural polyphenolic antioxidants, *Food Chem.* 125(2), 288-306.
- [4] Musialik, M., Litwinienko, G. (2005). Scavenging of dpph• radicals by vitamin E is accelerated by its partial ionization: the role of sequential proton loss electron transfer. *Org. Lett.* 7(22), 4951-4954.

POSTER PRESENTATION

STUDY OF THE HYPGLYCEMIC ACTIVITY OF THE COMMON FIGS OF THE REGION OF AIN DEFLA ALGERIA

Meliani Samiha

University of Medecine Departement of Pharmacy Saad Dahleb Blida , Algeria

Objective / Purpose: The fig tree is mentioned in the Quran in surah "At-Tine", which is the only one to have the title of a fruit or even a plant. Our work contributes to the study of the hypoglycemic activity of the common fig tree of the region of El Amra (Ain Defla) Algeria.

Material and Methods: To do this, 20 Wistar rats female, divided into 4 lots, were used: Lot 1: 5 normal controls; Lot 2: 5 normal controls treated with dry fig juice at 20%; Lot 3: 5 diabetic controls; Lot 4: 5 diabetic controls treated with dry fig juice at 20%. The rats are rendered diabetic by intra-peritoneal injection of a streptozotocin solution. The blood glucose is measured after 1 hour, 2 hours, 3 hours and after 4 hours of the administration of the fig juice; it's measured also on the 5th day, 8th day and 9th day of the beginning of the experiment.

Results and discussions: On the 9th day, we recorded a very significant decrease of the blood sugar level of diabetic rats treated with dry fig juice. This blood glucose level normalized for 3 rats / 5rats. In the short term (for 4 hours), an increase of blood sugar level, one hour after administration, for normal and diabetic rats. This increase is probably due to the high level of sugar content in the preparation. The blood glucose level is then corrected, four hours later. This may be the result of anti hyperglycemic effect of the active ingredients contained in the figs.

Conclusion/Discussion: The figs have despite their high sugar content a significant hypoglycemic activity. Whose mechanism of action still is unknown. The extract aqueous of the figs may imitate the action of insulin, or stimulate the synthesis of insulin.

Keywords: Figs, hypoglycemic activity

References:

[1] Azzi Rachid,,Contribution à l'étude de plantes médicinales utilisées dans le traitement traditionnel du diabète sucré dans l'Ouest algérien : enquête ethnopharmacologique; Analyse pharmaco-toxicologique de Figuier (*Ficus carica*) et de coloquinte (*Citrullus colocynthis*) chez le rat Wistar, departement of biology Telemcen Algeria, 2013

POSTER PRESENTATION

QUALITATIVE AND QUANTITATIVE DETERMINATION OF MONO- AND DICAFFEYOYLQUINIC ACIDS IN *INULA* SPECIES FROM BULGARIA

Antoaneta Trendafilova¹, Miroslav Rangelov¹, Milka Todorova¹

¹ *Institute of Organic Chemistry with Centre of Phytochemistry, Bulgarian Academy of Sciences, 1113 Sofia, Bulgaria; trendaf@orgchm.bas.bg*

Objective: Chlorogenic acid (CQA) and dicaffeoyl esters of quinic acid (DCQA) are widely distributed in Asteraceae family. These compounds are known to possess various biological activities - antioxidant, antiviral, antibacterial, anti-inflammatory activities and can reduce the relative risk of cardiovascular, diabetes type II and Alzheimer's diseases. The aim of this study was to compare HPTLC and HPLC methods for qualitative and quantitative analysis of CQA and four isomeric DCQA in the methanol extracts of six *Inula* species, growing in Bulgaria - *I. aschersoniana*, *I. conyza*, *I. ensifolia*, *I. oculus-christi*, *I. bifrons* and *I. germanica*.

Material and Methods: The aerial parts of the six *Inula* species were collected in full flowering stage from natural habitats in Bulgaria in 2016. Air-dried and powdered plant material was extracted with CH₃OH and the resulting extracts were further used for HPTLC and HPLC analyses. HPTLC was performed on HPTLC plates (Merck) on CAMAG system with toluene/ethyl acetate/formic acid/water (5:25:100:10:10, v/v/v/v) as mobile phase. Densitometric determination was performed at 325 nm. A gradient RP HPLC method was carried out at 325 nm.

Results: A preliminary HPTLC comparison of the methanol extracts with commercial standards showed presence of chlorogenic (CQA), 4,5-, 3,4-, 1,5- and 3,5-dicaffeoylquinic acids (DCQA). On the HPTLC plate, phenolic acids were observed under 366 nm UV light as blue bands at R_f values of 0.26, 0.49, 0.50, 0.63 and 0.66, respectively. Good separation was achieved for the main compounds CQA, 1,5- and 3,5- DCQA, which were further densitometrically determined at 325 nm. Additionally, HPLC method was applied for quantitative determination of CQA and DCQA isomers. The amount of CQA varied from 14.27 to 29.94 mg/g DE and the highest amount was detected in the methanol extract of *I. ensifolia*. 1,5- and 3,5-DCQA were the most abundant dicaffeoyl esters of quinic acid, but differed in their quantity in different species (20.32 - 65.0 and 1.91 - 22.92 mg/g DE, respectively). The content of 4,5- and 3,4-DCQA was relatively low (3.38-9.88 and 0.30-5.11 mg/g DE, respectively). The methanol extract of *I. germanica* was characterized with high amount of 1,5-DCQA. Similarly, *I. conyza* methanol extract was also rich in 1,5-DCQA, but contained subsequent amounts of 3,5-DCQA. Methanol extracts of *I. ensifolia*, *I. aschersoniana*, *I. oculus-christi* and *I. bifrons* contained almost equal amounts of 1,5-DCQA and 3,5-DCQA. There was no statistically significant difference between the mean values for CQA and 1,5- and 3,5-DCQA in studied extracts measured by HPTLC and HPLC.

Conclusion: HPTLC was found to be effective for qualitative analysis, while HPLC was more accurate for quantitative analysis. A combination of the two methods may be useful in a quality control setting, as it would allow rapid qualitative analysis of herbal material while maintaining accurate quantification of extract composition.

Acknowledgements: This work was supported by the NSF, Ministry of Education and Science, Bulgaria, Project DN 09/11.

Key words: *Inula*, Asteraceae, phenolic acids, HPTLC, HPLC-PDA

POSTER PRESENTATION

**TOTAL PHENOLIC CONTENT AND ANTIOXIDANT ACTIVITY OF THE
n-BUTANOL EXTRACT OF *MENTHA ROTUNDIFOLIA* FROM ALGERIA**

Ilhem Semra¹, Mehmet Öztürk², Ahmed Kabouche¹, Zahia Kabouche¹

¹Universite des Frères Mentouri Constantine 1, Laboratoire d'Obtention de Substances Thérapeutiques (LOST),
25000 Constantine, ALGERIA.

² Faculty of Sciences, Department of Chemistry, Mugla Sıtkı Koçman University, Mugla, TURKEY.

Objective / Purpose: The genus *Mentha* (mint) (Lamiaceae family) is represented by 25 to 30 species grown in different parts of the world. Fifteen *Mentha* species are distributed in Algeria [1]. *Mentha* has been applied in the traditional medicine as stimulant, carminative, antispasmodic, antitussive and diuretic. It has been used as antiseptic, antimicrobial and antioxidant agents [2]. Phenolics, flavonoids and terpenoids [3] have been identified from *Mentha* species. As part of our interest in Lamiaceae species [4-6], the objective of the present study was to evaluate total phenolic content and the antioxidant activity, of the *n*-butanol extract of *Mentha rotundifolia* L.

Materials and Methods: *Mentha rotundifolia* was collected at Hammam knif (North eastern Algeria). The powdered aerial part of the plant was extracted by a mixture of ethanol/water 80/20, v/v. The *n*-butanol extract was obtained by liquid-liquid extraction. Total phenolic content was determined with Folin–Ciocalteu reagent. Three antioxidant tests were carried out with DPPH Free Radical Scavenging assay, β -carotene/linoleic acid test and ABTS cation radical decolorization assay.

Results: The *n*-butanol extract of *Mentha rotundifolia* has a high content of polyphenols (356.75 ± 2.58 mg EAG./g). The greatest activity was obtained with the bleaching of β -carotene test (IC_{50} : 1.75 ± 0.37 μ g/mL). An excellent potential with the DPPH free radical scavenging test is also noted (IC_{50} : 9.33 ± 1.42 μ g/mL). This extract also possesses a good antiradical activity towards the ABTS⁺ radical (IC_{50} : 9.12 ± 0.74 μ g/mL).

Conclusion/Discussion: The *n*-butanol extract of *Mentha rotundifolia* L. possesses a great antioxidant potential with the three tests carried out (DPPH Free Radical Scavenging, β -Carotene/Linoleic Acid Bleaching and ABTS cation radical decolorization). The high content of polyphenols, in this extract, certainly contributes to this great antioxidant power.

Keywords: *Mentha rotundifolia*, antioxidant activity, DPPH, CUPRAC, ABTS⁺.

References:

- [1] Quezel P, Santa S.1963. ed. Centre National de la Recherche Scientifique. Tome II, Paris.
- [2] Ladjel S, Gherraf N, Hamada D. 2011. Journal of Applied Sciences Research. 7: 1665-7.
- [3] Phatak S.V, Heble M.R. 2002. Fitoterapia. 73:32- 39.
- [4] Kabouche A, Kabouche Z, Ozturk M, Kolak U, Topçu G. 2007. Food Chemistry 102 : 1281–1287.
- [5] Berrehal D, Boudiar T, Lakhel H, Khalfallah A, Kabouche A, Al-Freihat A, Ghannadi A, Sajjadi E, Mehrabani M, Safaei-Ghomi J, Kabouche Z. 2010. Natural Product Communications. 5 (6): 957-960.
- [6] Ferhat M, Kabouche Z, Fujimoto Y, Araya H. 2017. Natural Product Communications. 12(4): 483–486.

POSTER PRESENTATION

**RADICAL SCAVENGING AND ANTI-BACTERIAL ACTIVITIES OF
METHANOLIC EXTRACT FROM ALGERIAN FIR (*ABIES NUMIDICA*) LEAVES**

Inès Bellil, Djamila Benouchenne, Douadi Khelifi

*Laboratory of Genetic Biochemistry and Plant Biotechnologies, Faculty of Natural Sciences and Life,
University of Mentouri Brothers Constantine 1, 25000 Constantine, Algeria.
dbenouchenne1992@gmail.com, dkhelifi@yahoo.fr, bines07@yahoo.fr*

Objective / Purpose: Algerian forests are rich in plants due to its geographic position. It represents a high biodiversity that seems in different found species that are endemic in major, for example Algerian fir (*Abies numidica*) tree. It is a conifer belongs to *Pinaceae* family, used in folk medicine to treat cataplasm and respiratory system diseases. Nowadays, there is a huge interest to find new molecules from natural resources to prevent human diseases. For that, the objective of the present research is to investigate anti-oxidant and anti-bacterial activities of Algerian fir leaves crude extract.

Material and Methods: The air dried powdered leaves were extracted with methanol solvent using Soxhlet apparatus. The extract was evaporated under reduced pressure by rotary vacuum-evaporator type Buchi R-215 at 40°C. DPPH and ABTS assays were used to evaluate the anti-oxidant activity. The Anti-bacterial activity was also tested using the disc diffusion method against five bacterial strains; *Escherichia coli*, *Pseudomonas aeruginosa*, *Morganella morgani*, *Staphylococcus aureus* and *Bacillus subtilus*.

Results: The results showed that methanolic extract of Algerian fir leaves by DPPH method can reduce radicals ($IC_{50}=18.08\pm 0.21$) lower than ABTS ($IC_{50}=23.51\pm 0.54$). The methanolic extract exhibited no anti-bacterial activity against Gram negative bacteria; *Escherichia coli*, *Pseudomonas aeruginosa* as well as *Morganella morgani*. However, it represented a moderate activity against Gram positive; *Staphylococcus aureus* and *Bacillus subtilus*.

Conclusion / Discussion: *Abies numidica* is a threatened species in Algeria and according to the results obtained, it has a beneficial effect. So it is very important to protect it and exploit it in pharmaceutical mains and use it in food industry.

Keywords: methanolic extract, Algerian fir, radical scavenging, anti-bacterial activity.

POSTER PRESENTATION

CHEMICAL COMPOSITION AND BIOLOGICAL ACTIVITIES OF THE ESSENTIAL OILS FROM TWO CHEMOTYPES OF *CYMOPOGON SCHOENANTHUS* (L.) SPRENG. GROWING IN SOUTHERN ALGERIA.

Naima Hellali^{1,2}, Mahfoud Hadj Mohammed², Mebarka Bouziane²

¹ Faculty of Sciences and Technology, Department of process engineering, Ghardaïa University, BP 455, 47000 Ghardaïa, Algeria.

²Biogeochemistry Laboratory in Desert Environments, Faculty of Science, Kasdi Merbah University, BP 511, 30000 Ouargla, Algeria.

*Corresponding Author. E-mail: hellali.naima@univ-ghardaia.dz

The chemical composition of the essential oils of two populations of *Cymbopogon schoenanthus* collected in Ghardaïa and Illizi (Southern Algeria), were identified by (GC-MS), this study allowed us to identify the presence of two chemotypes of *C. schoenanthus* essential oil, the first one was characterized by the high content of piperitone, and the second group was characterized by other compounds. The screening of antimicrobial activity of oils was individually evaluated against representatives of gram-positive, gram-negative bacteria and fungi, using the agar diffusion method, whereas, the antioxidant activity of the essential oils was evaluated using the DPPH, ABTS and FRAP assay. The results of in vitro testing of the antimicrobial activities of *C. schoenanthus* essential oil of Ghardaïa region proved the existence of strong activity against all the tested strains, specially the growth of *E. faecium*, *S. aureus* and *S. agalactiae*. However, the same species growing in Illizi region shown less power in the inhibition of microorganisms. The essential oil obtained from *C. schoenanthus* growing in Illizi is markedly rich in non-phenolic constituents, due to this fact the essential oil is relatively weak antioxidant, which the same remark in the essential oil extracted from Ghardaïa.

Key words: *C. schoenanthus*, essential oil, chemotype, antimicrobial activity, antioxidant activity.

References:

- [1]. Alitonou GA, Avlessi F, Tchobo F, Noudogbessi JP, Tonouheva A., Yehouenou B, Menut C, Sohounhoute DK (2012). Chemical composition and biological activities of essential oils from the leaves of *Cymbopogon giganteus* Chiov. and *Cymbopogon schoenanthus* (L.) Spreng (Poaceae) from Benin. Int. J. Biol. Chem. Sci. 6(4):1819-1827.
- [2]. Avoseh O, Oyedeji O, Rungqu P., Nkeh-Chungag B and Oyedeji A., (2015). *Cymbopogon* Species; Ethnopharmacology, Phytochemistry and the Pharmacological Importance, *Molecules* 2015, 20, 7438-7453.
- [3]. Benzie IFF, Strain JJ (1996). The ferric reducing ability of plasma (FRAP) as a measure of "antioxidant power": the FRAP assay. Anal. Biochem. 239(1):70-76.
- [4]. Binsan W, Benjakul S, Visessanguan W, Roytrakul S, Tanaka M, Kishimura H (2008). Antioxidative activity of Mungoong, an extract paste, from the cephalothorax of white shrimp (*Litopenaeus vannamei*). Food Chem. 106(1):185-193.
- [5]. Bossou AD, Ahoussi E, Ruysbergh E, Adams A, Smaghe G, Kimpe ND, Avlessi F, Sohounhoute DCK, Mangelinckx S (2015). Characterization of volatile compounds from three *Cymbopogon* species and *Eucalyptus citriodora* from Benin and their insecticidal activities against *Tribolium castaneum*. Ind. Crops Prod. 76:306-317.
- [6]. Brand-Williams W, Cuvelier ME, Berset C (1995). Use of a free radical method to evaluate antioxidant activity. LWT-Food Sci. Technol. 28:25-30.
- [7]. Čavar S, Maksimović M, Vidica D, Parić A (2012). Chemical composition and antioxidant and antimicrobial activity of essential oil of *Artemisia annua* L. from Bosnia. Ind. Crops Prod. 37:479-485
- [8]. Chen Z, Berini R, Frolidi G (2013). EC50 estimation of antioxidant activity in DPPH assay using several statistical programs. Food Chem. 138:414-420.
- [10]. Gohari AR, Hajimehdipoor H, Saeidia S, Ajani Y, Hadjiakhoondi A (2011). Antioxidant Activity of some Medicinal Species using FRAP Assay. J. Med. Pla. 10(37):55-60
- [11]. Hadeif D, Saidi M, Youssi M, Moussacoti Y (2015). GC/MS Analysis of essential oils of *Cymbopogon schoenanthus* and *Origanum majorana* L. grown in eastern Algeria. Asian J. Chem. 27(10):3575-3578.
- [12]. Hammiche V, Maiza K (2006). Traditional medicine in Central Sahara: Pharmacopoeia of Tassili N'ajjer. J. Ethnopharmacol. 105:358-367.
- [13]. Hellali N, Hadj-Mahammed M (2008). Analyse par GC-MS de l'huile essentielle de la plante *Cymbopogon schoenanthus* de la région de Illizi, obtenue par hydrodistillation assistée par microondes. PhytoChem. BioSub J. 2(1):5.
- [14]. Hellali N, Hadj Mahammed M, Ramdane F, Talli A (2016). Antimicrobial and antioxidant activities of *Cymbopogon schoenanthus* (L.) spreng. essential oil, growing in Illizi – Algeria. J. Med. Pla. Res. 10(14):188-194.
- [15]. Ketoh GK, Koumaglo HK, Gliho IA (2005). Inhibition of *Callosobruchus maculatus* (F.) (Coleoptera: Bruchidae) development with essential oil extracted from *Cymbopogon schoenanthus* L. Spreng (Poaceae), and the wasp *Dinarmus basalis* (Rondani) (Hymenoptera: Pteromalidae). J. Stored Prod. Res. 41:363-371.
- [16]. Ketoh GK, Koumaglo HK, Gliho IA, Huiagnard J (2006). Comparative effects of *Cymbopogon schoenanthus* essential oil and piperitone on *Callosobruchus maculatus* development. Fitoterapia 77:506-510.
- [17]. Khadi A, Serralheiro MLM, Nogueira JMF, Neflati M, Smiti S, Arau' jo MEM (2008). Antioxidant and antiacetylcholinesterase activities of essential oils from *Cymbopogon schoenanthus* L. Spreng. Determination of chemical composition by GC-mass spectrometry and 13C NMR. Food Chem. 109: 630–637.
- [18]. Koba K, Sanda K, Raynaud C, Nonenene YA, Millet J, Chaumont J (2004). Activités antimicrobienne de trois *Cymbopogon* sp. Africains vis-à-vis de germes pathogènes d'animaux de compagnie. Ann. Med. Vet. 148:202-206.
- [19]. Lopes-Lutz D, Alviano DS, Alviano CS, Kolodziejczyk PP (2008). Screening of chemical composition, antimicrobial and antioxidant activities of Artemisia essential oils. Phytochemistry 69(8):1732-1738.
- [20]. Maiza K., Brac De La Perriere R. A., Hammiche V., Médicaments et aliments : l'approche ethnopharmacologique, (1993) p171.
- [21]. Mohd AAN, Husni S, Almajano PM, Gallego GM (2013). Solvent Effect on Antioxidant Activity and Total Phenolic Content of *Betula alba* and *Convolvulus arvensis*. Intern. J. Biol. Biom. Agri. Food Biotech. Engi.7(5) scholar.waset.org/1999.1/9996837.
- [22]. National Committee for Clinical Laboratory Standards (NCCLS) (1999). Approved standard M100-S9: performance standards for antimicrobial susceptibility testing. Wayne, Pennsylvania, USA.
- [23]. Neflati A, Bouhélil I, Ben Sghaier M, Boubakker J, Limem J, Bhouiri W, Le Dauphin J, Barillier D, Mosrati R, Chekir-Ghedira L, Ghedira K (2009). Antigenotoxic and antioxidant activities of *Pituranthos chloranthus* essential oils. Environ. Toxicol. Pharmacol. 27:187-194.
- [24]. Onadja Y, Ouédraogo A, Dorosso SA (2007). Chemical composition and physical characteristics of the essential oil of *Cymbopogon schoenanthus* (L.) Spreng of Burkina Faso. J. Appl. Sci. 7(4):503-506.
- [25]. Re R, Pellegrini N, Pannala A, Yang M, Rice-Evans C (1999). Antioxidant activity applying an improved ABTS radical cation decolorization. Free Radic. Biol. Med. 26(9-10):1231-7.
- [26]. Shi F, Jia X, Zhao C and Chen Y (2010). Antioxidant Activities of Various Extracts from *Artemisia selengensis* Turcz (LuHao). *Molecules* 15:4934-4946.
- [27]. Sousa EMBD, Câmara APC, Costa WA, Costa ACJ, Oliveira HNM, Galvão EL, Marques MMO (2005). Evaluation of the Extraction Process of the Essential Oil from *Cymbopogon schoenanthus* with Pressurized Carbon Dioxide. Brazilian Arch. Bio. Tech. 48:231-236.
- [28]. Trombetta D, Castelli F, Sarpietro M G, Venuti V, Cristani M, danielle C, Saija A, Mazzanti G, Bisignano G (2005). Mechanisms of antibacterial action of three monoterpenes, Antimic. Agen. Chemother. 49(6):2474–2478.
- [29]. Tyagi AK, Malik A (2010). In situ SEM, TEM and AFM studies of the antimicrobial activity of lemon grass oil in liquid and vapour phase against *Candida albicans*. Micron 41:797–805.

POSTER PRESENTATION

EVALUATION OF THE ANTIOXIDANT POTENTIAL, PHENOLIC AND FLAVONOID CONTENTS OF THE METHANOL EXTRACT OF *AMMOIDES ATLANTICA*

Mounira Benteldjoune¹, Tarek Boudiar², Abdelhakim Bakhouché³, María del Mar Contreras³, Jesús Lozano-Sánchez³, Chawki Bensouici², Antonio Segura-Carretero³, Zahia Kabouche¹

¹Université des frères Mentouri-Constantine 1, département de chimie, Laboratoire d'Obtention de Substances Thérapeutiques (L.O.S.T), 25000 Constantine, Algeria

²Biotechnology Research Centre (C.R.Bt), Ali Mendjli Nouvelle Ville BPE.73 Constantine, Alegria.

³Research and Development of Functional Food Centre (CIDAF), Bioregion building, PTS Granada, Avda. del

Objective / Purpose: The genus *Ammoides* (Apiaceae), tribe of Ammineae, includes two species in Algeria, from which the endemic species *A. atlantica* (Coss. Et Dur.) Wolf [1]. In Algerian traditional medicine, the aerial parts of *A. atlantica* are used in infusions for the treatment of headache, fever and diarrhea. It is also used in compresses, alone or soaked in alcohol or vinegar and mixed with henna, to treat children affected by mental debility [2].

Materials and Methods: The plant was collected during the flowering period in the North Eastern region of Algeria (Jijel) in May 2016. The aerial parts of *Ammoides* were extracted with methanol-water (80:20, v/v). *In-vitro* antioxidant activity was evaluated for the methanol extract using different antioxidant tests, namely, total antioxidant (DPPH scavenging, (ABTS⁺) scavenging, superoxide radical scavenging by alkaline DMSO, reducing power, cupric reducing antioxidant capacity (CUPRAC) and ferrous ions chelating assays compared to the synthetic antioxidants (BHA, BHT and ascorbic acid).

Results: The methanol extract of *Ammoides atlantica* exhibited a high antioxidant activity in DPPH assay (IC₅₀: 23.31 ± 0.99 µg/mL), ABTS⁺ (IC₅₀: 11.31 ± 2.49 µg/mL), O₂⁻ DMSO alkaline assay (IC₅₀: 3.19±0.17 µg/mL), Reducing power assay (A_{0.50} : 92.70±1.37 µg/mL), CUPRAC assay (A_{0.50}: 13.56±1.06 µg/mL) and Ferrous ions chelating assay (IC₅₀:102.35±2.91 µg/mL), respectively compared with BHA, BHT and ascorbic acid which were used as positive controls. Total Phenolic content (371.57±7.80 mg/g) and Total Flavonoid content 41.02±6.16 mg/g were also interesting

Conclusion / Discussion: From the results, *Ammoides atlantica* can be used as a source of antioxidants

Keywords: Apiaceae, *Ammoides atlantica*, antioxidant activity, DPPH, CUPRAC, ABTS⁺.

References:

- [1] Quezel, P., Santa, S. (1963). *Nouvelle Flore de l'Algérie et des régions désertiques méridionales*. Vol.2. Ed. CNRS, Paris France.
[2] Bellakhdar J. (1997). *La pharmacopée marocaine traditionnelle*. Casablanca/Paris : Éditions le Fennec /Ibis Press, p.739.

POSTER PRESENTATION

**EXTRACTION OF NATURAL PRODUCT FROM PLANT BY USING A
MICROWAVE**

Susmita Gudulkar & Usha Mukundan

*Department of Botany, Ramniranjan Jhunjhunwala College (Autonomous), Affiliated to University of Mumbai,
Ghatkopar (W), Mumbai 400 086, India.*

Authors email address: - gudulkar.2@gmail.com

Objective/Purpose: Plants have unique range of fine chemicals known as secondary metabolites. Shikimic acid is found widely in plants but generally occurs in low concentration. It is a hydro aromatic intermediate of common pathway of aromatic amino acid biosynthesis. This work basically discusses the extraction of natural product that is shikimic acid from plant material by using a microwave. Shikimic acid has several benefits including its use in treating fungal infection and act as antioxidant, antibacterial, antiviral medication.

Material and Methods: Plant materials like seeds of Chinese star anise (*Illicium verum*) can be a source of shikimic acid. Shikimic acid extracted for 10, 20, 30, 40, 50 and 60 seconds by using micro oven. It is cheapest and simplest method and less time consuming.

Results: Most plants contain water in more amounts so this is effective means of extraction. Plants with low water content take longer to heat in microwave. The temperature of the plant material heated in a microwave cannot exceed the boiling point of water (100°C). Hence, the shikimic acid is in stable form.

Conclusion: Microwave can be used to isolates flu drug raw material from star anise while minimising pigment contamination. Shikimic acid can be extracted easily, and optimization of base material for production of *oseltamivir (Tamiflu)*, which is an effective drug against the H5N1 influenza virus. Shikimate can also be used to synthesize *(6S)-6 Fluoroshikimic acid*, an antibiotic which inhibits the aromatic biosynthetic pathway.

Key words: Shikimic acid, star anise, microwave.

References:

- Quiroz, D. C. D., Carmona, S. B., Bolívar, F., & Escalante, A. (2014). Current perspectives on applications of shikimic and aminoshikimic acids in pharmaceutical chemistry. *Res Rep Med Chem*, 4, 35-46.
- M Estevez, A., & J Estevez, R. (2012). A short overview on the medicinal chemistry of (—)-shikimic acid. *Mini reviews in medicinal chemistry*, 12(14), 1443-1454.

POSTER PRESENTATION

URSOLIC ACID AS A BIOCHEMICAL MARKER IN FAMILY MYRTACEAE

Nidhi Pathak , Himanshu Dawda and Usha Mukundan

Department of Botany, Ramniranjan Jhunjhunwala college (Autonomous), Affiliated to University of Mumbai, Ghatkopar (W), Mumbai 400 086, India.

Authors email address: - pathaknidhi932087@gmail.com

Objective /Purpose:-Over the decades, biochemistry has become remarkably important to resolve the dispute of different taxon in plants. Different chemical markers can be used to differentiate plant families, genus and species, Ursolic acid a potentially important molecule from the family Myrtaceae can be used for the better understanding of different taxon. Apart from taxonomic possibilities, it also shows many pharmacological effects such as Reducing oxidative stress, anti-Inflammation, Anti-oncogenic, Inducing Apoptosis and hepato-protective nature.

Material and method: A detailed study on secondary metabolites was observed by screening different plants from *Myrtaceae*. HPTLC method was performed for the qualitative and quantitative analysis of different compounds mainly ursolic acid.

Results: HPTLC analysis resulted in presence of many secondary compounds in association with ursolic acid. Using pairing affinity for different molecules dendrogram was prepared to understand a phylogenetic relation between species of the family *Myrtaceae*.

Conclusion: Ursolic acid was found to be present in family *Myrtaceae* as a biochemical Marker.

Key words: Ursolic acid, *Myrtaceae*, Biochemical compound, HPTLC.

References:

Wink M., Waterman P., (1999). Chemotaxonomy in relation to molecular phylogeny of plants. In: "Biochemistry of plant secondary metabolism" (M. Wink, ed.), Sheffield Academic Press and CRC Press. Annual Plants Review 1999; 2:300-341.

POSTER PRESENTATION

DETERMINATION OF ESSENTIAL OIL COMPOSITION, PHENOLIC AND FLAVANOID CONTENTS OF *PHLOMIS LONGIFOLIA* VAR. *LONGIFOLIA* COLLECTED FROM HATAY DISTRICT OF TURKEY

Orçun Çınar¹, Murat Tunçtürk², Tuba Seçmen¹, Ayşe Erdoğan¹, Ünal Karık³

¹Department of Food Technology and Medicinal, Aromatic Plants, Bati Akdeniz Agricultural Research Institute, Antalya, Turkey

²Department of Field Crops, Faculty of Agriculture, Van Yüzüncü Yıl University, Van, Turkey, murattuncturk@yyu.edu.tr

³ Department Of Medicinal Plants, Aegean Agricultural Research Institute, İzmir, Turkey

Objective / Purpose: *Phlomis* is a large genus in the *Lamiaceae* family, with over 100 species distributed throughout Euro-Asia and North Africa countries. Their flowered parts are generally used as an herbal tea to treat gastrointestinal troubles and to promote good health by protecting the liver, kidney, bone and cardiovascular system. Phytochemical investigations of *Phlomis* species were the subject of several studies, and, consequently, essential oils, flavonoids, iridoids, phenylethylalcohol glycosides and other components were isolated.

Material and Methods: *Phlomis longifolia* var. *longifolia* was collected from Hatay district of Turkey. Its essential oil was obtained by using Clevenger apparatus. Essential oil composition was determined by using GC-MS/FID device. Total phenolic and flavanoid contents were determined by using spectrophotometer device.

Results: Essential oil rate was determined as 0,33%. The main essential oil components were determined as germacrene (34,27%), β -caryophyllene (20,97%), *trans*- β -farnesene (7,76%), *trans*-cadine-1,4-diene (6,79%) and α -copaene (5,61%). Total phenolic content was determined as 11,87 mg GAE/g plant. Total flavanoid content was determined as 10,26 mg CE/g plant.

Conclusion / Discussion: *Phlomis longifolia* essential oil can be used for cosmetic industry because of germacrene which is a good-scent main component.

Keywords: *Phlomis*, essential oil composition, total phenolic and flavanoid content

POSTER PRESENTATION

EFFECTS OF MULBERRY (*MORUS ALBA* L.) LEAF EXTRACT (Immunflex®) ON MMP-9, BMP-2 AND COMP IN HUMAN OSTEOARTHRITIC CHONDROCYTES *IN VITRO*

Zehra Aydin Bek¹, Zubeyir Elmazoglu¹, Berna Goker², Berivan Bitik³,
Cem Nuri Aktekin⁴, Cimen Karasu¹

1. Department of Medical Pharmacology, Faculty of Medicine, Gazi University, Ankara Turkey

2. Department of Rheumatology, Faculty of Medicine, Gazi University, Ankara Turkey

3. Department of Rheumatology, Faculty of Medicine, Baskent University, Ankara Turkey

4. Department of Orthopaedic Surgery, Faculty of Medicine, Yildirim Beyazıt University, Ankara Turkey

Objective: Osteoarthritis (OA) is a common disease of older individuals that is characterized by loss of articular cartilage. In OA, an inflammatory process occurs and cytokines and enzymes develop that further damage the cartilage. Matrix metalloproteinases (MMP-9, MMP-13 etc) are proteolytic enzymes, responsible for destruction of collagen and matrix components.¹ COMP, is structural protein of extracellular matrix, is expressed in higher level in OA patients.² Bone morphogenetic protein-2 (BMP-2) signaling is involved in maintaining the normal function of synovial joints.³ The redox modulatory regimens, anti-inflammatory and anti-oxidant rich diet, can help to improve OA. The mulberry plant contains a variety of phenolic compounds, steroids and triterpenes. *Morus alba* leaf extracts have been shown to have anti-inflammatory and antioxidant effects in a different cell type.⁴ Based on these findings, we aim to investigate the potential anti-osteoarthritis and chondroprotective effects of two different *Morus alba* leaf extracts in human OA chondrocytes.

Methods: Primary chondrocytes were isolated from cartilage tissue of OA patients (grade 4). Primary chondrocytes were treated with different concentrations of (1) aqueous and (2) ethanolic extract of *Morus alba* leaf extracts as Immunflex®. Cell viability (MTT assay, neutral red) and ROS levels (DCFH-DA assay) were tested. MMP-9, COMP and BMP-2 protein expression levels were examined by Western blot.

Results: We found that both ethanolic and aqueous extracts had no cytotoxicity at lower concentrations. When compared to untreated OA chondrocytes, ethanolic extract treated chondrocytes showed a reduced ROS level significantly ($p < 0.01$). Ethanolic extract also has a potential modulating effect on MMP-9, COMP and BMP-2.

Conclusion: The results provide evidence for the potential use of Immunflex® as an anti-osteoarthritis complex against OA pathology.

Keywords: *Morus alba*, mulberry leaf, osteoarthritis, primary chondrocytes.

Acknowledgements: This study was financially supported by the Scientific and Technological Research Council of Turkey (Grant No: 31S012)

References:

- Permatasari DA, *et al.*, Quercetin prevents proteoglycan destruction by inhibiting matrix metalloproteinase-9, matrix metalloproteinase-13, a disintegrin and metalloproteinase with thrombospondin motifs-5 expressions on osteoarthritis model rats. *J Adv Pharm Technol Res.* 2019 Jan-Mar;10(1):2-8.
- Tseng S, *et al.*, Cartilage Oligomeric Matrix Protein (COMP): A Biomarker of Arthritis. *Biomark Insights.* 2009 Feb;4:33-44.
- Ashraf S, *et al.*, Regulation of senescence-associated signaling mechanisms in chondrocytes for cartilage tissue regeneration. *Osteoarthritis Cartilage.* 2016 Feb;24(2):196-205.
- Choi EM and Hwang JK. Effects of *Morus alba* leaf extract on the production of nitric oxide, prostaglandin E2 and cytokines in RAW264.7 macrophages. *Fitoterapia.* 2005 Dec;76(7-8).

POSTER PRESENTATION

ASTRAGALOSIDE IV AND FORSYTHOSIDE B MODULATE HYPERGLYCEMIA-MEDIATED NEUROTOXICITY IN PRIMARY HIPPOCAMPAL NEURONAL CELLS

Zubeyir Elmazoglu¹, Zeynep Inan², Cansel Keskin², Naciye Bozkurt³,
İhsan Calis⁴, Cimen Karasu¹

¹ Department of Medical Pharmacology, Faculty of Medicine, Gazi University, Ankara, Turkey, ² Department of Biology, Gazi University, Ankara, Turkey

³ Faculty of Medicine, Gazi University, Ankara, Turkey,

⁴ Department of Pharmacognosy, Faculty of Pharmacy, Near East University, TRNC.

Background: Recent studies have revealed that cognitive deficits and hippocampal abnormalities are the complications of Diabetes Mellitus (DM) that is characterized by chronic hyperglycemia, hyperlipidemia, insulin resistance, oxidative stress and other alterations [1]. The well known mechanisms for the progression of hippocampal neuronal loss in DM, includes oxidative damage and mitochondrial dysfunction [2]. Unfortunately, current therapeutic strategies are not sufficient enough to overcome hyperglycemia induced neurotoxicity. So, we aimed to explore the effects of Astragaloside IV (AS-IV) and Forsythoside B (FST-B) on high glucose induced neurodegeneration in hippocampal neurons.

Methods: Primary hippocampal neurons were isolated from Wistar albino pups (E17-19). Cells were exposed high glucose (150 mM) for 16 h to induce glucotoxicity. For the neuroprotection experiments cells were pre-treated with various concentrations (1 nM-1 µM) of AS-IV or FST-B for 8 h, then post-treated with high glucose (150 mM) for 16 h. Cell viability was assessed both MTT and NRU assay. Moreover cytotoxicity and intracellular ROS were monitored using Trypan blue exclusion assay and fluorescent probe DCFHDA, respectively.

Results: Our results indicated that 150 mM glucose exposure caused significantly decreased cell viability ($p < 0.01$) along with a significant increased toxicity ($p < 0.05$) and high ROS generation levels ($p < 0.01$). On the other hand, pre-treatment with AS-IV and FST-B ameliorated the glucose induced neurotoxicity (> 50 nM) in terms of increased viability (1 nM-500 nM), improved membrane integrity and lowered the ROS generation capacity (≥ 50 nM).

Conclusion: We demonstrated that AS-IV and FST-B showed neuroprotective effects against hyperglycemia mediated neurotoxicity via inhibiting neuronal loss, strengthen membrane integrity and impairing oxidative damage.

Keywords: Hyperglycemia, hippocampal neurons, Astragaloside IV, Forsythoside B, neuroprotection.

Acknowledgements: This study was financially supported by the Scientific and Technological Research Council of Turkey (Grant No: 315S088 and 215S197)

References:

1. Zhao L, *et al.*, Characteristic Metabolic Alterations Identified in Primary Neurons Under High Glucose Exposure. *Front Cell Neurosci.* 2018 Jul 17;12:207.
2. Liu D, *et al.*, Effects of exposure to high glucose on primary cultured hippocampal neurons: involvement of intracellular ROS accumulation. *Neurol Sci.* 2014 Jun;35(6):831-7.

POSTER PRESENTATION

**THE ESSENTIAL OIL YIELD AND COMPOSITIONS OF LEMON VERBANA
(*LIPPIA CITRODARA* KUNTH.) CULTIVATED IN ORDU**

M. Muharrem Özcan¹ Ayşegül Kırılı¹ Özbay Dede¹ İrem Ayran² S.Ayşe Çelik² Yüksel Kan²

¹Ordu University, Agriculture Faculty, Department of Field Crops, Ordu/Turkey

²Selçuk University, Agriculture Faculty, Department of Medicinal Plants, Konya/Turkey

e-mail: irem.ayran@gmail.com, phone + 90-505 939 31 35

Objective / Purpose: In the study, essential oil yield and its components leaves obtained from lemon verbana (*Lippia citrodara* Kunth.) cultivated in Ordu were investigated.

Material and Methods: The air-dried leaves of lemon verbana were subjected to hydrodistillation for 3h using a Clevenger-type apparatus to produce essential oil. The GC-MS analysis was carried out with Agilent 7890 GC-MS system. The relative percentages of the separated compounds were calculated from total ion chromatograms. The identification of the oil components was based on the Wiley and NIST mass spectral library.

Results: The essential oil yield of lemon verbana was determined as 1.45%. The most important essential oil component of Geranial was 20.8 %. The major essential oil components of lemon verbana were determined as Limonen (18.1 %), Neral (13.8 %), respectively.

Conclusion / Discussion: According to the results of this study, it was determined that essential oil yield and components of lemon verbana (*Lippia citrodara* Kunth.) were determined. In addition to the use of essential oil, leaves of the plant are used as herbal tea. Lemon verbana is used appetizing, relaxing the stomach, colds, cough and digestion problems in folk medicine.

Keywords: Lemon verbana, *Lippia citrodara* Kunth., essential oil, composition

POSTER PRESENTATION

THE ESSENTIAL OIL YIELD AND COMPOSITIONS OF LAURIER (*LAURUS NOBILIS L.*) CULTIVATED IN SAMSUN

İrem Ayran¹ S.Ayşe Çelik¹ Asuman Kan² Yüksel Kan¹

¹Selçuk University, Agriculture Faculty, Department of Medicinal Plants, Konya/Turkey

²Konya Technical University, Technical Sciences Vocational School of Higher Education, Department of Food Processing, Konya/Turkey

e-mail: irem.ayran@gmail.com phone + 90-505 939 31 35

Objective / Purpose: In the study, essential oil yield and its components obtained from leaf of laurier (*Laurus nobilis L.*), cultivated in Samsun ecological conditions were investigated.

Material and Methods: The air-dried leaves of laurier were subjected to hydrodistillation for 3h using a Clevenger-type apparatus to produce essential oil. The GC-MS analysis was carried out with Agilent 7890 GC-MS system. The relative percentages of the separated compounds were calculated from total ion chromatograms. The identification of the oil components was based on the Wiley and NIST mass spectral library.

Results: Essential oil yield was determined as 0.4 % in dried leaves. The major essential oil component of 1,8-cineol was 45,7 %. The other major essential oil components of laurier leaf were determined as α -Terpinyl acetate (15.6 %), sabinen (8.4 %) , α -pinene (4.8 %) and β -pinene (4.1 %) , respectively.

Conclusion / Discussion: According to the results of this study, it was determined that essential oil yield and components of laurier (*Laurus nobilis L.*)leaf were determined. Essential oil yield and compositions of laurier leaf were one of the significant quality factors. Laurier leaf are used to antibacterial, diaphoretic, pain relieving, discomfort, antiseptic and stomach discomfort,diabetic, migraine, weakness in folk medicine. At the same time, Laurier is used as spices in the worldwide.

Keywords: Laurier, *Laurus nobilis L.*, essential oil, composition, 1,8-cineole

POSTER PRESENTATION

**A STUDY ON ESSENTIAL OIL YIELD and COMPONENTS OF LEMON BALM
(*MELISSA OFFICINALIS* L.) CULTIVATED DIFFERENT PROVINCES IN
TURKEY**

S.Ayşe Çelik¹, İrem Ayran¹, Asuman Kan², Emine Bİlginoğlu¹, Yüksel Kan¹

¹*Selçuk University, Agriculture Faculty, Department of Medicinal Plants, Konya/Turkey*

²*Konya Technical University, Technical Sciences Vocational School of Higher Education, Department of Food Processing, Konya/Turkey*

e-mail: sadiyeyse.sayar@gmail.com, phone + 90-539 799 33 63

Objective / Purpose: In the study, essential oil yield and its components obtained from Lemon balm (*Melissa officinalis*) cultivated different provinces (in Konya and Elazığ) were investigated.

Material and Methods: Lemon balm belonging to the family Lamiaceae were subjected to hydrodistillation for 3 h using a Clevenger-type apparatus to produce essential oil. The GC-MS analysis was carried out with Agilent 7890 GC-MS system. The relative percentages of the separated compounds were calculated from total ion chromatograms. The identification of the oil components was based on the Wiley and NIST mass spectral library.

Results: The essential oil yield of lemon balm was 0,34 % in Konya province, while It was 0.30 % in Elazığ province. The major essential oil components of lemon balm were determined as Geranial. The amount of geranial in Konya province and in Elazığ province were 33.4% and 46.0%, respectively.

Conclusion / Discussion: The aim of this research attempts to contribute to knowledge of differences essential oil yield and components of lemon balm produced in different provinces of our country were determined. In addition to used as essential oil, leaves of the plant are used as herbal tea. Lemon balm is used sedative, gastric, antitussive, diaphoretic and antiseptic effects in folk medicine.

Keywords: Lemon balm, *Melissa officinalis* L., essential oil, component

POSTER PRESENTATION

THE ESSENTIAL OIL YIELD and COMPOSITIONS of *LAVANDULA ANGUSTIFOLIA* MILLER and *LAVANDULA INTERMEDIA* EMERIC EX LOISEL. CULTIVATED IN BAYBURT ECOLOGICAL CONDITIONS

S.Ayşe Çelik¹ İrem Ayran¹ Asuman Kan² Yüksel Kan¹

¹Selçuk University, Agriculture Faculty, Department of Medicinal Plants, Konya/Turkey

²Konya Technical University, Technical Sciences Vocational School of Higher Education, Department of Food Processing, Konya/Turkey

e-mail: sadiveayse.savar@gmail.com phone + 90-539 799 33 63

Objective / Purpose: In this study, it was investigated essential oil yield and compositions of *Lavandula angustifolia* Miller and *Lavandula intermedia* Emeric ex Loisel., belong to Lamiacea, cultivated in Bayburt ecological conditions, Turkey.

Material and Methods: The air-dried flowers of lavender species were subjected to hydrodistillation for 3h using a Clevenger-type apparatus to produce essential oil. The GC-MS analysis was carried out with Agilent 7890 GC-MS system. The relative percentages of the separated compounds were calculated from total ion chromatograms. The identification of the oil components was based on the Wiley and NIST mass spectral library.

Results: Essential oil yield of lavender species was determined as 1.5 % in *Lavandula angustifolia* Miller and 6.5 % in *Lavandula intermedia* Emeric ex Loisel. The most important essential oil component of *Lavandula angustifolia* Miller were linalol (42.8 %) linalyl acetate (13.9 %) and borneol (2.6 %) and *Lavandula intermedia* were determined as linalol (45.4 %), linalyl acetate (22.2%) and borneol (4.6 %). The amount of camphor of *Lavandula intermedia* Emeric ex Loisel. was determined 2.4%.

Conclusion / Discussion: According to the results of this study, it was determined that significant amount essential oil yield and components of the lavender species (*Lavandula angustifolia* Miller and *Lavandula intermedia* Emeric ex Loisel. were determined. Lavender varieties are promoted primarily in essential oil as an linalol and linalil aasetat that give fragrance and also used antimicrobial, antioxidant and similar aims.

Keywords: Lavender, *Lavandula angustifolia* Miller, *Lavandula intermedia* Emeric ex Loisel., essential oil, composition.

POSTER PRESENTATION

THE ESSENTIAL OIL YIELD and COMPOSITIONS of FENNEL (*Foeniculum vulgare* MILL.) CULTIVATED in KONYA, TURKEY

Halil İbrahim Polat¹ Esra İnfal¹ Emine Saliha Toprak¹ S.Ayşe Çelik¹ İrem Ayran¹

¹*Selçuk University, Agriculture Faculty, Department of Medicinal Plants, Konya/Turkey*
e-mail: salihaemine@gmail.com, phone: + 90-541 712 80 95

Objective / Purpose: In this study, it was investigated essential oil yield and compositions of fennel (*Foeniculum vulgare* Mill.) cultivated in Medicinal and Aromatic Plants Farm of Selçuk University, in Konya, Turkey.

Material and Methods: The air-dried seeds of *Foeniculum vulgare* Mill. were subjected to hydrodistillation for 3 h using a Clevenger-type apparatus to produce essential oil. The GC-MS analysis was carried out with Agilent 7890 GC-MS system. The relative percentages of the separated compounds were calculated from total ion chromatograms. The identification of the oil components was based on the Wiley and NIST mass spectral library.

Results: The essential oil yield in fennel was determined as 2.16 % . The major essential oil components of *Foeniculum vulgare* Mill. were determined as (E)-anethole (83.7%) and limonene (5.5%).

Conclusion / Discussion: Fennel commonly is known as important medicinal and aromatic plant cultivated for its essential oil, spices herb, phytotherapy and aromatherapy applications. According to the results of this study, the observed significant differences in the major components of fennel essential oil cultivated in Konya.

Keywords: Fennel, *Foeniculum vulgare* Mill., Essential oil, Composition

POSTER PRESENTATION

THE POLYPHENOLS EXTRACTED FROM THE *ARTEMISIA ABSINTHIUM L* PLANT COMBINE BRAIN OXIDATIVE STRESS, NEURO-INFLAMMATION AND NEURODEGENERATION INDUCED BY HEAVY METALS (ALUMINUM AND MERCURY).

Hallal Nouria^{1,2} ; Benyettou Imène² ; Kharoubi Omar².

¹*Ahmed Ben Yahya El Wencharissi University Center, Tissemsilt.*

²*Experimental Biotoxicology Biodespollution and Phytoremediation Laboratory (BTE-BD-PR). University of Oran 1, Ahmed Ben Bella.*

hallalnouria@yahoo.fr

The objective of our study is the analysis of the polypholic extract of *Artemisia absinthium L* (stems and leaves) by HPLC and to evaluate the effect of its administration on the alterations of the cerebral structures (cortex, cerebellum hippocampus and striatum) in aluminum and mercury intoxicated rats [(30 mg AlCl₃ / kg body weight), (5 mg HgCl₂ / kg body weight)].

Thus, the analysis of extracts of *Artemisia absinthium L* allowed the identification of nine compounds of which: catechin which represents the dominant flavonoid (35,50%), followed by quercetin (31,37%), then the caffeic acid (14.49%), chlorogenic acid (10.18%), caticin (4.02%), vanillic acid (2.55%), hydroxycyanamic acid (0.18%) , rutin (0.72%) and gallic acid (0.36%).

In the rat, the results obtained demonstrate that treatment with the polyphenol extract of *Artemisia absinthium L* attenuates Al and Hg-induced alterations by decreasing lipid peroxidation and protein oxidation, increasing activity of antioxidant enzymes (SOD, CAT, GPx, GR, and trxR). In addition, the polyphenol extract of *Artemisia absinthium L* inhibits the release of pro-inflammatory cytokines (IL1, IL6 and TNF α), nitric oxide (NO) and also prevents neuronal apoptosis by inhibition of activation of caspase-3 and the reduction of mDNA oxidation.

These results indicate that treatment with *Artemisia absinthium L* extract may be part of an effective therapeutic strategy to reduce neuronal death and combat the oxidative stress, neuro-inflammation and neurodegeneration induced by certain heavy metals.

Key words: Polyphenols, *Artemisia absinthium L*, Neurodegeneration.

FULL-TEXT PAPERS

MESMAP-5 Draft Proceedings Book

STUDIES ON MEDICINAL AND AROMATIC PLANTS UNDER THE FARMERS CONDITIONS IN THE PROVINCE OF KAYSERİ

Ahmet Yenikalaycı^{1*}, Kemal Gül², Mahmut Güneş³

¹ *Muş Alparslan Üniversitesi, Faculty of Applied Sciences, Plant Production and Technologies Department, Muş, Turkey*

² *Directorate of Provincial Agriculture and Forestry, Plant Production and Plant Health Branch Directorate, Kayseri, Turkey*

³ *County Agriculture and Forestry Directorate, Kayseri, Turkey*

*Corresponding Author e-mail: a.yenikalayci@alparslan.edu.tr

Abstract: This study comprises the results of demonstrative studies on some medicinal and aromatic plants carried out by Kayseri Directorate of Provincial Agriculture and Forestry under the field conditions in 2015-2017 within the Project of Improving the Medicinal-Aromatic and Dye Plants Cultivations, funded by General Directorate of Plant Production (BÜGEM). In this project, five different plants consisting of lavandin (*Lavandula x intermedia*), lemon balm (*Melissa officinalis* L.), oregano (*origanum onites* L.), salep (*Orchis sancta* and *Serapias womeraceae*), and black cumin (*Nigella sativa*) were studied in 11 counties for three years. Desired results were obtained from lavandin, lemon balm, black cumin and oregano plants under the ecological conditions of Kayseri. For Lavandin species, dry flower yield was 300-1270 kg ha⁻¹, and essential oil rate in dry flowers was found 2.72-9.00 %. For Lemon Balm species, dry herbage yield and essential oil rate in dry leaf were 5250-6100 kg ha⁻¹, and 0.07-0.24 %, respectively. For Oregano species, essential oil rate was found 4.64 %. In terms of cultivation of black cumin, there aren't any considerable problems except for weeds affecting sufficient yields. The weed problem was primarily originated by not having herbicides authorized by the Ministry of Agriculture and Forestry. Although application of different planting times and mulching methods in salep plants, desired results were not obtained because of the harsh winter conditions and frost damage.

Key words: Field Conditions, Medicinal Plants, Cultivation

1. INTRODUCTION

Medicinal and aromatic plants are existing in nature for centuries and versatile plants used by human being for different aims. These plants, used as food, medicine, paint and spices since ancient ages, are utilized in different fields because of containing of divergent active ingredients. Although these plants, being talked about, lost their former significances, they have been nearly discovered again and become the center of attraction in recent years. These plants, undertaken in the scope of alternative, traditional or supplementary medicine and being in the position of herbal treatment, have many advantages. Being cultivated in marginal and high-altitude-areas, without irrigation, fertilizing, using pesticides are the preeminent of them. Also, having little soil selectiveness and a suitable species to every kind of geographical conditions because of the species richness and enabling us to cultivate economically with a very low inputs are the other main advantages.

This project has been carried out in the province of Kayseri since 2015 within the scope of the Project of Improving the Medicinal-Aromatic and Dye Plants Cultivations, sustained by BÜGEM (General Directorate of Plant Production), carried out on 5 plants (Lavandin, Lemon Balm, Oregano, Salep, and Black Cumin) in 11 counties and furthermore it has been expanding yearly by means of adding new plants. In this area by the means of increasing the plant species, determining the suitable alternative plants to the ecological conditions in Kayseri area providing more income, enabling the grower to learn and spread out their cultivation are aimed.

2. MATERIALS AND METHODS

2.1. Materials

The materials used in demonstrations were provided from the Directorate of Aegean Agricultural Research Institution/Menemen/İZMİR, the Directorate of Directorate of Fruit Research Institute/Eğirdir/İSPARTA, The Directorate of Transitional Zone Agricultural Research Institute/ESKIŞEHİR, and the Directorate of Horticultural Central Research Institute/YALOVA. The species used in the study; *Lavandula x intermedia*, *Melissa officinalis* L., *Origanum onites* L., *Orchis sancta*, *Serapias vomeraceae*, *Nigella sativa*.

2.2. Methods:

In the demonstrations; the distance on the line was 50 cm, between the lines were 120 and 150 cm for Lavandin. Seedlings of the seedlings were carried out in May. Lemon Balm seedlings were planted in May like Lavandin and the distances were 30 x 70 cm. For Oregano plants, seedlings were planted as 20-30 cm distance on the line, 40-60 cm distance between the line. Black Cumin seed were sown in April as 15-20 kg ha⁻¹. In the fields of Lavandin, Lemon Balm, Oregano, Salep, and Black Cumin fertilizer applications were carried out almost similarly as 20-20-0 Compound Fertilizer, 200-250 kg ha⁻¹ to the ground and 10-20 tons of fertilizer burnt were added in some locations. Salep plants were planted in October in the first year. Salep tubers were planted within a burnt manure-enriched soil bed, 25-30 cm in height and 80-100 cm in width, in 15-15 cm soil depth and 10-20 cm row spacing. Unsuccessful results were obtained in the first year. Therefore, planting time was changed to 3 different planting times as October (a month later than the first planting time), November and December to observe the result of different planting times. Also, different mulching method made of wheat straw and burnt manure, greenhouse and low tunnel methods were tried to see the developments under cover.

3. RESULTS AND DISCUSSION

3.1. Lavandin (*Lavandula x intermedia*): Lavandin seedlings were obtained in the years of 2015 and 2016. Demonstrative activity researches were carried out at totally 14 locations with a total of 7 locations first year and 7 locations second year. Harvestings were carried out in 2016-2017. Accordingly, while good results were taken from some locations in all plants, it was not possible to get any results from some of them since just first year or in the other years because of the reasons originated from the growers. In Lavandin, demonstrations were established in 3,35 decares in 7 locations in 6 counties in 2015, in 5,2 decares in 7 locations in 7 counties in 2016, totally in 8,55 decares. The harvest wasn't made in the first year because of plant growth being slow, however to enhance the tillering in flowering plants, flowers were severed from their stems.

In lavandin demonstrations, constituted in 2015, however even if just a bit desiccations in plants originated from cold weather and frost in some locations in 2015-2016, plants recovered again by tillering with the warming weather in Spring.

In three designated locations below, the results of dry flowers yield, essential oil rate and essential oil yield could be obtained.

Table 1. According to the 2. Years (2016) Harvests in Some Locations, Dry Flowers Yields, Essential Oil Rates and Essential Oil Yields.

Queue Number	Location Name	Dry Flowers Yield (kg ha ⁻¹)	Essential Oil Rate (%)	Essential Oil Yield (L ha ⁻¹)
1	Yahyalı/Mustafabeyli	1270	6,1	77
2	Yeşilhisar/Merkez	300	5,7	17
3	Yeşilhisar/Kayadibi	680	6,0	48

When examining Table 1, dry flowers yields, essential oil rates and essential oil yields are seen regarding Lavandin trials results on the grower basis in Mustafabeyli/Yahyalı, Yeşilhisar and Kayadibi/Yeşilhisar.

The highest dry flowers yield became in Yahyalı, Mustafabeyli Location as 1270 kg ha⁻¹, the lowest dry flowers yield became in Yeşilhisar location as 300 kg ha⁻¹. Essential oil rates in dry flower were very close to each other between 5,7-6,1%. While essential oil yields were 77 L ha⁻¹ because of high dry flowers yield in Mustafabeyli/Yahyalı location, it was stayed at a lower level of 17 L ha⁻¹ because of low dry flowers yield in Yeşilhisar. The reasons why dry flowers rate was so low were those grower retarded the harvest because of his beekeeping and shadows of the trees around flower yields increased more in the following years, but these values could not be measured.

Table 2. The Amount of Essential Oils Obtained in Dry Flower in 2017 Year Harvest in Lavandin Locations (%)

Queue Number	Location Name	Essential Oil Rate (%)	Queue Number	Location Name	Essential Oil Rate (%)
1	Yahyalı Karaköy	6,00	6	Felahiye Merkez	9,00
2	Yeşilhisar Merkez	5,70	7	Yahyalı Mustafabeyli	6,80
3	Develi Sindelhöyük	5,67	8	Tomarza Işıklar	4,50
4	İncesu Garipçe	5,20	9	Kocasinan Yazır	2,82
5	Kocasinan Akçatepe	5,80		Average Value	5,72

As examined in Table 2, the essential oil ratios of the locations in the dry flower ranged between 2,82-9,00%. The highest Felahiye Center is located at the lowest Kocasinan Yazır location. Average essential oil ratio of the locations in dry flowers was 5,72%. The emergence of a essential oil rate of 9% in the Felahiye Center is an important indicator for this location. The altitude of this location is 1330 m, the south open face, the sun is good and the climatic open days are high. It is a known fact that sunbathing increases the rate of essential oil. In Kocasinan Yazır and Tomarza Işıklar, essential oil ratios were lower than other locations because of the mixing of some lavender flower stalks in dry flower sample and lack of good sample preparation. These essential oil ratios in dry flowers are a good indicator. In the scope of the support given to young farmers in the province, the producers removed the essential oils from the products they obtained in the essential oil distillation system and placed them in small glass bottles and marketed them.

Arabaci and Bayram, (2005), (1340-4430 kg ha⁻¹), and Arslançan et al. (2014) (1620-3410 kg ha⁻¹) reported that low yields of lavender were lower in producers of lavender. the inadequacy, harvest method and the methods of removing the dry flowers from stalks, the lack of sufficient infrastructure in this issue has been due to. Deaf oil yields were also low. Otherwise, when we look at the essential oil ratios, we see a rate of 5.7% which is much higher than the values indicated by Arabaci and Bayram (2005) (1.54-2.34%).

3.2. Lemon Balm (*Melissa officinalis* L.): In 2015, 4 activities were carried out in 0.9 decares area, in 4 locations in 4 counties, in 2016 in 2.05 decares, in 3 locations and in 3 counties. In total, 2,95 decare area demonstrations have been established and followed up. In the demonstrations established in 2015, a harvest was made in the autumn plants before entering the winter, but there was not much yield. In the winter, although the above-ground parts of the plants completely desiccated in all locations with the arrival of the spring with the new shoots occurred in the plant development was not a problem. Due to winter and cold weather in Kayseri, there was no negative situation. Plant growth and yields were very good after the first year. A harvest in the flowering period in June, and another harvest before the winter in September-October, plants was harvested twice per year. While a high herbage yield was obtained in the first in June, a lower herbage yield was obtained in the second.

Table 3. Total Dry Herbage, Essential Oil Ratio and Essential Oil Yield Values of Some Locations According to the Harvest Results of the Lemon Balm

Queue Number	Location Name	Dry Herbage Yield (kg ha ⁻¹)	Essential Oil Rate (%)	Essential Oil Yield (L ha ⁻¹)
1	Yahyalı/Mustafabeyli	5250	0,08	4,2
2	Kocasinan/Yazır	6100	0,07	4,2

In Table 3, while the total dry herbage yield was 5250 kg ha⁻¹, the essential oil rate was 0.08% and the essential oil yield was 4,2 L ha⁻¹ in the Yahyalı Mustafabeyli location, these were 6100 kg ha⁻¹, 0.07% and 4,2 L ha⁻¹ in the Kocasinan Yazır location at the 2nd year harvest. The dry and fresh herbage yields for the second year were higher than in the first year, but these data could not be recorded since the yield values could not be weighed. According to this, it can be said that the lemon balm plant is in compliance with the ecological conditions in Kayseri and it has no climatic, disease or harmful problems.

Table 4. Essential Oil Ratio of Dry Leaf in Lemon Balm According to 2017 Harvest

Queue Number	Location Name	Essential Oil Rate (%)	Queue Number	Location Name	Essential Oil Rate (%)
1	Yahyalı Mustafabeyli	0,17	3	Tomarza Işıklar	0,09
2	Kocasinan Yazır	0,13	4	İncesu Garipçe	0,24
				Average Value	0,16

Table 4 shows the rates of essential oil in four locations in the lemon balm plant. According to this, the highest essential oil ratio was found at İncesu Garipçe location with the highest rate of 0,24% and the lowest value was found at 0,09% Tomarza Işıklar location. Values are close (0.1-0.35%), to the values reported by Uzun et al., (2014), some higher than the values reported by Uyanık and Gürbüz (2014) (0.03-0.08%) and Koç, H., (2002). According to the results of Abdellatif et al. (2014) (% 1,54-2,34), it is slightly lower. Essential oil ratios are close to the findings obtained in our country.

3.3. Oregano (*Origanum onites L.*): Demonstrations in oregano were established in 2016 in 5 locations and 5 counties in a total area of 2.5 decares. Plant growth was very good in locations. A superficial harvest was made before the first year of winter and no significant yield was obtained. In the second year flowering period, the herbage yields of the plants harvested in June were taken but could not be measured. In some locations 1 and 2 reaps were taken in 2 years, in second reaps in September and October, a very low herbage yield was mentioned. No disease, no harmful organism or no cold and frost damage in winter in the plants was observed. Producers in Yahyalı Kopçu and İncesu Garipçe locations sold their products in the markets and generated certain amounts of income. In the İncesu Garipçe location, the amount of essential oil detected in the 2nd year (2017) *Origanum onites L.* leaves was 4.64%. Since our producer is an organic farming producer, it has been able to pack and sell medicinal and aromatic plants (lavandin, lemon balm, oregano and sage) in the organic market. He is still doing this job.

3.4. Salep (*Orchis sp.*)

Demonstrations in the Salep plant were established in 2015 in 3 counties in 0.9 decares area, in 2016 in 6 locations in 6 counties in 3.2 decares, in a total of 4.1 decares. The first year plantings were made in October. In autumn there was no outflow of tubers. In the spring, there was some output in the Kocasinan location as in March, but they also disappeared with the effect of late frosts in spring. It was planned to investigate different applications such as different planting time, mulching methods and production conditions under cover for 2 years. In order to see the results of different planting time in the

2nd year, 3 different planting time applications were made in a way that the sowing times of the tubers were one month after the first planting time, from September to October-November. Different mulching methods were tried. To keep the pillows warm, burnt farm manure was laid in straw-straw and some were planted in the greenhouse and some were planted in the form of a low tunnel cover. After planting, demonstrative follow-ups were performed regularly. No improvement was observed in plants in autumn and winter Only in the greenhouse and under the cover planted some output was observed in winter, but in the later stage they disappeared. A few plants were found in the Felahiye location, which overlooks the southern slope of the open area.

It is thought that the winter conditions in Kayseri are hard and long, the damages of late frosts in spring and the salep species used in the spring are unsuccessful due to the cold and frost sensitive species *Orchis sancta* and *Serapias vomeraceae*. Salep plant is a difficult plant culture. Arabacı et al. (2014), in their study on the Effect of Different Cultural Practices in Salep Orchids, reported that many of the studies conducted for culturing Salep orchids were in vitro studies and that the plants were failing at the stage of adaptation to outdoor conditions. We are of the opinion that in the conditions of Kayseri cultivation of salep plants can be made in other species compatible with the Central Anatolia Region. Tutar et al. (2012) in our opinion in accordance with the salep each region of their own species and work with their ecological conditions can be achieved with successful results stated.

3.5. Black Cumin (*Nigella sativa* L.): The demonstrations in Black Cumin were implemented in 3 locations (13 decares) in 3 counties in 2015, and in 5 locations (14 decares) in 4 counties in 2016, totally in 27 decares. Seed of 15-20 kg per hectare were used for seed decantation in April and sprinkling irrigation was done. Sprouts and exits occurred in locations. However, due to the large number of weed populations, black cumin did not show any improvement. Since the Ministry is not a licensed herbicide and mechanically it is not suitable for anchoring, it is not possible to obtain a product at the level of economic efficiency in black cumin. Because the yield values of the decay remained at very low levels of 100-200 kg ha⁻¹, farmers did not even need to harvest because they could not remove them.

4. DISCUSSION AND CONCLUSION

The following conclusions can be drawn from these studies conducted in 45.1 decares area in 43 locations in 11 counties in 5 different medicinal and aromatic plant species (Lavandin, Lemon Balm, Oregano, Salep and Black Cumin) for 3 years between 2015-2017. As the studies were conducted under farmer conditions, the desired results could not be obtained from each demonstration depending on the farmers. In the trials and demonstrations carried out in the farmer's conditions, the choice of the wrong farmers, the lack of equipment and equipment infrastructure at the desired level, the willingness to appear at first, and then the difficulty in raising their work intensity or cultivation in these plants, the lack of interest and the lack of enough results could not be achieved. For these reasons, some demonstrations were disabled at the first stage. However, a certain result has been reached in the studies. We tried to reduce this risk by establishing the demonstrations at more locations. Under the conditions of Kayseri, these plants can be cultivated economically and they have an idea about their adaptation to Kayseri ecology, their efficiency and quality. These results shed light on local people for those who will work or invest in these plants. As a matter of fact, the Provincial Directorate of Agriculture and Forestry has a lot of people who are interested or not interested in agriculture and want to invest in these matters. These outputs are important in informing and referring them to the right direction.

Under the conditions of farmers, making and conducting these demonstrations cause the producers to recognize and learn these plants, and to recognize, see and be interested in the farmers of the neighboring farmers and other nearby villages. In the next stage, these plants gradually enter the crop rotation in this area and make it become agricultural. There are many examples of this, but this can take many years. Before the start of this project, none of the medicinal and aromatic plants in Kayseri were cultivated and

produced. Now some of the producers still continue to grow existing plants, new producers, new plants are added, they are included in municipalities under the leadership of Provincial Directorate of Agriculture and Forestry. According to this study; In the ecological conditions of Kayseri, positive results were obtained in the cultivation of lavandin, Lemon Balm and oregano plants. Although we do not see a problem in plant growth in the sowing of Black Cumin plants, weed problem has emerged as an important problem. Due to the lack of a medicinal product licensed from the Ministry of Agriculture and Forestry, it was not possible to obtain sufficient yield. In order to be successful in the agriculture of black cumin in Kayseri situations, we think it is necessary to plant more broadly (40-50 cm) in order to allow the weeding of the weeds mechanically, it would be appropriate to make different sowing times, including autumn, winter and early spring. Due to the fact that the winter conditions in Kayseri are hard and long, and the species used are not compatible with this region, appropriate results could not be obtained. Salep plant is a plant that is difficult to culture, but we believe that other types of studies compatible with the Central Anatolia region can be done.

Acknowledgments

As the Kayseri Project Team, we would like to thank the Ministry of Agriculture and Forestry, the General Directorate of Plant Production for the Project of Improving the Medicinal-Aromatic and Dye Plants Cultivations.

References

1. Abdellatif, F., Boudjella, Hadjira, Zitouni, A., Hassani, Aicha, 2014. Chemical composition and antimicrobial activity of the essential oil from leaves of Algerian *Melissa officinalis* L., *Exclı Journal*, v:13, s: 772–781.
2. Arabacı, O., Bayram, E., 2005. The Effect of Nitrogen Fertilization and Plant Density on Some Agronomic and Quality Traits of Lavender (*Mill.*) Under Ecological Conditions of Aydın, *Adü Journal of the Faculty of Agriculture*, 2005; 2(2): 13 – 19
3. Arabacı, O., Tutar, M., Öğretmen, N.G., Yaşar, F., Tan, U., 2014. Effects of Different Cultural Practices in Salep Orchids, II. Symposium of Medical and Aromatic Plants, Atatürk Horticultural Central Research Institute Directorate, Yalova, 23-25 september 2014, p.479-487.
4. Aslanca, H., Sarıbaş, R., Dağistanlıoğlu, C., Tınmaz, A.B., Arslan, D., 2014. Determination of Agricultural Properties of Some Lavender (*Lavandula x Intermedia Emeric Ex Loisel.*) Cultivars in Isparta Conditions, II. Symposium of Medical and Aromatic Plants, Atatürk Horticultural Central Research Institute Directorate, Yalova, 23-25 september 2014, p.344-347.
5. Koç, H., 2002. Healthy Living with Plants, Prof. Dr. Hüseyin KOÇ, Gaziosmanpaşa University Faculty of Agriculture Field Crops Department, Tokat.
6. Uyanık, M., Gürbüz, B., 2014. A Study on Determination of Diurnal Variability in Lemon Balm (*Melissa officinalis* L.), II. Symposium of Medical and Aromatic Plants, Atatürk Horticultural Central Research Institute Directorate, Yalova, 23-25 september 2014, p.432-437.
7. Uzun, A., Kevseroğlu, K., Gürbüz, B., 2014. Determination of Some Vegetal and Chemical Traits of Lemon Balm (*Melissa officinalis* L.) Ecotypes Presented in the Flora of Middle and East Black Sea Region, II. Symposium of Medical and Aromatic Plants, Atatürk Horticultural Central Research Institute Directorate, Yalova, 23-25 september 2014, p.377.
8. Tutar, M., Çiçek, F., Sarı, A.O., Bilgiç, A., Yıldız, Ö., 2012. The Culture of Some Orchid Species Distributed In Aegean Region, Turkey 2. Orchids and Sahlep Workshop Proceedings, Aegean Agricultural Research Institute, 25-26 April 2012, İzmir, p.301-320.

FULL PAPER

CURCUMIN, A POLYPHENOL DERIVED FROM *CURCUMA LONGA*: CLINICAL EVIDENCE AND MOLECULAR MECHANISMS IN PREVENTION AND MANAGEMENT OF CANCER

Nazim Sekeroglu^{a*} & Sevgi Gezici^b

^a Department of Food Engineering, Faculty of Engineering and Architecture, Advanced Technology Application and Research Center, Kilis 7 Aralik University, 79000 Kilis, Turkey

E-mail: nsekeroglu@gmail.com, www.nazimsekeroglu.com

^b Department of Molecular Biology and Genetics, Faculty of Science and Literature; Advanced Technology Application and Research Center, Kilis 7 Aralik University, 79000 Kilis, Turkey

E-mail: drsevgigezici@gmail.com, www.sevgigezici.com

Abstract: Cancer is the second leading cause of death globally, and it has been a major health problem in the past decades. Several strategies are involved in treatment and prevention of cancer cases such as chemotherapy, immune therapy, radiation, hormone therapy, targeted therapy and herbal therapy. Among them, herbal therapy with medicinal and aromatic plants has been gained great importance in recent years. Medicinal plants and their bioactive compounds may offer clinically useful anticancer agents, due to their rich anticarcinogenic and chemoprotective potentials. Curcumin, the rhizome of *Curcuma longa*, is widely used in therapeutic purposes because of its several biological properties such as anti-inflammatory, anticancer, antioxidant, and etc. From this point of view, this review is aimed to comprehensively highlight the effects of curcumin for the prevention and treatment of cancer. **Methods:** This report reviews the pharmacological and anticancer effects of curcumin with a focus on its molecular mechanisms in prevention and management of various cancer cases. For this purpose, electronic databases, including PubMed, Scopus, ScienceDirect, Cochrane library, Google Scholar and MedlinePlus, etc. were searched to summarize *in vitro*, *in vivo* and clinical studies on anticancer effects of *Curcuma longa* and curcumin, polyphenolic bioactive compound of the plant up-to-date. **Results and Conclusion:** To sum up, curcumin, the active ingredient of turmeric and exerts remarkable therapeutic potential for the treatment of various types of cancers. Curcumin is strong anti-oxidant and anti-inflammatory properties and therefore it possesses anticancer potentials through various molecular mechanisms. Based on documented data this natural compound could be a key source of anticancer agent in modern anticancer therapy after clinical confirmation.

Key Words: *Curcuma longa*; curcumin; anti-cancer agents; molecular mechanism; natural products; medicinal plants.



Graphical Abstract

INTRODUCTION

Cancer has become a growing health problem around the world with the high numbers of cases and deaths. Meanwhile, it is predicted that it will be more critical health problem in both developed and developing countries for the next years. More recently, the researchers have mainly focused on examining the use of plant-derived natural products with less or no side effects for prevention and treatment in the cancer cases, since an increase in the incidences of drug resistance and side effects of drugs used for cancer therapy (Gezici, 2018; Gezici and Sekeroglu, 2019). Curcumin (turmeric), a widely used spice, food preservative and coloring material, is isolated from *Curcuma longa* Linn. belonging to Zingiberaceae (ginger) family. This plant is originated from southeastern Asia, and distributed throughout tropical and subtropical regions. The rhizome of the plant is known as 'Golden Spice' because of its brilliant yellow color, and it is a very popular spice in Africa, Thailand, Pakistan and India (Verma et al., 2018; Rai et al., 2018; Li et al., 2019).




Fig. 1. Pictures of plant (A), rhizomes (B) and rhizome powder (C) of turmeric (*Curcuma longa* L.)

Chemical Structure of Curcumin: *C. longa* L. has included curcumin, demethoxycurcumin, and bisdemethoxycurcumin. Curcumin, the most important constituent of *C. longa* L., is responsible for its yellow color (Fig. 1). Chemically, curcumin is a bis- α , β -unsaturated β -diketone with keto-enol tautomerism and its chemical structure was given in the Fig. 2. (Akram et al., 2010; Rai et al., 2018; Li et al., 2019).




Fig. 2. Chemical structure of Curcumin

Properties and Health Benefits of Curcumin: Curcumin has a long history of use in traditional medicine for treatment of various disease and disorders including cough, diabetic ulcers, hepatic diseases, biliary disorders, rheumatism, sinusitis and anorexia (Li et al., 2019). It has been used to cure acne, heal wound, prevent skin damage, reduce cholesterol, treat diabetes, control blood pressure, prevent inflammation and cancer cell growth (Fig.3). It behaves as a free radical scavenger in the human body, thus it is able to prevent oxidative damage as blocking the free radicals. It also decreases the production of reactive oxygen species and stimulates antioxidant capacities in the body (Farzaei et al., 2018; Pagano et al., 2018). It has been used for treatment of various diseases, owing to its wide range of biological effects including not only antitumor activities but also as antioxidant, antimicrobial, anti-inflammatory, anticoagulant, antidiabetic and immunomodulatory activities (Aggarwal et al., 2007; Pan et al., 2010, Verma et al., 2018).




Fig. 3. General benefits of curcumin

Role of Curcumin in Carcinogenesis and Cancer Pathways: Cancer is a multifactorial disease that occurs when a normal cell loses its cellular division control. At the molecular level of carcinogenesis, curcumin targets numerous pathways at multiple levels including activator protein 1 (AP-1), Nuclear Factor Kappa B (NF- κ B), Peroxisome Proliferator-Associated Receptor Gamma (PPAR- γ), Signal Transducer and Activator of Transcription (STAT), Wnt/ β -catenin, transcription factor Nrf-2-Keap, Tumor Necrosis Factor-alpha (TNF- α), Interleukins (IL), and oncogenic kinases (Li et al., 2019; Shanmugam et al., 2015).




Fig. 4. Curcumin and cancer pathways (<http://www.cancermind.com/curcumin-anticancer-cure-cancer-review/>)

Curcumin are able to interact with molecular targets involved in inflammation, thus it modulates inflammatory response by reducing the activity of cyclooxygenase-2 and lipoxygenase, inducing nitric oxide synthase enzymes, inhibiting tumor necrosis factor-alpha and interleukins. Moreover, it interferes with NF- κ B, modulates growth of tumor cells through regulation of multiple cell signaling pathways including cyclin D1, c-myc, Bcl-2, Bcl-x, cFLIP, XIAP, c-IAP1, and caspase activation pathway (Jurenka, 2009; Yan et al., 2019). Curcumin related cancer pathways are figured out in the Fig. 4.

Curcumin effects carcinogenesis process through various mechanisms as increasing of growth suppressors and apoptosis, decreasing of proliferative signals and angiogenesis, and restricted replicative immortality (Fig. 5). Curcumin targets protein kinases and growth factors which are involved in carcinogenesis process. ErbB-2 (avian erythroblastosis oncogene B) or HER2/neu is a member of the epidermal growth factor receptor (EGFR) family is the main targets of curcumin. Moreover, curcumin is known as a potential anti-inflammatory agent that able to inhibit inflammatory enzymes namely nitric oxide synthase (iNOS), cyclooxygenase-2 (COX-2), and lipoxygenase (LOX) (Shanmugam et al., 2015; Farzaei et al., 2018; Yan et al., 2019).




Fig. 5. Role of curcumin in carcinogenesis

Clinical Trials and Cancer: Numerous clinical studies including pilot clinical studies and Phase I, II and III clinical trials are currently in progress for revealing of the therapeutic potentials of curcumin. The clinical trials have been summarized in Table 1. The anti-cancer potential of curcumin has been successfully showed in various types of cancers such as oral, lung, breast, prostate, pancreatic, colorectal, multiple myeloma, head and neck squamous cell carcinoma. Clinical trials with curcumin indicate safety, tolerability, non-toxicity (even up to doses of 8000 mg/day), and efficacy (Shanmugam et al., 2015; Yan et al., 2019). Researches showed that curcumin is capable of inhibition of carcinogenesis even at three stages including initiation, promotion, and progression (Fig. 5). Dhillon et al., [4], at the M.D. Anderson Cancer Center in Houston determined that curcumin has potential activity to inhibit of malignant cell proliferation in pancreatic cancer in phase II clinical trial, and also, found that it is utilized for chemoprevention of colon cancer (Johnson and Mukhtar, 2007). Moreover, Dai et al. [3] designed and synthesized curcumin mono-carbonyl analogs for cancer therapy against human lung cancer A549 cells (Gezici and Sekeroglu, 2017, 2019).

Table 1. Clinical trials the use of curcumin in cancer cases

Cancer Cases	Organization / Cancer Center	Trial Phase	Date
Colon cancer	Chao Family Comprehensive Center	Phase II	Unknown
Colon cancer	University of Pennsylvania	Phase II	2019, June
Rectal cancer	MD Anderson Cancer Center	Phase II	2010, July
Pancreatic cancer	Tel-Aviv Sourasky Medical Center	Phase III	Unknown
Pancreatic cancer	MD Anderson Cancer Center	Phase II	2009, December
Multiple myeloma	MD Anderson Cancer Center	Pilot study	2008, December
Osteosarcoma	Tata Memorial Hospital	Phase I and Phase II	2012, May
Familial adenomatous polyposis	John Hopkins University	Phase II	2013, March
Colorectal cancer	MD Anderson Cancer Center	Phase I	Unknown

Conclusions and Future Directions: To conclude, curcumin is a pharmaceutical agent that possess various therapeutic applications in cancer prevention and treatment. However, curcumin was limited in clinical use because of its low bioavailability, which might be led by poor absorption and rapid metabolism. To increase its bioavailability, several curcumin formulations have been developed namely powder, tablets, capsules, liposomal encapsulation, emulsions, and nanoparticles. Nonetheless, further clinical trials are required to validate their safety. It will be a great potential to be a dietary supplement with various formulations that is able to prevent tumorigenesis through altering different molecular pathways.

References

- [1] Aggarwal BB, Sundaram C, Malani N, Ichikawa H. Curcumin: The Indian solid gold. *Adv. Exp. Med. Biol.*, 2007, 595, 1–75.
- [2] Akram, M., Shahab-Uddin, A. A., Usmanghani, K., Hannan, A., Mohiuddin, E., Asif, M. (2010). Curcuma longa and curcumin: a review article. *Rom J Biol Plant Biol*, 55(2), 65-70.
- [3] Dai F.; Liu GY.; Li Y.; Yan WJ.; Wang Q.; Yang J.; Lu DL.; Ding DJ.; Lin D.; Zhou B. Insights into the importance for designing curcumin-inspired anticancer agents by a prooxidant strategy: The case of diarylpentanoids. *Free Radic. Biol. Med.*, 2015, 85, 127-37.
- [4] Dhillon N.; Aggarwal BB.; Newman RA.; Wolff RA.; Kunnumakkara AB.; Abbruzzese JL.; Ng CS.; Badmaev V.; Kurzrock R. Phase II trial of curcumin in patients with advanced pancreatic cancer. *Clin. Cancer Res.*, 2008, 14, 4491-9.
- [5] Farzaei, M., Zobeiri, M., Parvizi, F., El-Senduny, F., Marmouzi, I., Coy-Barrera, E., Abdollahi, M. (2018). Curcumin in liver diseases: A systematic review of the cellular mechanisms of oxidative stress and clinical perspective. *Nutrients*, 10(7), 855.
- [6] Gezici, S., Sekeroglu, N. (2017). Regulation of MicroRNAs by natural products and bioactive compounds obtained from common medicinal plants: novel strategy in Cancer therapy. *Indian J. Pharm. Educ. Res.* 51(3), 498-503.
- [7] Gezici, S. (2018). Promising anticancer activity of lavender (*Lavandula angustifolia* Mill.) essential oil through induction of both apoptosis and necrosis. *Annals of Phytomedicine*, 7(2), 38-45.
- [8] Gezici, S., Sekeroglu, N., (2019). Current perspectives in the application of medicinal plants against cancer: novel therapeutic agents. *Anti-Cancer Agent Med. Chem.* <https://doi.org/10.2174/1871520619666181224121004>.
- [9] Johnson JJ.; Mukhtar H. Curcumin for chemoprevention of colon cancer. *Cancer Lett.*, 2007, 255, 170-81.
- [10] Jurenka, J. S. (2009). Anti-inflammatory properties of curcumin, a major constituent of *Curcuma longa*: a review of preclinical and clinical research. *Alternative medicine review*, 14(2).
- [11] Li, H., Sureda, A., Devkota, H. P., Pittalà, V., Barreca, D., Silva, A. S., Nabavi, S. M. (2019). Curcumin, the golden spice in treating cardiovascular diseases. *Biotechnology advances.* <https://doi.org/10.1016/j.biotechadv.2019.01.010>.
- [12] Pagano, E., Romano, B., Izzo, A. A., Borrelli, F. (2018). The clinical efficacy of curcumin-containing nutraceuticals: An overview of systematic reviews. *Pharmacological research*, 134, 79-91.
- [13] Pan, L.; Chai, H.; Kinghorn, AD. The continuing search for antitumor agents from higher plants. *Phytochem. Lett.*, 2010, 3, 1-8.
- [14] Rai, M., Ingle, A., Pandit, R., Paralikar, P., Rehman, F., Anasane, N., Buxy, S. (2018). *Curcuma longa* L.: From Ethnomedicinal to Novel Biomedical Applications. *Ethnobotany: Application of Medicinal Plants*.
- [15] Shanmugam, M., Rane, G., Kanchi, M., Arfuso, F., Chinnathambi, A., Zayed, M., Sethi, G. (2015). The multifaceted role of curcumin in cancer prevention and treatment. *Molecules*, 20(2), 2728-2769.
- [16] Verma, R. K., Kumari, P., Maurya, R. K., Kumar, V., Verma, R. B., Singh, R. K. (2018). Medicinal properties of turmeric (*Curcuma longa* L.): A review. *IJCS*, 6(4), 1354-1357.
- [17] Yan, Z. J., Zhou, Q., Zhang, L. M., Xu, Z. J., & Chen, Y. Q. (2019). Curcumin regulates the proliferation and apoptosis of human oligodendrogloma cells through both the NF- κ B and AKT pathways. *Int J Clin Exp Med*, 12(1), 611-617.

FULL PAPER

**THE EFFECTS OF NITROGEN AND FARM MANURE
APPLICATIONS ON ESSENTIAL OIL CONTENT OF DILL
(*ANETHUM GRAVEOLENS* L.)**

Ferit Özen¹, Gülsüm Yaldiz², Mahmut Çamlica²

¹ Medicinal and Aromatic Plants Department, Mudurnu Süreyya Astarçı Vocational School, Mudurnu Bolu, Turkey

² Department of Field Crops, Faculty of Agriculture and Natural Sciences, Bolu Abant İzzet Baysal University, 14280 Bolu, Turkey. E-mail: g_yaldiz@hotmail.com

Abstract

Anethum graveolens L. or dill, belonging to Apiaceae (Umbelliferae) family, is an annual aromatic herb known for culinary and medicinal use since ancient times. It has been used in ayurvedic medicines since ancient times and it is a popular herb widely used as a spice and also yields essential oil. The present study aimed to determine the effects of different doses of chemical (ammonium nitrate) and organic fertilizer (farm manure) applications on the essential oil content of *Anethum graveolens* L. (dill). In this study, different doses of farm manure (750, 1000, 1250 and 1500 kg da⁻¹) and ammonium nitrate (3, 6, 9, 12 kg da⁻¹) with a control (no manure) were applied by sowing. The experiments were arranged in the Completely Randomized Blocks Design with three replicates. Each experimental plot consisted of five rows, with a distance of 0.3 m between rows and 0.2 m between plants, and a plot size of 3.3 m². The essential oil percentage was subjected to hydro-distillation for 3 hours using a modified Clevenger apparatus. The hydro-distilled essential oil content ranged from 0.26% to 6.18%. The highest essential oil content were obtained in the treatments with farm manure doses of 1250-1500 kg/da and control application, which were significantly different compared to the other treatments. Among all of the essential oils included in the study, dill seeds showed the highest essential oil yield, followed by dill leaves and dill herbs. As suggested by the present study, a suitable ratio of farm manure in the soil can increase the essential oil content. This organic manure could thus be considered as suitable substitutes for chemical fertilizers when growing medicinal plants that are gaining both an increased importance and demand.

Keywords: Dill, essential oil, farm manure

1. INTRODUCTION

Anethum graveolens L. (dill) has been used in ayurvedic medicines since ancient times and it is a popular herb widely used as a spice and also yields essential oil. It is an aromatic and annual herb of apiaceae family. There are various essential components of dill seeds and herb; carvone being the predominant odorant of dill seed and α -phellandrene, limonene, dill ether, myristicin are the most important odorants of dill herb. Other compounds isolated from seeds are coumarins, flavonoids, phenolic acids and steroids (Bailer et al., 2001). Dill is used as an ingredient in gripe water, given to relieve colic pain in babies and flatulence in young children (Pulliah, 2002). The seed is aromatic, carminative, mildly diuretic, galactagogue, stimulant and stomachic (Hornok, 1992; Sharma, 2004). The essential oil in the seed relieves intestinal spasms and griping, helping to settle colic (Fleming, 2000; Duke, 2001). The carminative essential oil improves appetite, relieves gas and aids digestion. Fertilization is one of the most significant agricultural practices used to improve the yield and quality of traditional crops. The need to use renewable forms of energy and reduce costs of fertilizing crops has revived the use of organic fertilizers worldwide (Seifritz, 1982). Furthermore, organic manure activates many species of living organisms which release phytohormones and may stimulate the plant growth and absorption of nutrient (Naguib and Aziz, 2004).

The present study aimed to determine the effects of different doses of chemical (ammonium nitrate) and organic fertilizer (farmyard manure) applications on the essential oil contents of dill. We have chosen dill for our study because it is one of the most popular culinary herbs worldwide, has a variety of cultivars available, and is often commercially produced in greenhouses, used by local people for medicinal purposes or as a nutrient.

2. MATERIAL AND METHODS

The present experiment was carried out during growing season in 2017 at Bolu Abant İzzet Baysal University, Bolu. The field experimental site was located at research and application area of Agriculture and Natural Sciences Faculty. Seeds of dill were obtained from Erzurum province. The experiment was designed in a randomized complete-block design with a split-plot arrangement, with three replications being conducted in open-field conditions. Farmyard manure (FYM) and ammonium nitrate (AN) were in the main plots, whereas different AN levels (3, 6, 9, 12 kg da⁻¹) and different FYM levels (750, 1000, 1250 and 1500 kg da⁻¹) with a control (no manure) were in sub-plots. Each experimental plot consisted of five rows, with a distance of 0.3 m between rows and 0.2 m between plants, and a plot size of 3.3 m². Average climatic data were recorded 16.08 °C temperature; 41.37 kg m⁻² rainfall; 69.2% humidity during the vegetation period for 2017 (Anonymous, 2017). Experimental area soils are clay-loam with a pH value of 7.8, organic matter content of 4.7%, phosphorus ratio of 10.3 ppm and potassium ratio of 235 ppm (Anonymous, 2016a). Dill was regularly irrigated to demonstrate good progress in its period vegetation since irrigation is a very important factor for cultivation of dill. The dill seeds were sown on the 20th day of april (2017). As experimental factors, the different doses of FYM (750, 1000, 1250, 1500 kg da⁻¹) was applied a week before sowing. Also, As a base fertilizer, 12 kg da⁻¹ diammonium phosphate (DAP) were applied with sowing to the conventional plants. AN (3, 6, 9, 12 kg da⁻¹) was applied to the conventional plants in two splits in April and July. It was uniformly spread on the plots and lightly worked into the soil with harrow. The nutrient composition of the soil and the FYM were analyzed before use. The properties of FYM are provided in Table 1. FMY has the highest organic matter, calcium, magnesium and iron content. No chemicals were used in the experiments.

Table 1. Chemical Analyses of Farmyard Manure (Anonymous, 2016b).

Analysis Parameters	Unit	Analysis results
Organic matter	%	40.68
Total Nitrogen (N)	%	1.80
Moisture	%	59.50
pH (potentiometric)		7.61
EC (1 10 ⁻¹)	mS cm ⁻¹	3.26
Total phosphorus (P)	%	0.52
Water soluble phosphorus (P ₂ O ₅)	%	0.32
Total potassium (K)	%	1.36
Water soluble potassium (K ₂ O ₅)	%	1.04
Total calcium (Ca, ICP EPA 3052)	ppm	7758.00
Total magnesium (Mg, ICP EPA 3052)	ppm	5099.00
Total iron (Fe, ICP EPA 3052)	ppm	10606
Total copper (Cu, ICP EPA 3052)	ppm	24.97
Total zinc (Zn, ICP EPA 3052)	ppm	93.53
Total manganese (Mn, ICP EPA 3052)	ppm	368.30
Total lead (Pb, ICP EPA 3052)	ppm	2.33
Total cadmium(Cd, ICP EPA 3052)	ppm	DLA
Total cobalt (Co, ICP EPA 3052)	ppm	DLA

Note: DLA: Detection limits Cd<0.03ppm, Co<0.08 ppm,Pb<0.09 ppm

The experiment harvest was started on 02 July and the final harvest made on 28 July. The data were analyzed using XLSTAT statistic software program and obtained results were subjected to analysis of variance and Least Significant Difference Test (LSD) in order to find differences among the dill populations and controls at p=0.05.

Isolation of essential oil: Essential oil analyses were carried out in accordance with method TS 8882. A sample of approximately 20 g was taken from the dried plants of each species and placed into glass Clevenger flasks. Approximately 200 mL (ten times the sample weight) of distilled water was added and samples were then subjected to hydro-distillation for 3h. The essential oil accumulated on top and separated from the rest of the sample. The amount obtained was recorded (in mL) from the graduated section of the flask and weights were then used to calculate percentage essential oil yields.

3. RESULT AND DISCUSSIONS

Organic FYM demonstrated statistically significant difference ($P < 0.05$) among all of the treated plots with respect to essential oil yield except herb essential oil (Table 2). The trend in essential oil yield was slightly different compared to that in FYM applications. In general, it was higher in all of the treated plots compared to AN fertilizer. The hydro-distilled essential oil content ranged from 0.26% to 6.18%. The essential oil content was found as 0.26% to 0.72% in dill herbs, 0.27% to 1.70% in dill leaves, 4.58% to 6.18% in dill seeds. The highest essential oil content were obtained in the treatments with FYM doses of 1250-1500 kg da⁻¹ and 3 kg da⁻¹ AN application, which were significantly different compared to the other treatments. Also, control application exhibited significantly higher essential oil content compared to the all AN applications in dill herbs.

Among all of the essential oils included in the study, dill seeds showed the highest essential oil yield, followed by dill leaves and dill herbs. The essential oil yield (0.26- 6.18%) in the present study was comparable to that in a study by Kruger and Hammer (1996) who found that essential oil percentage of different dill cultivars (fruits) vary from 1.91% to 7.25%. Similarly, Badoca and Lamartib (1991) observed that the oil content of European dill seeds varied from 1.75-5.8%. Also, Said-al and Omer (2016) reported that the essential oil content of Egyptian dill herbs varied from 1.933-3.267%. The essential oil content of dill herbs, leaves, seeds determined in the present study was within the range reported for dill in all of these previous studies.

By contrast, Stanojević et al., (2015) reported the yield of the dill essential oils from seeds and herbs to be 4% and 2.80%, respectively. Such variations in the essential oil content of dill across countries might be attributed to the varied agroclimatic conditions of the regions, as well as different fertilizer applications.

Table 2: Essential oil of herb, leaf and seed in dill.

Doses (kg da ⁻¹)	Herb EO (%)	Leaf EO (%)	Seed EO (%)
Control	0.65	1.47C	5.59AB
3	0.34	1.33D	5.98AB
6	0.26	0.23H	4.63C
9	0.26	0.30F	5.26BC
12	0.35	0.30F	5.23BC
750	0.30	0.27G	4.58C
1000	0.64	0.55E	5.56AB
1250	0.55	1.50B	5.91AB
1500	0.72	1.70A	6.18A
LSD (5%)	0.71	0.03	0.77




Figure 1. Essential oil of herb, leaf and seed in dill.

4. CONCLUSION

Anethum graveolens L. is a very popular and widely grown herb worldwide. Organic and inorganic fertilization studies have been carried out with respect to cultivation, breeding, high herbage yield and high essential oil content, etc., although few studies have been conducted concerning the organic FYM effect on essential oil content and the quality of dill. Especially, FMY doses 1250-1500 kg da⁻¹ had the highest essential oil content. As suggested by the present study, a suitable ratio of organic FYM in the soil can increase essential oil content. FMY could thus be considered as suitable substitutes for chemical fertilizers when growing medicinal plants that are gaining both an increased importance and demand.

REFERENCES

1. Anonymous 2016a. Data of Bolu Directorate of Provincial Food Agriculture and Livestock (<http://www.tarim.gov.tr>). Accessed, 12 December.
2. Anonymous 2016b. Soil, Fertilizer and Water Resources Central Research Institute. Ankara, Turkey: Republic of turkey ministry of agriculture and forestry Ankara.
3. Anonymous 2017. Bolu Meteorology General Directorate. Bolu, Turkey: Turkish State Meteorological Service.
4. Bailer J, Aichinger T, Hack G, Hueber KD, Dachler M. 2001. Essential Oil Content and Composition in Commercially Available Dill Cultivars in Comparison to Caraway. *Industrial Crops and Products*. 14:229-39.
5. Hornok L. 1992. Cultivation and Processing of Medicinal Plants: Academic publications. p. 338.
6. Sharma R. 2004. *Agrotechniques of Medicinal Plants*. New Delhi: Daya Publishing House, 3-8.
7. Duke JA. 2001. *Handbooke of Medicinal Herbs*. London: CRC Press, p. 42.
8. Fleming T. 2000. *PDR for Herbal Medicines*. New Jersey: Medical Economics Company, p. 252.
9. Nair R, Chanda S. 2007. Antibacterial Activities of Some Medicinal Plants of the Western Region of India. *Turkish Journal of Biology*. 31:231-6.
10. Seifritz W. 1982. Alternative and Renewable Sources of Energy in Optimizing Yield: The Role of fertilizers in proceedings of 12th IPI Congress, p. 153 -163.
11. Naguib NY, Aziz AA. 2004. Yield and Quality of *Hyoscyamus muticus* L. in Relation to Some Fertilizer Treatments. *Egyptian Journal of Horticulture*, 29:115-26.
12. Stanojević LP, Radulović NS, Djokić TM, Stanković BM, Ilić DP, Cakić MD, Nikolić VD. 2015: The Yield, Composition and Hydro Distillation Kinetics of the Essential Oil of Dill Seeds (*Anethii fructus*) Obtained by Different Hydrodistillation Techniques. *Industrial Crops and Products*, 65:429-436.
13. Said-Al Ahl H, Omer EA. 2016. Essential Oil Content and Chemical Composition of Eight Dill (*Anethum graveolens* L.) Cultivars Cultivated Under Egyptian Conditions. *International Journal of Pharmacy and Pharmaceutical Sciences* 8(5):227-231.
14. Kruger H, Hammer K. 1996. A New Chemotype of *Anethum graveolens* L. *Journal of Essential Oil Research*, 8:205-6.
15. Badoca A, Lamartib A. 1991. A Chemotaxonomic Evaluation of *Anethum graveolens* L. (dill) of Various Origins. *Journal of Essential Oil Research*, 3:269-78.

COMPOUNDS FROM AN AROMATIC ANNUAL PLANT GROWING IN ARID ZONE OF ALGERIA AND SCREENING OF ITS ANTIBACTERIAL ACTIVITY

Mebarka Bouziane^{1,2*}, Naima Hellali¹, Ferdous Cherifi^{1,2}, Bakir Abanou^{1,2}, Mahfoud Hadj
Mahammed^{1,2}

¹Laboratory of BioGeochemistry of Desert Environments,
Faculty of Mathematics and matter sciences, University of Ouargla. Ouargla 30000, Algeria.

²University of Ouargla. Ouargla 30000, Algeria.

*Corresponding author: bmebarka@yahoo.fr

Abstract: The Asteraceae *Matricaria pubescens* (Desf.) is a spontaneous species from arid regions of Algeria. To get its volatile fraction, the aerial parts of this plant were subjected to an hydrodistillation using a cleavenger apparatus. A yield of 0.1% was obtained. The sample was analyzed by GC-MS on a non polar capillary column. Among identified components in this volatile fraction were; spathuleol (10.59%), decanoic acid (5.03%), transisoelemicin (4.70%), cadinol (1.59%). Beside, the purpose of the present study was to continue the exploration of the biological effects of *M. pubescens*. Indeed, the prepared essential oil and its fragrant water have been the subject of an antibacterial investigation. However, most of the microorganisms tested were resistant to the tested samples.

Keywords: GC-MS, volatile composition, bioactivity, spontaneous plant, arid zones

INTRODUCTION

Matricaria pubescens (Desf.) belongs to the *Asteraceae* (Chemli R., 1997), it is an annual plant and in spring it begins to appear spontaneously in its native areas in the Sahara of Algeria. Guertoufa is its vernacular name; it is widely appreciated in these regions for its agreeable fragrance. Nomad and cities habitants used it to preserve and to give a good smell to their food; they collect it from its native areas and sale it in week-markets. For traditional medication, the species is used alone or with other medicinal plants to treat various ailments such as spasm, colic and digestive troubles. In previous studies, it has been demonstrated that the plant showed interesting biological activities such as antimicrobial anti-inflammatory and anticancer effects (Makhloufi A. et al., 2014, Gowsiya S. et al., 2014, Maiza K. et al., 2014). To the best of our knowledge, no previous investigations were performed about the volatile composition of this plant from Biskra city except some ones which reported a study of the essential oil of *M. pubescens* growing in other region of Algeria (Tadrent W. et al., 2014).

The present paper presents a continuance to our work on the spontaneous species of Algerian Sahara; it aims at the exploration of the volatile chemical composition of this plant. Furthermore, an investigation of the antimicrobial activity was performed for this fraction and the resulted fragrant water.

MATERIAL AND METHODS

Extraction of the essential oils: *M. pubescens* plant material was collected at the flowering stage from Biskra city (Algeria) where latitude longitude coordinates are respectively, 34°51'1.37"N, 5°43'40.98"E (<http://latitudelongitude.org/dz/biskra>). Dr O. Oulad Belkhir agronomist from the faculty of biology, university of Ouagla, carried out plant identification. The aerial parts of *M. pubescens* were air-dried at room temperature in the dark. The essential oil was obtained by hydrodistillation using a Clevenger apparatus for 3 h. The essential oil (sample1) was collected directly by simple physical separation of the distillate, while the odorant water (sample2) was the remaining aqueous phase after decantation. Both samples were stored at 4 °C.

GC-MS Analysis: The essential oil sample was analyzed using gas chromatography with MS detection. In fact, an Agilent HP6890 GC-MS system was used; it was equipped with a mass-selective detector using electron impact ionization. The separations were performed on HP-5MS a non polar capillary column (30 m x 0.25 mm; film thickness 0.25 μ m). The temperature program of the analysis started at 60°C that was maintained for 8 minutes and then increased to 220°C by 2°C / minute, secondly it ramps at 10°C to reach 295°C and held constant for 25 min. Injector temperature was set at 250 °C. Helium was the carrier gas at 1 ml/min flow rate and the Splitless mode was applied. Concerning MS detection, it was used 70 eV for the electron impact ionization energy and the scan mode ranged from 20 to 550 (m/z). Kovats indices (I_r) for all the compounds were determined according to Van Den Dool equation (Van Den Dool H. and Kratz P. D., 1963) using retention times of n-alkanes serie (C₅-C₃₁), that had been injected after the essential oil under the same chromatographic conditions. Detected components were confirmed by comparison of their retention indices with those reported in published works (Adams R.P., 1989, http://www.flavornet.org/d_kovats_ov101.html, Zhu M. et al, 2008, Bélanger A. et al., 2008) as well as by comparison of their mass spectra with those of standards in Wiley, NIST libraries. Quantitative data were obtained electronically from TIC % without the use of correction factors.

ANTIBACTERIAL ASSAY

Microbial strains: The antimicrobial activity was evaluated by paper disc diffusion and dilution method towards Gram-negative and Gram-positive referenced bacteria; *Escherichia coli* ATCC 25922, *Klebsiella pneumonia* ATCC 70060, *Proteus mirabilis* ATCC 49452, *Enterobacter cloacae* ATCC 13047, *Pseudomonas aeruginosa* ATCC 27853, *Staphylococcus aureus* ATCC 25923, *Enterococcus faecalis* ATCC 29212. Moreover, against the following microorganisms that were taken from patients; *E. coli*, *K. pneumonia*, *S. aureus*, *Streptococcus*, *Aspergillus niger* and *Candida albicans*.

Disc diffusion method (aromatograms): The qualitative antibacterial test was performed using culture growth at 37°C for 18 h and adjusted to approximately 10⁸ colony forming unit per milliliter (CFU/ml). The Mueller Hinton Agar (MH) medium was used for the bacteria culture and Sabouraud agar medium for yeasts. 500 μ l of the inoculums were spread over plates containing (MH) and a sterilized Whatman paper discs (0.5 cm) impregnated with 5 μ l of the essential oil (sample 1) or the aromatic water (sample 2) were placed on the surface of the well dried media. The plates were left for 20 min at room temperature to allow the diffusion of the samples. Plates were incubated at 37°C during 24 h for those containing bacteria and 5 days for those of yeasts. The inhibition zone observed around the antibiograms after incubation was measured. All experiments were repeated in triplicate. (Boussaada et al.(2008) and Yeşil-Çeliktaş et al., (2007).

Dilution method and determination of MIC: The following dilutions in DMSO were prepared from the essential; 25%, 50% and 75% and a disc containing only DMSO was the blank of this assay. The minimum inhibitory concentration (MIC) is the lowest concentration of the essential oil giving no visible growth in the naked eye (Kaloustian et al., 2008).

RESULTS AND DISCUSSION

GC-MS analysis: The hydrodistillation of *M. pubescens* aerial parts provided a yellow essential oil (sample 1) with an agreeable smell with a yield of 0.1 % v/w. From the data obtained from GC-MS analysis, we were able to identify the main composition of sample 1, as it is shown on table 1. Among them the following major compounds were revealed: spathulenol (10.59%), isoelmicin (4.70%), decanoic acid (5.03%), terpineol (3.09%), cadinol (1.59%), and in a second order it was present Z-ocimene (0.53%) and α -pinene (0.23%). As it can be seen on the table 1, other constituents were also present in the studied sample.

Table 1: Compounds from *Marticaria pubescens* essential oil.

Ir (exp)	Ir (ref)	Tic (%)	Compound	SM (exp)	SM (ref)
801	801	0,64	Hexanal	44(100), 41(99), 43(84), 56(83), 57(68), 27(50), 55(32)	44(100), 56(82), 41(66), 43(53), 29(38), 27(33), 72(19)
928	932	0,23	α -pinene	93(100), 91(36), 92(29), 77(28), 41(23), 79(22), 105(12), 121(11)	93(100), 91(45), 92(44), 77(36), 41(32), 79(27), 105(12), 121(11)
1040	1040	0,53	Z-ocimene	93(100), 91(52), 79(41), 92(39), 41(34), 77(33)	93(100), 79(50), 91(42), 41(40), 77(36)
1119	1121	3,09	Terpineol	43(100), 79(36), 139(34), 93(34), 71(32), 69(31), 111(30)	43(100), 93(46), 139(35), 41(33), 79(33), 1(32), 111(30)
1162	1162	0,43	Borneol	95(100), 41(25), 110(20), 43(19), 55(16), 39(15), 67(1), 93(11)	95(100), 41(30), 110(18), 43(16), 39(14), 55(13), 67(12), 93(10)
1193	1203	0,74	Trans- <i>p</i> -menth-2-en-1-ol	84(100), 83(39), 41(35), 139(35), 43(30), 91(20), 55(2)	84(100), 41(56), 43(38), 83(37), 55(34), 139(31), 91(28)
1281	1285	0,36	Bornylacetate	95(90), 43(70), 93(46), 121(39), 41(32), 55(20), 69(19)	95(100), 43(75), 93(45), 121(34), 41(27), 55(15), 69(13)
1399	1373	5.03	Decanoic acide	60(100), 73(91), 41(60), 43(54), 129(47), 55(45), 71(34), 29(30)	60(100), 73(88), 41(55), 43(55), 55(40), 129(38), 71(36), 29(34)
1415	1441	0.56	Ionon-5,6-epoxide	123(100), 43(57), 41(15), 39(10), 55(7)	123(100), 43(59), 41(25), 39(15), 55(14)
1571	1578	10.59	Spathulenol	43(100), 41(67), 91(63), 205(58), 119(52), 93(47), 79(44), 105(43), 159(34)	43(100), 41(69), 91(42), 119(42), 93(40), 105(34), 159(33), 205(33)
1573	1573	0,97	Caryophyllene oxide	43(100), 41(94), 79(30), 93(64), 91(60), 69(56), 95(42), 55(42), 67(41)	43(100), 41(93), 79(56), 93(66), 91(57), 95(42), 69(40), 55(39), 67(37)
1599	1600	0,83	Humulene epoxide2	43(100), 41(53), 55(35), 67(25), 109(21)	43(100), 109(44), 67(302), 55(23)
1608	1592	0,82	Calaren epoxide	41(100), 43(37), 69(34), 55(30), 91(26), 93(24)	41(100), 69(53), 93(41), 55(40), 91(39), 109(37)
1631	1612	0.69	Viridiflorol	43(100), 41(78), 81(50), 93(49), 55(47), 79(43)	43(100), 41(68), 81(33), 55(32), 93(30), 79(28)
1636	1640	1.59	Cadinol	161(100), 43(44), 204(36), 105(26), 41(25), 81(24), 119(19), 79(18)	161(100), 43(94), 41(47), 204(47), 105(39), 81(36), 79(29), 119(26)
1656	1654	4.70	Isoelemicin	208(100), 193(89), 77(20), 79(17), 133(15), 209(14), 165(13)	208(100), 193(69), 209(16), 79(10), 77(9), 133(7), 165(7)
1664	1675	0.54	Asaron	208(100), 193(48), 77(20), 91(19)	208(100), 193(49), 91(17), 77(15)
1981	1962	2,65	Palmitic acide	43(100), 41(87), 73(83), 55(75), 60(71), 57(69), 69(48), 71(36),	43(100), 41(75), 60(58), 55(53), 73(52), 57(48), 69(24), 71(19),
2303	--	0,55	Heptacosane	57(100), 43(79), 71(64), 85(47), 41(40), 55(39), 69(21), 56(19), 29(18)	57(100), 43(80), 71(62), 85(42), 55(28), 41(28), 69(18), 56(14), 29(13)

It can be noticed that the most of the relative percentages of other compounds are insignificant. But in spite of these low values, the chromatogram shows that they represent a great part of the separated compounds. In fact, these substances constitute also the essential oil of the studied plant and give it certainly the pleasant characteristic odor. More careful studies are required to clarify this chemical composition. As it can be seen on the table 1, more than 90 % of main composition of *M. pubescens* essential oil is constituted by oxygenated terpenes. Spathulenol (10.59%) was identified as the major constituent, in addition of isoelemicin (4.70%) and decanoic acid (5.03%). In previous study (Tadrent W. et al., 2014) spathulenol, with a close percentage (0.1%-19.4%) is the only common compound in the main composition of the essential oil of *M. pubescens*. Also, it was reported other main constituents in this oil: α -bisabolol oxide B (0.1%-7.0%), α -bisabolol (5.2%-56.9%) and α -bisabolol oxide A (10.2%-53.6%). That probably returns to the different collection origins of this plant, which are related to abiotic factors such as the specific region climate and geographical factors such as altitude and the nature of the soil of Biskra and its surroundings.

Antimicrobial assay: Essential oil of *M. pubescens* presented a variable efficiency against the tested germs; results are reproduced on table 2. In fact, *E. cloacae*, *E. coli* and *S. aureus* revealed the great sensitivity to sample 1 but *E. faecalis* and *K. pneumoniae* strains were more resistant towards this sample. Table 2 shows also that only the pure oil has presented medium to weak action against these last bacteria in addition of *P. mirabilis*, when the other dilutions from sample 1 were inactive. Generally, the absence of a zone of inhibition does not necessarily mean the inactivity of the tested sample, sometimes some product propagates slowly in the culture medium that delays their effect (Bensizerara et al., 2013). Otherwise, as it is shown on table 3 all bacteria isolated from patients were strongly resistant to both tested samples; except a moderate effect of sample 1 against *E. coli*. In comparison with the previous results, it is possible that this negative action is not due to the inefficiency of our samples but to these microbial strains themselves. In fact, they are frequently used in our hospitals where they show this strong resistance to the prescribed antibiotics. Concerning the inefficiency of the fragrant water, it returns probably to its hydrophilic composition that could not penetrate through the lipophilic membranes of microorganisms. Other studies such as (Mighri et al. (2001) reported that essential oils from an Asteraceae had an antibacterial activity against *S. aureus* and *P. aeruginosa*.

Table 2: Diameter (mm) of the inhibition zone.

% of E.O	<i>E. cloacae</i>	<i>P. mirabilis</i>	<i>E. faecalis</i>	<i>K. pneumoniae</i>	<i>E. coli</i>	<i>S. aureus</i>
100%	+++	++	+	+	++	++
75%	+++	-	-	-	++	++
50%	++	-	-	-	++	++
25%	++	-	-	-	++	++

(+++): 20mm; (++) $10 \leq D \leq 20$ mm; (+): $D \leq 10$ mm; (-): negative

Table 3: Results of microbial sensibility (microbial strains from patients).

Microbial strains	Samples	
	E.O	Odorant water
<i>E.coli</i>	S (++)	R
<i>Streptocoque</i>	R	R
<i>S. aureus</i>	R	R
<i>K. pneumoniae</i>	R	R
<i>Aspergillus niger</i>	R	R
<i>Candida albicans</i>	R	R

CONCLUSION

The composition of the essential oils of *M. pubescens* species growing in Biskra (Algeria) was partially determined. A screening of the antimicrobial activity of the isolated essential oil and its fragrant water were also determined. GC-MS analysis revealed the presence of spathuleol (10.59%), decanoic acid (5.03%), isoelemicin (4.70%), terpineol (3.09%), palmitic acid (2.65%), cadinol (1.59%). Microbial strains of reference were more sensitive to the essential oil however; microbial strains taken from patients were strongly resistant. In both cases, fragrant water was inactive against tested microorganisms. The specific chemical composition influenced clearly such activity.

REFERENCES

1. Adams, R.P., 1989. Retention indexes. Identification of Essential Oil Components by Gas chromatography/Quadrupole Mass Spectroscopy. Academic Press, New York, Appendix I,
2. Bélanger, A., Collin, G., Garneau, F-X., Gagnon, H., Pichette, A., 2008, Aromas From Quebec. II. Composition of the Essential Oil of the Rhizomes and Roots of *Asarum canadense* L., J. Essential Oil Res., 22, 164-169.
3. Boussaada, O., Ammar, S., Saidana, D., Chriaa, J., Chraif, I., Daami, M., Helal, A. N., Mighri, Z., 2008, Chemical composition and antimicrobial activity of volatile components from capitula and aerial parts of *Rhaponticum acaule* DC growing wild in Tunisia. Microbiol. Res., 163: 87-95.
4. Celikel, N., Kavas, G., 2008, Antimicrobial properties of some essential oils against some pathogenic microorganisms. Czech J. Food Sci., 26: 174–181.
5. Chemli R., 1997, plantes médicinales et aromatiques de la flore de Tunisie. CIHEAM - Options Méditerranéennes, 23, 119-125.
6. Gowsiya, S., Sujatha, N., Santosh, K. M., 2014, Anti-bacterial *K. pneumoneae* Medicinal plants as source of antibacterial agents to counter *Klebsiella pneumonia*, J. App. Pharm. Sci., 4 (01), 135-147.
7. <http://latitudelongitude.org/dz/biskra>. Accessed 27 / 02 / 2019.
8. http://www.flavornet.org/d_kovats_ov101.html, Accessed 27 / 02 / 2019.
9. Kaloustian, J., Chevalier, J., Mikail, C., Martino, M., Abou, L., Vergnes, M. -F., 2008, Étude de six huiles essentielles : composition chimique et activité antibactérienne, Phytotherapy, 6(3),
10. Maiza, K., Longeon, A., Hammiche, V., Guyot, M., Benabdesselam-Maiza, F., 2014. Biological Activities Of Plants Collected In The Algerian Sahara. Life Sciences Leaflets, 52, 52-56.
11. Makhloufi, A., Bouyahyaoui, A., Seddiki, N., Benlarbi, L., Mebarki, L., Boulanouar, A., 2014, Phytochemical screening and anti-listerial activity of essential oil and crude extracts from some medicinal plants growing wild in Bechar (south west of algeria), Int. J. Phytoth., 4 (2), 95-100.
12. Bouyahyaoui, A., Seddiki, N., Benlarbi, L., Mebarki, L., Boulanouar, A., 2014, Phytochemical screening and anti-listerial activity of essential oil and crude extracts from some medicinal plants growing wild in Bechar (south west of algeria), Int. J. Phytoth., 4 (2), 95-100.
13. Tadrent, W., Kabouche, A., Touzani, R., Kabouche, Z., 2014, Chemotypes investigation of essential oils of “Guertoufa” herbs, J. Mater Environment Sciences. 5 (4), 1200-1205.
14. Van Den Dool, H., Kratz, P. D., 1963, A generalization of the retention index system including linear temperature programmed gas-liquid partition chromatography, Journal of Chromatography, II, 463-471.
15. Yeşil-Çeliktaş, O., Hameş-Kocabaş E.E., Bedir E., Vardar-Sukan F., Özek T., Başer K.H.C., 2007, Antimicrobial activities of methanol extracts and essential oils of *Rosmarinus officinalis*, depending on location and seasonal variations. Food Chem., 100: 553–559.
16. Zhu, M., Li, E., He, H., 2008, Determination of volatile chemical constituents in tea by simultaneous distillation extraction, vacuum hydrodistillation and thermal desorption, Chromatographia, 68, 7(8), 603-610.

CHARACTERISATION OF VOLATILE COMPOUNDS OF *MENTHA PULEGIUM* VIA HS-SPME

Ela Nur Şimşek Sezer*, Meryem Bozkurt, Nihal İlhan, Tuna Uysal

Department of Biology, Faculty of Art and Science, University of Selçuk, Konya, Turkey,

*correspondence: elasimsek@selcuk.edu.tr

Abstract

Mentha pulegium L. is a herbaceous perennial plant belonging to the Lamiaceae family. The plant grows up in moist areas of the mountains and plains. In therapeutic applications, this plant and its active constituents have been used traditionally for several disorders. Their diuretic, antispasmodic, digestive, diaphoretic, sedative, antiseptic, stimulant and expectorant effects have been reported before. The aim of this study is the determination of the volatile and semi-volatile oil content of *Mentha pulegium*. The studied species was collected from its natural habitat. The powdered plant samples were analysed directly on a Shimadzu QP2010 system. Two SPME fibres preferred for analyses; 85 µm Carboxen/PDMS and 85 µm polyacrylate. The volatile oil compositions were identified in *Mentha pulegium* samples by SPME-GC/MS analysis. According to the analyses, significant differences were observed between the two SPME fibres. The major volatile component for 85 µm Carboxen/PDMS fiber were 2-Isopropyl-5-methylhex-2-enal (30.37%) and for 85 µm polyacrylate fiber 2(3H)-Benzofuranone, 3a, 4, 7,7a-tetrahydro-3a-methyl (48.59%), respectively. In addition, the most common and shared components to both fibers were, p-Menthone (14.15%- 8.82 %) and thymol (phenol, 5-methyl-2-(1-methyl ethyl)) (8.68 %- 7.67 %). In conclusion, *Mentha pulegium* is a plant that is rich in essential oils, and this study is the first study about the volatile content of plant with SPME method.

Keywords: Yarpuz, SPME-GC-MS, Turkey.

INTRODUCTION

Mentha is a member of the Lamiaceae family and is distributed throughout the world. *Mentha pulegium* L. is one of the *Mentha* species and is mostly known as pennyroyal (Chalchat et al., 2000). *Mentha* is a perennial herb and has a pungent odour, is grown over a wide area in Turkey (Eryigit, 2006). The plant is native to North Africa, Europe and the Middle East (Aghel et al., 2004; Debbab et al., 2009). The plant contains a variety of secondary metabolites, such as essential oils, resins, tannins, and pectins (Simon et al., 1984; Zargari, 1990). *Mentha pulegium* L. is a plant rich in terms of essential and volatile oils, especially. As far as we know the volatile content of pennyroyal have reported at several studies before and pulegone is the principal volatile component (Zargari, 1990; Lorenzo et al., 2002; El Fadl & Chtaina, 2010; Benayad et al., 2008; Boughdad, 2011). And also the chemical composition of pennyroyal oil is defined by various studies (Mkaddem et al., 2007; Benayad, 2008; Hajlaoui et al., 2009; Derwich et al., 2010; Boughdad, 2011). Pennyroyal oil is characterised by the preponderance of pulegone (70-90%) along with the other monoterpenic ketones such as menthone, isomenthone and piperitenone (Bruneton, 2009). The pennyroyal oil and its constituents exhibit interesting biological activities such as antibacterial, antioxidant, antifungal and insecticidal (Jazani et al., 2009; Derwich et al., 2010; Hmiri et al., 2011; Alpsy et al., 2011; Benayad et al., 2008; Boughdad, 2011).

Firstly, to understand the biological effect of the plants, it is necessary to know its volatile components. Gas chromatography–mass spectrometry (GC–MS) is commonly used to determine the volatile components of various aromatic and medicinal plant species. According to the literature, headspace solid phase microextraction (HS-SPME) and gas chromatography-mass spectrometry (GC-MS) has gained broad approval as an effective extraction technique for various samples in the last twenty years (Cuevas-Glory et al., 2007; Panighel, & Flamini, 2014; Xu et al., 2016). Even though the volatile

composition of *Mentha pulegium* has been studied before, there is not enough research about SPME-GC / MS in the literature. At present study aims to reveal the volatile constituent of *Mentha pulegium* species naturally grown in Turkey by using HS-SPME technique.

MATERIAL AND METHODS

Material: The *Mentha pulegium* samples were collected from its natural habitat. The plant samples are dried at room temperature without sunlight. Dried plants pulverised by helping a hand grinder. The ground samples (3gr) sealed in a 10 ml vial. The SPME fibres preferred for analysis were 85 µm Carboxen/PDMS and 85 µm polyacrylate. The SPME apparatus was directly injected into the upper space of the vial to adsorb volatile substances and then directly injected into the Shimadzu QP2010 ULTRA GC-MS apparatus using a Restek Rxi-5 MS capillary column.

HS-SPME-GC/MS Analysis: The volatile components were separated for about 15 minutes with different SPME fibers attached to a GC injector (the temperature of the injector was 250 °C). The plant material analysed on a Shimadzu QP2010 ULTRA FID GC-MS system. A Restek Rxi-5 MS column (30 m×0.25 mm id, film thickness 0.25 µm) used with helium as a carrier gas. The GC oven temperature was programmed to hold at 40 °C for 3 min and then to increase to 240 °C at 5 °C/min, finally holding at 240 °C for 3 min. The column flow rate was 1.8 mL/min. The ion source temperature was 200 °C, and the interface temperature set at 250 °C.

Evaluation of data: After the analysis, each sample was demonstrated with GC-MS chromatograms. Chromatograms of all samples were exposed to noise reduction before peak area integration, and then the peak areas of compounds in the chromatogram were combined. Compounds were identified by comparison with three libraries, W9N11, SWGDR4G4 and SWGDR4G5. Compounds were mostly found in the W9N11 library and specified with other libraries. Identification was performed based on the retention time (RT) of the components in the sample. The relative percentage of the volatile oil constituents was calculated from peak areas. All analyses were carried out triplicate, and all data are means of three independent analyses.

RESULTS AND DISCUSSION

The results obtained from the analysis of pennyroyal by two different SPME fibres are shown in **Table 1**. Totally 102 compounds were separated and identified from the studied samples. The major constituents of the investigated taxa with 85 µm Carboxen/PDMS fiber were 2-Isopropyl-5-methyl-2-hexenal (30.37 %), p-menthone (14.15 %) and thymol (8.68 %). For the other fiber 85 µm polyacrylate; 2(3H)-Benzofuranone, 3a, 4, 7, 7a-tetrahydro-3a-methyl (48.59 %), p-menthone (8.82 %) and thymol (7.67 %). 24 compound is common for two fibres. As reported earlier, the volatile components in *Mentha* taxa have various biological activities, usage areas and medical values (Su et al. 2014). One of the most common compounds is 2-isopropyl-5-methyl-2-hexenal, which is used as a flavouring component. Another compound menthone is a mono terpen, which naturally has a minty taste in some essential oils. Menthone is used in flavour, perfume and cosmetic products with its characteristic aromatic and tiny scent. According to our analyses, there were precious components available in the content. For example, α-pinene (0.12 %), β-pinene (0.11 %), Carvacrol (1.28 %), Limonene (0.13 %) and 1,8-Cineole (0.15 %). Our results are compatible with previous studies. Zekri et al., (2013) reported the volatile content of *Mentha pulegium* L. oils from from three regions of Morocco and they found similar results with us. Díaz-Maroto et al., (2007) studied the volatile composition of pennyroyal oil extracts via SDE method. In this study, 41 components were determined, and rates were a little more than our results. This difference may result from the used method. The SPME method which we preferred, is much more sensitive in identifying the compounds, and this sensitivity allows identification of small components in the content. Furthermore, the essential oil content of a plant can vary greatly depending on the plant's habitat, chemotype and process.

Table 1. The volatile composition of pennyroyal plant via HS-SPME (RT=Retention time, nd=not detected)

	Compound Name	Fibres			
		85 µm Carboxen/PDMS		85 µm polyacrylate	
		RT	% area	RT	% area
1.	Hexanal	4,357	0.53	nd	-
2.	Heptanal	7,834	0.14	nd	-
3.	Alpha pinene	9,009	0.12	nd	-
4.	Beta pinene	10,675	0.11	nd	-
5.	3-Octanol	11,596	2.31	11,620	1.18
6.	Octanal	11,843	0.07	nd	-
7.	Benzene, methyl(1-methylethyl)	12,645	1.51	nd	-
8.	Limonene	12,812	0.13	nd	-
9.	1,8-Cineole	12,897	0.15	nd	-
10.	2-Ethylhexanol	12,970	0.06	nd	-
11.	Ocimene	13,285	0.07	nd	-
12.	Cyclopentane, 1-methyl-2-propyl-	13,595	0.07	nd	-
13.	p-menth-2-en-1-ol	14,037	0.27	nd	-
14.	trans sabinene hydrate	14,368	0.40	14,388	0.11
15.	Cis-sabinene hydrate	15,599	0.26	nd	-
16.	Linalool	15,737	3.84	15,746	1.52
17.	Nonanal	15,888	0.12	nd	-
18.	1-(3,4-methylenedioxyphenyl) ethane-1-ol	16,374	0.06	nd	-
19.	Octyl cyclobutanecarboxylate	16,725	0.12	nd	-
20.	Cis-Verbenol	17,450	0.14	nd	-
21.	2-Isopropyl-5-methyl-2-hexenal	17,797	30.37	nd	-
22.	Nonenal	18,029	0.17	nd	-
23.	p-Menthone	18,166	14.15	17,772	8.82
24.	Borneol I	18,259	1.74	18,243	0.64
25.	Isopulegone	18,615	5.96	18,623	1.86
26.	Z-citral	19,160	0.23	nd	-
27.	1-alpha-terpinyl acetate	19,271	0.71	nd	-
28.	3-Decanol	19,595	0.06	nd	-
29.	1-Tetradecene	19,832	0.11	nd	-
30.	2-Cyclohexen-1-one, 2-(2-methyl-2-propenyl)	19,930	0.24	19,915	0.04
31.	Cis piperitone oxide	20,191	6.86	20,187	2.73
32.	Carveol	20,465	0.45	20,365	0.35
33.	Alpha sinensal	20,836	0.27	20,719	0.37
34.	1,2-cyclohexanediol, 1-methyl-, trans-	21,439	0.38	nd	-
35.	Piperitone	21,562	0.32	21,551	0.17
36.	2-Decenal	21,840	0.07	nd	-
37.	2,6,6-trimethyl-cyclohex-1-enecarboxylic acid	21,904	0.51	21,904	0.25
38.	Carvone	22,192	0.18	22,192	0.06
39.	2-Acetyl-4, 4-dimethyl-cyclopent-2-enone	22,616	0.12	22,619	0.12
40.	Bicyclo[3.2.0]heptan-2-one, 5-formylmethyl-6-hydroxy-3,3-dimethyl	22,824	3.04	22,828	2.40
41.	Thymol	23,050	8.68	23,058	7.67
42.	Linalyl butyrate	23,244	0.07	nd	-
43.	Carvacrol	23,366	1.28	23,376	1.13
44.	1-hexadecyl-3-ol, 3,7,11,15-tetramethyl	23,571	0.10	23,578	0.05
45.	2,4-Decadienal	23,795	0.30	nd	-
46.	Cyclohexasiloxane, dodecamethyl-	24,360	0.26	nd	-
47.	2-Cyclohexen-1-one, 3-methyl-6-(1-methylethylidene)	24,644	1.99	24,653	1.69
48.	Trans, cis,-3-ethylbicyclo[4.4.0]decane	24,840	0.40	24,947	0.30
49.	2-Heptyl furan	24,937	0.50	nd	-
50.	2-Hexen-1-ol, propanoate	25,060	0.20	nd	-
51.	2-Cyclohexen-1-one, 2-hydroxy-6-methyl-3-(1-methylethyl)	25,289	0.19	nd	-
52.	Cyclohexanone, 2-(1-methylethylidene)-	25,524	0.06	nd	-
53.	Butyric acid, dodecyl ester	25,770	0.07	nd	-

MESMAP-5 PROCEEDINGS BOOK
 April 24th - 26th, 2019 / Cappadocia – Turkey

54.	Copaene	25,894	0.15	nd	-
55.	Beta bourbonene	26,199	0.18	nd	-
56.	Beta funebrene	26,400	0.23	nd	-
57.	Cineralone	26,706	0.24	nd	-
58.	Caryophyllene	27,360	0.69	27,371	0.05
59.	Carane, 4,5-epoxy-, trans	27,855	1.44	nd	-
60.	Aromadendrene	28,007	0.53	28,018	0.05
61.	2(1H)-Naphthalenone, octahydro-3-methyl-, (3alpha,4abeta,8aalpha)	28,157	0.26	nd	-
62.	Alpha humulene	28,485	0.91	28,496	0.07
63.	Hotrienol	29,144	0.16	nd	-
64.	Gamma muurolene	29,238	0.18	nd	-
65.	Germacrene-D	29,381	0.34	nd	-
66.	2-hydroxy-4', 5'-dimethylacetophenone	29,762	0.37	29,772	0.42
67.	Beta bisabolene	30,248	1.89	30,255	0.46
68.	Naphthalene, 1,2,3,4,4a,5,6,8a-octahydro-7-methyl-4-methylene-	30,434	0.09	nd	-
69.	Delta cadinene	30,723	0.13	nd	-
70.	Mint furanone II	30,859	0.75	29,849	0.78
71.	Limonen-6-ol, pivalate	31,858	0.20	31,868	0.07
72.	Caryophyllene oxide	32,575	0.07	32,585	0.02
73.	12-Oxabicyclo[9.1.0]dodeca-3,7-diene, 1,5,5,8-tetramethyl-	33,371	0.12	nd	-
74.	Laurinsaeure, 4-octylester	46,362	0.07	nd	-
75.	Phospine oxide, trihenyl-	55,882	0.44	nd	-
76.	M-Cymene	nd	-	12,669	0.11
77.	Bicyclo[3.1.1.1.]hept-3-en-2-ol, 4, 6, 6-trimethyl	nd	-	17,444	0.06
78.	p-Menthan-3-one	nd	-	18,165	3.28
79.	Trans-p-Mentha-(7), 8-dien-2-ol	nd	-	19,110	0.05
80.	Alpha terpineol	nd	-	19,224	0.34
81.	Cyclohexanone, 2-isopropyl-2,5-dimethyl	nd	-	20,000	0.34
82.	Methanone, (1-hydroxycyclohexyl)phenyl	nd	-	20,296	0.30
83.	1,6,6-Trimethyl-8-oxabicyclo[3.2.1.]	nd	-	20,630	0.04
84.	2(3H)-Benzofuranone, 3a, 4, 7, 7a-tetrahydro-3a-methyl	nd	-	21,133	48.59
85.	benzene, 1-methoxy-4-methyl-2-(1-methylethyl)	nd	-	21,226	1.74
86.	1-(1,1-Dimethylethoxy)-6-methyl-1-cyclohexene	nd	-	21,414	0.25
87.	13-Octadecenal	nd	-	23,818	0.04
88.	Benzene, 2,4-diisocyanato-1-methyl-	nd	-	25,082	5.66
89.	2-cyclohexen-1-one, 3,4,4-trimethyl-	nd	-	25,534	0.09
90.	7-Isopropyl-7-methyl-nona-3,5-diene-2,8-dione	nd	-	26,730	0.54
91.	Pyrrrole-3-carbonitrile, 5-formyl-2,4-dimethyl	nd	-	26,793	0.56
92.	2-hydroxymethyl benzimidazole	nd	-	26,890	1.47
93.	Pulegone	nd	-	27,864	0.73
94.	Cyclohexane 1-methyl-4-(1-methylethylidene)	nd	-	28,166	0.21
95.	2,5-methano-1H-inden-7-ol, octahydro-	nd	-	29,154	0.04
96.	Beta bisabolene	nd	-	30,255	0.46
97.	Mint furanone I	nd	-	30,871	0.19
98.	Spathulenol	nd	-	32,410	0.02
99.	12-oxabicyclo[9.1.0]dideca-3, 7-diene	nd	-	33,392	0.06
100.	1(2H)-Quinolinecarboxylic acid, 6-amino	nd	-	36,197	0.11
101.	4-Isopropoxybutanol	nd	-	40,862	1.12
102.	2,5-Dimethyl-4-hydroxy-3-hexanone	nd	-	52,186	0.66

CONCLUSION

In this study, *Mentha pulegium* species from Turkey was analysed via HS-SPME-GC/MS method. As far as we know this study is the first study about the volatile composition of the plant with the SPME method. It is crucial to reveal the valuable natural compounds which have a large number of effects in the different plant species. Such studies, which are attempting to determine the active substances from plants, are of great importance in terms of more specific studies to be carried out in the future.

ACKNOWLEDGEMENTS

We want to thank the BAP (Scientific Researching Projects) Foundation of Selçuk University for their financial support (Project number 19704015).

REFERENCES

1. Aghel, N., Yamini, Y., Hadjiakhoondi, A., & Pourmortazavi, S. M. 2004. Supercritical carbon dioxide extraction of *Mentha pulegium* L. essential oil. *Talanta*, 62(2), 407-411.
2. Alpsoy, L., Sahin, H., Karaman, S. 2011. Anti-oxidative and anti-genotoxic effects of methanolic extract of *Mentha pulegium* on human lymphocyte culture. *Toxicol Ind Health*. 27(7), 647-654.
3. Benayad, N. 2008. Les huiles essentielles extraites des plantes médicinales marocaines: Moyen efficace de lutte contre les ravageurs des denrées alimentaires stockées. Research Project. University of Sciences, Rabat (Maroc), 63p.
4. Boughdad, A., Elkasimi, R. and Kharchafi, M. 2011. Activité insecticide des huiles essentielles de *Mentha* Sur *Callosobrochus maculatus* (F) (Coleoptera, Bruchidea). AFPP – Neuvième Conférence Internationale sur les ravageurs en Agriculture. Montpellier, 9p.
5. Bruneton, J. 2009. Pharmacognosie, Phytochimie et plantes médicinales. 4^{rt}Edt. Technique et Documentation., 1268p.
6. Chalchat, J.C., Gorunovic, M.S., Maksimovic, Z.A., Petrovic, S.D. 2000. Essential oil of wild growing *Mentha pulegium* L from Yugoslavia. *Journal of Essential Oil Res.*, 12, 598-600.
7. Cuevas-Glory, L. F., Pino, J. A., Santiago, L. S., & Sauri-Duch, E. 2007. A review of volatile analytical methods for determining the botanical origin of honey. *Food Chemistry*, 103(3), 1032–1043.
8. Debbab, A., Aly, A. H., Edrada-Ebel, R., Wray, V., Müller, W. E., Totzke, F., ... & Mosaddak, M. 2009. Bioactive metabolites from the endophytic fungus *Stemphylium globuliferum* isolated from *Mentha pulegium*. *Journal of Natural Products*, 72(4), 626-631.
9. Díaz-Maroto, M. C., Castillo, N., Castro-Vázquez, L., Ángel González-Viñas, M., & Pérez-Coello, M. S. 2007. Volatile composition and olfactory profile of pennyroyal (*Mentha pulegium* L.) plants. *Flavour and Fragrance Journal*, 22(2), 114-118.
10. Derwich, E. Benziane, Z. and Boukir, A. 2010. GC/MS Analysis and Antibacterial Activity of the Essential oil of *Mentha Pulegium*. *J.Agr.Biol.Sc.* 6(3), 191-198.
11. Eryigit, F. 2006. *Mentha pulegium* L. ve *Salvia tomentosa* Miller bitkilerinin metanol özütlerinin in vitro antioksidan aktivitelerinin belirlenmesi. Yüksek lisans tezi, Kimya Anabilim Dalı, Isparta.
12. El Fadl, A. and Chtaina, N., 2010. Etude de base sur la culture de la menthe du Maroc. Programme Régional de lutte intégrée contre les organismes nuisibles (Integrated Pest Management) au Proche Orient. Report, Office National de sécurité sanitaire des produits alimentaires (ONSSA).
13. Hajlaoui, H. Trabelsi, N. Noumi, E. Snoussi, M. Fallah, H. Ksouri, R. and Bakhrouf, A. 2009. Biological activities of the essential oils and methanol extract of tow cultivated mint species (*Mentha longifolia* and *Mentha pulegium*) used in the Tunisian folkloric medicine. *Springer Science. World. J. Microbiol. Biotechnol.* 25, 2227-2238.
14. Hmiri, S. Rahouti, M. Habib, Z. Satrani, B. Ghanmi M. and El Ajjour, M. 2011. Évaluation du potentiel antifongique des huiles essentielles de *Mentha pulegium* et d'*Eucalyptus Camaldulensis* dans la lutte biologique contre les champignons responsables de la détérioration des pommes en conservation. *Bulletin de la Société Royale des Sciences de Liège.* 80, 824 – 836.
15. Jazani, N.H. Ghasemnejad-Berenji, H. and Sadegpoor S. 2009. Antibacterial Effects of Iranian *Mentha pulegium* Essential Oil on Isolates of *Klebsiella* sp. *Pakistan J.Biol.Sc.* 12 (2), 183-185.
16. Lorenzo, D., Paz D., Dellacassa, E., Davies, P., Vila, R. and Canigüeral, S., 2002. Essential Oils of *Mentha pulegium* and *Mentha rotundifolia* from Uruguay. *Braz. Arch.Biol.Technol.* 45 (2), 519-524.
17. Mkaddem, M. Bousaid, M., and Ben Fadhel, N. 2007. Variability of volatiles in 9 Tunisian *Mentha pulegium* L. (Lamiaceae), *Journal of Essential Oil Research.* 19, 211-215.
18. Panighel, A., & Flamini, R. 2014. Applications of solid-phase microextraction and gas chromatography/mass spectrometry (SPME-GC/MS) in the study of grape and wine volatile compounds. *Molecules*, 19(12), 21291–21309.
19. Simon, J.E., Chadwick, A.F., Craker, L.E., 1984. An Indexed Bibliography: 1971–1980: The Scientific Literature on Select Herbs, and Aromatic and Medicinal Plants of the Temperate Zone. Elsevier, Amsterdam, New York.
20. Su X., Liu Y.P., Chen K.L. 2014. Analysis on Volatile Components in *Mentha haplocalyx* of Special Herbs in Qinghai Province. *Asian Journal of Chemistry.* 26: 555–558.
21. Xu, C.H., Chen, G.S., Xiong, Z.H., Fan, Y.X., Wang, X.C., & Liu, Y. 2016. Applications of solid-phase microextraction in food analysis. *Trac- Trends in Analytical Chemistry*, 80, 12–29.
22. Zargari, A., 1990. Herbal Medicines. 1st Edt, Publication of Tehran University, Tehran.14-18.
23. Zekri, N., Amalich, S., Boughdad, A., El Belghiti, M. A., & Zair, T. 2013. Phytochemical study and insecticidal activity of *Mentha pulegium* L. oils from Morocco against *Sitophilus Oryzae*. *Mediterranean Journal of Chemistry*, 2(4), 607-619.

EFFECTS OF OLIVE LEAF (*OLEA EUROPEA L.*) EXTRACT ON THE GROWTH PERFORMANCE AND SOME BLOOD PARAMETERS OF COMMON CARP (*CYPRINUS CARPIO*)

Ebru Yilmaz¹, Deniz Çoban², Murat Er³

¹Adnan Menderes University, Bozdoğan Vocational School, Aydın, Turkey

²Adnan Menderes University, Faculty of Agriculture, Department of Aquaculture, Aydın, Turkey

³Adnan Menderes University, Faculty of Veterinary Medicine, Department of Animal Nutrition and Nutritional Diseases, Aydın, Turkey

corresponding author: ebruyilmaz@adu.edu.tr

fact that oleuropein has many pharmacological qualities including antioxidant, antimicrobial, antiatherogenic and antiviral activities. In this study, the effects of olive leaf extract on growth performance and some blood parameters of common carp were evaluated. For this purpose, 3 different experimental diets containing rate of 0% (OLE0), 0.25% (OLE025) and 0.5% (OLE05) olive leaf extract were prepared. Olive leaf extract were obtained from Talya Herbal product. The olive leaf extract that we used in study include oleuropein at the rate of 2.3%. This study consisted of 3 repetitions for each group with 71.58±0.66 g and 200 L aquarium as 25 fish per aquarium for 60 days. The objective of this study was to determine the effects of olive leaf extract on the growth performance and some blood parameters in the common carp. At the end of the experiment, the weight increase, specific growth rate and feed conversion rate were found to be similar to the trial groups (p>0.05). The dietary supplementation of OLE did not affect the RBC count, hemoglobin concentration, hematocrit ratio, mean corpuscular hemoglobin concentration, mean corpuscular volume and mean corpuscular hemoglobin (p>0.05). In serum biochemical findings of fish, olive leaf extract was found to be a significant difference among the groups (p<0.05). Results in the present study showed that the olive leaf extract can use in common carp diets without adverse effects on growth performance, hematological and serum biochemical parameters of fish.

Key words: *Cyprinus carpio*, Olive leaf (*Olea europea L.*) extract, Blood parameter, Growth performance

INTRODUCTION

Olive (*Olea europaea L.*) and its derivatives are very important products in both Turkey and the surrounding Mediterranean and Middle East regions (Bircan, 2006). Olive leaf is the most important source of oleuropein known in nature. Several researchers revealed the fact that Oleuropein has many pharmacological qualities including antioxidant, antimicrobial, anti-inflammatory, antiatherogenic and antiviral activities (Visioli et al., 1998; Owen et al., 2000). In addition, it is stated that oleuropein has a strong antioxidant effect with its quality of bonding endogenous peptides in addition to its anti-inflammatory, anti-atherosclerotic and anticarcinogenic properties (Gikas et al., 2007). Recent studies determined that the medical plants are effective on the growth parameters of fishes (Yilmaz et al., 2017a; Yilmaz et al., 2017b). Also, It has been documented that natural products such as, *Azadirachta indica* extract (Harikrishnan et al., 2003), *Astragalus radix* and *Ganoderma lucidum* (Yin et al., 2009), stimulate the immune system and disease resistance in *Cyprinus carpio*.

MATERIAL AND METHODS

Olive leaf extract used in the present study was obtained from Talya Herbal product, and was supplemented in test diets at 0%, 0.25% and 0.5%, which were then named as OLE0,OLE025,OLE05 groups, respectively. In this experiment commercial fish diet (48% crude protein, Gümüşdoğa, Muğla) was used (Table 1). A control group was prepared without any supplements. The addition of olive leaf extract to feed was carried out by the spraying method. Feeds were prepared weekly and stored in capped glass bottles at +4°C to preserve the efficacy of the components of the olive leaf extract. During the experiment that lasted 60 days, fishes were manually fed 3 times a day in a ratio of 2% of their body weight.

Table 1. Proximate analysis of the commercial diet (2 mm) and olive leaf extract *

Nutrient	Diet percentage%
Crude Protein (%)	48
Crude Lipid (%)	18
Crude Cellulose (%)	3
Crude Ash (%)	12
Moisture (%)	12
Macro elements (%)	
Calcium (%)	1.2
Total Phosphore %	1.5
Meth+Cyst (%)	1.8
Lysine (%)	2
Compound	Olive leaf extract (OLE) percentage %
Oleuropein	2.3
Epicatechin	4.7
Quercitrin	5.2
Rutin	4.1
Quercetin	7.5
Vanillic acid	2.3

Ingredients: Fish oil, fish meal, soybean and by products, wheat and by products, yeast and by products, amino acids, vitamins and minerals. The ratios of phenolic compounds in the extracts of olive leaf was taken from Talya Herbal Company (Certificate of analysis 2018)

Cyprinus carpio fishes were obtained from a local trout fish farm in İzmir, Turkey. The US-EPA (United States Environmental Protection Agency) guidelines for qualitative assessment of fish health were followed for the visual and external inspection of each fish (Klemm et al., 1993). At the start of the trial, a total of 225 fish with initial mean weight of 71.58 ± 0.66 g were randomly distributed into nine 200-L aquariums (25 fish per aquarium). During the experimental period water quality identified as follows: temperature $24.40 \pm 0.29^\circ\text{C}$, pH 7.11 ± 0.39 , salinity 0.5 ppt, total dissolved solid (TDS) 11.90 mg/l and dissolved oxygen 8.48 ± 0.21 mg/L. Proximate analyses of the diets were performed using standard methods (AOAC, 2000). Moisture was detected after drying at 105°C until a constant weight was achieved. Crude protein was analyzed by the Kjeldahl method, and crude ash by incineration at 525°C in a muffle furnace for 12 h. Crude fat was analyzed by methanol/chloroform extraction. The haematological and biochemical analyses were conducted in the private Detay Analysis Laboratory. At the end of the 60-day feeding trial, blood was taken from a total of 9 fish/ group, 3 fish from each aquarium. After the fish were caught randomly from each aquarium, they were anesthetized with clove oil (Mylonas et al., 2005). They were well wiped and cleaned in order to avoid mucus mixing into the blood, and blood was taken from the fish through the caudal vein by a 5-mL plastic syringe without harming the fish (Val et al., 1998). The blood samples were put into the K_3EDTA tubes and haematological and biochemical analyses were done. Haematological parameters, such as RBC count, HCT, hemoglobin (Hb) and mean cell hemoglobin concentration (MCHC), were determined by a blok cell counter (Mindray BC-3000 Plus, China) within 2 h. Blood serum was separated by centrifugation (3000g, 20 min). The clear supernatant was aspirated carefully for the estimation of various blood chemical parameters in serum. The biochemical parameters like creatinine, total protein, glucose, albumin, globulin, uric acid, urea, glutamic oxaloacetic transaminase, glutamic pyruvic transaminase and cholesterol were assessed by the automatic analyzer apparatus (Reflotron plus, Roche, Germany) using commercially available Reflotlon kits (Roche Diagnostics, Germany) for the above mentioned blood chemical parameters (Bain et al., 2006). The analyses of the data obtained in the experiment were done by using the SPSS 17 statistics program. One way variance analysis (ANOVA) was applied to the data which were then subjected to Tukey multiple comparison test. The differences between groups were evaluated to be $p < 0.05$. Fish experiments were performed in accordance to the guidelines for fish research from the animal ethic committees at Adnan Menderes University (Protocol Number:2018/085).

RESULTS

It was determined that the specific growth rate (SGR) values among the growth parameters showed similarity for all groups. With regard to survival rate and feed conversion ratio (FCR), the groups were similar to each other. Nutrient component analysis results showed that there was no statistically difference between moisture, lipid, crude protein and crude ash amounts of all groups (Table 2.)

Table 2. The growth performance and proximate composition of common carp

	Control	OLE 0.25%	OLE 0.50%
Initial fish weight (g)	71.82±0.88	71.26±0.81	71.65±0.40
Final fish weight (g)	134.71±4.04	136.44±0.82	137.39±2.66
Weight gain (%)	62.88±3.16	65.18±0.80	65.74±2.87
FCR	0.81±0.05	0.80±0.00	0.80±0.01
SGR (%/day)	1.04±0.03	1.08±0.01	1.08±0.03
Survival (%)	100	100	100
Crude Protein	12.32±0.25	12.77±0.87	12.55±0.25
Crude Lipid	3.29±0.38	3.80±0.56	3.38±0.06
Crude Ash	2.44±0.83	2.58±0.58	2.10±0.34
Moisture	76.29±0.99	75.27±0.88	75.78±0.79

*The different letters on the lines indicate that differences are important (p<0.05)

The hematological parameters were presented in Table 3. Olive leaf (*Olea europea* L.) extract used in study did not significantly affect on the hematological parameters (P>0.05).

Table 3. Hematological parameters of *C. carpio* fed diets containing olive leaf extract (OLE)

Blood Parameter	Control	OLE 0.25%	OLE 0.50%
Hb (g/dL)	13.74±0.20	14.06±0.18	13.96±0.19
Hct (%)	22.53±3.05	24.64±1.24	26.13±4.65
RBC (10 ⁶ mm ³)	1.48±0.15	1.49±0.09	1.55±0.16
MCV (µm ³)	153.70±26.55	165.52±3.45	168.60±30.84
MCH (pg cell ⁻¹)	93.66±9.57	94.69±6.76	90.50±10.08
MCHC (%)	61.79±7.36	57.18±3.36	54.81±9.64

Values are mean ±SE (n=9) Hct: hematocrit; Hb: hemoglobin; MCV: mean cell volume; MCH: mean cell hemoglobin; MCHC: mean cell hemoglobin concentration

The serum biochemical parameters were presented in Table 4. It was found that the values of albumin, globulin, glucose, glutamic oxaloacetic transaminase, glutamic pyruvic transaminase, total protein, cholesterol, urea, uric acid and creatinine among serum biochemical parameters have statistical difference among groups (P<0.05). Also, urea and uric acid levels were similar in all the treatment groups (p>0.05).

Table 4. Serum biochemical parameters of *C. carpio* fed diets containing olive leaf extract (OLE)

Parameter	Control	OLE 0.25%	OLE 0.50%
ALB (g dL ⁻¹)	1.04±0.13 ^b	1.18±0.03 ^{ab}	1.24±0.05 ^a
GLO (g dL ⁻¹)	3.70±0.09 ^c	4.06±0.20 ^b	4.36±0.11 ^a
TPROT (g dL ⁻¹)	4.80±0.03 ^c	5.31±0.08 ^b	5.64±0.11 ^a
GLU (U l ⁻¹)	192.±8.74 ^a	174.20±7.94 ^b	160.20±8.49 ^b
GOT (U l ⁻¹)	60.92±5.82 ^a	52.32±6.24 ^{ab}	44.47±5.12 ^b
GPT (U l ⁻¹)	27.59±1.56 ^a	20.34±1.00 ^b	18.37±1.00 ^b
CHOL (mg dL ⁻¹)	138.40±3.84 ^a	122.40±2.30 ^b	116.80±4.65 ^b
Urea (mg dL ⁻¹)	0.16±0.01 ^a	0.16±0.01 ^a	0.14±0.00 ^a
Uric acid (mg dL ⁻¹)	1.32±0.05 ^a	1.30±0.08 ^a	1.32±0.10 ^a
Creatinine (mg dL ⁻¹)	0.39±0.02 ^a	0.28±0.02 ^b	0.22±0.02 ^b

Values are mean ± SE (n=9). Different letters represent the significant differences at P<0.05 ALB.albumin; GLO.globulin, TPROT.total protein; GLU.glucose; GOT=glutamic oxaloacetic transaminase; GPT=glutamic pyruvic transaminase; CHOL.cholesterol ment. Values are mean ± SEM (n=3)

DISCUSSION

In the present study, olive leaf (*Olea europea* L.) extract did not change significantly average weight gain, specific growth rate (SGR) and feed conversion rate (FCR) ($p>0.05$). Similarly, in a study by Baba et al. (2018), it was observed that there is no change in the final body weight (FW), specific growth rate (SGR) and feed conversion ratio (FCR) of rainbow trout fed by a diet containing 0.25% and 0.50% olive leaf (*Olea europea* L.) extract ($p>0.05$). Haematological analyses are important in determining the fish health and decreases in the amount of erythrocyte and haematocrit as well as haemoglobin destruction are considered to be indicator of anemia. The amount of erythrocyte sedimentation gives information about the existence of diseases. Total leucocyte amount and types of it are used in the diagnosis of diseases (Blaxhall and Daisley, 1973). In this study, the olive leaf (*Olea europea* L.) extract used in study did not significantly affect on the red blood cell count, hematocrit, hemoglobin, mean corpuscular hemoglobin concentration, the mean corpuscular volume and mean corpuscular hemoglobin level ($p>0.05$). In our study, the serum biochemical parameters, i.e. the values of albumin, globulin and total protein were found to be higher in the groups fed with olive leaf extract than the control group ($P<0.05$). In a different study, low utilization of garlic decreased total protein (Sahu et al. 2007a). In a different study which involved addition of mango to feed, a decrease of albumin in the ratio of 0.5 g/100g was reported (Sahu et al., 2007b). In our study, it was observed that the blood glucose level in fishes fed by olive leaf extract were lower in the trial groups than the control group. In the study of Baba et al., (2018), it was observed that the blood glucose level didn't change in fishes fed by olive leaf. When we look at the studies involving the use of vegetable resource, it is noticed that the plants of *Cynodon dactylon*, *Aegle marmelos*, *Withania somnifera*, ginger, mango and garlic decreased the blood glucose values. The serum enzymes including GOT and GPT are the indicators of damages that occurred in tissue and organs (Campbell, 2004). Our study shows that olive leaf extract causes decrease in the amounts of GOT and GPT. Similarly, the study by Baba et al. (2018) found that the amounts of GOT and GPT decreased in *O. mykiss* fishes fed by the olive leaf extract. Another study found decrease in the amounts of GOT and GPT in the *O. niloticus* fishes fed by garlic (Shalaby et al., 2006). The present study found that the olive leaf extract decreased the cholesterol amount in *Cyprinus carpio*. In a study which involved the addition of *Cynodon dactylon*, *Aegle marmelos*, *Withania somnifera* and ginger to the feeds of the *Oreochromis niloticus*, it was found that cholesterol significantly decreased (Immanuel et al., 2009). In another study involving the addition of olive leaf extract to the trout feeds, it was seen that there was no change in the cholesterol amount (Baba et al., 2018). This shows that different plant species may have different effects on fat metabolism. The reason of it may be explained by the fact that the components of plants and their mechanisms of action are different. Therefore, the increases of urea in blood are considered to be an indicator of gill and liver diseases rather than kidney (Stoskopf, 1993; Campbell, 2004). Creatinine in fishes consists of creatine and excreted by kidneys (Campbell, 2004). Uric acid in fishes consists of purine nucleotides as exogen and endogen. Later on, uric acid is converted to urea by liver. Kidneys have small effect in this conversion and urea is removed lastly from gills (Stoskopf, 1993). The present study demonstrated that the urea and uric acid amounts in *Cyprinus carpio* fed by olive leaf extract didn't change and that the creatine amounts decreased.

CONCLUSION

In conclusion, it was found that olive leaf extract has no negative effect on the metabolism of *Cyprinus carpio*. The positive effect of adding olive leaf extract to feed in the ratios of 0.25% and 0.50% on immunity shows that it can be used as a feed additive.

ACKNOWLEDGEMENTS

We would like to thank the Tarbiyomer and Kılıc Company for providing research facilities.

REFERENCES

- [1] AOAC, 2000. Official methods of analysis. Association of Official Analytical Chemists. (17th ed.) Washington, USA.
- [2] Baba, E., Acar, Ü., Yılmaz, S., Zemheri, F., and Ergün, S., 2018. Dietary olive leaf (*Olea europea* L.) extract alters some immune gene expression levels and disease resistance to *Yersinia ruckeri* infection in rainbow trout *Oncorhynchus mykiss*. Fish Shellfish Immunology, 79,28–33.

- [3] Bain, B.J., Lewis, S.M., and Bates, I., 2006. Dacie and Lewis Practical Haematology, Philadelphia, Elsevier Press., 600 pp.
- [4] Bircan, C., 2006. Determination of aflatoxin contamination in olives by immunoaffinity column using high-performance liquid-chromatography. *Journal of Food Quality*, 29(2), 126–138.
- [5] Blaxhall, P.C. and Daisley, K.W., 1973. Routine Hematological Methods for use with Fish Blood. *Journal of Fish Biology*, 5(6),771–781.
- [6] Campbell, T.W., 2004. Clinical Chemistry of Fish and Amphibians. In: Thrall, M.A., Baker, D.C., Campbell, T.W., DeNicola, D., Fettman, M.J., Lassen, E.D., Rebar, A., Weiser, G., Eds. *Veterinary Hematology and Clinical Chemistry*. Lippincott Williams & Wilkins, Pennsylvania. 499–517.
- [7] Gikas, E., Bazoti, F.N., and Tsarbopoulos, A., 2007. Conformation of Oleuropein, the major bioactive compound of *Olea europaea*. *Journal of Molecular Structure: Theochem*, 821(1-3), 125-132.
- [8] Harikrishnan, R., Nisha, R.M., and Balasundaram, C., 2003. Hematological and Biochemical Parameters in Common Carp, *Cyprinus carpio*, Following Herbal Treatment for *Aeromonas hydrophila* Infection. *Aquaculture*, 221 (1-4): 41–50.
- [9] Immanuel, G., Uma, R.P., Iyapparaj, P., Citarasu, T., Punitha, P.S.M., Michael, B.M., and Palavesam, A., 2009. Dietary Medicinal Plant Extracts Improve Growth, Immune Activity and Survival of Tilapia *Oreochromis mossambicus*. *Journal of Fish Biology*, 74(7), 1462–1475.
- [10] Klemm, D.J., Stober, Q.J., and Lazorchak, J.M., 1993. *Fish Field and Laboratory Methods for Evaluating the Biological Integrity of Surface Waters*. U.S. Environmental Protection Agency (EPA/600/R-92/111), Cincinnati, OH.
- [11] Mylonas, C.C., Cardinaletti, G., Sigelaki, I., and Polzonetti-Magni, A., 2005. Comparative Efficacy of Clove Oil and 2-phenoxyethanol as Anesthetics in the Aquaculture of European Sea Bass (*Dicentrarchus labrax*) and Gilthead Sea Bream (*Sparus aurata*) at Different Temperatures. *Aquaculture*, 246 (1–4), 467– 481.
- [12] Owen, R.W., Giacosa, A., Hull, W.E., Haubner, R., Würtele, G., Spiegelhalder, B., and Bartsch, H. (2000). Olive oil consumption and health: the possible role of antioxidants. *Lancet Oncology*, 1(2),107–112.
- [13] Sahu, S., Das, B.K., Mishra, B.K., Pradhan, J., and Sarangi, N., 2007a. Effect of *Allium sativum* on the immunity and survival of *Labeo rohita* infected with *Aeromonas hydrophila*. *Journal of Applied Ichthyology*, 23(1), 80–86.
- [14] Sahu, S., Das, B.K., Pradhan, J., Mohapatra, B.C., Mishra, B.K., and Sarangi, N., 2007b. Effect of Magnifera indica Kernel as a Feed Additive on Immunity and Resistance to *Aeromonas hydrophila* in *Labeo rohita* Fingerlings. *Fish & Shellfish Immunology*, 23 (1), 109-118.
- [15] Shalaby, A.M., Khattab, Y.A., and Rahman, A.M., 2006. Effects of Garlic (*Allium sativum*) and Chloramphenicol on Growth Performance, Physiological Parameters and Survival of Nile tilapia (*Oreochromis niloticus*). *Journal of Venomous Animals and Toxins Including Tropical Diseases*, 12(2), 172-201.
- [16] Stoskopf, M. 1993. *Fish Medicine* (1st Ed.). Philadelphia, Saunders Company, 882 pp.
- [17] Val, A.L., De Menezes, G.C. and Wood, C.M., 1998. Red Blood Cell Adrenergic Responses in Amazonian Teleost. *Journal of Fish Biology*, 52(1), 83-93.
- [18] Visioli, F., Bellosta, S., and Galli, C., 1998. Oleuropein, the bitter principles of olives, enhances nitric oxide production by mouse macrophages. *Life Sciences*, 62(6), 541–6
- [19] Yilmaz, E., Çoban, D., and Bayar, F., 2017a. The effects of the fig (*Ficus carica*) extract and onion (*Allium capa*) extract addition to *Cyprinus carpio* feeds in different amount on the growth performance. *Ege Journal of Fisheries and Aquatic Sciences*, 34(3), 321-326
- [20] Yilmaz, E., Çoban, D., Kırım, B., Güler, M., and Öztürk, S., 2017b. The Effects of the Carvacrol and Thymol Addition to *Pangasius Hypophthalmus* Feeds in Different Amount on the Growth Performance. *Aquaculture Studies*, 17(2), 157-162.
- [21] Yin, G., Ardo, L., Thompson, K.D., Adams, A., Jeney, Z., and Jeney, G., 2009. Chinese Herbs (*Astragalus radix* and *Ganoderma lucidum*) Enhance Immune Response of Carp, *Cyprinus carpio* and Protection against *Aeromonas hydrophila*. *Fish & Shellfish Immunology*, 26(1), 140-145.

EVALUATION OF THE RELIABILITY OF PLANTS USAGE IN FOODS: A REVIEW

Ezgi Şenol¹, Beraat Özçelik*^{1,2}

^{1*} *Department of Food Engineering, Faculty of Chemical and Metallurgical Engineering, Istanbul Technical University, Istanbul, Turkey, ozcelik@itu.edu.tr; ezgisenol23@hotmail.com*

² *BIOACTIVE Research and Innovation Food Manufacturing and Chemical and Metallurgical Engineering, Istanbul, Turkey*

Abstract

Functional foods is a concept that have been put forward in recent years. A food given an additional function (often one related to health-promotion or disease prevention) by adding new ingredients or more of existing ingredients is known as functional. Functional foods are products that provide significant benefits on the human physiology and metabolic functions beyond the body's need for essential nutrients, thus enabling to protect from diseases and achieve a healthier life. Although the concept of functional foods is new, its application is very old. As the side effects of the drugs have been learned and health problems have increased, the orientation to functional foods have increased and the products are diversified. Functional foods have been even more than normal food (Ozcelik, 2016). The functional food markets, whose trade volume has exceeded \$ 100 billion around the worldwide, has increased in terms of quantity and diversity in developed countries such as the USA, the European Union and Japan. In Turkey, the functional products market for 2015, it is stated that the ratio reached to 600 million Turkish Liras (Anonym,2014). The usage of plants for food industry to obtain functional food products has become a notable issue in recent years. A rising interest for medicinal and aromatic plants and the active substances obtained from them has led to studies to determine the effects of these plants. The purpose of his review is todetermine and evaluate the reliability of plants in terms of food and drugs effets.

Key Words: functional food, plant, reliability

1.INTRODUCTION

The function of functional, medically useful, regulating, special-purpose nutrients, which are not nutrients, but which have benefits and functions like nutrients, are called "phytochemicals" and the number of these chemicals is expressed in thousands. Basic functions of this type of foods are to collect the free radical-containing metabolites and inactivate them as well as to stimulate the immune system and to regulate antibacterial and antiviral effects in the cellular and intercellular environment activities. (Aksoy, 2007; Dunder, 2001). The purpose of this study is to look for the variety of plants or their ingredients for being used to produce functional food.

1.1. Evaluation In Terms of Food Industry: Approximately 1/3 of the 300 plant families grown in nature contain essential oil. The most known families with their volatile oils are: Pinaceae, Laureceae, Myrtaceae, Rutaceae, Laminaceae (Labiatae), Apiaceae (Umbelliferae), Zingiberaceae, Asteraceae (Compositae), Piperaceae, Irridaceae, Chenopodiaceae, Verbenaceae, Brassicaceae and Ranunculaceae. Some of these families have a special importance. For example, in the Labiatae family, Thymus, Lavandula, Melissa, Mentha species and some other plants found in many Mediterranean and European countries are valuable volatile oil sources (Ceylan, 1996). The amount and composition of volatile oils in Lamiaceae plants depends on the light (Johnson et al. 1999), the availability of nutrients from the plant (Skoula et al., 2000) and season (Kokkini et al., 1997).

The antioxidant activity of some aromatic plants is related to the amount of secondary components in plant structure and phenolic compounds in the structure (Skerget et al., 2005). The amount of these components varies from plant to plant due to individual factors (morphogenetic, ontogenetic, diurnal and ecological factors), genetic and genome differences (Ceylan, 1996). Among these compounds, the most common are flavonoids, phenolic acids and phenolic terpenes (Javanmardi et al., 2003). Antioxidant effect of phenolic compounds is to clean the free radicals (Rice-Evans et al., 1995;

Pekkarinen et al., 1999), to compound with the metal ions (metal chelating) and to prevent or reduce singlet oxygen formation (Rice-Evans et al. (1995).

Flavonoids and other phenolic compounds are mostly found in the leaves, flowers and woody parts of the plant (Kähkönen et al., 1999). For this reason, generally aromatic plants are used as a form of extracts of volatile oil obtained by methods such as extraction, drying. (Baytop, 1999) or distillation (Botsoglou et al., 2003a). There are many companies exporting products from plants in the Aegean Region, where has a great potential for aromatic plants. The primary criterias formed for many plants and plant products found in our food chain are important for researchers and firms. For example, the basic phytochemical groups found in fruits and vegetables are commonly used in daily life; Polyphenols, flavones, phenyloids, are the most powerful microchemicals in terms of antioxidant activity. Because of these properties, they protect cells by inhibiting LDL oxidation. At the same time, it has been determined that, primarily polyphenols, isoflavones and flavonoids have a change effect on metabolic profile of steroids and p450 substrates. (Fuhrman et al., 1997).

In vitro studies have shown that flavonoids have a wide spectrum of physiological, pharmacological activity ranges including the antiallergic, anti-inflammatory, antibacterial, antifungal, antiviral and anticancer. Carrot, citrus, strawberry, apple, raspberry, broccoli, ginkgo bloba, black and green tea, parsley, soybeans, cereals, cabbage, squash, potatoes, tomatoes, cucumbers, vegetables and fruits are rich sources of flavonoids (Van Hethof et al., 1997). There are many studies that test the efficacy of plant products or plant origin products against pathogens and saprophyte microorganisms; These studies have shown that microbial safety of foods can be protected in this way. For this purpose, the plants themselves (leaves, body, bud, flowers, fruits, seeds, onions and rhizomes) as well as the compounds obtained from plants (extracts, essential oils) are used.

1.1.1. Fruit and Vegetable Juices: It has been known that it has an inhibitory effect on microorganisms due to its low pH environment and / or its antimicrobial components that depend on the origins. Many fruit and vegetable juices, especially lemon juice, which is frequently used as acidifying and flavoring in foods, have antimicrobial activity on various microorganisms. Lemon juice obtained from lime; *Serratia spp.*, *Salmonella paratyphi*, *Shigella flexneri*, *Pseudomonas aeruginosa*, *Klebsiella pneumoniae*, *Citrobacter spp.*, *Escherichia coli*, *Staphylococcus aureus*, *Enterococcus faecalis*, *Clostridium spp.*, *Bacteroides spp.*, and *Porphyromonas spp.* It has been reported that it has inhibitory activity in varying proportions (Aibunu et al, 2007).

1.1.2. Spices: It is also known that it is used to extend the shelf life of foods by showing antimicrobial activity on pathogenic and saprophyte microorganisms as well as flavor and aroma enhancing properties. (Hsieh, 2001). According to the researchs, it was observed that he cinnamon (960,1, 0.2 and 0.3) added to the pasteurized apple juice decreased the *L. monocytogenes* by 4 - 6 log cfu / ml, and the number of *E. coli* O157: H7 was decreased by around 1.6-2 log cfu / ml; It was reported that the vanilla (260.10, 0.15 and 0.20) decreased the number of *L. monocytogenes* in orange juice by 2-3 log cfu / ml and the mint added to tomato juices decreased the microbial flora by 4.77 - 8.34 log cfu / ml.

Clove and garlic were tried against to *Bacillus sphaericus*, *Staphylococcus aureus*, *Si. epidermidis*, *Enterobacter aerogenes*, *E. coli*, *P. aeruginosa*, *Salmonella Typhi* and *S. flexneri* microorganisms and it was found that garlic nd cleve had he inhibitory effect on these microorganisms. When it comes to mint, 11 different gram negative bacteria strains were tried. (*E.coli*, *K. pneumoniae*, *P. aeruginosa*, *S Typhi*, *S. Paratyphi A*, *S. Paratyphi B*, *Proteus mirabilis*, *Proteus vulgaris*, *E. aerogenes*, *Shigella dysenteriae*, *Y. enterocolitica*). It was found to have inhibitory effect against these microorganisms.

Mustard flour used in mustard preparation was tested against pathogens of *S. typhimurium*, *E. coli* O157: H7 and *L. monocytogenes* at two different storage temperatures at a concentration of 0.10; At 5 * C on 1st, 3rd and 5th days, at 22 ° C, 12th and 24th hours, respectively, it was seen that it reduced the number or microorganisms below the detectable level (Oncul and Karabiyikli 2014).

1.1.3. Essential Oils: The antimicrobial activities do not depend on a single mechanism, but are based on the presence of many target points on the cell and different groups of chemical components. The

mechanisms of essential oils on the bacterial cell is degradation of the cell wall, separation of the cytoplasmic membrane and membrane proteins, leakage of cell components, coagulation of the cytoplasm, and loss of proton repellent force. The environmental and genetic factors such as species and / or sub-species of the plant, part of the plant used, isolation methods, harvesting time, growing conditions cause modifications in chemical composition (Oncul and Karabiyikli 2014).

1.1.4. Plant extracts Most of the plant extracts, which are evaluated as GRAS, are thought to have a wide range of application in the food industry as a flavoring agent and preservative, and many in vitro studies have been carried out on some extracts with high antimicrobial activity. Several studies have shown that extracts can offer alternative solutions to conventional food additives, preservatives and disinfectants. It was observed that the amount of 0,1%- 0,2% roseberry and its extracts had the inhibitory effect on the some food borne microorganisms (*Listeria innocua*, *L. monocytogenes*, *Lactobacillus sake*, *Leuconostoc mesenteroides*, *Leu. dexiranicum*, *Leu. carnosum*, *Lb. curvatus* *Brochothrix thermosphacla*, *Br. thermosphacta*, *Lactococcus lactis*). The effect of cranberry and cinnamon mixture (1: 1) was test on *L. Monocytogenes* and it has been reported that the mixture had inhibition effect on the microorganism. The antimicrobial activity of the sumac extracts were tested on different microorganisms (*B. cereus*, *B. subtilis*, *St. aureus*, *H. pylori*, *L. monocytogenes*, *E. coli*, *E. coli II*, *Salmonella spp.*) was tested and it was stated that *B. cereus* and *H. pylori* were highly sensitive among the tested microorganisms and inhibition value of the extracts containing sumac was higher than the others. The antimicrobial activity of the plant extracts prepared with ethanol was tested against *S. typhimurium*, *E. coli* 0157: H7, *L. monocytogenes* at various ratios of 6-7 log, and it was stated that clove was the most effective plant for inhibition among the others. At the next stage of the study, the lettuce samples were contaminated with pathogen cultures at different inoculation levels (6.80 log cfu / g, 6.70 log cfu / g and 5.31 log cfu / g) and then treated with 1, 3, 5 and 10 minutes with clove extracts. . It has been reported by researchers that clove extract provides 3 log inhibition within 3 minutes. The contents of these extracts obtained from different foods depend on the origin; It has been reported that antimicrobial activity is generally caused by compounds that have the polyphenol structure (Oncul and Karabiyikli 2014).

1.2. Evaluation In Terms of Pharmaceutical Industry: Nowadays, there is a tendency for the natural products, this idea is increasing in order to avoid the health problems caused by advancing technological processes. Pharmaceutical companies based on the use of traditional plants, have identified and synthesized the active ingredients of some of them and have been able to predict their effects in the body with the dose standardization.

Treatment with plants is known as complementary-salternative medicine (CAM). In spite of this widespread usage, the definition of side effects of phytotherapy have not been yet available in the clinical side of the modern medicine and it is estimated that few of the unwanted situations occurred due to plant based treatment are reported. There are also no obligation regarding explaining the side effects of these products (Gozum et al, 2007). There very few data about the interaction of plants used for treatment purposes with other drugs. For example, the interaction of some plants with anticoagulants has begun. Plants can change the effects of the drugs in the body and the bioavailability rates. Their interaction with nutrients is an issue that has never been evaluated (Bush et al, 2007).

1.3. The Side Effects of Medicinal Plants: There is an evidence that some endemic plants have toxic effects on the liver. For example, they may increase the liver enzyme levels. *Sauropus androgynus* and *Aristolochia* species are reported to cause hepatitis, bronchiolitis obliterans, renal failure and even death. The hepatotoxic effect of a plant taken with a drug may be higher than expected. *Echinacea purpurea* (echinacea), which can be used for colds, flu, and nausea, can cause vomiting and cogulation of blood cells. *Ephedra* (ephedra, ma huang) species, which are called as Herbal ectasi, are found in slimming teas, blends and energy drinks and they are used for alternative purposes against drugs such as fenfluramine, dexfenfluramine (TC Health Ministry, Turkish Hygiene and Experimental Journal of Biology, 2009).

Also, some plants cause unwanted side effects when they are consumed with drugs. For example, herbs used for depression (*Hypericum perforatum*, St. John's wort, yellow centaury) are sold as edible herbal

Prozac and may inhibit the enzyme monoamino oxidase (MAO) as well as increase serotonin, dopamine and norepinephrine levels. Therefore, they should not be used with prescribed antidepressants. In the literature, it is reported that this herb can cause adverse events such as gastrointestinal disorders, weakness, confusion, dizziness, dry mouth, hair loss, manic disorder, hyperactivity, irritation, allergy and sensitivity to light (Schey et al, 2000). Table 1 shows the information about some plants and part of plants known as positive (P) or negative (N) and used for medical purposes.

Table 1. Some plants and part of plants known as positive (P) or negative (N) and used for lots of purposes.

NO	LATIN NAME	USED PART	POZITIVE/ NEGATIVE	FUNCTION	USAGE AREA	SOURCE
1	<i>Abies sp.</i>	Whole plant	P	Containing volatile oil	Aroma Industry	Park J,2005
2	<i>Acer saccharum</i>	Plant Juice	P	High content of glucose	Glucose Syrup Production	Kork K et al,2002
3	<i>Achillea millefolium</i>	Flower	P	Effective on circulation system	Pharmaceutical Industry	Candan F et al,2003
4	<i>Acorus calamus</i>	Risome	N	Natural painkiller	Pharmaceutical Industry	Balakumbahan R et all 2010
5	<i>Acorus tatarinowii</i>	Risome	N	Effective on nervous system	Medicine	Liao W ve Zhang X,2005
6	<i>Actinidia chinensis</i>	Fruit	P	High content of Vitamin C	Food Extract Production	Huang S. et al,2013
7	<i>Adonsonia digitata</i>	Fruit	P	Antioxidant	Bakery Product Production	Kabbashi A.et al ,2014
8	<i>Aesculus hippocastanum</i>	Plant body shell	P	Microvaxulokinetic Activity	Pharmaceutical Industry	Gitelson A and Merzlyak M,1994
9	<i>Aframomum meleguetta</i>	Seed	N	Antimicrobial Activity	Pharmaceutical Industry (extracts are solved in the alcohol)	Okigbo RN and Ogbonnaya UO,2006
10	<i>Alchemilla alpina</i>	Plant Body	P	High content of nitrogen phosphorus	Transgenic Plant Research	Micheal D et al,1996
11	<i>Alchemilla vulgaris / A. arvensis</i>	Plant Body	P	High content of bioactive biochemicals	Herbal Medicine	Altameme H et all,2015
12	<i>Alkanna tinctoria</i>	Root	N	High content of alkaline esters	There is no enough research	Assimopoulou A et al,2004
13	<i>Allium cepa</i>	Seed	P	Antimicrobial	Food Industry as an addtive	Benkeblia B,2004
14	<i>Allium ursinum L.</i>	Leaf	P	High content of volatile compounds	Inhibitory agent	Djurdjevic L,2004
15	<i>Alpinia galanga</i>	Risome	P	Bio adhesive effect	Medicine	Onmetta-aree, J et al,2006

4. Conclusion and Suggestions: Nowadays, the production of functional foods containing the bioactive components found in the plants become more popular. Especially directly plant tea consumption is one of the most preferred way. A great deal of plants has beneficial extracts for health and using these extracts with a certain limits provide more healthy food products. However, the use of these products together with the drugs directly and inadequate information about these plants can lead to a chain of events that cause death. Unfortunately, the potential damages are not discussed in scientific environments. As a solution, academicians interested in this field should inform the people with their scientific based researches. Further studies must be done about plant mostly used for healing purposes as well.

References

- [1] Özçelik M., M. Bitkisel Kaynaklı Bazı Fonksiyonel Gıdalar. *Biyoloji Bilimleri Araştırma Dergisi*, (1), 57-68.
- [2] Anonym, 2014. <http://www.gidateknolojisi.com.tr/haber/2015/04/diyabetik-ve-fonksiyonel-gida-pazari-600-milyon-lira-buyuklugue-ulasti> Date: 12.10.2015.
- [3] Aksoy, M. (2007). Ansiklopedik beslenme, diyet ve gıda sözlüğü. *Hatipoğlu Yayınevi*, 323.
- [4] Dündar, Y. (2001). Fitokimyasallar ve sağlıklı yaşam.
- [5] Ceylan, A. (1996). Tıbbi Bitkiler II (Uçucu Yağ Bitkileri) Ege Üniv. *Zir. Fak. Yay.*, (481).
- [6] Johnson, C. B., Kirby, J., Naxakis, G., & Pearson, S. (1999). Substantial UV-B-mediated induction of essential oils in sweet basil (*Ocimum basilicum* L.). *Phytochemistry*, 51(4), 507-510.
- [7] Skoula, M., Abbes, J. E., & Johnson, C. B. (2000). Genetic variation of volatiles and rosmarinic acid in populations of *Salvia fruticosa* mill growing in Crete. *Biochemical systematics and ecology*, 28(6), 551-561.
- [8] Kokkini, S., Karousou, R., Dardioti, A., Krigas, N., & Lanaras, T. (1997). Autumn essential oils of Greek oregano. *Phytochemistry*, 44(5), 883-886.
- [9] Škerget, M., Kotnik, P., Hadolin, M., Hraš, A. R., Simonič, M., & Knez, Ž. (2005). Phenols, proanthocyanidins, flavones and flavonols in some plant materials and their antioxidant activities. *Food chemistry*, 89(2), 191-198.
- [10] Javanmardi, J., Stushnoff, C., Locke, E., & Vivanco, J. M. (2003). Antioxidant activity and total phenolic content of Iranian *Ocimum* accessions. *Food chemistry*, 83(4), 547-550.
- [11] Rice-evans, C. A., Miller, N. J., Bolwell, P. G., Bramley, P. M., & Pridham, J. B. (1995). The relative antioxidant activities of plant-derived polyphenolic flavonoids. *Free radical research*, 22(4), 375-383.
- [12] Pekkarinen, S. S., Heinonen, I. M., & Hopia, A. I. (1999). Flavonoids quercetin, myricetin, kaempferol and (+)-catechin as antioxidants in methyl linoleate. *Journal of the Science of Food and Agriculture*, 79(4), 499-506.
- [13] Kähkönen, M. P., Hopia, A. I., Vuorela, H. J., Rauha, J. P., Pihlaja, K., Kujala, T. S., & Heinonen, M. (1999). Antioxidant activity of plant extracts containing phenolic compounds. *Journal of agricultural and food chemistry*, 47(10), 3954-3962.
- [14] Baytop, T. (1999). *Türkiye'de bitkiler ile tedavi: geçmişte ve bugün*. Nobel Tıp Kitabevleri.
- [15] Botsoglou, N. A., Fletouris, D. J., Florou-Paneri, P., Christaki, E., & Spais, A. B. (2003). Inhibition of lipid oxidation in long-term stored chicken meat by dietary oregano essential oil and α -tocopherol acetate supplementation. *Food research international*, 36(3), 207-213.
- [16] Fuhrman, B., Buch, S., Vaya, J., Belinky, P. A., Coleman, R., Hayek, T., & Aviram, M. (1997). Licorice extract and its major polyphenol glabridin protect low-density lipoprotein against lipid peroxidation: in vitro and ex vivo studies in humans and in atherosclerotic apolipoprotein E-deficient mice. *The American journal of clinical nutrition*, 66(2), 267-275.
- [17] van het Hof, K. H., De Boer, H. S., Wiseman, S. A., Lien, N., Westrate, J. A., & Tijburg, L. B. (1997). Consumption of green or black tea does not increase resistance of low-density lipoprotein to oxidation in humans. *The American journal of clinical nutrition*, 66(5), 1125-1132.
- [18] Aibinu, I., Adenipekun, T., Adelowotan, T., Ogunsanya, T., & Odugbemi, T. (2007). Evaluation of the antimicrobial properties of different parts of *Citrus aurantifolia* (lime fruit) as used locally. *African Journal of Traditional, Complementary, and Alternative Medicines*, 4(2), 185.
- [19] Öncül, N., & Karabiyikli, Ş. (2015). Bitkisel Kaynaklı Antimikrobiyallerin Gıdalarda Bulunan Mikroorganizmalar Üzerindeki İnhibitif Etkinliği. *Academic Food Journal/Akademik GIDA*, 13(2).
- [20] Gozum, S., Arıkan, D., & Buyukavci, M. (2007). Complementary and alternative medicine use in pediatric oncology patients in eastern Turkey. *Cancer nursing*, 30(1), 38-44.
- [21] Bush, T. M., Rayburn, K. S., Holloway, S. W., Sanchez-Yamamoto, D. S., Allen, B. L., Lam, T., ... & Roth, L. W. (2007). Adverse interactions between herbal and dietary substances and prescription medications: a clinical survey. *Alternative therapies in health and medicine*, 13(2), 30-35.
- [22] TC Health Ministry, Turkish Hygiene and Experimental Journal of Biology, 2009.
- [23] Schey, K. L., Patat, S., Chignell, C. F., Datillo, M., Wang, R. H., & Roberts, J. E. (2000). Photooxidation of Lens α -Crystallin by Hypericin (Active Ingredient in St. John's Wort). *Photochemistry and photobiology*, 72(2), 200-203.
- [24] Park, J. H., Choi, G. J., Lee, H. B., Kim, K. M., Jung, H. S., Lee, S. W., ... & Kim, J. C. (2005). Griseofulvin from *Xylaria* sp. strain F0010, an endophytic fungus of *Abies holophylla* and its antifungal activity against plant pathogenic fungi. *J Microbiol Biotechnol*, 15(1), 112-117.
- [25] Saiya-Cork, K. R., Sinsabaugh, R. L., & Zak, D. R. (2002). The effects of long term nitrogen deposition on extracellular enzyme activity in an *Acer saccharum* forest soil. *Soil Biology and Biochemistry*, 34(9), 1309-1315.
- [26] Candan, F., Unlu, M., Tepe, B., Daferera, D., Polissiou, M., Sökmen, A., & Akpulat, H. A. (2003). Antioxidant and antimicrobial activity of the essential oil and methanol extracts of *Achillea millefolium* subsp. *millefolium* Afan. (Asteraceae). *Journal of ethnopharmacology*, 87(2-3), 215-220.
- [27] Balakumbahan, R., Rajamani, K., & Kumanan, K. (2010). *Acorus calamus*: An overview. *Journal of Medicinal Plants Research*, 4(25), 2740-2745.
- [28] Liao, W. P., Chen, L., Yi, Y. H., Sun, W. W., Gao, M. M., Su, T., & Yang, S. Q. (2005). Study of antiepileptic effect of extracts from *Acorus tatarinowii* Schott. *Epilepsia*, 46, 21-24.
- [29] Huang, S., Ding, J., Deng, D., Tang, W., Sun, H., Liu, D., ... & Yu, J. (2013). Draft genome of the kiwifruit *Actinidia chinensis*. *Nature communications*, 4, 2640.
- [30] Kabbashi, A. S., Koko, W. S., Mohammed, S. E. A., Musa, N., Osman, E. E., Dahab, M. M., ... & Mohammed, A. K. (2014). In vitro amoebicidal, antimicrobial and antioxidant activities of the plants *Adansonia digitata* and *Cucurbit maxima*. *Advancement in Medicinal Plant Research*, 2(3), 50-57.
- [31] Gitelson, A., & Merzlyak, M. N. (1994). Spectral reflectance changes associated with autumn senescence of *Aesculus hippocastanum* L. and *Acer platanoides* L. leaves. Spectral features and relation to chlorophyll estimation. *Journal of Plant Physiology*, 143(3), 286-292.
- [32] Okigbo, R. N., & Ogbonnaya, U. O. (2006). Antifungal effects of two tropical plant leaf extracts (*Ocimum gratissimum* and *Aframomum melegueta*) on postharvest yam (*Dioscorea* spp.) rot. *African Journal of Biotechnology*, 5(9).
- [33] Morecroft, M. D., & Woodward, F. I. (1996). Experiments on the causes of altitudinal differences in the leaf nutrient contents, size and $\delta^{13}C$ of *Alchemilla alpina*. *New Phytologist*, 134(3), 471-479.
- [34] Altameme, H. J., Hameed, I. H., & Abu-Serag, N. A. (2015). Analysis of bioactive phytochemical compounds of two medicinal plants, *Equisetum arvense* and *Alchemilla vulgaris* seed using gas chromatography-mass spectrometry and fourier-transform infrared spectroscopy. *Malays. Appl. Biol*, 44(4), 47-58.
- [35] Assimopoulou, A. N., Boskou, D., & Papageorgiou, V. P. (2004). Antioxidant activities of alkanin, shikonin and *Alkanna tinctoria* root extracts in oil substrates. *Food Chemistry*, 87(3), 433-438.
- [36] Benkeblia, N. (2004). Antimicrobial activity of essential oil extracts of various onions (*Allium cepa*) and garlic (*Allium sativum*). *LWT-food science and technology*, 37(2), 263-268.
- [37] Djurdjevic, L., Dinic, A., Pavlovic, P., Mitrovic, M., Karadzic, B., & Tesevic, V. (2004). Allelopathic potential of *Allium ursinum* L. *Biochemical systematics and ecology*, 32(6), 533-544.
- [38] Oonmetta-aree, J., Suzuki, T., Gasaluck, P., & Eumkeb, G. (2006). Antimicrobial properties and action of galangal (*Alpinia galanga* Linn.) on *Staphylococcus aureus*. *LWT-Food Science and Technology*, 39(10), 1214-1220.

STUDY OF THE VARIETAL BEHAVIOR OF 6 BASIL VARIETIES (*OCIMUM BASILICUM* L) GROWN FOR THE FIRST TIME IN ALGERIA UNDER ARID CLIMATE

Nabila Slougui^{1,2}, Naima Draoui³, Cherifa Khaddar³, Mahfoud Hadj Mahammed²

¹ Preparatory Class Department at National Polytechnic School of Constantine, Algeria

² Biogeochemistry Laboratory of the Desert Environments, Kasdi Merbah University Ouargla, Algeria

³ Agronomy Department, Kasdi Merbah University Ouargla, Algeria

Email: slougui.nabila@gmail.com

Abstract

The objective of the present research is to study the behaviour of six varieties of basil, *Ocimum basilicum* L., in the edaphic-climatic conditions of Ouargla region. **Methods:** To achieve the plotted objectives, we have taken a number of parameters of growth, bloom, and yield. **Results:** The ANOVA analysis of all parameters showed that our tests were significant to very highly significant with the F computed above the theoretical F of 1%. **Conclusion:** The climate of the Algerian Sahara is adapted to the cultivation of new varieties of basil.

Keywords: Varietal behaviour, *Ocimum basilicum* L, growth parameters, yield parameters, arid climate.

INTRODUCTION

The history of aromatic and medicinal plants is associated with the evolution of civilizations with the aim of emphasizing the growing economic importance of the cultivation of aromatic and medicinal plants in the industrial investigation. And as part of the valorisation of the natural resources of arid zones, we have chosen to study the varietal behaviour of 6 varieties of basil of European origin in a Saharan climate, Hassi Ben Abdellah-Ouargla south-east of Algeria. Few agronomic studies have been carried out on basil. On the site, the test of Carron et al, which is a test of culture of basil in Mountain, ~~this test~~ was carried out in Belgium on five varieties of basil: great green, lettuce, green, purple, and exotic in 1996 and 1997 [1]. The objective of our trial is to study the behavior of six varieties of basil *Ocimum basilicum* L. in the edaphic-climatic conditions of the Ouargla region. In order to achieve the plotted objectives, we have taken a number of parameters of growth, bloom, and yield. We evaluated the obtained results using the statistical study of ANOVA.

MATERIAL AND METHODS

1. Climatic Data of the Agricultural Companion: The maximum temperature during the cycle of the crop does not exceed 33.8°C, and the minimum temperature is in the order of 4.2 °C. Humidity is high in January (68.1%) and with a significant evaporation in May [2].

Study Material:

Soil: The soil is characterized by a sandy texture, that is a basic pH with a very low organic matter rate. Soil salinity is low, C. E varies from 0.43 to 0.41 DS/m.

Irrigation Water: Irrigation water is pumped from the Alwelle aquifer which is at a depth of 1450 m. According to the Riverside classification, irrigation water belongs to the C₄S₄ class. The use of this water is very delicate and it requires a very permeable soil, well drained, and washed out soil.

Characteristics of Organic Fertilization: Organic fertilization is characterized by an acid pH (6.11) and a high electrical conductivity of the order of 10.10 DS/M.

Experimental Protocol

Experimental Device: The adopted experimental device is in complete random blocks, consisting of 6 treatments and 4 repetitions. Each block contains 6 elementary plots of 2 m in length on 1.5 wide.

Plant Material: The material studied is composed of six (06) Varieties of basil *Ocimum basilicum* L. (c.f. table 1)

Table 1: Presentations of Plant Material

Variety	Common name	Scientific name
V1	Great Green Basil	<i>Ocimum basilicum</i> L. Genovese
V2	Purple Basil	<i>Ocimum basilicum</i> L. Purpurescens
V3	Dwarf Basil Compact	<i>Ocimum basilicum</i> L. Minimum
V4	Basil Marseille	<i>Ocimum basilicum</i> L. Marceillais
V5	Basil Cinnamon	<i>Ocimum basilicum</i> L. Cinnamon
V6	Lemon basil	<i>Ocimum basilicum</i> L. Citriodora

Conduct of the Cultivation: **1. Tillage:** The tillage is carried out on a ploughing of 25 to 30 cm depth using a hoe. **2. Fertilization:** The organic and mineral bottom fertilization is applied to standards of 9 kg of manure from cattle per elemental parcel and 3 qx/ha of 15-15-15 per ha; therefore, (45 u N, 45 u P₂O₅ and 45 u K₂O) with the addition of phosphorus in the form of TSP: 55U/ha. **3. Nursery Seedlings:** Seedlings are made in the “nursery area” with containers filled with peat at a weight of 1.20 g of seeds to monitor our culture well. The irrigation is done on a daily basis. **4. Pre-irrigation:** Pre-irrigation has a dual purpose, to wash the salts and moisten the soil with its holding capacity. **5. Transplanting:** The plants we have planted have been of good vigour and biometrics of a height of 10 cm. Planting density was 11 plants with a spacing between the 30-cm planes and between the 40-cm lines per elemental plot.

Statistical Analysis of the Results: The obtained results for the studied parameters of the six varieties of *Ocimum Basilicum* L. were subjected to an analysis of ANOVA. This analysis allows testing the similarity of variables in statistical terms. The calculated F study and its comparison to the theoretical F, in particular the treatments, makes the meaning of the test out.

RESULTS AND DISCUSSION

Duration of the Cycle: The date of flowering is an important parameter as it allows judging the maturity of the varieties. Two periods are considered, the sowing-flowering period and the planting-flowering period. The duration of the seedling-flowering cycle of the experimented plants varies between 74 days for Purple (V2), which has the longest duration and 100.5 days for the Compact Dwarf (V3) with the most limited duration (Figure 1). Similar results are obtained for the parameter (transplanting-flowering cycle time) (Figure 2).

Plant Height: Based on the reported results in Figure 3, plant height varies very highly significance as a result of variety. The Great Green (V1) gave the best height (40.92 cm) against Marseille (V4) which is the variety that gave the lowest height (18 cm). The works of Philip and Staci [3] show that the height of the great green can reach 60 to 90 cm. According to another study, the height of basil varies from 20 to 50 cm, which reinforces the obtained results in the experimental range [4].

Mortality Rates : In general, plants with high mortality rates are the least suitable for edaphic-climatic conditions or those with low vigour at the time of transplantation. The analysis of variance reveals a very highly significant difference in the mortality rate parameter between the studied varieties. The highest rate is presented by Marseille (V4) with 40.62% which proves to be the least adapted to the Saharan conditions. The other varieties gave values below 3.12% (large green) or even zero (purple, dwarf compact and cinnamon) (Figure 4).

Total Biomass (leaves and stems): Total plant biomass is influenced by a number of factors including climatic conditions, fertilization and variety. The results show a very highly significant variation in the total biomass of the different tested varieties, varying between 82.84 g and 19.70 G. The best weight is

obtained with the Dwarf Compact (V3) 82.84 G; however, the lowest weight is recorded by the Purple (V2) 19.7 g (Figure 5).

Number of Branches per Plant: The number of branches per plant is variety function and water supply and fertilization. The analysis of variance reveals a very highly significant difference between treatments. The highest number of ramifications is presented by the Lemon variety (V6) with 13.24 branches per plant but the lowest value is recorded by the Dwarf Compact (V4) and cinnamon (V5) with 7.47 and 9.6 ramifications per consecutive plant (Figure 6).

Leaf Weight/Total Biomass Ratio: The variance analysis reveals a highly significant difference between the varieties. The results presented in Figure 7 show that Purple (V2) and Marseille (V4) provided the most important ratio (0.87) while the Lemon (V6) gave the lowest ratio (0.74).

Number of Leaves per Plant: The number of leaves is an important parameter because the more important the plant has a large number of leaves, the more important the vegetative weight is. The results shown in Figure 8 show that the Compact Dwarf (V3) recorded the most important number of leaves (2861.12) while Purple (V2) provided the lowest number (176.37). The analysis of variance reveals a highly significant difference in this variable.

Leaf Area: The analysis of variance reveals a very highly significant difference in the tested varieties. Leaf area of leaves varied between a minimum of 9 cm² obtained with the Compact Dwarf and a maximum of 17.62 cm² obtained for the large green (V1) (Figure 9). According to Carron et al [1], the Great Green leaves are 3 to 4 cm in length.

Branch Width: The results shown in Figure 10 show that the width of the variety ramifications varies between 36.55 and 19.02 cm. The Great Green (V1) recorded a significant value; however, the basil Marseille (V4) provided the lowest width. The analysis of variance reveals a very highly significant difference between the studied varieties.

CONCLUSION

The statistical study of the varietal behaviour of six varieties of *Ocimum basilicum* L., 4 of them grown for the first time in Algeria under the edaphic-climatic conditions of the Ouargla region revealed the success of our trial. The lack of similar agronomic work has not allowed us to develop a comparative study and in particular to draw distinctive conclusions. This test was performed by studying some parameters of yield, morphological, and early maturity. The analysis of all the studied parameters allowed us to conclude that the climate of the Sahara is adapted to the cultivation of new varieties of basil. These results remain preliminary but encouraging for the importance that this culture can present on the economic level given the diversity of the use of basil in the fields of aromatherapy, the food and cosmetics industry, and in culinary use as a condiment plant.

REFERENCES

- [1] Carron, C. A., Rey, C. H. and Brutin, (2017). Essai de variétés de basilic en montagne. *Revue suisse vitic. Arboric. Hortic*, 38(1), 51-55.
- [2] I. T. D. A. S., Ouargla, 2009.
- [3] Philippe et Delwaulle, *Revue suisse vitic. Arboric. Hortic*, 17, 1-8, 2006.
- [4] P. Schauenberg, guide des plantes médicinales, Paris, 287, 2005.




Figure 1: Effect of the variety on the duration of the sowing-flowering cycle (days)




Figure 2: Effect of the variety on the duration of the transplant-flowering cycle (days)




Figure 3: Variety effect on plant height (cm)




Figure 4: Effect of variety on mortality rate (%)




Figure 5: Effect of variety on total biomass




Figure 6: Effect of variety on the number of branches




Figure 7: Effect of variety on total leaf/biomass ratio




Figure 8: Effect of the variety on the number of leaves per plant




Figure 9: Effect of variety on leaf surface




Figure 10: Effect of variety on branch width

ETHNOMEDICINAL STUDY OF *CORNUS MAS L.* FROM THE FOLK AND TRADITIONAL MEDICINE OF REPUBLIC OF MACEDONIA

Biljana Bauer

Department of Pharmacognosy, Faculty of Pharmacy, University of Ss Cyril and Methodius, Majka Tereza No 47, 1000 Skopje Republic of Macedonia
biba@ff.ukim.edu.mk

Abstract

This study was carried out concerning ethnomedicine on preparations obtained from *Cornus mas L.* from the folk and traditional medicine of Republic of Macedonia, an area so far less frequently studied from the perspective of folk traditional medicinal, local food and handicraft and other uses. European cornel acts as astringent and antidiarrhoeic. Tea, juice, fresh and dry fruits because of its antimicrobial, anti-inflammatory action are used at diarrhoea, colitis and other intestine disorders. Dry fruits are used directly as a medicine against diarrhoea. It acts chemo static and protective on capillary. Cornels as mountainous fruits were used by Macedonians as a food long time ago. Fruits are used dry as a food. Dry fruits are used for preparing compote or directly in nutrition in winter and spring period. From the fruit jam, marmalade, compote and juice are prepared at home and in the food industry. Dry fruits are kept in rakiya and sugar for obtaining liqueur. Alcohol beverage *rakiya* - dryankoitsa was prepared from cornels. Manufacture of hammer in Ohrid or stick in Mariovo (koledashki) for Christmas Eve (koleda) is made from a cornel branch. Handmade handle for Macedonian traditional instrument *gajdarka*, in Kumanovo is manufactured from cornel because it is smooth and not cracked. Cornel is a symbol of durability, solidity, and with its hardness influences the health. The district, from the ethnobotanical point of view, shows traces of the influences of the neighboring regions. Some medicinal uses are linked to beliefs or residual forms of magic prescriptions.

Key words: European cornel, medicinal, food, handicraft, use

INTRODUCTION

During the past decade, the Balkans have been the focus of a number of ethnobotanical studies that have revealed the complex relationships of humans and plants at the location of a geographical cross roads where the complex features of their history and the structure of their ethnical and religion meet. Most of the previous studies were mainly aimed at medicinal and edible plants and traditional handicrafts in the countries on the Balkans [1-8] and shed new light on existing ethnological and anthropological data. Republic of Macedonia has old ethnomedicine traditions, consisting of many recipes with herbal, animal, and mineral original ingredients. The folk and traditional medicine of Republic of Macedonia pays special attention to disease prevention. This study was carried out concerning ethnomedicine on preparations obtained from *Cornus mas L.* from the folk and traditional medicine of Republic of Macedonia, an area so far less frequently studied from the perspective of plant folk traditions. In this study local medicinal uses, local food uses and local handicraft and other uses of European cornel are described.

LOCAL NAMES

Republic of Macedonia: дрен, дренка, дренушка, петров цвет, тајане, равнец, месечник [9] [10].

Yörüks: Karamluk [11].

English: Cornelian cherry, European cornel, dogwood [10].

BOTANY AND ECOLOGY

European cornel is a deciduous shrub or small tree, reaching 3-8 m tall or slightly larger, shape is rounded. *Bark* is dark gray to reddish brown with scaly, exfoliating patches. Each node has two oppositely positioned flower buds on one-year old shoot. Leaves are ovate or ovate-elliptic up to lanceolate and narrowly elliptic, with a sharp top and round base dark green (4-11 cm long and 3-5 cm

wide). Leaves are entire, and with distinctively parallel lateral *veins* (up to 6 pairs of side) arching strongly as they approach the margins. Inflorescences resembled the umbel structure. Each umbel is surrounded at the base by small, yellowish, petaloid bracts. Yellow flowers on short stalks bloom in early spring before the leaves emerge. The flowers have 4 rudimentary sepals, 4 petals, 4 stamens, and one pistil. Fruits are ellipsoid, fleshy, single-seeded, stone fruit type (drupes to 2-3 cm long) which mature to cherry red in mid-summer. Fruits (*Corni fructus*) are collected from September to October. European cornel grows most in the dry and stony places exposed to the sun, at the end of the deciduous forest, among the shrubbery throughout the entire territory of the Republic of Macedonia [9]. European cornel is a wide spread plant, beside the bushes, the roads, in the oak and beech forests and other habitats up to 1200 m above the sea level [12]. It is not indicate the exact degree of danger of the European cornel in Republic of Macedonia.

LOCAL MEDICINAL USES

Cornus mas act as astringent and antidiaric. Tea, juice, fresh and dry fruits because of its antimicrobial, anti-inflammatory action are used at diarrhea, colitis and other intestine disorders. Dry fruits are used directly as a medicine against diarrhea [13]. It acts chemo static and protective on capillary [9].

Yörüks in Republic of Macedonia collected primarily for medicinal herbal teas (infusion) for internal cleansing [11].

LOCAL FOOD USES

Cornels as mountainous fruits were used by Macedonians as a food long time ago [14]. Fruits are also used dry as a food. Dry fruits are used for preparing compote or directly in nutrition in winter and spring period [13]. From the fruit jam, marmalade, compote and juice are prepared at home and in the food industry. Dry fruits are kept in brandy (*rakiya*) and sugar for obtaining liqueur [9]. Alcohol beverage *rakiya* - *dryankoitsa* (дренкојца) is prepared in region of Struga [14].

LOCAL HANDICRAFT AND OTHER USES

Manufacture of hammer in Ohrid or stick in Mariovo (коледашки) for Christmas Eve (коледе) was reported [15]. In Kumanovo from *Cornus mas* handmade handle for Macedonian traditional instrument *gajdarka*, because it is smooth and not cracked is manufactured [16].

Cornus mas is a symbol of durability, solidity, and with its hardness influences the health. The cornel tree is considered as the strongest and most healthy tree and as such is a symbol of health. Sometimes it is considered as the longest living tree, and it is an imaginable symbol of the long life. The cornel tree blooms the earliest, and because of that is regarded as a symbol of joy, youth and strength. *Cornus mas* is used during wedding rituals in order to cause agreement in marriage and fertility among spouses. *Cornus mas* is used for the health of the cattle, especially for the lambs. Early in the morning they hit the lambs, and with cornel stick cows are forced to the bulls. It is believed that such a cow will host a male calf [15].

In some areas in Republic of Macedonia people carry with them cornels stick, because they believed that the vampires run away from it. They are sheltered from witches and other unclean strengths. Gates and windows are decorated with cornel twigs, which sheltered from thunders [15]. In Porechje people collect *Cornus mas* on St. George's day (Gjurgovden) on 6th of May, because of its durability, to be given to them [17]. When there are many cornels (cornel fruits) there will be a great winter (long winter with lots of snow). Cornel tree is small because part of it was used for the cross on which Jesus Christ was crucified, or from other traditional forecasting the bear walked out of its winter bearing earlier, after noticing that the cornels blossomed. Bear was irritated and broken the cornel tree and from than it doesn't grow high [15].

Ritual acts with cornel twigs are carried out with the assurance that they will bring health to the participants. Such practices are connected with some holidays, mainly with Gjurgjovden (Ѓурѓовден) Saint George's Day, when people are decorated with cornel twigs, or they are sealed with the twigs and are swinging on *Cornus mas* to be health like it [18] [19]. Swinging on *Cornus mas* twigs is performed

on Easter in Kumanovo and in Shopsko-bregalnica ethnographic entity as a symbol of new beginning on Gjurgjovden, to be healthy as cornel [20]. Also, swinging on swings tied up on green tree i.e. cornel, walnut etc is done on Gjurgjovden in the same entity [21]. People are bathing in the warm water in which cornel twigs are placed [15].

Surova or Vasilica (on 14th of January) is a holiday that marks the beginning of the Old New Year according to the Julian calendar in Shopsko-bregalnica ethnographic entity. The stick named *surovica* [суровица] or *surovaska* [суроваска] made of oak branch, or of cornel or hazelnut, served as a ritual requisite through which the magical power of the tree was transferred with the hit. Everyone in the house is hit to be healthy as a cornel. On Četerepe (St. Forty), 40 мученици, Младенац (22nd of March) from 39 twigs from *Vitis vinifera* and one cornel branch fire burns, and with the smoke the house is fumigated for moths and other animals to get away [21]. Domestically produced and hand-made brooms consisting of *Cornus mas*, nettle and *Euphorbia* are made in March and they are used for sweeping early in the morning before the sun rises [19]. Fruits from cornels are used as a nafora (antidors prosfor) [15]. Flower buds in several areas in Republic of Macedonia are used on Easter. Holy Communion was practiced with cornel flower buds and wine [15]. In the village Gorna Pcinja because of its great distance from the church, sacrament named as komka “комка” is given with cornel flower buds, nettle, *Geranium* and wine in the house [20]. Ritual act is done with cornel flower buds, nettle, wine and bread for giving the sacrament to all in the family, on the great Thursday on Easter, for God to give health [19].

DISCUSSION AND CONCLUSION

The usage of preparations obtained from *Cornus mas* L. from the folk and traditional medicine of Republic of Macedonia, from the ethnobotanical point of view, shows traces of the influences of the previous nations who lived in this area [22]. Some medicinal uses are linked to beliefs or residual forms of magic prescriptions. The use of cornels remains well alive nowadays.

REFERENCES

- [1] Dogan, Y., and Nedelcheva, A., 2015. Wild plants from open markets on both sides of the Bulgarian-Turkish border. *Indian Journal of Traditional Knowledge*, 14(3), 351–358.
- [2] Ivancheva, S., Stancheva, B., 2000. Ethnobotanical Inventory of medicinal plants in Bulgaria. *J Ethnopharmacol*, 69(2), 165-172.
- [3] Nedelcheva, A.M., Dogan, Y., Guarrera, P.M. 2007. Plants traditionally used to make brooms in several European countries. *J Ethnobiol Ethnomed*, 3: 20.
- [4] Nedelcheva, A., Dogan, Y., Obratov-Petkovic, D., Padure, I.M., 2011. The traditional use of plants for handicrafts in Southeastern Europe. *Hum Ecol*, 39 (6), 813-828.
- [5] Nedelcheva, A., Dogan, Y., 2011. Usage of plants for weather and climate forecasting in Bulgarian folk traditions. *Indian J Tradit Know*, 10(1), 91-95.
- [6] Nedelcheva, A., 2013. An ethnobotanical study of wild edible plants in Bulgaria. *Eur Asian Journal of BioSciences*. 7: 77–94.
- [7] Redzic, S., 2006. Wild edible plants and their traditional use in the human nutrition in Bosnia-Herzegovina. *Ecol Food Nutr*. 45(3), 189-232.
- [8] Šoukand, R., Pieroni, A., Biró, M., Denes, A., Dogan, Y., Hajdari, A., Kalle, R., Reade, B., Mustafa, B., Nedelcheva, A., Quave, C.L., Luczaj, L., 2015. An ethnobotanical perspective on traditional fermented plant foods and beverages in Eastern Europe. *J Ethnopharmacol*. 170, 284–296.
- [9] Dervendzi, V., 1992. Contemporary medication with medicinal herbs. Tabernakul, Skopje, 203-205. (in Macedonian)
- [10] Dimitrovski, T., 1990. Bulletin of the Macedonian terminology development committee. Manu, Skopje, 61-85. (in Macedonian)
- [11] Nedelcheva, A., Pieroni, A., Dogan, Y., 2017. Folk food and medicinal botanical knowledge among the last remaining Yörüks of the Balkans. *Acta Soc Bot Pol*. 86(2), 3522.
- [12] Matevski, V., 2005. The Flora of the Republic of Macedonia. Vol.1, Book 6., MANU, Skopje, 1526-1527. (in Macedonian)
- [13] Mojsoski, P., 2008. Natural antibiotics. Iris-P, Struga, 1-80 (in Macedonian)
- [14] Konstantinov, M., 1992. Macedonians, Maring, Skopje, 1-615 (in Macedonian)
- [15] Vrazinovski, T., 2000. Dictionary of the folk mythology of the Macedonians, Matica makedonska, Skopje, 1-350 (in Macedonian)
- [16] Dzimrevski, B., 2000. Instrumental musical tradition in Kumanovo, in Folklore in Kumanovo and surroundings, Centar za kultura “Trajko Prokopiev”, Kumanovo, 289-313 (in Macedonian)
- [17] Obrembski, J., 2001. Macedonian sociological studies, Matica makedonska, Skopje, 1-281 (in Macedonian)
- [18] Matijashevic-Pokupec, V., 2000. Contemporary situation of the ritual calendar songs in the Kumanovo. In: Folklore in Kumanovo and surroundings. Centar za kultura “Trajko Prokopiev”, Kumanovo, 163-190 (in Macedonian)
- [19] Vrazinovski, T., 2002. Macedonian folk mythology of the Macedonians. Matica makedonska, Skopje, 1-272 (in Macedonian)
- [20] Petreska, V., 2000. Spring customs, rituals and beliefs in Kumanovo. In: Folklore in Kumanovo and surroundings. Centar za kultura “Trajko Prokopiev”, Kumanovo, 289-313 (in Macedonian)
- [21] Malinov, Z., 2006. Traditional folk calendar in Shopsko-bregalnica ethnographic entity. Special Editions Volume 68, Institute of folklore “Marko Cepenkov”, Skopje, 1-353 (in Macedonian)
- [22] Nikolovski, B. 1995. Contributions to the history of the health culture of Macedonia. Makedonsko farmacevtsko drustvo, Skopje, 1-399 (in Macedonian)

DEVELOPMENT OF MICROEMULSION CONTAINING *PLANTAGO MAJOR* EXTRACTS: FORMULATION AND EVALUATION OF TOPICAL ANTI-INFLAMMATORY ACTIVITIES

Asih Triastuti,^{1*} Oktavia Indrati,² Farida Hayati³

¹ Laboratory of Pharmaceutical Biology, Department of Pharmacy UII, 55584 Yogyakarta, Indonesia

² Laboratory of Pharmaceutical Technology, Department of Pharmacy UII, 55584 Yogyakarta, Indonesia

³ Laboratory of Pharmacology, Department of Pharmacy UII, 55584 Yogyakarta, Indonesia

Corresponding author: asih.triastuti@uii.ac.id

Abstract

Plantago major L. (Plantaginaceae) is widely used to treat various diseases. It is reported to have anti-microbial, anti-diabetic, anti-spasm, anti-inflammatory activity and wound healing properties. *P. major* contains biologically active compounds such as polysaccharides, flavonoids, iridoid glycosides, terpenoids, and alkaloids.¹ Despite several publications supporting health benefits of this plant, not much work has been done to develop an anti-inflammatory formulation containing extracts of *P. major*. Therefore, the aim of this work was to formulate extract of *P. major* into microemulsion to increase economical value of this plant and to support the society with affordable dosage form. **Material and Methods:** Microemulsions are clear, optically isotropic and thermodynamically stable systems generally composed of a blend of oil, water and surfactant(s).² Three formulas were developed with variation of surfactants tween 80: lecithin, Formula 1 = 10:1, Formula 2 = 11:1 and Formula 3 = 12:1 respectively, along with cosurfactant (propylene glycol) and extract of *P. major* (0.3%). Microemulsions were stored at 4°C, 25°C, and 40°C for 4 weeks, and were characterized regarding their organoleptic, homogeneity, pH, and viscosity. Particle size was determined on day 0 and week 4 using Horiba@SZ-100 nano particle analyzer. Formula 3 was investigated for its anti-inflammatory activity using croton-oil induced ear edema in mice.³ Percentage of inflammatory inhibition was calculated and confirmed by histological section of ears. **Results:** During 4 weeks of storage, all formula showed homogeneity and stability and met the particle size requirements for microemulsions. Topical application of microemulsions reduced ear edema and pro-inflammatory cells in the tissue and its activity was similar to hydrocortisone 1% ($p > 0.05$). **Conclusion / Discussion:** Our data indicated that *P. major* extracts can be developed into modern formulation with potent anti-inflammatory activity.

Keywords: *Plantago major*, microemulsion, topical anti-inflammation, croton oil-induced ear edema

INTRODUCTION

Ethnopharmacological studies show that *Plantago major* is used in the treatment of various diseases. It is reported to have anti-inflammatory,¹⁻³ anti-microbial⁴ anti-diabetic,⁵ anti-spasm activity,⁶ and wound healing properties.⁷ *P. major* contains biologically active compounds such as polysaccharides, lipids, caffeic acid derivatives, flavonoids, iridoid glycosides, terpenoids, alkaloids and some organic acids.^{6,8-11} *P. major* shows anti-inflammatory properties by reducing pro-inflammatory cytokine levels¹, inhibiting cyclooxygenase-2⁹, and inhibiting neutrophil ROS production.¹² Despite several publications supporting health benefits of this plant, not much work has been done to develop the formulation of this plant into anti-inflammatory dosage form.

One of the practical dosage forms is microemulsion, which was first introduced by Hoar and Schulman in 1943. Microemulsion is a dispersion system comprising aqueous phase, oil, surfactant, and co-surfactant. It forms a single liquid solution that is clear, thermodynamically stable and optically isotropic having a droplet diameter usually ranging from 10 to 100 nm. Several studies have proved the potential of microemulsion to rise cutaneous delivery of hydrophilic and lipophilic drugs. Microemulsion has such advantages as the ability to increase the solubilization, dissolution rate, and bioavailability of poorly soluble drugs, the good thermodynamic stability and the easiness to produce.¹³⁻¹⁶ Due to these advantages, it is promising to deliver the *P. major* extract into microemulsion.

MATERIALS AND METHODS

General: Unless explicitly stated otherwise, all chemicals used were of “pro analysis” quality. Ethanol, isopropyl myristate, Tween 80, Span 80, propylene glycol, methyl paraben, and propyl paraben were supplied from Sigma-Aldrich. Distilled water was used in the preparation of microemulsion. Ethanol, croton oil, ether, Phosphate Buffered Saline (PBS) and Hematoxyline-Eosin (HE), were obtained from Sigma-Aldrich (St. Louis, MO, USA). Hydrocortisone (1%) was obtained from Kimia Farma.

Ethics: All animals used in this study were obtained from the Laboratory of Pharmacology, Department of Pharmacy Universitas Islam Indonesia. Male Balb/c mice (32-36 g) were used and allowed one week to adapt to environment before experiments. Animals were maintained on a 12 h light/dark cycle in an air-conditioned room and given standard chow and water *ad libitum*. All animal experiments were approved by Medical and Health Research Ethics Committee (MHREC) Faculty of Medicine Universitas Gadjah Mada, Yogyakarta, Indonesia with reference number KE/FK/345/EC and exactly performed according to the “Principles of Laboratory Animal Care” (NIH Publication No. 86-23).

Preparation of plant extract: *P. major* dried herbs was purchased from Directorate Research and Development of Medicinal Plants and Traditional Medicine (B2P2TO2T Tawangmangu, Indonesia). *P. major* were macerated for 3 days using ethanol as the solvent. Extract was evaporated using rotary evaporator (Heidolph L4000) until most of its solvent evaporated.

Preparation of microemulsion: Three formulations were produced for the same percentage but different ratios of Tween 80 and Span 80, with Formulation 1 (8:1), Formulation 2 (9:1), and Formulation 3 (10:1). First, the aqueous phase, consisting of ethanol extract of *P. major*, Tween 80, methyl paraben, and distilled water, were mixed using magnetic stirrer at 50°C. Separately, the oil phase that consisted of isopropyl myristate, propyl paraben, and Span 80 was stirred at the same condition. After that, the oil phase was added to the aqueous phase using magnetic stirrer at 50°C to produce microemulsion.

Microemulsion characterization: All three formula were then analyzed for its characteristics including the homogeneity, viscosity, globule size, pH, and stability. Particle Size Analyzer SZ100 (Horiba, Japan) was used to measure the average size of microemulsion droplet by loading it into a clear zeta cell at 25°C. The pH values for microemulsion were determined at 25°C by pH meter (Horiba LaquaAct, Japan). The viscosity of optimized microemulsion was determined using Brookfield viscometer (Brookfield DV2T, USA) with spindle number 61 at 60 rpm. The physical stability of microemulsion was evaluated by visual inspection such as by phase separation, transparency, drug precipitation and color change. Three batches of microemulsion were stored at 4°C, 25°C and 40°C for 4 weeks and examined for physical stability. All measurements were carried out in triplicate.

Croton-oil induced ear edema test: Formula 3 was investigated for its anti-inflammatory activity using croton-oil induced ear edema in mice according to Patel et al., 2012. Adult male Balb-c mice (25-35 g) were randomly divided into 5 groups of 7 animals. Ear edema was induced on the anterior and the posterior surfaces of the right ear of the mice by topical application of 2 µl croton oil dissolved in 20 µl of acetone. Microemulsion based, microemulsion formula 3, and hydrocortisone (1 mg) were also dissolved in acetone and were applied with croton oil. The inflammations were measured on 4 hours after the application of croton oil. Animals were euthanized and two ear punches (6 mm diameter) were taken from each mouse. Percentage of inflammatory inhibition was then calculated and confirmed by histological section of ears.

Ear biopsies from croton oil-induced mouse ear edema: Ear biopsies were collected and fixed in 70% ethanol for 24 h and then preserved in 10% formalin. Subsequently, the ears were dehydrated, blocked in paraffin, and then sectioned with a microtome (4 µm). The cross-sections were stained with HE for the evaluation of leukocyte infiltration, vasodilatation, and edema intensity. A representative area was selected for qualitative light microscope analysis (40 and 400 x magnification).

RESULTS AND DISCUSSION

P. major microemulsion characteristic and stability: All the formulations with various ratios of surfactants formed transparent microemulsion. The microemulsion was brown and has specific odor due to its ethanolic extracts of *P. major*. The clarity of the microemulsion is caused mainly by the activity of surfactants and co-solvent in the formulation. The surfactants as well as the co-solvent (propylene glycol) lower the interfacial tension between the oil phase (isopropyl myristate) and the aqueous phase, so the oil droplets can disperse in the aqueous solution and yield a clear microemulsion system. The various characteristics of each formulation can be seen in table 1.

Table1. Evaluation of various characteristics of *P. major* microemulsion

Characteristics	Formulation1	Formulation 2	Formulation3
Organoleptic	Brownish color, specific odor, transparent	Brownish color, specific odor, transparent	Brownish color, specific odor, transparent
Homogeneity	Homogeneous	Homogeneous	Homogeneous
Viscosity (cP)	85.2 ± 25.3	115.2 ± 12.4	122.8 ± 6.0
pH value	6.85 ± 0.01	6.86 ± 0.02	6.88 ± 0.03
Droplet size	93.3 ± 3.06 nm	75.8 ± 1.93 nm	53.6 ± 4.73 nm
Polydispersity index	0.382	0.428	0.402

Since previous studies have found that rheological property of microemulsion is of Newtonian type, the viscosity measurement in this study was performed using one shear rate for all the formulations. This resulted in the viscosity of the microemulsion that is also of Newtonian type and suitable for topical use (Table 1). In addition, there are differences in the viscosity among the three formulations due to different ratio of surfactants. The more amount of Tween, the more viscous microemulsion.

The droplet size shown in Table 1 indicated that the microemulsion had a droplet size ranging from 10 to 100 nm. The ratio of standard deviation to the mean of droplet size is shown as polydispersity index. It indicates the uniformity of microemulsion droplet size within the formulation. The lower the polydispersity index, the better the uniformity of the droplet size in the microemulsion. The polydispersity values of all formulations are about 0.4 indicating that uniformity of droplet size within each formulation is good. From stability test, the visual observation confirmed that all the formulations did not significantly change in their appearance over the storage period at 4°C and 25°C although there was a slight change in their viscosity, pH value, droplet size, and polydispersity index. There was a tendency for the droplet size to become smaller over 4 weeks storage. This phenomenon possibly occurred due to the solubilization of some of the extract and/or of the oil phase in surfactant micelles during the storage.

Meanwhile, Formulation 1 and 2 stored at 40°C were not stable during the 4 weeks storage. There was phase separation in these microemulsions, which did not occur in Formulation 3 because the high storage temperature affected the activity of Tween 80. Previous study reported that in emulsions stabilized by non-ionic surfactants (e.g. Tween 80) elevated temperatures raise droplet coalescence through dehydration of non-ionic head groups. This process alters the solubility of the surfactant in the oil and water phases that can lead to phase separation.

Anti-inflammation activities of P. major microemulsion: In the topical anti-inflammatory test using croton oil induced ear edema, the negative control group demonstrated the greatest degree of edema, while the mice group treated with microemulsion of *P. major* extracts demonstrated significant reduction of the edema (Figure 1).




Figure 1. Photomicrograph of transverse sections of mice ears (treatments: normal group (A), croton oil (B) hydrocortisone 1% (C), microemulsion base (D) and microemulsion of *P. major* extracts (E)), stained with hematoxylin-eosin and examined under light microscopy (magnification 400x).

Croton oil (contains 12-*o*-tetracanoilphorbol-13-acetate (TPA)) topical application triggered local inflammation with edema formation and polymorphonuclear leucocytes infiltration hence suitable for studying anti-inflammatory agent that act on acute phase of inflammation^{15,17} It induces an inflammatory response by activating phospholipase A2 and initiating arachidonic acid metabolites, followed by COX-1 and COX-2-induced prostaglandin production and 5-lipoxygenase-induced leukotriene production, the inflammatory mediators involved in edema and leukocyte migration.^{18,19} Application of croton oil increased ear edema in negative control group strengthening the right design of this models. The histopathological analysis from ear edema exhibited dermal inflammatory reactions noted by decreasing of infiltration of leukocyte and vascular dilatation. Inflammatory cells were also counted (data not published) from the histological sections. The inflammatory cells were decreased in *P. major* microemulsion treatment group significantly different with croton oil group.

CONCLUSION

P. major microemulsion had been successfully produced by combining Tween 80 and Span 80 as the surfactants. Compared to Formulation 1 and Formulation 2, Formulation 3 containing Tween 80-Span 80 ratio of 10:1 was the most stable microemulsion after four weeks of storage at various temperatures. Furthermore, *P. major* microemulsion possess topical anti-inflammatory activities strengthened the notation that this plant may develop into a new anti-inflammatory agent.

ACKNOWLEDGEMENTS

The authors wish to thank the *Indonesian Directorate General of Higher Education (DIKTI)* and Directorate of Research and Community Service (DPPM) Islamic University of Indonesia for providing the financial support.

REFERENCES

1. Hussan F, Mansor AS, Hassan SN, Kamaruddin NTTNE, Budin SB, Othman F. Anti-Inflammatory Property of Plantago major Leaf Extract Reduces the Inflammatory Reaction in Experimental Acetaminophen-Induced Liver Injury. *Evidence-based Complementary and Alternative Medicine*. 2015. doi:10.1155/2015/347861

2. Hussan F, Osman Basah RH, Mohd Yusof MR, Kamaruddin NA, Othman F. Plantago major treatment enhanced innate antioxidant activity in experimental acetaminophen toxicity. *Asian Pacific Journal of Tropical Biomedicine*. 2015;5(9):728-732. doi:10.1016/j.apjtb.2015.06.013
3. Mello JC, Gonzalez MVD, Moraes VWR, et al. Protective effect of plantago major extract against t-BOOH-induced mitochondrial oxidative damage and cytotoxicity. *Molecules*. 2015;20(10):17747-17759. doi:10.3390/molecules201017747
4. Metİner K, Ozkan O, Seyyal A. Antibacterial Effects of Ethanol and Acetone Extract of Plantago major L . on Gram Positive and Gram Negative Bacteria. *Kafkas Univ Vet Fak Derg*. 2012;18(3):503-505. doi:10.9775/kvfd.2011.5824
5. Abdulghani MA, Hamid I, Al-Naggar RA, Osman MT. Potential antidiabetic activity of Plantago Major leaves extract in Streptozocin-induced diabetic rats. *Research Journal of Pharmaceutical, Biological and Chemical Sciences*. 2014;5(2):896-902.
6. Samuelsen AB. A Review: The Traditional Uses, Chemical Constituents and Biological Activities of Plantago major L. *Journal of Ethnopharmacology*. 2000;71:1-21. doi:10.1016/S0378-8741(00)00212-9
7. Amini M, Kherad M, Mehrabani D, Azarpira N, Panjehshahin MR, Tanideh N. Effect of Plantago major on Burn Wound Healing in Rat. *Journal of Applied Animal Research*. 2010;37(1):53-56. doi:10.1080/09712119.2010.9707093
8. J. Jamilah, AA. Sharifa NS. GC-MS Analysis of Various Extracts from Leaf of Plantago major Used as Traditional Medicine. *World Applied Sciences Journal*. 2012;17(February):67-70.
9. Ringbom T, Segura L, Noreen Y, Perera P, Bohlin L. Ursolic acid from Plantago major, a selective inhibitor of cyclooxygenase-2 catalyzed prostaglandin biosynthesis. *Journal of Natural Products*. 1998;61(10):1212-1215. doi:10.1021/np980088i
10. Kolak U, Boĝa M, Uruřak EA, Ulubelen A. Constituents of Plantago major subsp. intermedia with antioxidant and anticholinesterase capacities. *Turkish Journal of Chemistry*. 2011;35(4):637-645. doi:10.3906/kim-1102-990
11. Nazarizadeh A, Mikaili P, Moloudizargari M, Aghajanshakeri S, Javaherypour S. Therapeutic Uses and Pharmacological Properties of Plantago major L . and its Active Constituents. 2013;3(9):212-221.
12. Reina E, Al-Shibbani N, Allam E, Gregson KS, Kowolik M WL. The Effects of Plantago major on the Activation of the Neutrophil Respiratory Burst. *Journal of Traditional and Complementary Medicine*. 2013;3(4):268-272. doi:10.4103/2225-4110.119706
13. Fanun M. Microemulsions as delivery systems. *Current Opinion in Colloid and Interface Science*. 2012;17(5):306-313. doi:10.1016/j.cocis.2012.06.001
14. Lopes LB. Overcoming the cutaneous barrier with microemulsions. *Pharmaceutics*. 2014;6(1):52-77. doi:10.3390/pharmaceutics6010052
15. Patel M, Murugananthan, Gowda S. In Vivo Animal Models in Preclinical Evaluation of Anti- Inflammatory Activity- A Review. *International Journal of Pharmaceutical Research & Allied Sciences*. 2012;1(2):1-5.
16. Wang H, Chen Z. Microemulsion-based novel transdermal delivery system of tetramethylpyrazine : preparation and evaluation in vitro and in vivo. 2011:1611-1619.
17. Trivellatograssi L, Malheİros A, Meyre-Silva C, et al. From popular use to pharmacological validation: A study of the anti-inflammatory, anti-nociceptive and healing effects of Chenopodium ambrosioides extract. *Journal of Ethnopharmacology*. 2013;145(1):127-138. doi:10.1016/j.jep.2012.10.040
18. Tubaro a., Dri P, Delbello G, Zilli C, Loggia R Della. The Croton oil ear test revisited. *Agents and Actions*. 1986;17(3-4):347-349. doi:10.1007/BF01982641
19. Call DR, Nemzek J a, Ebong SJ, Bolgos GL, Newcomb DE, Remick DG. Ratio of local to systemic chemokine concentrations regulates neutrophil recruitment. *The American journal of pathology*. 2001;158(2):715-721. doi:10.1016/S0002-9440(10)64014-X

FULL PAPER

ANTIBACTERIAL ACTIVITY AND COMPOSITION OF ESSENTIAL OILS OF LEAVES AND BASIL FLOWERS OF THREE DIFFERENT GROWING SEASONS UNDER ARID CLIMATE

Nabila Slougui^{1,2}, Ahlem Tlili², Hadjer Bentayeb², Mahfoud Hadj Mohammed²

¹ Preparatory Class Department at National Polytechnic School of Constantine, Algeria,

² Biogeochemistry Laboratory of the Desert Environments, Kasdi Merbah University, Ouargla, Algeria
Email: slougui.nabila@gmail.com

Abstract

We wanted to study antibacterial activity of essential oils in relation to the growing seasons of the plant as the first parameter of study, and what part of plant that will present a better antibacterial activity as the second parameter of study. The relationship between this activity and the majority composition of the essential oil has been developed. **Material and Methods:** The antibacterial activity of 6 essential oils was evaluated on three two Gram-negative strains: *Escherichia coli* ATCC 25923 and *Pseudomonas aeruginosa* ATCC 27853 and one gram positive: *Staphylococcus aureus* ATCC 25922. The diffusion method in Agar medium (Aromatogram) is used for the study of antibacterial activity which allows determining the sensitivity of the different bacterial species towards our essential oils. The majority composition of essential oils was made by a Shimadzu QP2020 GC-MS. **Results:** According to our results, the essential oil of basil autumn flowers inhibits bacterial growth with inhibition diameters of 21.5 and 25 mm against *Escherichia coli* and *Staphylococcus aureus*, respectively. This has a direct relationship with its major composition: Linalool (34.99%), methylcinnamate (21.09%) and Methylchavicol (14.53%). **Conclusion / Discussion:** The essential oils of the flowers showed antibacterial activities better than that of the leaves and this in the three study seasons, summer, autumn, and winter.

Keywords: antibacterial activity, *Ocimum basilicum* L, essential oil, growing seasons.

INTRODUCTION

The south of Algeria is rich in its natural resources which are not well exploited. The plants with medicinal character constitute most of these resources. Spontaneous plants and aromatic plants in Algeria present a growing interest for researchers in the world. The traditional uses of plants for medicine were studied in south Algeria. Until now, 85 spontaneous medicinal plants have been identified [1]. To our knowledge, only two varieties of *Ocimum basilicum* are cultivated and marketed on a national scale in Algeria. These are: *Ocimum basilicum* and *Ocimum basilicum* minimum. The genus *Ocimum* belonging to the Lamiaceae comprises annual and perennial herbs and shrubs native to the tropical and subtropical regions of Asia, Africa, and South America [2]. Apart from culinary use, basil has traditionally been employed as a medicinal herb in the treatment of headaches, cough, diarrhoea, and constipation. It is also considered to be a source of aroma compounds and essential oils containing biologically active constituents that possess insect repellent, nematocidal and antibacterial activity. Many *Ocimum basilicum* varieties contain primarily phenol derivatives, such as eugenol, methyl eugenol, chavicol, methylcinnamate often combined with various amounts of linalool [3]. In the realm of the natural resources valorisation of arid regions in Algeria, our study, first interested in evaluating the antibacterial activities of essential oils of *Ocimum basilicum* from three different seasons and the relationship between this activity and the majority composition of these oils.

MATERIAL AND METHODS

Plant Material: The plant material used was harvested in the state of Ouargla exactly in the village of Oum Erraneb in southern east of Algeria. Oum Erraneb, agricultural zone, is a village in the municipality of Sidi Khouiled, located 18 km northeast of the Ouargla city, at latitude of 32° 03 ' 53.24 north and at a longitude of 5 ° 22 ' 33.34 east, elevated to 129 m in relation to sea level. We proceeded to separate the different organs, leaves and flowers.

Essential Oils Extraction: Samples, 1 kg of fresh material, were extracted by steam distillation for 2.5 hours. The essential oils were stored in dark glass bottles at 4°C [4] until the beginning of the analyses.

Antibacterial Activity Determination: The antibacterial activity was evaluated on three strains: two gram-negative, *Escherichia coli* ATCC 25923 (digestive tract infections) and *Pseudomonas aeruginosa* ATCC 27853 (the commensal flora of the skin in humans), and a Gram positive, *Staphylococcus aureus* ATCC 25922 (nosocomial infections). They are maintained by replanting on nutrient agar (NI) favourable to their growth for 24 hours, in darkness and at 37 °C [5]. The microorganisms used are pathogenic bacteria of the American type culture collection (ATCC) frequently involved in the contamination and alteration of food and often responsible for infections contracted in Hospitals [6]. The method of diffusion in agar medium (Aromatogram) is used for the study of the antibacterial activity which allows determining the sensitivity of the different bacterial species towards the essential oils studied. Petri dishes (90 mm) containing Muller-Hinton agar (MHA) are inoculated, sterile discs (6 mm diameter filter paper) impregnated with different essential oils (5µl per disc) are then gently deposited on the surface of the agar using a clamp. The whole is incubated at 37 °C in the oven for 24 hours [7]. After 24 hours of incubation, an area or a clear Halo is present around the disc. This inhibition diameter reflects the antibacterial activity of the essential oil. The result is expressed by measuring the diameter of the inhibition zone where the bacteria could not develop. The latter will be qualified: Not sensitive (-) or resistant, diameter < 8mm; sensitive (+), diameter from 9 to 14 mm; very sensitive (+ +), diameter from 15 to 19 mm; and extremely sensitive (+ + +), diameter > 20mm [8].

Determination of Essential Oils Composition by GC-MS: GC-MS analysis was performed in National Polytechnic School of Constantine, Algeria using a Shimadzu System comprising a model QP 2020. GC was equipped with non-polar capillary column HP-5 (30m * 0.25 mm, film thickness 0.25 µm). Helium was the carrier gas at a flow rate of 1.0 ml/min. The oven temperature was held at 45°C for 10 min, and increased from 45°C to 180°C at a rate of 3°C/min to be maintained at 180°C for 5 min, and then increased from 180°C to 280°C at a rate of 5°C/min to be maintained at 280°C for 5 min. For GC-MS detection, an electron ionization system with ionization energy of 70 eV, was used. Injector temperature was 280°C. Diluted samples, in dichloromethane of 1 µl, were injected in the split/splitless (100:1 split) mode. Oil identification of components analysed was accomplished based on comparison of their Arithmetic index (AI), calculated from GC-FID analysis, with those of literature [9], and by comparison of their mass spectral fragmentation patterns with those of databases.

RESULT AND DISCUSSION

Antibacterial Activity: The results of antibacterial tests with essential oils (different parts each season) summarized in table 1. The results obtained showed that the essential oil of the different parts of the *Ocimum basilicum* L. had no antibacterial activity marked against the strain of *Pseudomonas aeruginosa* (ATCC 27853), the least susceptible to antibiotics tested. The microorganisms most susceptible to these essential oils were *Staphylococcus aureus* (ATCC 25923) and *Escherichia coli* (ATCC 25922). According to Bassol, gram-positive bacteria are more sensitive than gram-negative bacteria with essential oils [10]. The zones of inhibition of the antibiotics tested are around 19 and 30 mm. This sensitivity varies from one antibiotic to another. This activity is very strong compared to that of the oils studied. Indeed, the Autumn E.O. (essential oil) of *Ocimum basilicum* L. showed an important inhibitory effect against the microorganisms studied. According to our results, it is concluded that autumn promotes this strong activity compared to the other seasons. The inhibition diameters ranged from 8 to 18 mm for the leaves and from 12 to 25 for the flowers. According to our results, the essential oil of the flowers of *Ocimum basilicum* L. inhibits bacterial growth more than that of leaves. The results of Ijaz Hussain demonstrate that the autumn and winter E.O. are more active than in the spring and summer months: *Staphylococcus aureus* (24.4 ± 1.1; 24.0 ± 1.0; 23.2 ± 1.4; 22.2 ± 1.3 mm) and *Escherichia coli* (13.2 ± 0.8; 16.2 ± 1.0; 13.5 ± 0.8; 11.4 ± 0.6 mm) [7]. The activity of the essential oils tested could be related to their chemical compositions. According to other studies, the analysis of essential oils, flowers, and leaves of *O. basilicum* L. showed a variation in the chemical composition of one organ to another. In other words, the study of the antibacterial activity of certain constituents of essential oils made it possible to distinguish that phenolic compounds have strong antimicrobial activity, such as thymol, carvacrol, and eugenol. Constituents with low antibacterial activity are pulegone, menthone, 1.8-cineole,

p-cymene, isomenthone, myrcene, α -pinene, piperitone, limonene, linalool, terpinene, sesquiterpenes, and non-terpene compounds [7]. According to these literature data, the variation in the essential oils' antibacterial activity of *Ocimum basilicum* L's leaves and flowers is due to their difference in the content of active compounds which must be determined by GC-MS.

Table 1: Diameters of inhibition zones (including 6 mm disk) of *O. basilicum* L. essential oils (E.O)

The strains	Summer		Fall		Winter	
	Leaves	flowers	Leaves	flowers	Leaves	flowers
<i>Escherichia coli</i> (G-) ATCC 25922	8 mm -	12 mm +	16 mm ++	21.5 mm +++	6 -	13 mm +
<i>Staphylococcus aureus</i> (G+) ATCC 25923	10 mm +	14 mm +	18 mm ++	25 mm +++	6 -	20 mm +++
<i>Pseudomonas aeruginosa</i> (G-) ATCC 27853	6 -	6 -	6 -	6 -	6 -	6 -

Essential Oils Major Composition: The GC-MS Chromatograms of the 6 essential oils are illustrated in Figure 1 where very different profiles are observed. The identified major compounds as well as their respective outages are summarised in table 2.





Figure 1: GC-MS chromatograms of leaves and flowers essential oils in three seasons

Table 2: Major composition of essential oils

Majority Composition	Summer		Fall		Winter	
	Leaves	Flowers	Leaves	Flowers	Leaves	Flowers
linalool	30,26%	32,14%	30,75%	34,99%	29,13%	26,66%
methylchavicol	17,58%	08,05%	44,08%	14,53%	20,43%	18,00%
linalyl acetate	17,03%	-	-	-	16,15%	15,18%
methylcinnamate	-	18,66%	17,06%	21,09%	-	-

The essential oils of the autumn season consist of the three major compounds: linalool, methylchavicol, and methylcinnamate. The absence of linalylacetate is clearly noted by comparing it with the essential oils of the other seasons (c.f. table 1).

CONCLUSION

The study of the essential oils' antibacterial activity of *O. Basilicum* L. on three bacterial strains (*Escherichia coli* ATCC 25922, *Pseudomonas aeruginosa* ATCC 27853, and *Staphylococcus aureus* ATCC 25923). And according to the results of the Aromatogram, our E.O. presented an important antibacterial activity against two bacterial strains, with the exception of *Pseudomonas aeruginosa* ATCC 27853 which is known from literature by its extensive antibiotic resistance, flowers' essential oil inhibited bacterial growth more than that of leaves. Besides that autumn promotes this activity. This variability is due to the impacts of environmental factors on the chemical composition of E.O. and their antibacterial activities. The study of essential oils' composition by GC-MS has allowed the identification of the major compounds: linalool, methyl chavicol, linalylacetate, and methylcinnamate with rates that change from one season to another and from one part to another.

REFERENCES

- [1] Jirasek, V. J., 1983. List of Botanical Gardens and Institutes Offering Seeds of Plants Collected from Indigenous Habitats. *Taxon*, 32, 584-597.
- [2] Darrah, H. H., 1988. *The cultivated basil*. Missouri Buckeye Printing Company, MO, Independence.
- [3] Weker, E., Putievsky, E., Ravid, U., Dudai, N., Katzir, I., 1993 Glandular hairs and essential oil in developing leaves of *Ocimum basilicum* (lamiaceae). *Ann. Bot.* 71, 43-50.
- [4] Marotti, M., Piccaglia, R., Giovanelli, E., 1996. Differences in essential oil composition of basil (*Ocimum basilicum* L.) Italian cultivars related to morphological characteristics. *J. Agric. Food Chem.* 44, 3926-3929.
- [5] Branger, J., Knapp, S., Weijer, S., Leemans, J. C., Pater, J. M., Speelman, P., Florquin, S., van der Poll, T., 2004, Role of Toll-like receptor 4 in gram-positive and gram negative pneumonia in mice. *Infect Immun.* 72:788–794.
- [6] Nabrdalik, M., Grata, K., 2016, Antibacterial activity of *Ocimum basilicum* L. essential oil against Gram-negative bacteria, *Post Fitoter.* 17(2), 80-86.
- [7] Hussain, A. I., Anwar, F., Hussain Sherazi, S. T., Przybylski, R., 2008. Chemical composition, antioxidant and antimicrobial activities of basil (*Ocimum basilicum*) essential oils depends on seasonal variations. *Pakistan, Food Chem.* 108(3), 986-95.
- [8] Bougurra, A., Himed, L., Barakat, M., 2014. Étude de l'activité antimicrobienne de l'huile essentielle extraite des écorces de *Citrus reticulata*. *SAN*, 03(1), 32-39.
- [9] Adams, R. P., 2007. *Identification of Essential Oils Components by Gas Chromatography/Quadrupole Mass Spectrometry*, 4th Edition, Allured Publishing Corporation, Carol Stream, IL, USA.
- [10] Bassole, I. H. N., Ouattara, A. S., Nebie, R., *et. al.* 2011. Composition chimique et activités antibactériennes des huiles essentielles des feuilles et des fleurs de *cymbopogon proximus* (stapf.) et d'*ocimum canum* (SIMS). *Pharm. Méd. Trad.* 11, 37-51.

IMPACT OF SOME HERBICIDES AND HERBICIDE COMBINATIONS ON SOWING CHARACTERISTICS OF CORIANDER (*CORIANDRUM SATIVUM* L.)

Grozi Delchev

Department of Plant Production, Faculty of Agriculture, Trakia University, Stara Zagora, Bulgaria
delchevgd@dir.bg

Abstract

The research was conducted during 2013 - 2015 on pellic vertisol soil type. Under investigation was Bulgarian coriander cultivar Lozen 1 (*Coriandrum sativum* L.). Factor A included no treated control, 6 soil-applied herbicides – Tender EC (S-metolachlor) - 1.5 l/ha, Silba SC (metolachlor + terbuthylazine) - 3.5 l/ha, Sharpen 33 EC (pendimethalin) - 5 l/ha, Merlin flex 480 SC (isoxaflutole) - 420 g/ha, Smerch 24 EC (oxyfluorfen) - 1 l/ha, Raft 400 SC (oxidiargil) - 1 l/ha and 5 foliar-applied herbicides – Kalin flo (linuron) - 2 l/ha, Eclipse 70 DWG (metribuzine) - 500 g/ha, Sultan 500 SC (metazachlor) - 2 l/ha, Corrida 75 DWG (tribenuron-methyl) - 20 g/ha, Lontrel 300 EC (clopyralid) - 500 ml/ha. Factor B included no treated control and 1 antigraminaceous herbicide – Tiger platinum 5 EC (quizalofop-P-ethyl) - 2.5 l/ha. Soil-applied herbicides were treated during the period after sowing before emergence. Foliar-applied herbicides were treated during rosette stage of the coriander. Herbicides Merlin flex and Lontrel and herbicide combinations Merlin flex + Tiger platinum and Lontrel + Tiger platinum proven decrease germination energy of the coriander seeds. Laboratory seed germination and lengths of primary germ and primary root are decreased by herbicide Merlin flex and herbicide combination Merlin flex + Tiger platinum only. Herbicide Merlin flex and herbicide combination Merlin flex + Tiger platinum do not proven decrease waste grain quantities. High yields of coriander seeds are obtained by foliar treatment with antigraminaceous herbicide Tiger platinum after soil-applied herbicides Raft, Smerch, Sharpen, Silba and Tender. Tank mixtures of Tiger platinum with foliar herbicides Kalin flo, Eclipse, Sultan, Corrida and Lontrel also lead to obtaining of high seed yields. The use of the soil-applied herbicide Merlin flex does not increase the seed yield, due to its higher phytotoxicity against coriander. Alone use of soil-applied or foliar-applied herbicides leads to lower yields due to they must to combine for full control of weeds in coriander crops.

Key words: coriander, herbicides, herbicide combinations, seed yield, sowing characteristics

INTRODUCTION

Coriander (*Coriandrum sativum* L.) is an important essential-oil plant, referring to the *Apiaceae* family. The technology for coriander growing over time needs to be updated, as a result of ever-changing weather conditions. Specify and significantly improve crop rotations, tillage, timings of sowing and harvesting (Delibaltova et al., 2012). All of this leads to changes in the conditions for the weeds development, to the emergence of resistant or new weed species that were previously marginal or were not harmful.

The weeds occurring in coriander crops of are mainly from the group of the winter-spring species and from the group of early spring species. The most frequent winter-spring weeds are: *Alopecurus myosuroides* L., *Apera spica-venti* P.B., *Bromus arvensis* L., *Avena ludoviciana* Durien., *Lolium multiflorum* L., *Anthemis arvensis* L., *Chamomilla recutita* Rauchert, *Consolida regalis* Gray, *Viola tricolor* L., *Lithospermum arvense* L., *Centaurea cyanus* L., *Papaver rhoes* L., *Agrostemma githago* L. The most frequent early spring weeds are: *Avena fatua* L., *Galium aparine* L., *Sinapis arvensis* L., *Falopia convolvulus* Leve, *Myagrimum perfoliatum* L. (Mitchell and Abernethy, 1993; Atanasova et al., 2012). Biological peculiarity of coriander is its slow growth rates in the first stages and therefore it is highly vulnerable to the competitive impact of weeds at the beginning of its vegetation and weed infestation significantly (Atanasova and Gospodinov, 2005, Gramatikov et al., 2005). Effective weed control is one the important conditions for full realization of the biological potential of culture (Macko, 1990; Lugo and Santiago, 1996; Vaculik, 2007; Meena and Mehta, 2009; Delchev and Georgiev, 2015,

2015a; Delchev, 2018). The presence of weeds in coriander crops requires carrying out of their chemical control to preserve coriander yield at a maximum level (Zheljazkov and Zhalnov, 1995; Smart et al., 1996; Mathukia et al., 2014). The aim of this investigation was to establish the influence of some soil-applied herbicides, foliar-applied herbicides and herbicide combinations on sowing characteristics of the coriander seeds and the quantity of waste grain.

MATERIALS AND METHODS

The research was conducted during 2013 - 2015 on pellic vertisol soil type. Under investigation was Bulgarian coriander cultivar Lozen 1 (*Coriandrum sativum* L.). Two factors experiment was conducted under the block method, in 4 repetitions; the size of the crop plot was 15 m². Factor A included no treated control, 6 soil-applied herbicides – Tendar EC, Silba SC, Sharpen 33 EC, Merlin flex 480 SC, Smerch 24 EC, Raft 400 SC and 5 foliar-applied herbicides – Kalin flo, Eclipse 70 DWG, Sultan 500 SC, Corrida 75 DWG, Lontrel 300 EC. Factor B included no treated control and 1 antigraminaceous herbicide – Tiger platinum 5 EC. Active substances of herbicides and their doses are shown in Table 1.

Table 1. Investigated variants

№	Variants	Active substance	Doses	Treatment period
Antibroadleaved herbicides				
1	Control	-	-	-
2	Tendar EC	S-metolachlor	1.5 l/ha	ASBE
3	Silba SC	metolachlor + terbuthylazine	3.5 l/ha	ASBE
4	Sharpen 33 EC	pendimethalin	5 l/ha	ASBE
5	Merlin flex 480 SC	isoxaflutole	420 g/ha	ASBE
6	Smerch 24 EC	oxyfluorfen	1 l/ha	ASBE
7	Raft 400 SC	oxidiargil	1 l/ha	ASBE
8	Kalin flo	linuron	2 l/ha	rosette
9	Eclipse 70 DWG	metribuzine	500 g/ha	rosette
10	Sultan 500 SC	metazachlor	2 l/ha	rosette
11	Corrida 75 DWG	tribenuron-methyl	20 g/ha	rosette
12	Lontrel 300 EC	clopyralid	500 ml/ha	rosette
Antigraminaceous herbicides				
1	Control	-	-	-
2	Tiger platinum 5 EC	quizalofop-P-ethyl	2.5 l/ha	rosette

ASBE – after sowing, before emergence

Soil-applied herbicides were treated during the period after sowing before emergence. Foliar-applied herbicides were treated during rosette stage of the coriander. Introduction of herbicide combinations during rosette stage is done as herbicide tank mixtures – the mixing is done in the spray tank. All of herbicides, herbicide combinations and herbicide tank-mixtures were applied in a working solution of 200 l/ha. Mixing of foliar-applied herbicides was done in the tank on the sprayer.

The grain gained after every variant was cleaned through a sieves and the quantity of the waste grain was defined (siftings). All version seeds for sowing were defined for their germination energy and lab seed germination. It was studied intensity of early growth of seeds, expressed by the lengths of primary germ and primary root definite on the eighth day after setting the samples. Each index was determined in two repetitions of the year. Averages in each of the years of experience were used as repetitions in mathematical data processing were done according to the method of analysis of variance.

RESULTS AND DISCUSSION

One of the important conditions for obtaining a normal crop and a good harvest is the use of quality seeds. Apart from the high-yield cultivar which is resistance to diseases and pests, it must have the necessary sowing properties, the main of which are high germination energy and seed germination.

Germination energy is one of the most important characteristics of the sowing properties of the seed. The low germination energy is the reason for slower development of primary germ and primary root after seed germination and is associated with later germination in field conditions, less tempering of plants and a higher risk of frost in the winter. Its lead to lower seed yields. The obtained results show that the treatment of the coriander with herbicides Merlin flex and Lontrel and herbicide combinations Merlin flex + Tiger platinumium and Lontrel + Tiger platinumium lead to the decrease in the germination energy (Table 2). Analysis of variance, in which the years have taken for replications, shows that these decreases are mathematically proven.

Table 2. Influence of some herbicides and herbicide combinations on sowing characteristics of the coriander seeds (mean 2013 - 2015)

Herbicides		Germinative energy, %	Germination %	Length, cm		Waste grain, %
Antibroadleaved	Antigraminaceous			Primary germ	Primary root	
-	-	76.5	87.0	7.8	8.2	16.9
	Tiger platinumium	90.0	95.5	10.0	12.0	14.1
Tendar	-	90.0	95.0	10.1	12.1	14.4
	Tiger platinumium	90.5	95.0	10.4	12.5	14.0
Silba	-	90.0	96.0	10.6	12.5	14.5
	Tiger platinumium	90.0	95.5	10.3	12.7	14.0
Sharpen	-	90.0	96.0	10.7	12.3	14.4
	Tiger platinumium	90.5	96.5	10.4	12.5	14.4
Merlin flex	-	75.0	88.0	7.0	8.6	15.6
	Tiger platinumium	75.0	87.5	7.2	8.3	15.7
Smerch	-	89.5	94.5	10.3	11.9	13.9
	Tiger platinumium	90.5	95.0	10.2	11.7	14.1
Raft	-	91.0	96.0	10.2	11.5	14.3
	Tiger platinumium	91.0	95.0	10.2	11.4	14.0
Kalin flo	-	90.5	96.5	10.8	12.7	14.1
	Tiger platinumium	89.5	95.5	10.0	12.8	14.4
Eclipse	-	91.0	96.5	10.3	11.3	14.0
	Tiger platinumium	91.0	96.0	10.4	12.0	13.8
Sultan	-	90.5	95.0	10.5	11.8	13.0
	Tiger platinumium	91.5	96.0	10.3	11.5	13.6
Corrida	-	89.5	95.5	10.1	11.3	14.1
	Tiger platinumium	90.0	96.5	10.3	11.6	14.0
Lontrel	-	76.0	94.5	9.1	11.3	14.8
	Tiger platinumium	75.5	94.0	9.3	11.0	14.1
LSD 5 %		6.2	5.3	1.8	2.1	2.0
LSD 1 %		8.3	7.0	2.6	2.9	3.3
LSD 0.1 %		10.5	9.2	3.7	4.0	4.8

Germination is the most important index who characterizing the sowing properties of the seeds. At low laboratory germination sowing should be done with higher sowing rate, which increases the cost production. Laboratory germination of the seeds at all variant during the three years of study is above the requirements of the standard, although in different years account for some variation of its values. This is the positive effect of their use, because it is not necessary to increase the sowing rate (in kg/ha) and the cost of necessary seeds. At herbicide Merlin flex and herbicide combination Merlin flex + Tiger platinumium seed germination is lower than untreated control. The coriander seeds germinate normally by influence of the herbicide Lontrel and herbicide combination Lontrel + Tiger platinumium, although the initial rate of development is lower due to lower germination energy. Other soil-applied herbicides, foliar-applied herbicides and their combinations increase the indexes germination energy and seed germination. This means that they help for joint and fast germination of the coriander sowing-seeds. The obtained results for germination energy and seed germination are a prerequisite continue to investigate the effect of herbicides and their combinations on initial intensity of the growth of seeds, expressed by the lengths of primary germs and roots. It was found that the lengths of primary germ and primary root

of coriander are decreased by herbicide Merlin flex and herbicide combination Merlin flex + Tiger platinum. These decreasing are proven by analysis of variants. This two variants difficult young plants developments, reduces their resistance to cold and increase risk of frost damages during winter months. Herbicide Lontrel and herbicide combination Lontrel + Tiger platinum does not increase proven length of primary germ, but increase proven length of primary root. Other tank mixtures between soil-applied and foliar-applied herbicides led to increase of the lengths of primary germ and primary root of the coriander and recommended for use in seed production crops of coriander.

Table 3. Influence of some herbicide and herbicide combinations on seed yield of coriander (2013 - 2015)

Herbicides		2013		2014		2015		Mean	
Antibroadleaved	Antigraminaceous	kg/ha	%	kg/ha	%	kg/ha	%	kg/ha	%
-	-	2150	100	2333	100	2060	100	2181	100
	Tiger platinum	2227	103.6	2433	104.3	2142	104.0	2267	103.9
Tendar	-	2288	106.4	2492	106.8	2182	105.9	2321	106.4
	Tiger platinum	2417	112.4	2678	114.8	2348	114.0	2481	113.8
Silba	-	2288	106.4	2499	107.1	2188	106.2	2325	106.6
	Tiger platinum	2423	112.7	2685	115.1	2348	114.0	2485	113.9
Sharpen	-	2333	108.5	2550	109.3	2245	109.0	2376	108.9
	Tiger platinum	2526	117.5	2753	118.0	2408	116.9	2562	117.5
Merlin flex	-	2008	93.4	2240	96.0	1840	89.3	2029	93.0
	Tiger platinum	2176	101.2	2384	102.2	2017	97.9	2192	100.5
Smerch	-	2311	107.5	2522	108.1	2227	108.1	2353	107.9
	Tiger platinum	2498	116.2	2734	117.2	2396	116.3	2543	116.6
Raft	-	2356	109.6	2563	110.7	2250	109.2	2390	109.6
	Tiger platinum	2537	118.0	2767	118.6	2433	118.1	2579	118.2
Kalin flo	-	2376	110.5	2592	111.1	2268	110.1	2412	110.6
	Tiger platinum	2537	118.0	2755	118.1	2414	117.2	2569	117.8
Eclipse	-	2301	107.0	2522	108.1	2217	107.6	2347	107.6
	Tiger platinum	2481	115.4	2702	115.8	2385	115.8	2523	115.7
Sultan	-	2301	106.0	2508	107.5	2200	106.8	2336	107.1
	Tiger platinum	2473	115.0	2690	115.3	2377	115.4	2513	115.2
Corrida	-	2270	105.6	2492	106.8	2184	106.0	2315	106.1
	Tiger platinum	2457	114.3	2685	115.1	2369	115.0	2504	114.8
Lontrel	-	2258	105.0	2473	106.0	2167	105.2	2299	105.4
	Tiger platinum	2451	114.0	2676	114.7	2365	114.8	2497	114.5
LSD 5 %		80	3.7	85	3.6	79	3.8		
LSD 1 %		104	4.8	110	4.7	100	4.7		
LSD 0.1 %		133	6.2	147	6.3	128	6.2		

At the evaluation of the sowing characteristics we have to consider not only the characteristics of the sowing seeds but also the quantity of the waste grain (siftings) which are gained at the preparation of these seeds. Bigger quantity screenings lead to higher cost of the seed and reduce the economic effect of seed production of coriander. Herbicide Merlin flex and herbicide combination Merlin flex + Tiger platinum do not lead to are mathematically proven increase in the quantity of waste grain. Other herbicides and herbicide combinations lead to decreasing in the quantity of waste grain. Differences between them and untreated control are mathematically proven. Decreases in the values of germination energy and laboratory seed germination, changes in the intensity of the initial growth, expressed by the lengths of the primary root and primary germ at germination and changes in the quantity of waste grain by the influence of the relevant herbicides and herbicide combinations are explained by the depressing effects on growth and development of the coriander during its vegetative period.

To done a full evaluation of the sowing characteristics needed to establish not only the quality of seeds, but also the quantity of grain which will be received this seeds. Data for the influence of investigated herbicides and herbicide combinations on seed yield show that the lower yield is obtained by alone use of herbicide Merlin flex, followed by the untreated control (Table 3). The use of soil-applied herbicide Merlin flex do not increases seed yield despite its very good herbicidal effect against both graminaceous and broadleaved weeds. The reason for this is its higher phytotoxicity against coriander. Alone use of foliar-applied herbicides Kalin flo, Eclipse, Sultan, Corrida and Lontrel and as well as alone use of soil-applied herbicides Tendar, Silba, Sharpen, Smerch and Raft increases seed yields because the big numbers of weeds are destroyed by these herbicides. The increases of the yield by alone use of antigraminaceous herbicide Tiger platinum is less than the increase by other herbicides, because prevailing weeds in the trail are broadleaved.

Foliar treatment with Tiger platinum after soil-applied herbicides Tendar, Silba, Sharpen, Smerch and Raft increases the seed yields during the three years of the investigation. Tank mixtures of Tiger platinum with herbicides Kalin flo, Eclipse, Sultan, Corrida and Lontrel also lead to bigger increase in seed yields compared to yield at alone use of respective herbicides. Herbicide combination Merlin flex + Tiger platinum increases seed yield compared to alone use of Merlin flex, but not proven increase yield compared to no treated control.

CONCLUSION

Herbicides Merlin flex and Lontrel and herbicide combinations Merlin flex + Tiger platinum and Lontrel + Tiger platinum proven decrease germination energy of the coriander seeds.

Laboratory seed germination and lengths of primary germ and primary root are decreased by herbicide Merlin flex and herbicide combination Merlin flex + Tiger platinum only.

Herbicide Merlin flex and herbicide combination Merlin flex + Tiger platinum do not proven decrease waste grain quantities.

High yields of coriander seeds are obtained by foliar treatment with antigraminaceous herbicide Tiger platinum after soil-applied herbicides Raft, Smerch, Sharpen, Silba and Tender.

Tank mixtures of Tiger platinum with foliar herbicides Kalin flo, Eclipse, Sultan, Corrida and Lontrel also lead to obtaining of high seed yields.

The use of the soil-applied herbicide Merlin flex does not increase the seed yield, due to its higher phytotoxicity against coriander.

Alone use of soil-applied or foliar-applied herbicides leads to lower yields due to they must to combine for full control of weeds in coriander crops.

REFERENCES

1. Atanasova, D., Gospodinov, G., 2005. Studies on herbicides for weed control in coriander (*Coriandrum sativum* L.). Balkan Conference. Breeding and farming practices on field crops, Part II, 540-542. Institute of Agriculture – Karnobat.

2. Atanasova, D., Maneva, V. Dacheva, S., 2012. Phytosanitary Monitoring of Coriander Crops in South-East Bulgaria. *Plant Science*, 49 (2) 36-40.
3. Delchev, Gr., 2018. Chemical control of weeds and self-sown plants in eight field crops. Monograph, ISBN: 978-613-7-43367-6, LAP LAMBERT Academic Publishing, Saarbrücken, Germany, pp. 397.
4. Delchev, Gr., Georgiev, M., 2015. Achievements and problems in the weed control in oilseed canola (*Brassica napus* L.). *Scientific Papers. Series A. Agronomy*, Vol. LVIII, 174-180.
5. Delchev, Gr., Georgiev, M., 2015. Achievements and problems in the weed control in common wheat (*Triticum aestivum* L.) and durum wheat (*Triticum durum* Desf.). *Agricultural Science and Technology*, 7 (3), 281-286.
6. Delibaltova, V., Yanchev, I. Ivaylova, V., 2012. Effect of the date and density of sowing on the seed yield and yield components of coriander – (*Coriandrum sativum* L.) *Agrarni nauki*, 16 (4) 45-50.
7. Gramatikov, B., Koteva, V. Penchev, P. Atanasova, D., 2005. Technology for growing of coriander. Sofia.
8. Lugo, M. de; Santiago, L., 1996. Herbicide screening in cilantro and spiny coriander. *Journal of Agriculture of the University of Puerto Rico*, 80 (1-2) 73-75.
9. Macko V., 1990. Chemical control of weeds in coriander (*Coriandrum sativum* L.) at intensive occurrence of the disease in this crop. *Bilten za hmelj, sirak i lekovito bilje, Yugoslavia* v.41-45, 60-61.
10. Mathukia, R., Sagarka, B. Panara, D., 2014. Efficient and Economical Weed Management in Coriander (*Coriandrum sativum* L.). *Journal of Crop Science and Technology*, 3 (3) 30-33.
11. Meena, S., Mehta, R., 2009. Integrated weed management in coriander (*Coriandrum sativum*). *Indian Journal of Agricultural Sciences*, 79 (10) 824-826.
12. Mitchell, R. B., Abernethy, R. J., 1993. Tolerance of clary sage, coriander, and caraway to herbicides applied pre- and post-emergence 24–29. *Proceedings of the Forty-Sixth New Zealand Plant Protection Conference, Christchurch, New Zealand, 10–12 August 1993*.
13. Smart J., Makus, D. Coleman, R., 1996. Trifluralin for weed control in coriander, dill, and dandelion. *Horticulture Science*, 31 (5), 757-760.
14. Vaculík, A., 2007. The influence of the herbicides treatment on the yield and essential oil content of spice plants growth in the Czech Republic. Dissertation thesis. Department of Crop Science, Breeding and Plant Medicine, Brno, Czech Republic.
15. Zheljazkov, V., Zhalnov, I., 1995. Effect of herbicides on yield and quality of *Coriandrum sativum* L. *Journal of Essential Oil Research*, 7 (6) 633-639.

FULL PAPER

CHANGES IN THE WEED SPECIES IN FIELDS OF MILK THISTLE, CORIANDER, CHICKPEA AND FORAGE PEA WHICH ARE SOWN ON DAMAGED BY FROST AREAS OF WINTER OILSEED CANOLA

Grozi Delchev

Department of Plant Production, Faculty of Agriculture, Trakia University, Stara Zagora, Bulgaria
delchevgd@dir.bg

Abstract

During 2016 - 2018 was conducted a field experiment. On areas with damaged by frost winter oilseed canola, at the spring were sowed and investigated 4 spring crops: 1 milk thistle cultivar - Silmar (*Silybum marianum* Gaertn.); 1 coriander cultivar - Lozen (*Coriandrum sativum* L.); 1 chickpea cultivar - Kabule (*Cicer arietinum* L.); 1 forage pea cultivar - Mir (*Pisum sativum* L.). The same variants were planted on areas under conventional soil cultivation for each of these crops. Weed control was carried out by different herbicide combinations and herbicide tank mixtures. Coriander and milk thistle are suitable crops for sowing on areas after damaged by frost winter oilseed canola, when using herbicide combination Stomp aqua + Stratos ultra or herbicide tank mixture Zencor + Shadow. After plowing of canola crops, it is more appropriate to sow chickpea in which weed control is carried out by soil treatment with herbicide Merlin flex, followed by foliar treatment with herbicide tank mixture Challenge + Shadow. Herbicide combination Dual gold + Listego should not be used in chickpea. On plowed areas with damaged by frost winter canola hybrids without any problems can be sown forage pea, in which weed control is carried out by herbicide combinations Stomp aqua + Korum or Dual gold + Listego.

Keywords: milk thistle, coriander, chickpea, forage pea, weeds, herbicides

INTRODUCTION

Weed control is very important in the initial stage of milk thistle, coriander, chickpea and forage pea developments, when they are less competitive. The use of herbicides creates favorable conditions for germination, growth and development of these crops and also for the creation of well topped and high-yielding crops (Dann et al., 1987; Andrzejewska and Lamparski, 2006; Vaculík, 2007; Khan et al., 2009; Wágner, 2015; Delchev, 2018). In the crops of milk thistle, coriander, chickpea and forage pea occurring weeds of various biological groups. The implementation of the biological potential of these crops is closely related to the removal of the harmful effects of weeds (Meena and Mehta, 2009; Tanveer et al., 2010; Ratnam and Rao, 2011; Drapalova and Pluhackova, 2014; Mathukia et al., 2014; Tidemann et al., 2014). The aim of this experiment is to investigate the changes in the weed species in fields of 4 crops - milk thistle, coriander, chickpea and forage pea which are sown on damaged by frost crops of winter oilseed canola.

MATERIAL AND METHODS

During 2016 - 2018 was conducted a field experiment on pellic vertisol soil type. It was carried out a field experiment as a block method in 4 repetitions, on a 15 m² harvesting area. On areas with damaged by frost winter oilseed canola, at the spring were sowed and investigated 4 spring crops: 1 milk thistle cultivar - Silmar (*Silybum marianum* Gaertn.); 1 coriander cultivar - Lozen (*Coriandrum sativum* L.); 1 chickpea cultivar - Kabule (*Cicer arietinum* L.); 1 forage pea cultivar - Mir (*Pisum sativum* L.). These variants have been sown also on the areas with traditional for each of those crops soil tillage.

Weed control in milk thistle was carried out with herbicide combination Stomp aqua (pendimethalin) - 3 l/ha + Stratos ultra (cycloxydim) - 2 l/ha and herbicide tank mixture Zencor 70 WG + Shadow 3 EC (metribuzine + clethodim) - 500 g/ha + 1.6 l/ha. Soil-applied herbicide Stomp aqua was treated during

the period after sowing before emergence. Foliar-applied herbicides Stratos ultra, Zencor and Shadow were treated during rosette stage of the milk thistle.

Weed control in coriander was carried out with herbicide combination Stomp aqua (pendimethalin) - 3 l/ha + Stratos ultra (cycloxydim) - 2 l/ha and herbicide tank mixture Zencor 70 WG + Shadow 3 EC (metribuzine + clethodim) - 500 g/ha + 1.6 l/ha. Soil-applied herbicide Stomp aqua was treated during the period after sowing before emergence. Foliar-applied herbicides Stratos ultra, Zencor and Shadow were treated during rosette stage of the coriander. Weed control in chickpea was carried out with herbicide combinations Dual gold 960 EC (S-metolachlor) - 1.5 l/ha + Listego 40 (imazamox) - 1.2 l/ha and Merlin flex 480 SC (isoxaflutole) - 420 g/ha + herbicide tank mixture Challenge 600 SC + Shadow 3 EC (aclonifen + clethodim) - 4 l/ha + 1.6 l/ha. Soil-applied herbicides Dual gold and Merlin flex were treated during the period after sowing before emergence. Foliar-applied herbicides Listego, Challenge and Shadow were treated during 6 - 8 real leaf stage of the chickpea.

Weed control in forage pea was carried out with herbicide combinations Dual gold 960 EC (S-metolachlor) - 1.5 l/ha + Listego 40 (imazamox) - 1.2 l/ha and Stomp aqua (pendimethalin) - 3 l/ha + Korum (bentazone + imazamox) - 1.25 l/ha. Soil-applied herbicides Dual gold and Stomp aqua were treated during the period after sowing before emergence. Foliar-applied herbicides Listego and Korum were treated during 6 - 8 real leaf stage of the pea. Due to of low adhesion of the herbicides Listego and Korum were used in addition with adjuvant Dash HC – 1 l/ha. It was investigated the changes occurring in the weed species in 4 crops - forage pea, chickpea, coriander and milk thistle. It was appointed the efficacy of the herbicides according to 100 % visual scale of EWRS (European Weed Research Society). It was followed the selectivity of the herbicides by the scale of EWRS (rating 1 - without damages, rating 9 - culture is completely destroyed).

RESULTS AND DISCUSSION

On the area of frozen winter canola was sown milk thistle cultivar Silmar. The cultivar was sowed as soon as possible to enter in the field. The same cultivar was sown at the same time and on area with traditional coriander soil tillage. Weed control was done with herbicide combination Stomp aqua + Stratos ultra and with herbicide tank mixture Zencor + Shadow (Tables 1 and 2). In both weed control variants, the efficacy of the used herbicide combinations is very good. A small reduction in the efficacy of herbicide combination Stomp aqua + Stratos ultra against some annual broadleaf weeds such as *Galium aparine* L., *Papaver rhoeas* L. and *Sinapis arvensis* L., when sowing milk thistle after frozen canola has been found. Used herbicide combination and herbicide tank mixture enable effective weed control during crop growing against graminaceous and broadleaf weeds. This makes the milk thistle a suitable crop for sowing on areas after frozen winter oilseed canola.

On the area of frozen winter canola was sown coriander cultivar Lozen. The cultivar was sowed as soon as possible to enter in the field. The same cultivar was sown at the same time and on area with traditional coriander soil tillage. Weed control was done with herbicide combination Stomp aqua + Stratos ultra and with herbicide tank mixture Zencor + Shadow (Tables 3 and 4). A small reduction in the efficacy of herbicide combination Stomp aqua + Stratos ultra against some annual broadleaf weeds such as *Galium aparine* L., *Papaver rhoeas* L. and *Sinapis arvensis* L., when sowing coriander after frozen canola has been found. In both weed control variants, the efficacy of the herbicide combinations used is very good. This means that coriander is a suitable crop for sowing on areas after damaged by frost winter oilseed canola.

Table 1. Efficacy of herbicide combination and herbicide tank mixture against broadleaved weeds at milk thistle according to the 100 % visual scale of EWRS (mean 2016 - 2018)

Variants	Weeds									
	<i>Galiu n aparin e</i>	<i>Ch am omi lla rec utit a</i>	<i>Papav er rhoes</i>	<i>Sina pis arve nse</i>	<i>Raph anus raph anist rum</i>	<i>Ant he mis arv ensi s</i>	<i>Capse lla bursa pastor is</i>	<i>Cirsiu m arvens e</i>	<i>Co nvo lvul us arv ensi s</i>	<i>Card aria drab a</i>
Milk thistle – Sowing after canola										
Stomp aqua + Stratos ultra	98	100	99	98	100	100	100	0	0	0
Zencor + Shadow	15	100	98	98	100	100	100	0	0	0
Milk thistle – Normal sowing										
Stomp aqua + Stratos ultra	100	100	100	100	100	100	100	0	0	0
Zencor + Shadow	15	100	100	100	100	100	100	0	0	0

Table 2. Efficacy of herbicide combination and herbicide tank mixture against graminaceous weeds and self-sown plants at milk thistle according to the 100 % visual scale of EWRS and selectivity according to the 9-rate scale of EWRS (mean 2016 - 2018)

Variants	Weeds									
	<i>Avena fatua</i>	<i>Ave na ludo vici ana a</i>	<i>Alop ecur us myo soro ides</i>	<i>Apera spica venti</i>	<i>Bromus arvensis</i>	<i>Triticum durum *</i>	<i>Sorg hum hele pens e</i>	<i>Cyn odo n dact ylon</i>	<i>Agr opyr um repe ns</i>	Sele cti vity
Milk thistle – Sowing after canola										
Stomp aqua + Stratos ultra	100	100	100	100	100	100	100	100	100	1
Zencor + Shadow	100	100	100	100	100	100	100	100	100	1
Milk thistle – Normal sowing										
Stomp aqua + Stratos ultra	100	100	100	100	100	100	100	100	100	1
Zencor + Shadow	100	100	100	100	100	100	100	100	100	1
* - self-sown plants of durum wheat										

On the area of frozen winter canola was sown chickpea cultivar Kabule. Sowing was done in late February. The same cultivar was sown at the same time and on area with traditional chickpea soil tillage. Weed control was done with two herbicide combinations: Dual gold + Listego and Merlin flex + herbicide tank mixture Challenge + Shadow (Tables 5 and 6). It was found a stronger phytotoxicity of herbicide Listego to the chickpea plants, although Listego provides better control of perennial broadleaf weeds compared to the herbicide tank mixture Challenge + Shadow. Combining of soil-applied herbicide Merlin flex with herbicide tank mixture Challenge + Shadow does not lead to phytotoxicity at chickpea plants.

Table 3. Efficacy of herbicide combination and herbicide tank mixture against broadleaved weeds at coriander according to the 100 % visual scale of EWRS (mean 2016 - 2018)

Variants	Weeds									
	<i>Galiu n aparine</i>	<i>Ch am omi lla rec utit a</i>	<i>Papav er rhoes</i>	<i>Sina pis arve nse</i>	<i>Raph anus raph anist rum</i>	<i>Ant he mis arv ensi s</i>	<i>Capse lla bursa pastor is</i>	<i>Cirsiu m arvens e</i>	<i>Co nvo lvul us arv ensi s</i>	<i>Card aria drab a</i>
Coriander – Sowing after canola										
Stomp aqua + Stratos ultra	98	100	98	98	100	100	100	0	0	0
Zencor + Shadow	14	100	98	97	100	100	100	0	0	0
Coriander – Normal sowing										
Stomp aqua + Stratos ultra	100	100	100	100	100	100	100	0	0	0
Zencor + Shadow	15	100	100	100	100	100	100	0	0	0

Table 4. Efficacy of herbicide combination and herbicide tank mixture against graminaceous weeds and self-sown plants at coriander according to the 100 % visual scale of EWRS and selectivity according to the 9-rate scale of EWRS (mean 2016 - 2018)

Variants	Weeds									
	<i>Avena fatua</i>	<i>Aven a ludo vicia naa</i>	<i>Alop e cur us myso roi des</i>	<i>Apera spica venti</i>	<i>Bromus arvensis</i>	<i>Triticum durum *</i>	<i>Sorg hum hele pens e</i>	<i>Cyno don dact ylon</i>	<i>Agro pyru m repe ns</i>	Sel ecti vit y
Coriander – Sowing after canola										
Stomp aqua + Stratos ultra	100	100	100	100	100	100	100	100	100	1
Zencor + Shadow	100	100	100	100	100	100	100	100	100	1
Coriander – Normal sowing										
Stomp aqua + Stratos ultra	100	100	100	100	100	100	100	100	100	1
Zencor + Shadow	100	100	100	100	100	100	100	100	100	1

* - self-sown plants of durum wheat

At these two technologies, the high efficiency of the using herbicides is the reason for a good control of weeds in both sowing - after frozen canola and after deep plowing. These results lead to the conclusion that after plowing canola crops is appropriate to sow chickpea in which weed control should be carried out by soil treatment with herbicide Merlin flex, followed by foliar treatment with the herbicide tank mixture Challenge + Shadow. On the area of frozen winter canola was sown in early spring forage pea cultivar Mir. The cultivar was sowed as soon as possible to enter in the field. It was done a pre-sowing cultivation accompanied with harrowing. Weed control was done with two herbicide combinations: Dual gold + Listego and Stomp aqua + Korum (Tables 7 and 8). Herbicide Korum has a better effect than herbicide Listego.

Table 5. Efficacy of herbicide combinations against annual broadleaved weeds and self-sown plants at chickpea according to the 100 % visual scale of EWRS (mean 2016 - 2018)

Variants	Weeds									
	Galiun aparine	Cha mo mill a recu tita	Papaver rhoes	Consolid a regalis	Am aran thus retro flex us	Ant hem is arve nsis	Falo pia con volv ulus	Ver onic a hede rifol ia	Lam ium purp ureu m	He lia nth us an nu us *
Chickpea – Sowing after canola										
Dual gold + Listego	93	100	94	100	92	92	93	100	100	0
Merlin flex + Challenge + Shadow	100	100	100	100	98	99	100	100	100	100
Chickpea – Normal sowing										
Dual gold + Listego	98	100	100	100	100	100	100	100	100	0
Merlin flex + Challenge + Shadow	100	100	100	100	100	100	100	100	100	100
* - self-sown plants of ClearField and ExpressSun sunflowers										

Table 6. Efficacy of herbicide combinations against perennial weeds, annual graminaceous weeds and self-sown plants at chickpea according to the 100 % visual scale of EWRS and selectivity according to the 9-rate scale of EWRS (mean 2016 - 2018)

Variants	Weeds									
	<i>Cirsium arvense</i>	<i>Con volv ulus arv ensi s</i>	<i>Sorgum helepen s</i>	<i>Cynodo n dactylon</i>	<i>Avena fatua</i>	<i>Loli um mul tiflo rum</i>	<i>Alo pec uru s myo sor oide s</i>	<i>Bromus arvensis</i>	<i>Triticum durum*</i>	Se lec tiv ity
Chickpea – Sowing after canola										
Dual gold + Listego	98	100	94	95	100	100	100	98	100	3
Merlin flex + Challeng e + Shadow	0	0	100	100	100	100	100	100	100	1
Chickpea – Normal sowing										
Dual gold + Listego	98	100	97	100	100	100	100	98	100	3
Merlin flex + Challeng e + Shadow	0	0	100	100	100	100	100	100	100	1
* - self-sown plants of durum wheat										

Korum has longer control over the secondary-emerged weeds until the pea plants cover the whole soil surface, competes with weeds and almost prevents secondary weed infestation. These results showed that after plowing of areas with frozen canola hybrids without any problems can be planted forage pea.

Table 7. Efficacy of herbicide combinations against annual broadleaved weeds at forage pea according to the 100 % visual scale of EWRS (mean 2016 - 2018)

Variants	Weeds									
	<i>Galiun aparine</i>	<i>Chamomilla recutita</i>	<i>Papaver rhoeas</i>	<i>Consolida regalis</i>	<i>Sinapis arvensis</i>	<i>Raphanus raphanistrum</i>	<i>Anthemis arvensis</i>	<i>Falopia convolvulus</i>	<i>Veronica hederaefolia</i>	<i>Lamium purpureum</i>
Forage pea – Sowing after canola										
Dual gold + Listego	93	100	94	100	95	98	100	93	100	100
Stomp aqua + Korum	100	100	100	100	100	100	100	100	100	100
Forage pea – Normal sowing										
Dual gold + Listego	100	100	100	100	100	100	100	100	100	100
Stomp aqua + Korum	100	100	100	100	100	100	100	100	100	100

Table 8. Efficacy of herbicide combinations against perennial broadleaved, annual graminaceous weeds and self-sown plants at pea according to the 100 % visual scale of EWRS and selectivity according to the 9-rate scale of EWRS (mean 2016 – 2018)

Variant s	Weeds									
	<i>Cirsium arvense</i>	<i>Convolvulus arvensis</i>	<i>Cardaria draba</i>	<i>Sonchus arvensis</i>	<i>Avena fatua</i>	<i>Lolium multiflorum</i>	<i>Alopecurus myosuroides</i>	<i>Bromus arvensis</i>	<i>Triticum durum*</i>	Selec tivity
Forage pea – Sowing after canola										
Dual gold + Listego	99	100	100	100	100	100	100	98	100	1
Stomp aqua + Korum	100	100	100	100	100	100	100	98	100	1
Forage pea – Normal sowing										
Dual gold + Listego	100	100	100	100	100	100	100	98	100	1
Stomp aqua + Korum	100	100	100	100	100	100	100	98	100	1
* - self-sown plants of durum wheat										

CONCLUSIONS

Coriander and milk thistle are suitable crops for sowing on areas after damaged by frost winter oilseed canola, when using herbicide combination Stomp aqua + Stratos ultra or herbicide tank mixture Zencor + Shadow.

After plowing of canola crops, it is more appropriate to sow chickpea in which weed control is carried out by soil treatment with herbicide Merlin flex, followed by foliar treatment with herbicide tank mixture Challenge + Shadow. Herbicide combination Dual gold + Listego should not be used in chickpea.

On plowed areas with damaged by frost winter canola hybrids without any problems can be sown forage pea, in which weed control is carried out by herbicide combinations Stomp aqua + Korum or Dual gold + Listego.

REFERENCES

1. Andrzejewska J., Lamparski L., 2006. Assessing the extent of damage caused by *Cleonus piger* Skop. and other entomofauna in the cultivation of milk thistle (*Silybum mariannum* Gaertn.). Journal of Plant Protection Research, No. 46, 49-59.
2. Dann P.R., Thomas A.G., Cunningham R.B., Moore P.H., 1987. Response by wheat, rape and field peas to pre-sowing herbicides and deep tillage. Australian Journal of Experimental Agriculture, v. 27(3) p. 431-437.
3. Delchev, Gr., 2018. Chemical control of weeds and self-sown plants in eight field crops. Monograph, ISBN: 978-613-7-43367-6, LAP LAMBERT Academic Publishing, Saarbrücken, Germany, pp. 397.
4. Drapalova, I., Pluhackova, H., 2014. Effect of herbicide treatment on milk thistle (*Silybum marianum* Gaertn.) germination. Conference MendelNet, 19-20 November 2014. Mendel University in Brno, Czech Republic, 33-38.
5. Khan, M.A., Blackshaw, R.E., Marwat, K.B., 2009. Biology of milk thistle (*Silybum marianum*) and the management options for growers in north-western Pakistan. Weed Biology and Management, 9 (2) 99–105.
6. Mathukia, R., Sagarka, B., Panara, D., 2014. Efficient and Economical Weed Management in Coriander (*Coriandrum sativum* L.). Journal of Crop Science and Technology, 3 (3) 30-33.
7. Meena, S., Mehta, R., 2009. Integrated weed management in coriander (*Coriandrum sativum*). Indian Journal of Agricultural Sciences, 79 (10) 824-826.
8. Ratnam, M.M., Rao, A.S., 2011. Integrated weed management in chickpea (*Cicer arietinum* L.). Indian Journal of Weed Science, 43 (1-2) 70-72.
9. Tanveer, A., Imran, S., Ayub, M., Yasin, M., 2010. Response of chickpea (*Cicer arietinum*) and *Euphorbia dracunculoides* to pre- and post-emergence herbicides. Pakistan Journal of Weed Science Research. 16 (3) 267-277.
10. Tidemann, B.D., Hall, L.M., Johnson, E.N., Beckie, H.J. Sapsford, K.L. Raatz, L.L., 2014. Efficacy of Fall- and Spring-Applied Pyroxasulfone For Herbicide-Resistant Weeds in Field Pea. Weed Technology 28(2) 351-360.
11. Vaculík, A., 2007. The influence of the herbicides treatment on the yield and essential oil content of spice plants growth in the Czech Republic. Dissertation thesis. Department of Crop Science, Breeding and Plant Medicine, Brno, Czech Republic.
12. Wágner, G., 2015. Competition and Herbicide Effect Studies with Green Pea. Thesis of PhD Dissertation. Keszthely, Hungary.

CORRELATION BETWEEN CHEMICAL COMPOSITION AND ANTIOXIDANT ACTIVITY OF ESSENTIAL OILS FROM SIX AROMATIC MEDICINAL PLANTS GROWING IN ILLIZI AND GHARDAÏA (SOUTHERN ALGERIA)

Naima Hellali^{1,2}, Mebarka Bouziane², Mahfoud Hadj Mahammed²

¹ Faculty of Sciences and Technology, Department of process engineering, Ghardaïa University, BP 455, 47000 Ghardaïa, Algeria, hellali.naima@univ-ghardaia.dz

² Biogeochemistry Laboratory in Desert Environments, Faculty of Science, Kasdi Merbah University, BP 511, 30000 Ouargla, Algeria.

Abstract

This study provides an assessment to the relationship between the chemical composition of the essential oils and the antioxidant activity of six aromatic medicinal plants growing in southern Algeria, namely *Cymbopogon schoenanthus*, *Cymbopogon Citratus*, *Artemisia judaica*, *Artemisia herba alba*, *Pituranthos chloranthus* and *Pituranthos scoparius*. The essential oils were obtained by hydrodistillation and analyzed by GC/MS. Their antioxidant activity was determined by using three methods of testing (DPPH, ABTS and iron reduction test (FRAP)), the results showed that the essential oil of *A. judaica* has the best antioxidant capacity in all tests. The statistical study allowed us to show a correlation between the chemical composition of the studied essential oils and the antioxidant activity and our results indicate that the high content of cyclic ether sesquiterpenes compounds (such as: Davana ether), the compounds have furan rings, or the existence of non-terpene alkenes/esters in the essential oil, increases its antioxidant effect.

Keywords: *C. schoenanthus*, *C. Citratus*, *A. judaica*, *A. herba alba*, *P. chloranthus*, *P. scoparius*, essential oil, antioxidant activity.

INTRODUCTION

Essential oils (EOs) are complex mixtures that can contain 20–60 compounds in different concentrations, and are characterized by two or three main components with higher concentrations compared to others present in lower concentrations. In general, the major compounds that are present in EOs are responsible for the biological activities (Budel et al., 2018), Antioxidant properties of essential oils from many plants have also been of great interest to the food processing industry, since their possible use as natural additives has emerged from a growing tendency to replace synthetic antioxidants with natural ones (khadri et al., 2008; Djeridane et al., 2006).

The medicinal plants contain various types of antioxidants, mostly polyphenols and flavonoids which exhibit high antioxidant activity (Gohari et al., 2011). As part of evaluation of the biological effectiveness of the essential oil from the medicinal plants, this paper presented a study of the antioxidant activity associated with the chemical composition of essential oils isolated from *Cymbopogon schoenanthus* (Poaceae), *Cymbopogon Citratus* (Poaceae), *Artemisia judaica* (Asteraceae), *Artemisia herba alba* (Asteraceae), *Pituranthos chloranthus* (Apiaceae), *Pituranthos scoparius* (Apiaceae).

MATERIALS AND METHODS

Plant material and extraction of EOs: The aerial parts of each plant were collected as follows: *C. schoenanthus* and *P. chloranthus* were collected from two Southern regions of Algeria, the first is Ghardaïa (weddi Cebceb, on April, 2014) and the second is Illizi (weddi tasset, on April, 2014). *A. herba alba* and *P. scoparius* were collected from weddi Cebceb in Ghardaïa (Algeria), on April 2014. *C. Citratus* were collected from Djanet in Illizi (Algeria), on December 2011. And concerning *A. judaica* were collected from weddi Tasset in Illizi (Algeria), on April 2014. Extraction of the EOs was carried

out by hydrodistillation in a modified Clevenger-type apparatus for 3 h, The oil collected from each plant was dehydrated with Na₂SO₄ and stored in sealed glass vials at 4°C, prior to analyses.

Essential oil chromatographic analysis: The EOs were injected on a gas chromatography (Hewlett-packar computerized system, Agilent model 6890 GC coupled to a Agilent 5973 N mass selective detector , and equipped with an Agilent technologies capillary HP-5MS column (30 m, 0,25 mm I.d, 0.25µm thickness), temperature program and conditions were described by Hellale et al., (2016).

Antioxidant activity determinations: Three methods have been applied for the antioxidant activity of extracted EOs in this study: the 1,1-diphenyl- 2-picrylhydrazyl radical (DPPH[·]) assay, the 2,20-azino-bis(3-ethylbenzothiazoline-6-sulphonate) radical cation (ABTS^{·+}) assay and the ferric reducing antioxidant power (FRAP) assay. All chemicals, antioxidant activity essays and calculation of Trolox equivalent antioxidant capacity (TEAC) were described by Hellali et al., (2016).

Statistical analysis: Correlation analysis of antioxidant activity versus the chemical composition of EOs were carried out using the correlation and regression program in Excel 2007 program.

RESULTS

The compounds identified by GC-MS of the EOs extracted and their relative proportions are listed in the Table 1.

Table 1: Chemical composition of the studied essential oils

Compound	Taux %							
	Cyc	CyII	CyGh	Ps	PcGh	PcII	Arh	ArJ
1 cis-Salvene	-	-	-	-	-	-	0.31	-
2 β-thujene	-	-	-	-	1.55	-	-	-
3 α-thujene	-	-	-	0.29	-	1.50	-	-
4 α-Pinene	-	-	-	10.53	12.95	14.24	-	0.41
5 Unknown	-	-	-	-	-	-	-	0.35
6 Camphene	-	-	-	0.11	-	-	1.24	-
7 Sabinene	-	-	-	11.97	5.17	-	0.94	-
8 β-Pinene	-	-	-	-	3.23	3.80	-	-
9 β-Myrcene	5.52	-	-	0.65	1.01	-	-	-
10 α-Phellandrene	-	-	1.48	17.50	21.96	1.22	-	-
11 2-Carene	-	4.923	-	-	-	-	-	-
12 4-Carene	-	-	19.76	-	0.19	-	-	-
13 O-Cymene	-	-	-	-	-	1.90	-	-
14 p-Cymene	-	-	-	-	3.51	-	-	0.84
16 Limonene	-	1.413	7.39	36.90	31.55	3.85	-	-
17 Eucalyptol	-	-	-	-	-	-	4.11	-
18 Geraniol ester	-	-	-	-	-	-	0.61	-
19 cis-b-Ocimene	-	0.014	-	-	-	-	-	-
20 trans-b-Ocimene	-	0.151	1.79	-	-	-	-	-
21 trans-Ocimene	-	-	-	-	-	3.33	-	-
22 γ-terpinene	-	-	-	1.57	0.96	-	-	-
23 Terpinolene	-	-	-	0.17	-	-	-	-
24 α-terpinolene	0.49	-	-	-	-	-	-	-
25 α-Thuyone	-	-	1.41	3.14	-	-	73.33	-
26 β-Thuyone	-	-	-	0.44	-	-	11.96	-
27 cis-p-menth-2-en-1-ol	-	-	29.21	-	-	-	-	-
28 (4E,6E)-2-Methyl-2,4,6-octatriene	-	-	-	-	-	-	-	5.32
29 Camphor	-	-	-	0.32	-	-	5.92	-
30 trans-p-menth-2-en-1-ol	-	-	15.62	-	-	-	-	-
31 Cyclohexanecetaldehyde, 2-Methylene	-	-	-	-	-	-	-	1.08
32 trans-sabinenhydrate	-	-	-	-	-	-	0.97	-
33 D-fenchone	-	0.037	-	-	-	-	-	-
34 1-terpineol	-	0.429	-	-	-	-	-	-
35 4-terpineol	-	-	-	0.93	0.88	-	-	-
36 α-terpineol	-	1.447	1,71	-	-	-	-	-
37 cis piperitol	-	0.106	5.81	-	-	-	-	-
38 trans piperitol	-	0.029	5.05	-	-	-	-	-

39	Neral	37.06	-	-	-	-	-	-	-
40	Piperitone	-	63.35	-	-	-	-	-	79.04
41	Geraniol	0.54	-	-	-	-	-	-	-
42	Geranial	54.91	-	-	-	-	-	-	-
43	α -Cubebene	-	-	-	0.10	-	-	-	-
44	O-Methyleugenol	-	-	-	-	0.79	-	-	-
45	β -elemene	-	0.616	-	-	-	-	-	-
46	β -caryophyllene	-	0.060	-	-	-	-	-	-
47	Calarene	-	0.094	-	-	-	-	-	-
48	α -caryophyllene	-	0.032	-	-	-	-	-	-
49	Germacrene-D	-	0.210	-	0.92	-	-	0.62	1.41
50	β -selinene	-	0.269	-	-	-	-	-	-
51	α -selinene	-	0.242	-	-	-	-	-	-
52	α -muurolene	-	0.562	-	-	-	-	-	-
53	β -Cubebene	-	-	-	-	1.15	1.19	-	-
54	Unknown	-	-	-	0.23	-	-	-	-
55	Bicyclogermacrene	-	-	-	0.27	-	-	-	0.77
56	Davana ether (isomer 1)	-	-	-	-	-	-	-	0.40
57	Davana ether (isomer 2)	-	-	-	-	-	-	-	1.18
58	γ -cadinene	-	0.345	-	-	-	-	-	-
59	δ -cadinene	-	0.946	-	0.76	-	-	-	-
60	Myristicin	-	-	-	-	12.58	65.15	-	0.95
61	Davana ether	-	-	-	-	-	-	-	0.56
62	Elemol	-	6.915	3.48	-	-	-	-	-
63	Spathulenol	-	-	-	-	-	1.49	-	0.47
64	Davanone	-	-	-	-	-	-	-	7.23
65	Selina-6-en-4-ol	-	-	1.82	-	-	-	-	-
66	Apiol	-	-	-	-	0.97	1.18	-	-
67	α -Cadinol	-	-	-	0.44	-	-	-	-
68	10-epi- γ -eudesmol	-	1.266	-	-	-	-	-	-
69	β -eudesmol	-	9.305	1.39	-	-	1.15	-	-
70	α -eudesmol	-	1.875	-	-	-	-	-	-
71	Globulol	-	-	1.89	-	-	-	-	-
72	Butylidene phthalide	-	-	-	0.18	-	-	-	-
73	2-Isopropenyl-4a,8-dimethyl-1,2,3, 4,4a,5,6,7-octahydronaphthalene	-	-	2.19	-	-	-	-	-
74	(Z)-Ligustilide	-	-	-	12.26	1.74	-	-	-
75	3-Butyl-2-benzofuran-1(3H)-one	-	-	-	0.13	-	-	-	-
Monoterpene hydrocarbons (%)		6.01	6.501	30.42	79.88	81.89	29.8	2.18	1.25
Oxygenated monoterpenes (%)		92.51	65.398	53.76	4.83	0.88	0	96.3	79.04
Sesquiterpene hydrocarbons (%)		0	3.376	2.19	2.05	1.15	1.19	0.62	2.18
Oxygenated sesquiterpenes (%)		0	19.361	13.63	0.44	0	2.64	0	9.84
Composes oxygenes non terpeniques (%)		0	0	0	12.57	16.08	66.3	0	0.95
Other compounds (%)		0	0	0	0	0	0	0.92	6.4

Cyc: *C. Citratus*, CyII: *C. schoenanthus* from Illizi, CyGh: *C. schoenanthus* from Ghardaïa, Ps : *P. scoparius* , PcGh: *P. chloranthus* from Ghardaïa, Pcll: *P. chloranthus* from Illizi , Arh : *A. herba alba* , ArJ : *A. judaica*,

Results from the antioxidant activities assays of extracted EOs are summarized in Table 2.

Table 2. Antioxidant properties against DPPH, ABTS and FRAP assays of the studied EOs

Essential oil	Harvest area	DPPH assay IC ₅₀ (mg/ml)	ABTS assay IC ₅₀ (mg/ml)	FRAP assay TEAC (μ moles / g)
<i>C. schoenanthus</i>	Illizi	44.219 \pm 3.78	18.91 \pm 0.17	3.33 \pm 0.3
	Ghardaïa	21.68 \pm 5.1	15.16 \pm 4.42	2.94 \pm 0.6
<i>C. citrates</i>	Illizi	15.584 \pm 0.72	53.74	6.64 \pm 2.3
<i>A. judaica</i>	Illizi	3.63\pm0.17	2.09\pm0.17	86.28\pm 3.2
<i>A. herba alba</i>	Ghardaïa	11.478 \pm 0.66	6.36 \pm 1.43	17.02 \pm 0.7
<i>P. chloranthus</i>	Illizi	8.24 \pm 1.11	6.10 \pm 1.37	24 .27 \pm 2.2
	Ghardaïa	10.747 \pm 3.24	5.9 \pm 1.72	25.96 \pm 3.9
<i>P. scoparius</i>	Ghardaïa	5.812 \pm 0.88	7.47 \pm 2.64	49.18 \pm 3.8

In order to correlate the antioxidant activity with the chemical composition of the extracted essential oils, and to identify compounds in the EOs potentially associated with antioxidant activities, the coefficients of correlation " R^2 " between the abundances of the compounds and the antioxidant activities were calculated and determined by Excel 2007. The results are summarized in Table 3.

Table 3. Correlation coefficients between antioxidant activity and chemical classes that exist in extracted EOs

Grouped componets	Correlation coefficient " R^2 "		
	DPPH assay	FRAP assay	ABTS assay
Monoterpene hydrocarbons	0,000	0,001	0,032
Oxygenated monoterpenes	0,009	0,011	0,001
Monoterpenes cyclic ether	0,013	0,019	5E-05
Monoterpene ketones	0,052	0,078	0,214
Monoterpene Alcohols	0,104	0,118	0,066
Monoterpene Aldehydes	0,052	0,083	0,149
Sesquiterpene hydrocarbons	0,392	0,375	0,293
Oxygenated sesquiterpenes	0,042	0,017	2E-06
Sesquiterpenes cyclic ether	0,678	0,709	0,840
Sesquiterpenes Alcohols	0,304	0,249	0,153
Phenylpropene	0,002	0,000	0,002
Furan cycle compounds	0,711	0,682	0,845
Alkenes and esters non-terpenics	0,658	0,682	0,859
Hyterocycles	0,622	0,655	0,326

DISCUSSION AND CONCLUSION

The chemical composition of essential oils obtained from *C. schoenanthus* of Ghardaïa and of Illizi was different. The *C. schoenanthus* EOs of Illizi, for instance, was characterized by the high content of piperitone (Hellali et al., 2016), whereas the major compounds found in *C. schoenanthus* EO of Ghardaïa are cis-p-menth-2-en-1-ol (29.21%), trans-p-menth-2-en-1-ol (15.62 %), 4-Carene (19.76%), cis piperitol (5.81%) and trans piperitol (5.05 %).

P. chloranthus EO of Illizi is a chemotype characterized by absence of oxygenated monoterpenes content and high myristicin content (65.15 %). The second *P. chloranthus* chemotype was found in Ghardaïa region, which differs from the first at the presence of high content of monoterpene hydrocarbons (81.89%), Limonene (31.55 %), α -Phellandrene (21.96%) and α -pinene (12.95%) were the main compounds in this chemotype. Limonene, α -Phellandrene and α -pinene they also the major compounds in *Pituranthos scoparius* EO of Ghardaïa (36.90% and 17.50% and 10.53 respectively), in addition to the presence of a significant percentage of **Sabinène 11.97%** in *Pituranthos scoparius* EO. Hellali et al., (2017) have shown that for the *A. judaica* EO, the oxygenated monoterpenes had the most important proportions (79.04%), which represents the proportion of piperitone. The oxygenated monoterpenes have been found 96.29% in *A. herba alba* EO. α -thuyone (73.33%) and β -thuyone (11.96%) were detected as major compounds in *A. herba alba* EO. Finally, *C. citratus* was mainly constituted by oxygenated monoterpenes (92,51%), the results showed that Néral (37.06%) and Géranial (54.91%) were the two chemical compounds, taking part of the major compounds of *C. citratus* EO.

Antioxidant activities of studied essential oils were analyzed using free radical scavenging (DPPH and ABTS) and reducing power (FRAP) assays; the results are presented in Table 2. In the DPPH assay, the

radical scavenging activity of *A. judaica essential oil* was stronger than *P. scoparius essential oil* and followed by the *P. chloranthus essential oil* of Illizi, *P. chloranthus essential oil* of Ghardaïa, *A. herba alba*, *C. citratus*, *C. schoenanthus* of Ghardaïa and *C. schoenanthus* of Illizi. In the ABTS assay, the most potent radical scavenging activity was also found in *A. judaica*, which was followed by *P. chloranthus essential oil* of Ghardaïa. The results obtained by the method of FRAP confirm the important antioxidant potential of *A. judaica essential oil*. Antioxidant activities of essential oils from aromatic plants are mainly attributed to the active compounds present in them. This can be due to the high percentage of main constituents, but also to the presence of other constituents in small quantities or to synergy among them (Politeo et al., 2006). Some study indicate that the antioxidative effectiveness of essential oil depends on the content of phenolic compounds and the reaction activity of the phenol towards the chain-carrying peroxy radicals and on the stability of the phenoxyl radical formed in the reaction (Ćavar et al., 2012). The antioxidant activity may be ascribed to the presence of the several chemical components. Monoterpenes found in these essential oils may act as radical scavenging agents. It seems to be a general trend that the essential oils, which contain monoterpene hydrocarbons, oxygenated monoterpenes and/or sesquiterpenes, have high antioxidative properties (Khadri et al., 2008). The statistical analyzes allowed us to highlight a possible correlation between the antioxidant activity and the chemical composition of the essential oils. Through these results of Table 3, we can be noted that there are a strong correlations between the antioxidant capacity of an EOs and cyclic ether sesquiterpene compounds, furane ring compound and Alkenes and esters non-terpenics with correlation coefficients 0.840, 0.45, 0.859 (in ABTS assay) respectively. This result suggests that the antioxidant capacity of essential oils is not related to the presence of some phenolic compounds or to the major compounds, it can be due to other minor ingredients in the EOs might contribute to bioactivity through synergistic effects.

REFERENCES

- [1] Budel, J.M., Wang, M., Raman, V., Zhao, J., Khan, S.I., Rehman, J.U., Techen, N., Tekwani, B., Monteiro, L. M., Heiden, G., Takeda, I. J. M., Farago, P.V., Khan, I. A., (2018). Essential Oils of Five Baccharis Species: Investigations on the Chemical Composition and Biological Activities, *Molecules*, 23, 2620; doi:10.3390/molecules 23102620.
- [2] Ćavar, S., Maksimović, M., Vidica, D., Parić A., (2012). Chemical composition and antioxidant and antimicrobial activity of essential oil of *Artemisia annua* L. from Bosnia. *Ind. Crops Prod.*, 37, 479-485.
- [3] Djeridane, A., Yousfi, M., Nadjemi, B., Boutassouna, D., Stocker, P., Vidal, N., (2006). Antioxidant activity of some algerian medicinal plants extracts containing phenolic compounds, *Food Chemistry*, 97(4), 654-660.
- [4] Gohari AR, Hajimehdipoor H , Saeidnia S, Ajani Y, Hadjiakhoondi A (2011). Antioxidant Activity of some Medicinal Species using FRAP Assay, *J. Med. Pla.*, 10(37), 55-60
- [5] Hellali, N., Hadj Mahammed ,M., Ramdane, F., Talli, A., (2016). Antimicrobial and antioxidant activities of *Cymbopogon schoenanthus* (L.) spreng. essential oil, growing in Illizi – Algeria. *Journal of Medicinal Plants Research*. 10(14), 188-194.
- [6] Hellali, N., Hadj mahammed, M., Masrouk, H., (2017). Physico-chemical Study and Evaluation of Antimicrobial and Antioxidant Activities of *Artemisia judaica* L. Essential Oil, Growing in Illizi, Algeria. *Asian Journal of Chemistry*, 29(1), 181-186.
- [7] Khadri, A., Serralheiro, M.L.M., Nogueira, J.M.F., Neffati, M., Smiti, S., Araujo, M.E.M., (2008). Antioxidant and antiacetylcholinesterase activities of essential oils from *Cymbopogon schoenanthus* L. Spreng. Determination of chemical composition by GC–mass spectrometry and ¹³C NMR. *Food Chem.*, 109, 630–637.
- [8] Politeo, O., Juki, M., and Milo, M., (2006). Chemical Composition and Antioxidant Activity of Essential Oils of Twelve Spice Plants, *CROATICA CHEMICA ACTA* , 79 (4), 545-552.

CHEMICAL COMPOSITION AND BIOLOGICAL ACTIVITIES OF THE ESSENTIAL OILS FROM TWO CHEMOTYPES OF *CYMOPOGON SCHOENANTHUS* (L.) SPRENG. GROWING IN SOUTHERN ALGERIA.

Naima Hellali^{1,2}, Mahfoud Hadj Mohammed², Mebarka Bouziane²

¹ Faculty of Sciences and Technology, Department of process engineering, Ghardaïa University, BP 455, 47000 Ghardaïa, Algeria.

² Biogeochemistry Laboratory in Desert Environments, Faculty of Science, Kasdi Merbah University, BP 511, 30000 Ouargla, Algeria.

*Corresponding Author. E-mail: hellali.naima@univ-ghardaia.dz

Abstract

The chemical composition of the essential oils of two populations of *Cymbopogon schoenanthus* collected in Ghardaïa and Illizi (Southern Algeria), were identified by (GC-MS), this study allowed us to identify the presence of two chemotypes of *C. schoenanthus* essential oil, the first one was characterized by the high content of piperitone, and the second group was characterized by other compounds. The screening of antimicrobial activity of oils was individually evaluated against representatives of gram-positive, gram-negative bacteria and fungi, using the agar diffusion method, whereas, the antioxidant activity of the essential oils was evaluated using the DPPH, ABTS and FRAP assay. The results of in vitro testing of the antimicrobial activities of *C. schoenanthus* essential oil of Ghardaïa region proved the existence of strong activity against all the tested strains, specially the growth of *E. faecium*, *S. aureus* and *S. agalactiae*. However, the same species growing in Illizi region shown less power in the inhibition of microorganisms. The essential oil obtained from *C. schoenanthus* growing in Illizi is markedly rich in non-phenolic constituents; due to this fact the essential oil is relatively weak antioxidant, which the same remark in the essential oil extracted from Ghardaïa.

Key words: *C. schoenanthus*, essential oil, chemotype, antimicrobial activity, antioxidant activity.

INTRODUCTION

Cymbopogon genus is a member of the family of Poaceae (Gramineae) which are herbs known worldwide for their high essential oil content. They are widely distributed across all continents where they are used for various purposes (Avoseh et al., 2015). *Cymbopogon schoenanthus* L. belongs to this family, it is widely distributed from the tropical region through North Africa and Asia. In Algeria it grows as wild plant in dry areas and valleys. It is widely spread in the Algerian Sahara. *C. schoenanthus* is an aromatic and medicinal plant rich in essential oil, this plant is widely used in traditional medicine in the sahara of Algeria specially in Tassili n'ajjer. The traditional use of this species was varied from one region to another, for example in Algeria, it was used as: aches, toothaches, myalgia, anuria in Ghardaïa area and as an strengthening in Beni Abbes, but it was used to cooling and to treat anuria in Ouargla (Maiza et al., 1993). In addition to this Hammiche et al., (2006) reported that this plant is used as traditional medicine to treat digestive diseases (flatulence, urinary decrease), analeptic drink for the new mother after childbirth, bad breath, gumboils and urinary incontinence. *In vitro* *C. schoenanthus* essential oil showed different activities for example Ketoh et al., 2004; Koba et al., 2004 were reported that this plant has sedative, digestive and aromatic properties, with a strong and characteristic aroma, with some reports on its insecticidal activity.

MATERIALS AND METHODS

Plant material: The aerial parts of *C. schoenanthus* were collected from weddi Cebceb in Ghardaïa (Algeria), on April, 2014.

Essential oil extraction: The fresh aerial parts of *C. schoenanthus* were subjected to hydrodistillation in a modified Clevenger-type apparatus for 3 h. The essential oil obtained was dried over anhydrous sodium sulfate and stored in sealed glass vials at 4 to 6°C prior to analyses.

Essential oil chromatographic analysis: 0.2 µL of sample was injected on a gas chromatography (Hewlett-packar computerized system, Agilent model 6890 GC coupled to a Agilent 5973 N mass selective detector , and equipped with an Agilent technologies capillary HP-5MS column (30 m, 0,25 mm I.d, 0.25µm thickness), a split/splitless injector used in the split mode (200:1), using Helium (20 mL/min). The initial temperature of the column was 40°C which was heated gradually to 325°C with a 2°C /min. High purity helium was used as carrier gas at 36 cm/s. The quadrupole, source and transfer line temperatures were maintained at 150, 230 and 280°C, respectively. A solvent delay of 3 min. Identification of components was assigned by matching their mass spectra of peaks with those obtained from authentic samples.

Antioxidant activity determinations: The difference in their chemical composition of extracted *C. schoenanthus* essential oil between the two regions (Ghardaïa and Illizi), urged us to study the antioxidant activity of *C. schoenanthus* essential oil again in the same conditions as cited by Hellali et al. (2016), in order to evaluate and compare the capacity antioxidant of this herb. Three methods have been applied for the antioxidant assessment of the *C. schoenanthus* essential oil in this study: the 1,1-diphenyl- 2-picrylhydrazyl radical (DPPH[·]) assay (Brand-Williams et al.,1995), the 2,20-azino-bis(3-ethylbenzothiazoline-6-sulphonate) radical cation (ABTS^{·+}) assay (Miller et al.,1993) and the ferric reducing antioxidant power (FRAP) assay (Benzie and Strain, 1996). All measurements were performed in duplicate.

Antimicrobial screening: The antimicrobial activities were determined by using the drop agar diffusion method (Lopes et al., 2008). The microorganisms tested were the fungi *Candida albicans* ATCC 10231, and the bacteria *Escherichia coli* ATCC 8739, *Salmonella typhimurium* ATCC 14028, *Staphylococcus aureus* ATCC 6538, *Enterococcus faecium* ATCC 19434, and *Streptococcus B* (*Streptococcus agalactiae*).

RESULTS

Chemical composition of essential oil: The aerial parts of *C. schoenanthus* essential oils from Ghardaïa area, were analyzed by GC–MS. Fifteen compounds were identified showing high amounts in monoterpenes (84,18%), with mainly oxygenated monoterpene (53,76%) (Table 1). The main components identified were cis-p-menth-2-en-1-ol (29.21%), trans-p-menth-2-en-1-ol (15.62 %), 4-Carene (19.76%), cis piperitol (5.81%) and trans piperitol (5.05 %).

The results completed by literature search, the comparison show a significant difference in the content of the volatile oil of the aerial parts of this species collected in Illizi area which characterized by the presence remarkable of piperiton in their composition (Hellali et al., 2016).

Antioxidant activities: In this piper we report the antioxidant activity of *C. schoenanthus* essential oil growing in Ghardaïa area (Table 2), In our previous studies, the antioxidant investigation of the *C. schoenanthus* essential oil from Illizi region, were previously evaluated using various antioxidant tests (DPPH, ABTS and FRAP assay).

The results of Table 2 indicated that the *C. schoenanthus* essential oil from Ghardaïa region strongly transformed the ABTS radical into its reduced form. It has an IC₅₀ value of 15.16 mg/ml. These measurements are compared with our previously results, its showed that the essential oil obtained from *C. schoenanthus* growing in Illizi is markedly rich in non-phenolic constituents, due to this fact the essential oil is relatively weak antioxidant, which the same remark in the essential oil extracted from Ghardaïa.

Table 1. Chemical composition of essential oil of *C. schoenanthus*

N	Tr	Component	Percentage (%)
1	19.76	4-Carene	19.76
2	13.33	α -Phellandrene	1.48
3	14.78	Limonene	7.39
4	15.60	trans- β -Ocimene	1.79
5	19.62	α -Thujone	1.41
6	20.91	cis-p-menth-2-en-1-ol	29.21
7	22.10	trans-p-menth-2-en-1-ol	15.62
8	25.43	α -terpineol	1,71
9	25.70	cis piperitol	5.81
10	26.65	trans piperitol	5.05
11	47.67	Elemol	3.48
12	51.39	Selina-6-en-4-ol	1.82
13	53.11	β -eudesmol	1.39
14	53.47	Globulol	1.89
15	55.56	2-Isopropenyl-4a,8-dimethyl-1,2,3,4,4a,5,6,7-octahydronaphthalene	2.19
-	-	monoterpenes hydrocarbons	30,42
-	-	Oxygenated monoterpenes	53,76
-	-	sesquiterpenes hydrocarbons	2,19
-	-	Oxygenated sesquiterpenes	13,63

Table 2. Antioxidant activity of *C. schoenanthus* essential oil expressed in TEAC and IC50.

Variables	DPPH	ABTS	FRAP
TEAC (μ mol/g)	2.105	14.577	2.94 \pm 0.6
IC ₅₀ (mg/mL)	21.68 \pm 5.1	15.16 \pm 4.42	-

Averages \pm Standard Deviation were obtained from two different experiments.

Antimicrobial activity: The results of *in vitro* testing of the antimicrobial activities of *C. schoenanthus* essential oil from Ghardaïa region are presented in the Table 3, we used antibiotic ampicillin and antifungal nystatin as positives probe. The qualitative test was to determine the presence or absence of inhibition zones around the disks.

The results obtained proved the existence of strong activity against all the tested strains, specially the growth of *E. faecium*, *S. aureus* and *S. agalactiae*. The results obtained with same species growing in Illizi region (Hellali et al., 2016) shown less power in the inhibition of microorganisms, and that antimicrobial activities have been mainly explained through the high proportion of piperitone.

Table 3. Antimicrobial activity of *C. schoenanthus* essential oil.

Name of bacterial strain	Inhibition diameter (mm)*		
	Ampicillin/ Nystatin 10µg/100µg	<i>C.</i> <i>schoenanthus</i> <i>Of Ghardaïa</i>	<i>C. schoenanthus</i> <i>Of Illizi</i> (Hellali et al., 2016)
<i>Escherichia coli</i> ATCC 8739	12±1	25±1.4	15±1.4
<i>Salmonella typhimurium</i> ATCC 14028	17,33±1.5	16,5±0.7	10.5 ±0.7
<i>Staphylococcus aureus</i> ATCC 6538	44.3±0.5	34±2.8	19.5 ±0.7
<i>Enterococcus faecium</i> ATCC 19434	45.3±1.5	41±1.4	21±1.4
<i>Streptococcus B (Streptococcus agalactiae)</i>	34.3±0.5	31±1.4	12.75 ±0.3
<i>Candida albicans</i> ATCC 10231	42±1	21±1.4	12±1.4

*Including disc diameter of 6 mm, Averages ± Standard Deviation were obtained from two different experiments.

DISCUSSION AND CONCLUSION

To the best of the study knowledge, this is the first study providing data on chemical composition of the essential oil of *C. schoenanthus* from Ghardaïa especially and evaluating the influence of the climatic and geographic condition on their chemical composition. As a conclusion we can say that according to the chemical composition and geographical origin, two chemotypes were defined for *C. schoenanthus* for the first time in southern Algeria: the type of Illizi area (piperitone) and the type of Ghardaïa area (cis-p-menth-2-en-1-ol, trans-p-menth-2-en-1-ol, cis piperitol and trans piperitol).

The essential oil of Ghardaïa showed an activity antioxidant more important than the Illizi origin. We can explain this result by the ability of some terpenes (such as the monoterpene alcohols) to give up a hydrogen atom or more to the free radicals, but on the other hand, these compounds give another compounds are more stable. For example: we can obtain the piperitone by the dehydrogenation of trans or cis piperitol. But generally, *C. schoenanthus* essential oil can be considered as a relatively weak antioxidant. The antibacterial of this oil was studied *in vitro* on five bacterial strains and one fungal strain, the results obtained proved the existence of strong activity against all the tested strains. *C. schoenanthus* essential oil of Ghardaïa region was characterized by the presence of cis-p-menth-2-en-1-ol, trans-p-menth-2-en-1-ol, cis piperitol and trans piperitol, these components are alcohols monoterpene. *C. schoenanthus* essential oil of Illizi region (Hellali et al., 2016) shown an important activity antimicrobienne due to the presence of high proportion of monoterpenes specially the high content of piperitone, and we supposed that the piperitone, cis-p-menth-2-en-1-ol, trans-p-menth-2-en-1-ol, cis piperitol and trans piperitol, all these compounds have the same cyclic intermediate in the biosynthesis in *C. schoenanthus*, and the presence of hydroxyl function in place of the ketone function increases the inhibition efficiency of the essential oil of *C. schoenanthus* collected from Ghardaïa region Against the microorganisms tested.

REFERENCES

- [1] Avoseh, O., Oyedeji, O., Rungqu, P., Nkeh-Chungag, B., and Oyedeji, A., (2015). *Cymbopogon* Species; Ethnopharmacology, Phytochemistry and the Pharmacological Importance. *Molecules*. 20, 7438-7453.
- [2] Benzie, I.F.F., and Strain, J.J., (1996). The ferric reducing ability of plasma (FRAP) as a measure of "antioxidant power": the FRAP assay. *Anal. Biochem.* 239(1),70-76.
- [3] Brand-Williams, W., Cuvelier, M.E., and Berset, C., (1995). Use of a free radical method to evaluate antioxidant activity. *LWT-Food Sci. Technol.* 28, 25-30.
- [4] Hammiche, V., and Maiza, K., (2006). Traditional medicine in Central Sahara: Pharmacopoeia of Tassili N'ajjer. *J. Ethnopharmacol.* 105, 358-367.
- [5] Hellali, N., Hadj Mohammed, M., Ramdane, F., and Talli, A., (2016). Antimicrobial and antioxidant activities of *Cymbopogon schoenanthus* (L.) spreng. essential oil, growing in Illizi – Algeria. *J. Med. Pla. Res.* 10(14), 188-194.
- [6] Ketoh, G.K., Koumaglo, H.K., and Glitho, I.A., (2005). Inhibition of *Callosobruchus maculatus* (F.) (Coleoptera: Bruchidae) development with essential oil extracted from *Cymbopogon schoenanthus* L. Spreng (Poaceae), and the wasp *Dinarmus basalis* (Rondani) (Hymenoptera: Pteromalidae). *J. Stored Prod. Res.* 41, 363-371.
- [7] Ketoh, G.K., Koumaglo, H.K., Glitho, I.A., and Huignard, J., (2006). Comparative effects of *Cymbopogon schoenanthus* essential oil and piperitone on *Callosobruchus maculatus* development. *Fitoterapia*. 77, 506-510.
- [8] Koba, K., Sanda, K., Raynaud, C., Nononene, Y.A., Millet, J., and Chaumont, J., (2004). Activités antimicrobienne de trois *Cymbopogon* sp. Africains vis-à-vis de germes pathogènes d'animaux de compagnie. *Ann. Med. Vet.* 148, 202-206.
- [9] Lopes-Lutz, D., Alviano, D.S., Alviano, C.S., and Kolodziejczyk, P.P., (2008). Screening of chemical composition, antimicrobial and antioxidant activities of *Artemisia* essential oils. *Phytochemistry*. 69(8), 1732-1738.
- [10] Maiza, K., Brac De La Perriere, R. A., and Hammiche, V., (1993). Pharmacopée traditionnelle saharienne : Sahara septentriona, Médicaments et aliments : l'approche ethnopharmacologique, Actes du 2e colloque européen d'Ethnopharmacologie et de la 11e conférence internationale d'Ethnomédecine, Heidelberg, 24-27 mars 1993, 169-171.
- [11] Miller, N.J., Rice-Evans, C., Davies, M.J., Gopinathan, V., and Milner, A., (1993). A novel method for measuring antioxidant capacity and its application to monitoring the antioxidant status in premature neonates. *Clinical Science*, 84(4), 407-412.

ASSESSMENT OF CHOLINESTERASE AND TYROSINASE ENZYMES INHIBITORY, PHYTOCHEMICAL COMPOSITION AND ANTIOXIDANT PROPERTIES OF *MENTHA SPICATA* L.

Sevgi Gezici^{a,*}, Murat Tuncurk^b, Lutfi Nohutcu^b, Nazim Sekeroglu^c

^a Department of Molecular Biology and Genetics, Faculty of Science and Literature; Advanced Technology Application and Research Center, Kilis 7 Aralik University, 79000 Kilis, Turkey,

E-mail: drsevgigezici@gmail.com, sevgigezici@kilis.edu.tr

^b Department of Field Crops, Faculty of Agricultural Engineering, Yuzuncu Yil University, 65080 Van, Turkey,

E-mail: murattuncurk@hotmail.com

^c Department of Food Engineering, Faculty of Engineering and Architecture; Advanced Technology Application and Research Center, Kilis 7 Aralik University, 79000 Kilis, Turkey,


E-mail: nsekeroglu@gmail.com, sekeroglu@kilis.edu.tr

Abstract

Because oxidative damage is one of the major factor contributing to neurodegeneration, combination of antioxidant activity and enzyme inhibition potential of the herbal extracts is closely related to combat neurodegenerative diseases and disorders efficiently. Spearmint (*Mentha spicata* L., Lamiaceae), an edible aromatic plant, are of important food sources used as flavoring, tea infusions and spicing in the world. Besides its importance in food, beverage, cosmetic and confectionary industries, there has been huge ethnobotanical information on the curative properties of *M. spicata* L. against common diseases. Even though biological properties of spearmint have been documented by previous scientific reports, one less known biological activity of spearmint is its neuroprotective activity. **Material and Methods:** In this concern, neuroprotective potentials, antioxidant capacities, mineral contents and total polyphenolic compositions of aqueous and methanol extracts of the leaves from *M. spicata* L. were aimed to examine in the current research. Neuroprotective potentials of the extracts were determined against inhibition of AChE, BChE, and tyrosinase enzymes, which are associated with pathogenesis of Alzheimer's disease. *In vitro* assays including DPPH, ABTS, FRAP, and CUPRAC were carried out to reveal antioxidant capacities of the extracts. The total phenolic and total flavonoid contents were analyzed spectrophotometrically. Mineral contents (K, Ca, Mg, Fe, Zn, Cu, Mn, Cd, and Pb) in spearmint leaves were also determined by Atomic Absorption Spectrometry. **Results:** A wide variation was observed in the mineral contents of the plant that Fe was found the major mineral with the values of 1471.41±12.28 mg/kg, whilst Cu and Pb were determined as minor minerals. With regard to total polyphenolic components, the methanol extract possessed higher phenols and flavonoids contents (3.158±0.52 mg/g extract as GAE and 1.31±0.08 mg/g extract as QE, respectively). A significant correlation was found between total polyphenolic compositions and antioxidant capacities of the spearmint leaves. Correlation analysis indicated that the methanol extract, which possessed higher total phenols and flavonoids components than the water extract, was also found the more efficient in terms of antioxidant capacities and neuroprotective potentials. Both of the extracts demonstrated powerful inhibition on BChE, whereas they showed moderate inhibition on AChE and tyrosinase enzymes with the values ranging from 24.06±0.12% to 57.68±0.18%. **Conclusions:** Consequently, our results indicate that *M. spicata* L., a natural antioxidant and neuroprotective agent, can provide considerable benefits against oxidative stress-related diseases.

Keywords: *Mentha spicata* L.; Alzheimer's Disease; neuroprotection; antioxidant; polyphenolic; mineral content

Graphical Abstract



1. INTRODUCTION

Spearmint (*Mentha spicata* L., Lamiaceae), an edible aromatic plant, are of important food sources used as flavoring, tea infusions and spicing in the world. The plant is indigenous to northern England and is cultivated in areas with climate ranging from tropical to temperate, such as America, Europe, China, South Africa and Brazil. Nowadays, spearmint is widely grown throughout all regions in the world such as America, Europe, China, South Africa and Brazil. Since ancient times, both fresh and dried spearmint plants have been practiced as medical and aromatic plants in western and eastern cultures (Fitsiou et al., 2016). Spearmint has been used broadly in cosmetic and soap, as well as toothpaste, breath freshener and antiseptic mouth rinse. It is widely used in food preservation and imparts food taste and aroma, confectionery, and chewing gum industries. Additionally, it has also been used in the treatment of several ailments in Turkish folk medicine. In term of medical uses, spearmint is considered as an herbal medicine in folkloric remedies for treating of colds and flu, respiratory tract problems, gastralgia, hemorrhoids, and stomachache (Brahmi et al., 2017; Kee et al., 2017).

Because oxidative damage is one of the major factor contributing to neurodegeneration, combination of antioxidant activity and enzyme inhibition potential of the herbal extracts is closely related to combat neurodegenerative diseases and disorders efficiently. The formation of Reactive Oxygen Species (ROS), such as superoxide ion (O_2^-), hydroxyl radical (OH) and Hydrogen peroxide (H_2O_2), have often been reported to induce DNA damage, protein carboxylation, and lipid peroxidation, causing a variety of chronic health disturbances and diseases, including cancer, ageing, Parkinson's disease, Alzheimer's disease, and cardiovascular diseases. Recent research indicates that several herbal plants can offer alternative sources of dietary ingredients to promote human health and might open promising opportunities for the treatment of a wide range of troublesome diseases and infections (Sun et al., 2016; Sahoo et al., 2018). Although, there are so many methods for combatting neurodegenerative diseases and disorders, they cannot always provide effective treatments and mediations. Hence, an extensive research on developing new treatment strategies against these disorders are still needed to cure nowadays. Since ancient times, natural products (NPs), originated from natural sources such as plants, have been used for the cure and treatment of many diseases in Anatolian folk medicine 'herbal therapies' (Gezici and Sekeroglu, 2019a; Awasthi et al., 2016; Godyń, et al, 2016).

So far, we have investigated various of medicinal plants using in vitro antioxidant, anticancer, antiproliferative, anticholinesterase, etc. experiments, which aimed to contribute to the finding of new herbal products for prevention and treatment of cancer and neurodegenerative diseases (Gezici, 2019; Gezici and Sekeroglu, 2019a; Gezici and Sekeroglu, 2019b; Sekeroglu et al., 2018a; Sekeroglu et al., 2018b; Gezici, 2018; Senol et al., 2018; Gundogdu et al., 2018; Karik et al., 2018; Belkhodja et al.,

2017; Gezici et al., 2017; Sekeroglu et al., 2017; Akgunlu et al., 2016; Orhan et al., 2013; Sekeroglu et al., 2012; Orhan et al., 2012, etc.). Taking our previous researches on medicinal plants and plant-derived natural products, the current study was undertaken to examine total polyphenolic and mineral contents, *in vitro* antioxidant capacities, neuroprotective and enzyme inhibitory activities of the extracts from leaves of *Mentha spicata* L. Besides its importance in food, beverage, cosmetic and confectionary industries, there has been huge ethnobotanical information on the curative properties of *M. spicata* L. against common diseases. As far as our knowledge, no detailed studies on combined biological activities have been performed with extracts spearmint growing under special conditions. Hence, this study may help us to find new potential sources as natural antioxidant and neuroprotective agents.

2. MATERIAL AND METHODS

2.1. Collection of Plant Material: The plant was growing by Prof. Dr. Murat Tuncturk at Medicinal and Aromatic Plants Garden, Van Yuzuncu Yil University, and the voucher specimen was deposited in the herbarium Van Yuzuncu Yil University, Van-Turkey. Taxonomic classification of the plant was given in the Table 1.

Table 1. Taxonomic classification of the plant and plant sample in the Medicinal and Aromatic Plants Garden

Kingdom:	<i>Plantae</i>
Subkingdom:	<i>Tracheobionta</i>
Division:	<i>Magnoliophyta</i>
Class:	<i>Magnoliopsida</i>
Subclass:	<i>Asteridae</i>
Order:	<i>Lamiales</i>
Family:	<i>Lamiaceae</i>
Genus:	<i>Mentha</i>
Species:	<i>spicata</i> L.



2.2. Extract Preparation: To prepare crude extracts, air dried samples (50 g) of the aerial parts of the plant were individually extracted with 250 ml methanol (70%) and distilled water for 2 days at the room temperature, as described in our previous research (Gezici and Sekeroglu, 2019b). The extracts yields (w/w%) are given in the Figure 1. Extraction yields of the methanol and water extracts of the aerial parts of the plant were determined as 21.370% and 11.517% (w/w), respectively.




Fig. 1. Extraction yield (w/w) of the plant

Determination of Total Polyphenolic Contents: Phenolic compounds in total were determined in accordance with slightly modified Folin-Ciocalteu's method (Singleton and Rossi, 1965; Gezici and Sekeroglu, 2019). Absorption was measured at 760 nm at a using a 96-well microplate reader (VersaMax Molecular Devices, USA). Total flavonoid content of the extracts was calculated by aluminum chloride colorimetric method (Woisky and Salatino, 1998; Gezici and Sekeroglu, 2019b). A number of dilutions of quercetin were obtained to prepare a calibration curve. Absorbance of the reaction mixtures was measured at wavelength of 415 nm with a using a 96-well microplate reader (VersaMax Molecular Devices, USA). The total phenol and flavonoid contents of the extracts were expressed as gallic acid and quercetin equivalents (mg g⁻¹ extract), respectively.


2.3. Determination of Mineral Contents: The presence of mineral elements such as potassium (K), calcium (Ca), magnesium (Mg), iron (Fe), zinc (Zn), copper (Cu), manganese (Mn), cadmium (Cd), and lead (Pb) in the aerial parts of the plant were determined by Atomic Absorption Spectrometry (AAS, Perkin-Elmer 2280) (Sekeroglu et al., 2017).

2.4. Antioxidant Activity Assays: Since oxidative damage is one of the major factors contributing to both cancer and neurodegeneration, *In vitro* methods including 2,2-diphenyl-1-picrylhydrazyl (DPPH), 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulphonic acid) (ABTS), ferric reducing antioxidant power (FRAP), and cupric ion reducing capacity (CUPRAC) were performed to reveal antioxidant activity of the extracts (Gezici et al., 2017; Sekeroglu et al., 2017; Gundogdu et al., 2018; Gezici and Sekeroglu, 2019b).

2.5. Enzyme Inhibition Assays: Neuroprotective activities of the extracts on AChE, BChE, and tyrosinase were evaluated in the current study. AChE and BChE inhibitory activity of the samples was measured by slightly modified spectrophotometric method of Ellman et al. (1961). Electric eel AChE (EC 3.1.1. Sigma, St. Louis, MO, USA) and horse serum BChE (EC 3.1.1. Sigma, St. Louis, 7 MO, USA) were used, while acetylthiocholine iodide and butyrylthiocholine chloride (Sigma, St. Louis, MO, USA) were employed as substrates of the reaction. 5,5'-Dithio-bis(2-nitrobenzoic)acid (DTNB, Sigma, St. Louis, MO, USA) was used for the measurement of the anticholinesterase activity. All reagents, conditions and calculations were same as described in our previous publication (Senol et al., 2018; Gezici and Sekeroglu, 2019b). The measurements and calculations were evaluated by using Softmax PRO 4.3.2.LS software. The experiments were done in quadruplicate. Galanthamine hydrobromide (Sigma, St. Louis, MO, USA) was used as the reference drug. Inhibition of tyrosinase (EC 1.14.1.8.1, 30 U, mushroom tyrosinase, Sigma) was determined using the modified dopachrome method with L-DOPA as substrate (Sekeroglu et al., 2012).

3. RESULTS AND DISCUSSION

Total polyphenolic compositions of the extracts were identified spectrophotometrically in the current research. With regard to total polyphenolic components, the methanol extract possessed higher phenols and flavonoids contents (3.158±0.52 mg/g extract as GAE and 1.31±0.08 mg/g extract as QE, respectively). Gallic acid and quercetin equivalent as commercial standards for total phenolic phenolic and flavonoid contents were shown in the Figure 2.






Fig. 2. Gallic acid and Quercetin Equivalent

K, Ca, Mg, Fe, Zn, Cu, Mn, Cd, Pb mineral contents of the plant were also determined in the presented research. A wide variation was observed in the mineral contents of the plant and the mineral values were given as mean±Sd in the Table 2.

Table 2. Mineral contents (mg/kg) of *M. spicata* L.

K (g/kg)	Ca (g/kg)	Mg (g/kg)	Fe (mg/kg)	Zn (mg/kg)
14,84±0,95	15,20±0,83	6,64±0,40	1471,31±12,28	19,53±1,08
Cu (mg/kg)	Mn (mg/kg)	Cd (mg/kg)	Pb (mg/kg)	
6,69±0,72	110,46±1,74	0,045±0,05	0,024±0,01	

Antioxidant activity of the extracts obtained from the aerial parts of the plant were evaluated by using radical scavenging against DPPH and ABTS radicals, and ion reducing antioxidant power on FRAP and CUPRAC. Regarding of antioxidant assays, both methanol and water extracts obtained from leaves of *M. spicata* L. exhibited excellent scavenging activities on DPPH, FRAP, ABTS, and CUPRAC, comparing the standard antioxidants. A significant correlation was found between total polyphenolic compositions and antioxidant capacities of the spearmint leaves. Correlation analysis indicated that the methanol extract, which possessed higher total phenols and flavonoids components than the water extract, was also found the more efficient in terms of antioxidant capacities and neuroprotective potentials.

Neuroprotective activity of the extracts was assessed through enzyme inhibition assays on cholinesterase enzymes. The spearmint leaves were extracted with water and methanol solvents, and subjected to enzyme inhibitory assays on acetylcholinesterase (AChE), butyrylcholinesterase (BChE), and tyrosinase which are closely related to pathogenesis of neurodegenerative disease. According to the results, both of the extracts demonstrated powerful inhibition on BChE, whereas they showed moderate inhibition on AChE and tyrosinase enzymes with the values ranging from 24.06±0.12% to 57.68±0.18%.

4. CONCLUSION

Consequently, a strong correlation was observed between the phytochemical and mineral constituents of the plant and its biological properties. In the light of the findings, our data indicate that *M. spicata* L., a natural antioxidant and neuroprotective agent, can provide considerable benefits against oxidative stress-related diseases. Moreover, it appears to be a natural source having promising inhibitory molecules, are worth to conduct further *in vivo* investigations that is under further investigation by our group.

Acknowledgments: The authors would like to thank Genetics Laboratory, Advanced Technology Application and Research Center, Kilis 7 Aralik University for technical support.

REFERENCES

- [4] Akgunlu, S., Sekeroglu, N., Koca-Caliskan, U., Ozkutlu, F., Ozcelik, B., Kulak, M., Gezici, S., 2016. Research on selected wild edible vegetables: Mineral content and antimicrobial potentials. *Ann. Phytomed.* 5(2), 50-57. <https://doi.org/10.21276/ap.2016.5.2.6>.
- [5] Awasthi, M., Singh, S., Pandey, V. P., Dwivedi, U. N. 2016. Alzheimer's disease: An overview of amyloid beta dependent pathogenesis and its therapeutic implications along with in silico approaches emphasizing the role of natural products. *Journal of the neurological sciences*, 361, 256-271. <https://doi.org/10.1016/j.jns.2016.01.008>.
- [6] Brahmi, F., Khodir, M., Mohamed, C., Pierre, D. 2017. Chemical composition and biological activities of Mentha species. *Aromatic and Medicinal Plants—Back to Nature*; InTech: London, England, 47-80. <https://doi.org/10.5772/67291>.
- [7] Fitsiou, E., Mitropoulou, G., Spyridopoulou, K., Tiptiri-Kourpeti, A., Vamvakias, M., Bardouki, H., Pappa, A. 2016. Phytochemical profile and evaluation of the biological activities of essential oils derived from the Greek aromatic plant species *Ocimum basilicum*, *Mentha spicata*, *Pimpinella anisum* and *Fortunella margarita*. *Molecules*, 21(8), 1069. <https://doi.org/10.3390/molecules21081069>.
- [8] Gezici, S. 2018. Promising anticancer activity of lavender (*Lavandula angustifolia* Mill.) essential oil through induction of both apoptosis and necrosis. *Annals of Phytomedicine*, 7(2), 38-45. <https://doi.org/10.21276/ap.2018.7.2.5>.
- [9] Gezici, S., 2019. Anticancer, Antiproliferative, Lysosomal and Lactate Dehydrogenase Inhibitory Effects of Fruit Extracts from Sumac (*Rhus coriaria* L.) on Human Lung Cancer Cells. *Acta Oncol Tur.* 2019; 52(1): 160-168. <https://doi.org/10.5505/aot.2019.09326>.
- [10] Gezici, S., Sekeroglu, N., 2019a. Neuroprotective potential and phytochemical composition of acorn fruits. *Ind. Crop. Prod.* 128, 13-17. <https://doi.org/10.1016/j.indcrop.2018.10.082>.
- [11] Gezici, S., Sekeroglu, N., 2019b. Current perspectives in the application of medicinal plants against cancer: novel therapeutic agents. *Anti-Cancer Agent Med. Chem.* <https://doi.org/10.2174/1871520619666181224121004>.
- [12] Gezici, S., Sekeroglu, N., Kijjoa, A., 2017. In vitro Anticancer Activity and Antioxidant Properties of Essential Oils from *Populus alba* L. and *Rosmarinus officinalis* L. from South Eastern Anatolia of Turkey. *Indian J. Pharm. Educ. Res.* 51(3), 498-503. <https://doi.org/10.5530/ijper.51.3s.74>.
- [13] Godyń, J., Jończyk, J., Panek, D., & Malawska, B. 2016. Therapeutic strategies for Alzheimer's disease in clinical trials. *Pharmacological Reports*, 68(1), 127-138. <https://doi.org/10.1016/j.pharep.2015.07.006>.
- [14] Gundogdu, M., Tunçtürk, M., Berk, S., Şekeroğlu, N., Gezici, S., 2018. Antioxidant Capacity and Bioactive Contents of Mulberry Species from Eastern Anatolia Region of Turkey. *Indian J. Pharm. Educ. Res.* 52(4), 96-101. <https://doi.org/10.5530/ijper.52.4s.82>.
- [15] Karik, U., Çınar, O., Tunçtürk, M., Sekeroglu, N., Gezici, S. 2018. Essential Oil Composition of Some Sage (*Salvia* spp.) Species Cultivated in İzmir (Turkey) Ecological Conditions. *Ind. J. Pharm. Educ. Res.* 52(4), 102-107. <https://doi.org/10.5530/ijper.52.4s.83>.
- [16] Kee, L. A., Shori, A. B., Baba, A. S. 2017. Bioactivity and health effects of *Mentha spicata*. *Integr Food Nutr Metab*, 5(1), 1-2. <https://doi.org/10.15761/IFNM.1000203>.
- [17] Orhan, I.E., Atasu, E., Senol, F.S., Ozturk, N., Demirci, B., Das, K., Sekeroglu, N., 2013. Comparative studies on Turkish and Indian *Centella asiatica* (L.) Urban (gotu kola) samples for their enzyme inhibitory and antioxidant effects and phytochemical characterization. *Ind. Crop. Prod.* 47, 316-322. <https://doi.org/10.1016/j.indcrop.2013.03.022>.
- [18] Orhan, I.E., Senol, F.S., Gulpinar, A.R., Sekeroglu, N., Kartal, M., Sener, B., 2012. Neuroprotective potential of some terebinth coffee brands and the unprocessed fruits of *Pistacia terebinthus* L. and their fatty and essential oil analyses. *Food Chem.* 130(4), 882-888. <https://doi.org/10.1016/j.foodchem.2011.07.11>.
- [19] Sahoo, A.K., Dandapat, J., Dash, U.C., Kanhar, S. 2018. Features and outcomes of drugs for combination therapy as multi-targets strategy to combat Alzheimer's disease. *J. Ethnopharmacol.* 215, 42-73. <https://doi.org/10.1016/j.jep.2017.12.015>.
- [20] Sekeroglu, N., Karaoglan, M., Gezici, S., Kulak, M., Ozkutlu, F., Kacar, O., Gul, F., 2018. Variation in the composition of the essential oils, hypericin and mineral elements in aerial parts, stem and flower of *Hypericum capitatum* (CHOISY) growing in Turkey with oxidative DNA damage protective activity. *J. Pharm. Res.* 17, 67-77. <https://doi.org/10.18579/jprckc/2018/17/2/123613>.
- [21] Sekeroglu, N., Senol, F.S., Orhan, I.E., Gulpinar, A.R., Kartal, M., Sener, B., 2012. In vitro prospective effects of various traditional herbal coffees consumed in Anatolia linked to neurodegeneration. *Food Res. Int.* 45, 197-203. <https://doi.org/10.1016/j.foodres.2011.10.0088>.
- [22] Sekeroglu, N., Urlu, E., Kulak, M., Gezici, S., Dang, R., 2017. Variation in Total Polyphenolic Contents, DNA Protective Potential and Antioxidant Capacity from Aqueous and Ethanol Extracts in Different Plant Parts of *Hypericum perforatum* L. *Indian J. Pharm. Educ. Res.* 51, 1-7. <https://doi.org/10.5530/ijper.51.2s.43>.
- [23] Senol, F. S., Sekeroglu, N., Gezici, S., Kilic, E., Orhan, I. E. 2018. Neuroprotective potential of the fruit (acorn) from *Quercus coccifera* L. *Turkish Journal of Agriculture and Forestry*, 42(2), 82-87. <https://doi.org/10.3906/tar-1711-18>.
- [24] Singleton, V. L., & Rossi, J. A. 1965. Colorimetry of total phenolics with phosphomolybdic-phosphotungstic acid reagents. *American Journal of Enology and Viticulture*, 16(3), 144-158.
- [25] Sun, J., Ren, X., Qi, W., Yuan, D., Simpkins, J.W. 2016. Geissoschizine methyl ether protects oxidative stress-mediated cytotoxicity in neurons through the 'Neuronal Warburg Effect'. *J. Ethnopharmacol.* 187, 249-258. <https://doi.org/10.1016/j.jep.2016.04.034>.
- [26] Woisky, R. G., & Salatino, A. 1998. Analysis of propolis: some parameters and procedures for chemical quality control. *Journal of apicultural research*, 37(2), 99-105. <https://doi.org/10.1080/00218839.1998.11100961>.

DETERMINATION OF VOLATILE AND SEMI-VOLATILE OIL CONTENT OF *ISATIS TINCTORIA* L. VIA SPME-GC-MS

Meryem Bozkurt*, Ela Nur Şimşek Sezer, Nihal İlhan, Tuna Uysal

Department of Biology, Science Faculty, Selçuk University, Konya, Turkey,

**E-mail address of the corresponding author: mbozkurt@selcuk.edu.tr*

Abstract

Isatis tinctoria L. is a species of the genus Brassicaceae and generally distributed in Central Asia. The plant is also used for medical purposes as well as being used as a colorant. The aim of this study is to determine and characterize the volatile oil components found in the leaves and flowers of *Isatis tinctoria* via the SPME-GC-MS method. The plant material was collected from the Ardıçlı Village (Konya Province) and preserved under suitable conditions. 65 µm PDMS/DVB and 85 µm polyacrylate fibers were used to determine the volatile and semi volatile oil composition of the samples. Analyzes were performed using the Shimadzu QP2010 ULTRA FID GC-MS system. According to the data obtained from the analysis of *Isatis tinctoria*, in leaf content 148 compounds and in flower content 156 compounds were detected respectively. According to the analysis, with 65 µm PDMS/DVB fiber, the most amount of components in the leaf Benzeneethanol (4.67%) and 5-Methylundecane (3.93%), the most common components in flower, 4-Octadecylmorpholine (18.68%), 4-Amino-N-(3-morpholin-4-ylpropyl)-1,2,5-oxadiazole-3-carboxamide (11.91%). With 85 µm polyacrylate fiber, the most amount of components in the leaves, benzene, 2,4-diisocyanato-1-methyl (33.41%) and pyrrole-3-carbonitrile, 5-formyl-2,4-dimethyl (14.47%), in the flower content 2,2,3,3,4,4,4-Heptafluoro-N-(2-morpholin-4-yl-ethyl)-butyramide (19.67%) and 4-Octadecylmorpholine (18.07%), respectively. This study is the first study in terms of characterization and comparison of volatile and semi volatile oil components by SPME-GC-MS analysis from leaves and flowers of *Isatis tinctoria* L. species from Turkey.

Keywords: Çivitotu, solid phase micro-extraction, Turkey.

INTRODUCTION

One of the rare colors used as dyestuff in natural dyeing is blue. *Isatis tinctoria* L. (çivit otu, Brassicaceae) is one of the primary sources of blue color in nature. Also, it is recommended for the treatment of wounds, ulcers and tumors, hemorrhoids, snake bites and various inflammatory diseases. For this reason, *I. tinctoria* has a long and well documented history as an indigo dye plant and a medicinal plant (Mohn, 2009). In recent years, a large number of phytochemical, biological and pharmacological studies have been conducted. More than 100 secondary metabolites have been described in *I. tinctoria*. These are like isatin, tryptanthrin, deoxyvasicinone, isaindigotone, isaindigotidione, quinazolines, indolinone, benzodiazepine, the indigo dyes indigo and indirubin, or their natural precursors. *I. tinctoria*, a rich source for glucosinolates represented by glucoraphanin, progoitrine and glucobrassicine compounds, has been reported to contain compounds such as aromatic and aliphatic carboxylic acids (Honda et al., 1980; Goetz and Schraudolf, 1983; Lockwood and Belkhiri, 1991; Hartleb and Seifert, 1994, 1995; Wu et al., 1997a,b; Zhu, 1998; Frécharde et al., 2001).

The species was used as a medical plant in both Europe and East Asia. However, Hamburger (2002) reported that *Isatis tinctoria* is a plant that has been neglected in Europe today but could be returned to the European phytotherapy list with continue systematic investigations. Heinemann (2004) and Recio et al. (2006a,b) reported that lipophilic *Isatis tinctoria* extracts have anti-inflammatory activity in skin erythema, acute and chronic inflammation, contact allergy and rheumatoid arthritis. Although several approaches can be pursued in the assessment of a plant's pharmacological potential, depending on the preparation mode, the composition of an extract can vary greatly. Therefore, in the evaluation of the volatile and semi-volatile profile of *Isatis*, we directly analyzed the flowers and leaves of plant pulverized by drying with SPME-GC-MS.

Solid phase micro extraction (SPME) is a simple, fast and economical extraction method that has gained popularity in foods and especially in the determination of volatile aroma profile in recent years. SPME has been successfully used to identify aroma compounds with different functional groups (Koufogianni, 2006). The aim of this study is to determine the volatile content of wild *Isatis tinctoria* and to put the economic importance of the identified components for this species of our town.

MATERIAL AND METHODS

Material: The *Isatis tinctoria* samples were collected from its natural habitat. The powdered samples (3gr) sealed in a 10 ml vial. The SPME fibres preferred for analysis were 65 μm PDMS/DVB and 85 μm polyacrylate. The SPME apparatus was directly injected into the upper space of the vial to adsorb volatile substances and then directly injected into the Shimadzu QP2010 ULTRA GC-MS apparatus using a Restek Rxi-5 MS capillary column.

GC-MS Analysis: The volatile components separated for 15 min by inserting the SPME fibre into a GC injector (the injector temperature was 250 °C). The extracts from SPME procedure analysed on a Shimadzu QP2010 ULTRA FID GC-MS system. A Restek Rxi-5 MS column (30 m \times 0.25 mm id, film thickness 0.25 μm) used with helium as a carrier gas. The GC oven temperature was programmed to hold at 40 °C for 3 min and then to increase to 240 °C at 5 °C/min, finally holding at 240 °C for 3 min. The column flow rate was 1.8 mL/min. The ion source temperature was 200 °C, and the interface temperature set at 250 °C.

Evaluation of data: After the GC-MS analysis, each sample was demonstrated with GC-MS chromatograms. Chromatograms of all samples were subjected to noise reduction prior to peak area integration, and then the peak areas of compounds in the chromatogram were integrated. Compounds were identified by comparison with three libraries, W9N11, SWGDR4G4 and SWGDR4G5. Identification of components in the sample was based on the retention time (RT). The relative percentage of the volatile oil constituents was calculated from peak areas. All analyses were carried out three times.

RESULT AND DISCUSSION

Totally, 148 component identified from the leaves and 156 compound identified from the flowers during our analyses. As a result of all analyzes with flowers and leaves, some common components were detected. These are 1,3-Cyclohexadiene-1-carboxaldehyde, 2,6,6-trimethyl; Dimethyl sulfone; 5,9-Undecadien-2-one, 6,10-dimethyl; 3-Methylbutanoic acid; 2-Methylbutanoic acid; Benzaldehyde; 3,5-Octadien-2-One; Naphthalene; Tetradecane; Benzene, 2,4-diisocyanato-1-methyl; Dihydroactinidioides; 4-Octadecylmorpholine; Dimethyltetradecylamine, Phytone, 2-Ethylhexanol and Phytane (Table 1). With 85 μm polyacrylate fiber, the Benzen, 2,4-diizosiyano-1-metil compound is 33.41% on the leaves and 10.81% on the flower. With 65 μm PDMS/DVB fiber, the Benzen, 2,4-diizosiyano-1-metil compound is 0.81% on the leaves and 0.31% on the flower. Allyl isothio cyanate, which is an essential component in the leaves of species, was identified by 65 μm PDMS/DVB fiber. These compounds are responsible for preventing the proliferation of cancer cells and promoting apoptosis or programmed cell death of tumor cells. The compounds such as isothiocyanate and D-limonene were reported to be related to cancer. Isothiocyanates are characteristic flavour compounds in *Brassica*, and the cancer chemoprotective attributes are recently responsible for their growing interest (Condursa et al., 2006; Hashimoto et al., 2019).

The other common compound is 4-Octadecylmorpholine. This compound is the highest in the flower of species, and it has been previously reported to possess antibacterial and antifungal potential. 4-Sec-Butyl-2,6-di-tert-butylphenol, an industrial chemical, is known as DTBSBP. DTBSBP may be used as an antioxidant in mineral/vegetable oils to prevent oil thickening and the oxidation process, which can lead to sediment formation. This compound has a value of 4.18% (65 μm PDMS/DVB) and 5.17% (85 μm polyacrylate) in the flowers of the species and 0.27% in the leaves (85 μm polyacrylate fiber). Parallel to these pharmacological investigations, advanced phytochemical and analytical studies of the plant have been carried out (Hamburger 2002). Condurso et al. (2006) were reported that leaf volatile in *I. tinctoria* were analyzed by 50/30 mm DVB/CAR/PDMS fiber of SPME/GC-MS, and they were characterized by acids, alcohols, aldehydes, ethers, furans, hydrocarbons, isothiocyanates, ketons,

nitriles, terpenes and sesquiterpens. Furthermore, these components were detected by Verzera et al. (2010). These data are consistent with our results.

Table 1. Volatile oil and semi-volatile oil of flowers and leaves from *Isatis tinctoria*

Flowers 65 µm PDMS/DVB Volatile compounds	Area %	Flowers 85 µm polyacrylate Volatile compounds	Area %	Leaves 65 µm PDMS/DVB Volatile compounds	Area %	Leaves 85 µm polyacrylate Volatile compounds	Area %
Hexanal	0.24	3-Methylbutanoic acid	0.06	Butyric acid	0.18	3-Methylbutanoic acid	0.58
Hexamethylcyclotrisiloxane	0.94	2-Methylbutanoic acid	0.33	Hexanal	0.61	2-Methylbutanoic acid	1.29
3-Methylbutanoic acid	0.52	Oxime-, methoxy-phenyl	0.13	4-Methyl-1-pentanol	1.21	Valeric acid	0.24
2-Hexenal	0.20	Dimethyl sulfone	0.04	3-Methylbutanoic acid	1.17	Dimethyl sulfone	0.11
3-Hexen-1-ol	0.17	Benzaldehyde	0.08	2-Hexenal	0.67	Benzaldehyde	0.25
Ethylbenzene	0.55	Glycerin	0.04	3-Hexen-1-ol	1.12	Glycerol	0.07
2-Methylbutanoic acid	0.25	Hexanoic acid	0.98	2-Methylbutanoic acid	2.06	Hexanoic acid	0.55
p-Xylene	0.25	Benzene, 1,3-dichloro	0.07	4-Methyloctane	0.43	Benzene, 1,3-dichloro	0.20
Styrene	0.29	2-Ethylhexanol	0.40	Allyl isothiocyanate	0.12	4-Methyldecane	0.14
Xylose	0.30	Benzyl alcohol	0.28	Valeric acid	0.56	2-Ethylhexanol	0.16
4-Heptenal	0.15	Acetophenone	0.07	Dimethyl sulfone	0.33	1,5-Heptadien-4-one, 3,3,6-trimethyl	0.57
Oxime-, methoxy-phenyl	1.13	3,5-Octadien-2-one	0.12	Methyl hexanoate	1.29	Tridecane	0.36
Dimethyl sulfone	0.18	Nonanal	0.05	alpha-Pinene	0.11	3,5-Octadien-2-One	0.40
Benzaldehyde	0.79	3-Acetyl-2,6-heptanedione	0.04	4-Methylpentanoic acid	0.11	2,6-Dimethylcyclohexanol	0.31
Nonane, 2-methyl	0.33	Benzeneethanol	0.95	Benzaldehyde	1.08	Propan-2-yl tetradecyl sulfite	0.04
Heptanoic acid	1.07	Naphthalene	0.17	4-Methylnonane	0.26	5-Butylnonane	0.03
2-Pentylfuran	0.28	1-Tridecene	0.34	Nonane, 2-methyl	0.68	3-Methylpyrazole	0.07
Decane	0.35	1,3-Cyclohexadiene-1-carboxaldehyde, 2,6,6-trimethyl	0.09	Nonane, 2,5-dimethyl	1.8	2,4,4-Trimethylhexane	0.05
Octanal	0.12	2-Ethoxyethyl acrylate	0.25	octenol	0.12	Naphthalene	0.29
Cyclotetrasiloxane, octamethyl	0.74	Anethole	0.10	6-Methyl-5-hepten-2-one	1.20	1,3-Cyclohexadiene-1-carboxaldehyde, 2,6,6-trimethyl	0.10
1,4-Dichlorobenzene	0.83	Decanoic acid, methyl ester	0.21	Butyl acetate	1.06	Dodecane	0.12
2,4-Heptadienal	0.42	2-Methyl-5-nitro-3-hexanol	1.4	Hexanoic acid	0.85	2-Methyldecane	0.12
5-Methyldecane	0.47	Benzene, 2,4-diisocyanato-1-methyl	10.81	Ethyl hexanoate	2.43	Dodecane, 4-methyl	0.22
Nonane, 2,5-dimethyl	0.56	Propanoic acid, 2-methyl-, 3-hydroxy-2,4,4-trimethylpentyl ester	0.16	1,4-Dichlorobenzene	2.88	1H-Pyrrole-2,5-dione, 3-ethyl-4-methyl	0.75
4-Methyldecane	1.5	1,3-Benzenediamine, 4-methyl	0.67	Undecane	2.06	3,7-Dimethyldecane	0.03
D-Limonene	0.49	myristyl alcohol	0.35	4-Methyldecane	3.53	4,6-Dimethyldodecane	0.13
2-Ethylhexanol	2.25	Decanoic acid, ethyl ester	0.38	1-Limonene	0.47	Tetradecane	0.21
1,5-Heptadien-4-one, 3,3,6-trimethyl	0.73	4-Pyrrolidinopyridine	2.27	Eucalyptol	0.58	2-Ethoxyethyl acrylate	0.38
Benzeneacetalddehyde	0.18	Pyrrole-3-carbonitrile, 5-formyl-2,4-dimethyl	4.81	2-Ethylhexanol	0.37	4-Propylbenzaldehyde	0.49
Undecane, 4,7-dimethyl	1.80	.alpha.-Ionone	0.09	1,5-Heptadien-4-one, 3,3,6-trimethyl	1.23	Phytane	0.21
3,5-octadien-2-one	1.4	2-Heptanol, 6-methyl	0.06	2-Bromo-4-methylpentane	0.27	Undecane, 4,7-dimethyl	0.08
Dodecane, 5-methyl	0.10	5,9-Undecadien-2-one, 6,10-dimethyl	0.17	5-Methylundecane	3.93	4-Methyltridecane	0.04
Undecane	0.41	Cetyl alcohol	0.10	5-(2-Methylpropyl)nonane	1.23	5-Methyltetradecane	0.12
2,3,6,7-Tetramethyloctane	1.44	4-(2,6,6-Trimethylcyclohexa-1,3-dienyl)but-3-en-2-one	0.39	3,5-Octadien-2-one	3.15	2,6,11-Trimethyldecane	0.20
Nonanal	0.55	Tetradecane	0.30	7-Methyl-1-undecene	0.11	Benzene, 2,4-diisocyanato-1-methyl	33.41
3-Acetyl-2,6-heptanedione	0.77	pentadecane	0.06	Methyl 6-methylheptanoate	0.71	1,3-Benzenediamine, 4-methyl	0.67
4,6-Dimethyldodecane	1.65	N,N-Dimethyldodecylamine	0.31	2,3,6,7-Tetramethyloctane	0.51	4-Pyrrolidinopyridine	6.63
4-Methylene-1,3-dioxolane	0.14	2,4-Di-tert-butyl-1-methoxybenzene	0.05	Octane, 6-ethyl-2-methyl	2.52	Pyrrole-3-carbonitrile, 5-formyl-2,4-dimethyl	14.47
3,7-Dimethyldecane	0.12	Dicyclohexyl ketone	0.15	2,6-Dimethylcyclohexanol	1.03	5,9-Undecadien-2-one, 6,10-dimethyl	0.65
6-Methylundecane	0.13	Dodecanoic acid, methyl ester	0.57	2,6,10-Trimethyldodecane	0.89	4-(2,6,6-Trimethylcyclohexa-1,3-dienyl)but-3-en-2-one	0.11
Decamethylcyclopentasiloxane	0.95	Dihydroactinidioides	0.35	Benzeneethanol	4.67	beta-Ionone	1.90
Isododecane	0.64	Megastigmatrienone 2	0.19	Methyl octanoate	0.40	Octadecane	0.14
2,6-Octadiene, 2,6-dimethyl	0.13	Ethyl laurate	0.85	Isododecane	1.23	Dihydroactinidioides	5.01
2,4-Dimethylundecane	0.38	Hexadecane	0.27	Ethyl palmitate	1.78	Methanone, diphenyl	0.34
2,5-Dimethylundecane	1.32	Hexadecanal	0.15	4a,5-Dimethyl-6,7,8,8a-tetrahydro-5H-benzo[d][1,3]dioxin-4-one	0.40	Decanoic acid, decyl ester	0.56
Naphthalene	0.85	Methanone, diphenyl	0.40	2,5-Dimethylundecane	1.94	4-Sec-Butyl-2,6-di-tert-butylphenol	0.27
alpha-Terpineol	0.21	2H-1,3-Oxazine-2-thione, tetrahydro-4,4,5,6-tetramethyl	5.55	Naphthalene	1.19	4-Octadecylmorpholine	2.52
1,3-Cyclohexadiene-1-carboxaldehyde, 2,6,6-trimethyl	0.32	4-Sec-Butyl-2,6-di-tert-butylphenol	5.07	1,3-Cyclohexadiene-1-carboxaldehyde, 2,6,6-trimethyl	1.57	Dimethyltetradecylamine	0.49
Dodecane	0.69	4-Octadecylmorpholine	18.07	Dodecane	1.44	Citronellol	0.13
Tetradecane	2.1	Naphthalene, 2,6-bis(1-methylethyl)	0.12	Hexadecane	1.03	Phytone	2.46
Dodecane, 4-methyl	0.52	Phytane	0.11	2,4-Dimethylundecane	0.44	5H-1,4-Dioxepin, 2,3-dihydro	5.79
1H-Pyrrole-2,5-dione, 3-ethyl-4-methyl	0.25	Dimethyltetradecylamine	2.44	Dodecane, 4-methyl	0.84	Dimethyl palmitamine	4.21

MESMAP-5 PROCEEDINGS BOOK
 April 24th - 26th, 2019 / Cappadocia – Turkey

2,6-Dimethylundecane	0.37	Benzoic acid, 2-ethylhexyl ester	0.72	Tetradecane	1.3	4-(Hexyloxy)-1-butanol	0.18
3,3-Dimethyloctane	0.14	Methyl myristate	0.23	2,6-Dimethylundecane	0.50	1,3,5-Triazine-2,4(1H,3H)-dione, 6-(2-methylphenylamino)	3.19
2,4-Dimethyldodecane	0.80	Dibutyl adipate	0.13	4,6-Dimethyldodecane	1.76	Laurinsaeure, 4-Octylester	0.39
N-Formyl-DL-phenylalanine	0.19	Ethyl nonadecanoate	0.18	N-Formyl-DL-phenylalanine	0.18	2,5-Dimethyl-4-hydroxy-3-hexanone	4.58
Phytane	1.57	2,3,7-trimethyloctanal	0.18	Phytane	2.25		
Estragole	0.31	Phytol	0.37	3-Methyl-5-propylnonane	0.82		
pentadecane	0.76	Phytone	0.66	Tridecane	0.19		
Tridecane	0.30	2,2,3,3,4,4,4-Heptafluoro-N-(2-morpholin-4-yl-ethyl)-butyramide	19.67	2,3,7-Trimethyldecane	0.62		
2,3,7-Trimethyldecane	0.43	Nonadecane	0.33	decanoic acid, methyl ester	1.96		
Butanoic acid, 2-ethylhexyl ester	0.26	1-Hexadecanaminium, N,N,N-trimethyl-, bromide	11.47	Dodecacyclohexasiloxane	0.47		
2,6,11-Trimethyldodecane	0.85	Methyl palmitate	0.11	2,6,11-Trimethyldodecane	0.21		
Dodecacyclohexasiloxane	0.63	Dodecyl propan-2-yl sulfite	0.21	Benzene, 2,4-diisocyanato-1-methyl	0.81		
Sulfurous acid, pentyl tridecyl ester	0.10	13- α -Methylandrosta	0.12	2-Methyltridecane	0.14		
Benzene, 2,4-diisocyanato-1-methyl	0.30	Eicosane	0.28	Decanoic acid	0.26		
10-Methylnonadecane	0.13	1,3,5-Triazine-2,4(1H,3H)-dione, 6-(2-methylphenylamino)	1.4	1-Oxaspiro[4.5]decan-2-one, 3-methylene	0.32		
Propanoic acid, 2-methyl-, 3-hydroxy-2,4,4-trimethylpentyl ester	0.20	Heneicosane	0.27	Decanoic acid, ethyl ester	1.71		
Cadinol	0.26	Dodecanoic acid, isooctyl ester	0.28	6,10-Dimethylundecan-2-one	0.24		
Dodecanal	0.43	Tetracosane	0.19	Pentamethylbenzaldehyde	0.20		
Heptadecane	0.21	Methoxyacetic acid, 4-tridecyl ester	0.07	Nonadecane	0.18		
5,9-Undecadien-2-one, 6,10-dimethyl	0.55			α -Ionone	0.19		
Nonadecane	0.09			5,9-Undecadien-2-one, 6,10-dimethyl	0.40		
beta-Ionone	1.12			4-(2,6,6-Trimethylcyclohexa-1,3-dienyl)but-3-en-2-one	1.08		
N,N-Dimethyldodecylamine	0.19			Methyl 10-methylundecanoate	1.45		
bis(2-Ethylhexyl) ether	0.09			Eicosane	0.36		
Dihydroactinidioid	1.98			Pentadecane	0.28		
Hexadecane	0.54			Dicyclohexyl ketone	0.31		
Isopropyl dodecanoate	0.10			Dodecanoic acid, methyl ester	2.58		
Decanoic acid, decyl ester	4.51			Dihydroactinidioid	0.90		
4-Sec-Butyl-2,6-di-tert-butylphenol	4.18			Lauric acid	0.71		
4-Octadecylmorpholine	18.68			Megastigmatrienone	0.75		
Dimethyltetradecylamine	1.51			Ethyl laurate	2.66		
Benzoic acid, 2-ethylhexyl ester	0.49			Decanoic acid, decyl ester	0.20		
Octadecane	0.17			Dimethyltetradecylamine	0.27		
Hexadecanal	0.18			Methyl myristate	0.17		
Phytone	1.00			Ethyl tetradecanoate	0.20		
4-Amino-N-(3-morpholin-4-ylpropyl)-1,2,5-oxadiazole-3-carboxamide	11.91			Phytone	1.46		
Hexadecanyldimethylamine	6.32			4-Octadecylmorpholine	1.12		
Eicosane	0.71			hexadecanyldimethylamine	2.12		
Hexacosane	0.14			Docosane	0.16		
laurinsaeure, 4-octylester	0.72			Laurinsaeure, 4-Octylester	2.16		
2-Linoleoylglycerol	0.24						
Linoleoyl chloride	0.67						
1,3-Di(trans-9-octadecenoyl)glycerol	0.56						

CONCLUSION

In conclusion, the volatile oil and semi-volatile oil contents of *Isatis tinctoria* have been examined for the first time by SPME/GC-MS analysis. When considering the substances contained in *I. tinctoria*, we can say that the species has an antifungal, antibacterial, therapeutic effect. Although the volatile and semi-volatile oils such as Anethole, Limonene, Cetyl alcohol, Decamethylcyclopentasiloxane, Terpinol, Estragole, Valeric acid, Glycerol, Citronellol, Butyric acid, Eucalyptol, Ethyl palmitate and Cadinol are found at low levels in leaves and flowers, these compounds are flavoring, aroma enhancer, food additive, It is used for perfume, cosmetics, pharmacological and medical purposes.

ACKNOWLEDGEMENTS

We want to thank the BAP (Scientific Researching Projects) Foundation of Selçuk University for their financial support (Project number 19704017).

REFERENCES

- [1] Condursa, C., Verzera, A., Romeo, V., Ziino, M., Trozzi, A., and Ragusa, S., 2006. The Leaf Volatile Constituents of *Isatis tinctoria* by Solid-Phase Microextraction and Gas Chromatography/Mass Spectrometry. *Planta Medica*, 72, 924 and 928.
- [2] Fréchar, A., Fabre, N., Pean, C., Montaut, S., Fauvel, M.T., Rollin, P., and Fouraste, I., 2001. Novel indole-type glucosinolates from woad (*Isatis tinctoria* L.). *Tetrahedron Lett.*, 42, 9015 and 9017.
- [3] Hamburger, M., 2002. *Isatis tinctoria*-From the rediscovery of an ancient medicinal plant towards a novel anti-inflammatory phytopharmaceutical. *Phytochemistry Reviews*, 1, 333 and 344.
- [4] Hartleb, I., and Seifert, K. A., 1994. Novel anthranilic acid derivative from *Isatis tinctoria*. *Planta Medica*, 60, 578 and 579.
- [5] Hartleb, I., and Seifert, K., 1995. Acid constituents from *Isatis tinctoria*. *Planta Medica*, 1995; 61, 95-96.
- [6] Hashimoto, Y., Yoshimura, M., and Huang, R.N., 2019. *Appl Entomol Zool*.
- [7] Heinemann, C., Schliemann-Willers, S., Oberthür, C., Hamburger, M., and Elsner, P., 2004. Prevention of experimentally induced irritant contact dermatitis by extracts of *Isatis tinctoria* compared to pure tryptanthrin and its impact on UVB-induced erythema. *Planta Medica*, 70, 385 and 390.
- [8] Honda, G., Tosirisuk, V., and Tabata, M., 1980. Isolation of an antidermatophytic, tryptanthrin, from the indigo plants, *Polygonum tinctorium* and *Isatis tinctoria*. *Planta Medica*, 38, 275 and 276.
- [9] Goetz, J.K., and Schraudolf, H., 1983. Two natural indole glucosinolates from Brassicaceae. *Phytochemistry*, 22, 905 and 907.
- [10] Lockwood, G.B., and Belkhiri, A., 1991. Glucosinolate spectrum of some Algerian Cruciferae. *Plant Systematics and Evolution*, 176, 11 and 20.
- [11] Koufogianni, A., 2006. Comparison of the aroma profile of different Mediterranean olive oil cultivars, Master Thesis, Ghent University.
- [12] Mohn, T., 2009. A comprehensive metabolite profiling of *Isatis tinctoria* leaf extracts, Philosophisch-Naturwissenschaftlichen Fakultät der Universität Basel, Basel. 1-157.
- [13] Recio, M.C., Cerdá-Nicolás, M., Hamburger, M., and Ríos, J.-L., 2006a. Anti-arthritis activity of a lipophilic woad (*Isatis tinctoria*) extract. *Planta Medica*, 72, 715–720.
- [14] Recio, M.C., Cerdá-Nicolás, M., Poterat, O., Hamburger, M., and Ríos, J.-L., 2006b. Anti-inflammatory and antiallergic activity in vivo of lipophilic *Isatis tinctoria* extracts and tryptanthrin. *Planta Medica*, 72, 539–546.
- [15] Wu, X.Y., Liu, Y., Sheng, W., Sun, J., and Qin, G., 1997a. Chemical constituents of *Isatis indigotica*. *Planta Medica*, 63, 55 and 57.
- [16] Wu, X.Y., Qin, G.W., Cheung, K.K. and Cheng, K.F., 1997b. New alkaloids from *Isatis indigotica*. *Tetrahedron*, 53, 13323 and 13328.
- [17] Zhu, Y.P., 1998. *Chinese Materia Medica*. Harwood Academic, Amsterdam.

FULL PAPER

IN VITRO SUSCEPTIBILITY TESTING OF TEA TREE OIL ACNE CREAM IN COMBINATION WITH SALICYLIC ACID AND NICOTINAMIDE

Saiqa Ishtiaq*, Rabia Arshad, Sairah Hafeez Kamran

Punjab University College of Pharmacy, University of the Punjab, Allama Iqbal Campus – 54000, Lahore Pakistan.

Abstract

Acne vulgaris, is a commonly occurring skin disease caused by *Staphylococcus aureus* and *Propionibacterium acnes* (Gollnick *et al.*, 2008). Tea Tree Oil (TTO) is derived mainly from the Australian native plant *Melaleuca alternifolia* (Maiden & Betche) Cheel, belongs to family Myrtaceae. The oil is very much effective in treating a broad range of infectious conditions, especially fungal skin conditions, as well as warts, acne, and vaginal yeast infection (Edris, 2007). The present study is focused to evaluate the *in-vitro* susceptibility and effectiveness of tea tree oil acne cream against acne culprits; *Staphylococcus aureus* and *Propionibacterium acnes*. **Materials and Methods:** Tea tree oil acne cream is a compounding and dispensing product of the *Nigehban Pharmaceuticals (Pvt.) Ltd.* Lahore Pakistan. The bacterial samples were collected from the skin portion of forehead and cheeks of the patients from outdoor of Mayo Hospital Lahore. The formulation was tested for its susceptibility by microwell dilution method on 15 samples of *S. aureus* and five samples of *P. acnes*. After finding the susceptibility, minimum inhibitory concentration of the cream was estimated to establish its effectiveness, by 96 well plate method. SPSS Ver. 21 was employed for data analysis obtained from experiments. All the results were presented in terms of \pm SEM. For analysis of variance, two-way ANOVA was applied with post-hoc Bonferroni test. $p < 0.05$ was considered as statistically significant.

Key words: tea tree oil, acne, salicylic acid and nicotinamide.

RESULTS

Antimicrobial assay of tea tree oil acne cream against *S. Aureus*

Table 1. Antimicrobial assay of tea tree oil acne cream against *S. aureus* by microwell dilution assay

Sr. No.	Antimicrobial assay of tea tree oil acne cream against <i>S. aureus</i>					
	1:5	1:10	1:20	1:50	1:100	1:1000
SMP 1	12.4±0.22	12.2±0.22	8.26±0.17*	12.16±0.17	8.3±0.25*	2.46±0.27*
SMP 2	12.5±0.35	4.26±0.22*	12.9±0.22	12.5±0.22	4.43±0.32*	3.36±0.32*
SMP 3	12.6±0.17	1.43±0.20*	9.53±0.17*	8.93±0.17*	4.26±0.17*	3.36±0.25*
SMP 4	13.3±0.26	3.36±0.32*	12.4±0.32*	9.83±0.32*	3.5±0.26*	1.43±0.26*
SMP 5	12.9±0.18	8.26±0.22*	8.93±0.22*	13.3±0.26	1.5±0.29*	9.53±0.22*
SMP 6	13.4±0.26	2.46±0.34*	9.3±0.26*	8.6±0.31*	3.36±0.32*	4.26±0.228
SMP 7	8.26±0.22	8.93±0.22	9.53±0.22	9.33±0.26	12.4±0.32	9.83±0.32
SMP 8	13.3±0.26	12.5±0.33	13.4±0.26	12.9±0.18	12.6±0.22	13.3±0.26
SMP 9	14.2±0.18	3.5±0.26*	12.6±0.22	13.5±0.29	2.46±0.34*	3.5±0.26*
SMP 10	13.5±0.30	14.2±0.18	13.3±0.26	12.9±0.14	8.9±0.23*	8.5±0.32*
SMP 11	9.53±0.22	9.3±0.26	14.2±0.18*	13.3±0.26*	9.33±0.28	8.26±0.22
SMP 12	13.5±0.29	8.26±0.22*	9.83±0.32*	13.4±0.26	12.1±0.18	8.1±0.18*
SMP 13	12.1±0.18	13.4±0.26	12.9±0.22	8.1±0.18*	9.53±0.22*	9.3±0.26*
SMP 14	8.1±0.18	9.5±0.18	14.2±0.18*	9.83±0.32	10.1±0.18*	8.4±0.26
SMP 15	9.83±0.32	8.1±0.18	10.1±0.18	2.43±0.32*	4.2±0.18 β	3.3±0.26*
Vancomycin	13.67±0.50	13.67±0.50	13.67±0.50	13.67±0.50	13.67±0.50	13.67±0.50

All the results are presented as \pm SEM. For statistical analysis two-way ANOVA with Bonferroni test was applied. Multiple comparisons were done using $p < 0.05$. *represents comparison among 1:5 with all other dilutions.

Antimicrobial assay of tea tree oil acne cream against *P. acnes*

Table 2: Antimicrobial assay of tea tree oil acne cream against *P. acnes* by microwell dilution assay

Sr. No.	Antimicrobial assay of tea tree oil acne cream against <i>P. acnes</i>					
	1:5	1:10	1:20	1:50	1:100	1:1000
Sample 1	2.46 \pm 0.34	9.6 \pm 0.26*	8.26 \pm 0.23*	8.6 \pm 0.32*	3.5 \pm 0.26	4.26 \pm 0.23*
Sample 2	8.23 \pm 0.23	9.3 \pm 0.26	8.53 \pm 0.26	9.1 \pm 0.18	8.2 \pm 0.18	8.2 \pm 0.26
Sample 3	4.26 \pm 0.23	13.3 \pm 0.26*	2.6 \pm 0.18*	3.3 \pm 0.18	10.13 \pm 0.14*	2.6 \pm 0.38*
Sample 4	1.2 \pm 0.15	1.1 \pm 0.22	1.6 \pm 0.15	1.2 \pm 0.22	1.3 \pm 0.22	1.6 \pm 0.15
Sample 5	No effect	No effect	No effect	No effect	No effect	No effect
Vancomycin	13.67 \pm 0.71	13.67 \pm 0.71	13.67 \pm 0.71	13.67 \pm 0.71	13.67 \pm 0.71	13.67 \pm 0.71

All the results are presented as \pm SEM. For statistical analysis two-way ANOVA with Bonferroni test was applied. Multiple comparisons were done using $p < 0.05$. *represents comparison among 1:5 with all other dilutions.

Minimum Inhibitory Concentration (MIC) of tea tree oil acne cream against *S. Aureus*

Table 3. Minimum Inhibitory Concentration (MIC) of tea tree oil acne cream against *S.aureus*

No of obs.	Conc. (mm)	Bacteria + Compound				Compound			Minimum Inhibitory Concentration	Avg.
1	12.5	0.132	0.127	0.149	0.058	0.074 ^a	0.069 ^b	0.091	0.078	
2	6.25	0.399	0.553	0.479	0.089	0.31	0.464	0.39	0.388	
3	3.125	0.465	0.424	0.341	0.075	0.39	0.349	0.266	0.335	
4	1.87	0.440	0.507	0.483	0.068	0.342	0.356	0.273	0.323	
5	0.935	0.452	0.486	0.491	0.061	0.379	0.446	0.422	0.415	
6	0.467	0.569	0.486	0.471	0.058	0.511	0.428	0.418	0.452	
7	0.233	0.488	0.492	0.450	0.062	0.426	0.367	0.388	0.393	
8	0.116	0.512	0.448	0.478	0.056	0.455	0.391	0.422	0.422	
9	0.05	0.553	0.443	0.459	0.057	0.496	0.386	0.402	0.428	
10	0.02	0.459	0.457	0.469	0.070	0.389	0.387	0.399	0.391	

a = shows MIC at 12.5 conc.

b = shows MIC at 6.25 conc.

Minimum Inhibitory Concentration (MIC) of tea tree oil acne cream against *P. Acnes*

Table 7. Minimum Inhibitory Concentration (MIC) of tea tree oil acne cream against *P. acnes*

No of obs.	Conc. (mm)	Bacteria + Compound				Compound			Minimum Inhibitory Concentration	Avg.
1	12.5	0.359	0.090	0.081	0.058	0.301	0.032	0.023 ^a	0.118	
2	6.25	0.626	0.155	0.096	0.089	0.537	0.066	0.007 ^b	0.203	
3	3.125	0.444	0.273	0.148	0.075	0.369	0.198	0.073 ^c	0.2130	
4	1.87	0.535	0.307	0.352	0.068	0.467	0.239	0.284	0.33	
5	0.935	0.635	0.461	0.365	0.061	0.574	0.4	0.304	0.426	
6	0.467	0.560	0.380	0.312	0.058	0.502	0.322	0.254	0.359	
7	0.233	0.527	0.511	0.388	0.062	0.465	0.449	0.326	0.413	
8	0.116	0.568	0.462	0.472	0.056	0.512	0.406	0.416	0.4446	
9	0.05	0.536	0.521	0.423	0.057	0.479	0.464	0.366	0.436	
10	0.02	0.384	0.517	0.142	0.070	0.314	0.447	0.072 ^d	0.277	

a = shows MIC at 12.5 conc.

b = shows MIC at 6.25 conc.

c = shows MIC at 3.125 conc.

d = shows MIC at 0.02 conc.

DISCUSSION/ CONCLUSION

A wide variety of essential oils are known to possess antimicrobial properties, and in many cases this activity is due to the presence of active monoterpene constituents (Delaquis *et al.*, 2002). Similarly, in the present study we employed the essential oil of *Melaleuca alternifolia* to establish its efficacy in the form of a topical preparation in combination with salicylic acid and nicotinamide. The formulations showed promising effects against these both bacteria strains. The results suggest that the tea tree oil acne cream is effective in a narrow dilution range and shows potent anti-acne effect. The results of the MIC reveal that the formulation has profound action on the bacterial strains. Tea tree oil (5%) had a significant effect in ameliorating the patient's acne by reducing the inflamed and non-inflamed lesions (Carson, 2006). It was reported, that exposing these organisms to MIC and MBC of tea tree oil, inhibited respiration and increased the permeability of bacterial cytoplasmic membrane by uptake of propidium iodide. In the case of *S. aureus*, tea tree oil also causes potassium ion leakage (Trombetta *et al.*, 2005). The tea tree oil acne cream was prepared with salicylic acid and nicotinamide. Salicylic acid is a mild keratolytic and anti-inflammatory agent. (Jacquet, 1988). Nicotinamide is reported to be effective as antibiotic, and provides potent anti-inflammatory action without the risk of inducing bacterial resistance (Shalita, 1995). The ingredients will not only improve the product's susceptibility against the strains but will also show patient compliance.

REFERENCES

- [1] Carson, C. F., Hammer, K. A., & Riley, T. V. (2006). *Melaleuca alternifolia* (tea tree) oil: a review of antimicrobial and other medicinal properties. *Clinical microbiology reviews*, 19(1), 50-62.
- [2] Delaquis, P. J., Stanich, K., Girard, B., & Mazza, G. (2002). Antimicrobial activity of individual and mixed fractions of dill, cilantro, coriander and eucalyptus essential oils. *International journal of food microbiology*, 74(1-2), 101-109.
- [3] Edris, A. E. (2007). Pharmaceutical and therapeutic potentials of essential oils and their individual volatile constituents: a review. *Phytotherapy Research: An International Journal Devoted to Pharmacological and Toxicological Evaluation of Natural Product Derivatives*, 21(4), 308-323.
- [4] Gollnick, H. P., Finlay, A. Y., & Shear, N. (2008). Can we define acne as a chronic disease?. *American journal of clinical dermatology*, 9(5), 279-284.
- [5] Jacquet, Y. F. (1988). The NMDA receptor: central role in pain inhibition in rat periaqueductal gray. *European journal of pharmacology*, 154(3), 271-276.
- [6] Shalita, A. R., Smith, J. G., Parish, L. C., Sofman, M. S., & Chalker, D. K. (1995). Topical nicotinamide compared with clindamycin gel in the treatment of inflammatory acne vulgaris. *International journal of dermatology*, 34(6), 434-437.
- [7] Trombetta, E. S., & Mellman, I. (2005). Cell biology of antigen processing in vitro and in vivo. *Annu. Rev. Immunol.*, 23, 975-1028.

HEADSPACE SOLID PHASE MICROEXTRACTION (HS-SPME) ANALYSIS OF *SIDERITIS OZTURKII* AYTAC & AKSOY

Tuna Uysal*, Ela Nur Şimşek Sezer, Meryem Bozkurt

Department of Biology, Faculty of Art and Science, University of Selçuk, Konya,
Turkey, * tuysal@selcuk.edu.tr

Abstract

The genus *Sideritis* (Lamiaceae) is represented by more than 150 bushy species and annual or perennial plants. Many studies have been conducted to elucidate the chemical composition and pharmacological activities of plants of the genus. *Sideritis ozturkii* Aytac & Aksoy is an endemic plant to Turkey and used as herbal tea and folk medicine in central Anatolia. The purpose of this study was the detection of the volatile oil components of *S. ozturkii* by solid phase microextraction/gas chromatography/mass spectrometry (SPME-GC-MS) technique. Dried and powdered samples were used directly. Two SPME fibres, 50/30 µm DVB/CAR/PDMS and 85 µm Carboxen/PDMS were preferred for analysis. SPME procedure was analysed on a Shimadzu QP2010 ULTRA FID GC-MS system. The volatile oil compositions were identified in *Sideritis ozturkii* samples by SPME-GC/MS analysis using two different fibres. The major constituents of dried leaves detected with 50/30 µm DVB/CAR/PDMS fiber were octanoic acid (11.70 %) and nonanoic acid methyl ester (16.68 %) for 85 µm Carboxen/PDMS fiber, respectively. On the other hand, the major constituents of flowers were detected with both fibres were hexanoic acid (24.32 %- 26.34 %) and hexanal (18.67 % -16.68 %) respectively. The comparative volatile oil composition of *S. ozturkii* leaves and flowers were revealed via the SPME method for the first time with this study.

Keywords: Paşa çayı, Solid-phase microextraction, Turkey.

INTRODUCTION

Sideritis (L.) is located within the Lamiaceae family. There are more than 160 species of the genus in all over the world. *Sideritis* is represented by 53 taxa, 40 of which are endemic, in Turkey (Duman, 2012). Endemism rate is 73%. *Sideritis* spp., known as Adaçayı (Sage) and Dağçayı (Ironwort), are widely used in Turkey. Although *Sideritis* species which are widely used as herbal tea in the folk have not been studied extensively in terms of the components which they contain, there are some studies (Topçu et al., 2001; Kırimer et al., 2004; Gonzalez-Burgos et al., 2011). The chemical components of the plant extracts of the species *Sideritis* are flavonoids, terpenoids, lignans, coumarins and iridoids (Gonzalez-Burgos et al., 2011; Koutsaviti et al., 2013; Çarıkçı et al., 2018). Moreover, it was reported that some components of the plant have anti-inflammatory, antirheumatic, antistress, analgesic, antioxidant, antispasmodic, antibacterial, digestive, diuretic, and antimicrobial activities (Gonzalez-Burgos et al., 2011; Koutsaviti et al., 2013; Topçu et al., 2001; Topçu et al., 2002; Goren et al., 2002; Tunalier et al., 2004; Fraga 2012).

Sideritis ozturkii, which is an endemic species to our country, is known as Dağçayı. In Central Anatolia, it is used as herbal tea and spices. There is a limited number of studies related to this species. There are some studies including anti-inflammatory and antinociceptive effects as well as phenolic content and antioxidants associated with antimicrobial activities of *S. ozturkii* (Küveli et al., 2007; Sagdic et al., 2008). SPME is an increasingly common method for rapidly ejecting volatile compounds from plants (Zhu et al., 2013). SPME is a pre-concentration technology based on direct sample analysis without the use of any solvent and preventing the production of artefacts. Conjunction with gas chromatography (GC)-mass spectrometry (MS) analysis, it has been successfully applied for profiling the metabolomic pattern of environmental, food, forensic and pharmacological samples, and as a powerful technique for the extraction of urinary potential biomarkers of various diseases, such as diabetes, cancer and autism syndrome (Gonçalves et al., 2012; Cozzolino et al., 2014).

In this study, solid phase microextraction/gas chromatography/mass spectrometry (SPME-GC-MS) technique was used for profiling the volatile oils of dried leaves and flowers from Turkish endemic sage species.

MATERIAL AND METHODS

Material: The plant material was collected from Konya-Derebucak province in 2016. The collected plant material washed and dried without sunlight. Dried and powdered leaf and flower samples were used directly. The ground samples (3gr) sealed in a 10 ml vial. Two SPME fibres, 50/30 μm DVB/CAR/PDMS and 85 μm Carboxen/PDMS were used for analysis. The SPME apparatus was directly injected into the upper space of the vial to adsorb volatile substances and then directly injected into the Shimadzu QP2010 ULTRA GC-MS apparatus using a Restek Rxi-5 MS capillary column.

HS-SPME-GC/MS Analysis: The volatile components were separated for about 15 minutes with different SPME fibres attached to a GC injector (the temperature of the injector was 250 °C). The plant material analysed on a Shimadzu QP2010 ULTRA FID GC-MS system. A Restek Rxi-5 MS column (30 m \times 0.25 mm id, film thickness 0.25 μm) used with helium as a carrier gas. The GC oven temperature was programmed to hold at 40 °C for 3 min and then to increase to 240 °C at 5 °C/min, finally holding at 240 °C for 3 min. The column flow rate was 1.8 mL/min. The ion source temperature was 200 °C, and the interface temperature set at 250 °C.

Evaluation of data: After the analysis, each sample was demonstrated with GC-MS chromatograms. Chromatograms of all samples were exposed to noise reduction before peak area integration, and then the peak areas of compounds in the chromatogram were combined. Compounds were identified by comparison with three libraries, W9N11, SWGDR4G4 and SWGDR4G5. Compounds were mostly found in the W9N11 library and specified with other libraries. Identification was performed based on the retention time (RT) of the components in the sample. The relative percentage of the volatile oil constituents was calculated from peak areas. All analyses were carried out triplicate, and all data are means of three independent analysis.

RESULTS AND DISCUSSION

The results obtained from the analysis of *Sideritis ozturkii* leave and flower by two different SPME fibres are shown in **Table 1**. Totally, 172 component identified from the leaves and 133 compound identified from the flowers during our analyses. However, we preferred to give 45 main components and their quantities in leaf and flower content are given in the table, respectively. When the content of the leaf was evaluated, it was found that octanoic acid, methyl ester (11,7- 9,73 %) and nonanoic acid, methyl ester (11,25- 16,68 %) ratios were high in both fibres. These compounds are also found in floral content but less in quantity. The most common found components in the flower content are hexanoic acid (24,32- 26,34 %) and hexanal (18,67- 14,29 %). 9 components are common to both leaf and flower content and were detected with both fibres. These are; nonanoic acid, methyl ester, Cyclopentasiloxane, decamethyl-, hexanoic acid, nonanal, octanoic acid, cyclohexasiloxane, dodecamethyl-, 1-octanol, heptanoic acid, nonanoic acid respectively. The phenolic composition, essential oils and flavonoids of *Sideritis* species were studied by many researchers before (Gergis et al., 1989; Gomez-Serranillos et al., 1998; Tabanca et al., 2001; Aboutabl et al., 2002; Kilic et al., 2003; Tunalier et al., 2004; Gabrieli et al., 2005). According to the previous study, it was reported that the main components of *S. ozturkii* essential oils were α -pinene and β -pinene via GC/MS analysis in three samples (Kirimer et al., 2001). In our analyses, α -pinene and β -pinene were present but in minor amounts. In this study, the analyses were performed after extraction, but in our study, the sample was used directly without any procedure. One another study, anti-inflammatory and antinociceptive effects of the *Sideritis ozturkii* acetone extract were investigated, and it was reported that the extract has anti-inflammatory and antinociceptive activity (Küpelı et al., 2007). According to the analyses, the components in the content have different effects. For example, octanoic acid and nonanoic acid methyl ester used as a flavouring agent and is present in many plants. On the other hand, hexanal which is one of the most compounds in flower is potentially useful as a natural extract that prevents fruit spoilage (Sharkey, 2016).

Table 1. The most common compounds identified in *S. ozturkii* leaf and flowers via SPME method

		Fibres			
		50/30 µm DVB/CAR/PD MS	85 µm Carboxen/PDMS	50/30 µm DVB/CAR/PD MS	85 µm Carboxen/PD MS
		Leaf		Flower	
	Compound Name	% area	% area	% area	% area
1.	Octanoic acid, methyl ester	11,7	9,73	0,76	-
2.	Nonanoic acid, methyl ester	11,25	16,68	0,86	0,86
3.	Cyclopentasiloxane,decamethyl-	4,52	1,4	1,75	1,16
4.	Hexanoic acid	4,3	5,1	24,32	26,34
5.	Nonanal	4	3,46	3,88	2,79
6.	Octanoic acid	3,89	5,7	0,93	0,73
7.	Benzene, 1-methoxy-4-(1-propenyl)-	3,78	-	1,62	5,66
8.	Cyclohexasiloxane,dodecamethyl-	3,54	9,73	2,54	5,08
9.	1-octanol	2,99	1,85	1,83	0,94
10.	Heptanoic acid, methyl ester	2,74	-	-	-
11.	Octanal	2,51	1,04	-	1,67
12.	Hexanoic acid, methyl ester	2,5	-	0,66	0,30
13.	1,2 Benzenedi carboxylic acid, diethyl ester	2,07	-	-	-
14.	1,2-Benzenedicarboxylic acid, dinonyl ester	2,05	-	-	-
15.	Pyridine, 2-methyl-4,6-dipropyl-	1,8	-	-	-
16.	1-heptanol	1,75	-	0,73	-
17.	Diethyl Phthalate	1,69	-	-	-
18.	Cyclotrisiloxane, octamethyl-	1,55	-	-	-
19.	Hexanal	1,47	-	18,67	14,29
20.	6-methyl-5-hepten-2-one	1,44	-	-	-
21.	Heptanoic acid	1,38	1,06	1,51	1,54
22.	Nonanoic acid	1,28	2,08	0,69	0,80
23.	Benzene, methyl(1-methylethyl)-	1,04	-	-	-
24.	Anethole	-	7,7	-	-
25.	2-furancarboxaldehyde	-	4,58	0,47	2,20
26.	Estragole	-	1,59	0,66	0,51
27.	Cycloheptasiloxane,tetradecamethyl	-	1,27	0,48	0,91
28.	Decanoic acid, methyl ester	-	1,07	-	-
29.	Decanal	-	1,02	0,63	0,70
30.	3-Methyl-hepta-1,6-dien-3-ol	-	-	-	2,71
31.	Pentanoic acid	0,86	-	1,70	2,07
32.	5-pentyl-2(5H)-furanone	-	-	-	1,73
33.	2-Octenal	-	-	3,70	2,56
34.	Dodecanal	-	-	0,60	1,28
35.	1-Decanol, 2-hexyl-	-	-	-	1,20
36.	Pentanoic acid, 2-methyl-, anhdride	-	-	-	1,15
37.	Furan, 2-pentyl-	-	-	6,21	-
38.	Butyric acid, 2,2-dimethyl, vinyl ester	-	-	2,58	-
39.	D-Limonene	-	-	1,95	0,48
40.	3-Octen-2-one	-	-	1,59	-
41.	2-Heptanone	-	-	1,52	-
42.	2-Heptenal	-	-	1,41	-
43.	Heptanal	-	0,23	1,20	0,25
44.	1-Pentanol	0,16	-	1,13	0,18
45.	1-Octen-3-ol	0,18	-	1,07	-

CONCLUSION

In this study, the endemic *Sideritis ozturkii* species from Turkey was analysed via HS-SPME-GC/MS method. According to the findings of this study, the leaf and flowers of *S. ozturkii* had a broad, volatile oil content. It is essential to identify the biological effects of a plant and determine the content for the determination of active ingredients. If the plant is an endemic species, its content is much more valuable. This study is a pioneering study, and our research will continue to determine the active phytochemicals and their effects in the content.

ACKNOWLEDGEMENTS

We want to thank Assist Prof.Dr. Hakkı DEMİRELMA for the collected plant material and the BAP (Scientific Researching Projects) Foundation of Selçuk University for their financial support (Project number 19704016).

REFERENCES

1. Aboutabl, E. A., Nassar, M. I., Elsakhawy, F. M., Maklad, Y. A., Osman, A. F., & El-Khrisy, E. A. M. 2002. Phytochemical and pharmacological studies on *Sideritis tautica* Stephan ex Wild. Journal of Ethnopharmacology, 82, 177–184.
2. Cozzolino R, De Magistris L, Saggese P, Stocchero M, Martignetti A, Di Stasio M, Malorni A, Marotta R, Boscaino F, Malorni L. 2014. Use of solid-phase microextraction coupled to gas chromatography-mass spectrometry for determination of urinary volatile organic compounds in autistic children compared with healthy controls. Anal Bioanal Chem. 406: 4649–4662.
3. Çarıkcı S., Özer Z., Dereli S., Açar D., Gören A. C., Kılıç T., 2018. Essential Oil Composition of Five *Sideritis* Species Endemic to Turkey. Süleyman Demirel University Journal of Natural and Applied Sciences, 22: 301-305.
4. Duman, H. 2012. *Sideritis*. In: Güner, A., Aslan, S., Ekim, T., Vural, M., Babaç, M.T. (edlr.), Türkiye Bitkileri Listesi (Damarlı Bitkiler). Nezahat Gökyiğit Botanik Bahçesi ve Flora Araştırmaları Derneği Yayını, İstanbul, pp. 585-588.
5. Fraga, B. M. 2012. Phytochemistry and Chemotaxonomy of *Sideritis* Species from the Mediterranean Region. Phytochemistry. 76: 7-24.
6. Gabrieli, C. N., Kefalas, P. G., & Kokkalou, E. L. 2005. Antioxidant activity of flavonoids from *Sideritis raeseri*. Journal of Ethnopharmacology, 96, 423–428.
7. Gergis, V., Argyriadou, N., & Poulos, C. 1989. Composition of the essential oils of *Sideritis cladestina* ssp. *cylleneu* and *Sideritis sipylea*. Journal of the Science of Food and Agriculture, 47, 501–507.
8. Gomez-Serranillos, P., Carretero, E., Slowing, K., Palomino, O. M., Villarrubia, A. I., & Villar, A. 1998. HPLC quantitative analysis of diterpenoids in *Sideritis* (Labiatae) species. Phytotherapy Research, 12, 101–103.
9. Gonçalves J, Figueira J, Rodrigues F, Câmara JS. 2012. Headspace solid-phase microextraction combined with mass spectrometry as a powerful analytical tool for profiling the terpenoid metabolomic pattern of hop-essential oil derived from Saaz variety. J Sep Sci. 35:2282–2296.
10. Gonzalez-Burgos, E., Carretero, M. E. Gomez-Serranillos, M. P. 2011. *Sideritis* spp.: Uses, Chemical Composition and Pharmacological Activities—A Review. Journal of Ethnopharmacol. 135: 209–225.
11. Goren, A. C., Kılıç, T., Yıldız, Y. K. Tümen, G. 2002. Diterpenes from *Sideritis trojana*. Natural Product Letters. 16:33-37.
12. Kırimer, N. Baser, K. H. C., Demirci, B., Duman, H. 2004. Essential Oils of *Sideritis* Species of Turkey Belonging to the Section Empedoclia, Chemistry of Natural Compounds. 40, 19-23.
13. Kılıç, T., Yıldız, Y. K., Goren, A. C., Tümen, G., & Topcu, G. 2003. Phytochemical analysis of some *Sideritis* species of Turkey. Chemistry of Natural Compounds, 39, 453–456.
14. Kirimer, N., Tabanca, N., Demirci, B., Baser, K. H. C., Duman, H., & Aytac, Z. 2001. The essential oil of a new *Sideritis* species: *Sideritis ozturkii* Aytac and Aksoy. Chemistry of Natural Compounds, 37, 234–237.
15. Koutsaviti, A., Bazos, I., Milenkovic, M., Pavlovic-Drobac, M., Tzakou, O. 2013. Antimicrobial Activity and Essential Oil Composition of Five *Sideritis* Taxa of Empedoclia and Hesiodia sect. from Greece. Records of Natural Products. 7:1- 6.
16. Küpeli, E., Şahin, F. P., Çalış, İ., Yeşilada, E., & Ezer, N. 2007. Phenolic compounds of *Sideritis ozturkii* and their in vivo anti-inflammatory and antinociceptive activities. Journal of ethnopharmacology, 112(2), 356-360.
17. Sağdıç O, Aksoy A, Ozkan G, Ekici L, Albayrak S, 2008. Biological activities of the extracts of two endemic *Sideritis* species in Turkey. Innovative food science & emerging technologies, 9, 1, 80-4.
18. Sharkey, Jackie. 2016. "Fruit spray developed by Guelph prof extends shelf life by 50 per cent". *CBC News*. Retrieved 21 August 2017.
19. Tabanca, N., Kirimer, N., & Baser, K. H. C. 2001. The composition of essential oils from two varieties of *Sideritis erythrantha* var. *erythrantha* and var. *cedretorum*. Turkish Journal of Chemistry, 25, 201–208.
20. Topcu, G., Gören, A. C., Kılıç, T., Yıldız, Y. K. Tümen, G. 2002. Diterpenes from *Sideritis sipylea* and *S. dichotoma*. Turkish Journal of Chemistry. 26: 189-194.
21. Tunalier, Z., Kosar, M., Ozturk, N., Baser, K. H. C., Duman, H., Kirimer, N. 2004. Antioxidant Properties and Phenolic Composition of *Sideritis* Species. Chemistry of Natural Compounds. 40, 206-210.
22. Zhu F, Xu J, Ke Y, Huang S, Zeng F, Luan T, Ouyang G. 2013. Applications of in vivo and in vitro solid-phase microextraction techniques in plant analysis: a review. Anal Chim Acta. 794:1–14.

FROM A TRADITIONAL REMEDY TO MODERN THERAPY; *IN VIVO* ANTIHEMORRHOIDAL STUDY OF *Malva sylvestris* L.

Ceylan Aka-Dönmez^{1a}, Özge Boyacıoğlu^{2a, 2b}, Petek Korkusuz³,
Ufuk Koca-Caliskan^{1b*}

^{1a}Department of Pharmacognosy, Faculty of Pharmacy, Gazi University, 06330, Etiler-Ankara, Turkey,
email:ceylanaka@gazi.edu.tr

^{1b}Department of Pharmacognosy, Faculty of Pharmacy, Gazi University, 06330, Etiler-Ankara, Turkey,
email:ukoca@gazi.edu.tr

^{2a}Bioengineering Department, Graduate School of Science and Engineering, Hacettepe University, 06800,
Ankara, Turkey, email:ozge.boyacioglu@hacettepe.edu.tr

^{2b}Department of Medical Biochemistry, Basic Sciences Division, Faculty of Medicine, Atılım University, 06830,
Ankara, Turkey, email:ozge.boyacioglu@atilim.edu.tr

³Department of Histology and Embryology, Faculty of Medicine, Hacettepe University, 06100, Ankara, Turkey,
email:petek@hacettepe.edu.tr

Abstract

It is estimated that about 50% of the world population is suffering from haemorrhoids, which is known as the most common anorectal disease. Patients have been trying to cope with the hemorrhoids by utilizing both herbal and chemical preparations. Leaves of *Malva sylvestris*, which is known as mallow in English, Ebegümeci in Turkish, belongs to Malvaceae family, have been utilized as a folk medicine to treat haemorrhoids. The studies demonstrated that the leaves have potent anti-inflammatory, antioxidant, anti-complementary, anti-cancer and skin tissue integrity activities. In this study, *in vivo* antihemorrhoidal activity and supporting *in vitro* activities of *M. sylvestris* leaf extract were examined in order to find a scientific base for a folkloric usage. **Material and Methods:** Hemorrhoid model was induced by applying croton oil externally in anal part of the rats. After the occurrence of hemorrhoids *M. sylvestris* leaf (50%) extract and a commercial herbal product (Pilex®) as a positive control were implemented externally in different rat groups. 7 days after the treatment, the animals were anesthetized in order to collect a large blood sample via intracardiac bleed and the serums were obtained to determine the Tumor Necrosis Factor alpha (TNF- α), Vascular Endothelial Growth Factor (VEGF) and Superoxide Dismutase (SOD) levels. Sacrificed rats were weighed and 2 cm long anorectal tissues were cut and weighed in a sequence to calculate the Anorectal Coefficients (ARC) for each rat. Moreover, the excised tissues were fixed in formaldehyde (10%) for histopathological evaluations. **Results:** According to the ARC scores, biochemical and histopathological results, the leaf extract of *M. sylvestris* showed significant biological activity compared to the controls. The value of ARC was 1.86 \pm 0.09 of the plant extract, while serum TNF- α and VEGF levels were 163.22 \pm 11.15 ng/L and 246.37 \pm 12.64 ng/L, respectively. SOD inhibition of the plant extract was 91.58 \pm 6.17%. **Conclusion / Discussion:** The data demonstrated that the antihemorrhoidal activity is directly proportional to the effect on capillary permeability. The effectiveness of the traditional usage has been verified by scientific studies.

Keywords: hemorrhoid, antihemorrhoidal, *in vivo*, croton oil, mallow

INTRODUCTION

Hemorrhoids term is originated from the Greek words "haema (blood)" and "rhoos (flow)". Hippocrates (BC 460) used to describe it as a vascular bleeding in the anus (Leff, 1987). There are limited epidemiological studies on the prevalence of hemorrhoid disease (Lohsiriwat et al., 2012; Everhart et al., 2008). Since the patients suffers from hemmorrhoids, refrain from consulting physicians and do consent from the operations, they try to treat themselves with alternative, mostly natural products. Even if a curative preparation is demanded not only in Turkey but also in all over the world, the number of effective drugs derived from natural products is limited. To heal early stages of hemorrhoids, herbals are quite effective, but they are not sufficient to demonstrate the desired performance in more severe stages. The most popular plants/extracts used against hemorrhoids are *Ruscus aculeatus* (butcher's broom), *Aesculus hippocastanum* (horse chestnut), and *Hammamelis virginiana* (witch hazel) (Abascal et al., 2005). Although there are several plants that have ethnomedicinal usage for hemorrhoids, pharmacological efficacy of many of them, particularly *Malva sylvestris*, have not been proven

scientifically. In traditional medicine, leaves and flowers of the plant are used for the treatment of hemorrhoids, inflammatory and respiratory problems, dermatological and urologic symptoms and gastrointestinal diseases such as heartburn, diarrhea/constipation, and stomachache (Gasparetto et al., 2012). Additionally, experimental studies on leaves revealed that they have an anti-inflammatory, antioxidant, anti-complementary, anticancer effects and improving of damaged skin activity (Baytop, T., 1999; Gasparetto et al., 2012).

The aim of this study is to evaluate antihemorrhoidal activity of the traditionally used *M. sylvestris* leaf extract based on ethnobotanical usage via *in vivo* experimental method.

MATERIALS AND METHODS

Chemicals: All the chemicals were purchased from Sigma Aldrich (St. Louis, MO, USA), and solvents were obtained from Merck at HPLC grade. Croton oil was purchased Alfa Aesar®. Rat Tumor necrosis factor- α (TNF- α) ELISA Kit (YL Biont), Rat Vascular Endothelial Growth Factor (VEGF) ELISA Kit (YL Biont) and Superoxide Dismutase (SOD) Assay Kit 500 test (Sigma Aldrich) were used for biochemical tests.

Plant material and preparation of the extract: The whole flowering cultured plant material was collected during August-September 2017 at Beysehir-KONYA region in Turkey. The leaves of plants were dried in shade and then grounded to powder. Powdered samples (500 g) were extracted with methanol (6×1500 mL) by maceration for 6 days. Filtered methanolic extracts were collected and evaporated until dryness.

Animals: The application for the ethics committee with protocol number 176 was approved by Kobay Experimental Animals Laboratory Local Ethics Committee on 28 April 2016. Healthy adult male Wistar albino rats (2 months old and 180±20 g weight) were used for this study. The rats were maintained at Kobay Experimental Animals Laboratory at 23-25 °C, under standard experimental conditions, at 50% relative humidity and 12 h light/dark cycle. The animals were fed with standard rodent chow and water *ad libitum*.

Induction of hemorrhoid: Hemorrhoid was induced by croton oil solution (deionized distilled water, pyridine and 6% croton oil in diethyl ether) in rats (Nishiki et al., 1988; Azeemuddin et al., 2014; Dey et al., 2016). According to literature cotton swabs (0.4 cm diameter) soaked in 200 μ L of croton oil solution were inserted into the anus and ano-rectal section, 1.5 cm from anal opening) of all the studied animals and kept for exactly 30 seconds. Hemorrhoid induction was repeated once a day for the following three days. The presence of edema was denoted at the end of the third day after the induction.

Treatment of hemorrhoid: After the observation of edema at the anus, treatment was started. The dose of Pilex cream was chosen as reference control according to the previous literature (Azeemuddin et al., 2014).

Cold cream was preferred as a base cream. The aqueous phase containing *M. sylvestris* extract dissolved in water, was mixed with an oily phase at a temperature of 70-80 °C. The cream was applied to the animals twice a day for 7 days (Gurel et al., 2013).

Table 1. Animal experimental groups

Animal Groups	Applications
Control	Rectal 500 μ L croton oil solution,
Vehicle control (base cream)	Rectal 500 μ L croton oil solution, 2x7days, externally 0.5 mL base cream
Reference control (Pilex® cream)	Rectal 500 μ L croton oil solution, 2x7days, externally 0.5 mL pilex cream
<i>Malva sylvestris</i> (50%) extract	Rectal 500 μ L croton oil solution, 2x7days, externally 0.5 mL <i>M. sylvestris</i> cream

On the day 8th, 5 mL blood samples were collected with intracardiac technique from rats under anesthesia for evaluation of the biochemical parameters. After the animals were sacrificed, both animals and their isolated rectoanal tissues (2 cm in length) were weighed.

Anorectal coefficient (ARC): On order to calculate the Anorectal Coefficient (ARC), animals and their isolated recto-anal tissues (2 cm in length) were weighed. ARC was calculated by the following formula: $ARC = \text{Anorectal tissue weight (miligram)} / \text{Animal weight (gram)}$.

Biochemical parameters: Rat Tumor Necrosis Factor- α (TNF- α) ELISA Kit (YL Biont), Rat Vascular Endothelial Growth Factor (VEGF) ELISA Kit (YL Biont), Superoxide Dismutase (SOD) Assay Kit 500 test (Sigma Aldrich) were used for evaluating serum TNF- α and VEGF levels in order to interpret the inflammation degree. SOD is associated with the antioxidant capacity.

Histomorphometric Analysis: The anorectal tissues were fixed in 10% formaldehyde at room temperature. After washing the tissues, they were dehydrated in a graded series of ethanol and cleared in xylene solution. Then the tissues were embedded in paraffin by using a tissue processor with vacuum (Leica, Germany). Thin sections of 3-5 micrometre thickness were obtained with a microtome and subsequently stained with Haematoxylin-Eosin. A histomorphometric examination was carried out in order to define the edema, necrosis, inflammation, congestion, haemorrhage, thickened mucosa, degenerative changes and fibrosis in lesion areas (Dey et al., 2016). Histomorphometric parameters are shown in Table 2. Sections referring to lesion areas were evaluated under the bright field microscope (Leica DMB6 B (Wetzlar, Germany) and attached digital camera Leica DFC7000 T (Wetzlar, Germany) using an image processing software (Leica, LAS Germany). Average scores per group were evaluated by two different researchers.

Table 2. Histomorphometric score parameters are indicated.

Parameter	Lesion Degree in Microscopic Area (%)		Histomorphometric Score
*Edema,	No lesion (0%)	= -	0
*Necrosis,	Minimal (5-10%)	=+	1
*Inflammation,	Mild (10-25%)	=++	2
*Congestion/Thickened,	Moderate (25-50%)	=+++	3
*Mucosa Haemorrhage, *Degenerative Changes *Fibrosis	Severe (more than 50%)	=++++	4

Immunohistochemical Analysis: Paraffin sections were labelled with myeloperoxidase using indirect immune peroxidase method. Anti-Myeloperoxidase Antibody (Boster Bio USA, PB9057, 1%) was used for immunohistochemical analysis. Myeloperoxidase immune reactivity in sections were scored in lesion areas and scaled in a range from – (0%), + (<10%), ++ (10-50%), +++ (>50%) (Krawisz et al., 1984; Sakai et al., 2002). Circumference of veins and stromal cells were taken into consideration in the analysis.

Statistical Analysis: Experiment and control groups are independent variables, whereas histomorphometric and immunohistochemical scoring parameters are dependent variables. The statistical analysis was performed in SPSS 25.0 statistical software and all data were evaluated for the normal distribution Shapiro-Wilk test. Then the total score values were analysed by Kruskal-Wallis non-parametrical analysis followed by Bonferonni Correction. Values were evaluated under 95% confidence interval and $p < 0.05$ was considered significant in all groups.

RESULTS

Anorectal coefficient (ARC) values: As a result, Among the treated groups, the lowest ARC value was observed in *M. sylvestris* cream administered group (Table 3).

Table 3. ARC values of (Mean ± SD) experimental groups

Animal Groups	ARC (Mean±SD)
Control	3.50±0.03
Vehicle control (base cream)	3.43±0.05
Reference control (Pilex® cream)	2.26±0.04***
<i>Malva sylvestris</i> (50%) extract	1.86±0.09***

*:p<0.05; **:p<0.01; ***:p<0.001; SD: Standart error mean

Serum TNF-α, VEGF and SOD levels: Serum TNF-α and VEGF levels of the plant showed a significant difference compared to negative control (Table 4). All the detected TNF-α and VEGF serum levels of mallow were less than the reference control group Pilex®. SOD inhibition of mallow was greater than the reference control group Pilex®.

Table 4. Serum TNF-α, VEGF and SOD experimental groups

Animal Groups	TNF ALPHA (Mean ± SD)(ng/L)	VEGF (Mean ± SD)(ng/L)	SOD (Mean % inhibition ±SD)
Control	217.30±1.35	400.20±6.39	84.16±1.73
Vehicle control (base cream)	192.80±3.65***	319.40±6.70***	86.74±1.88
Reference control (Pilex® cream)	186.50±1.42***	282.20±6.55***	89.51±1.30
<i>M. sylvestris</i> (50%) extract	163.22±11.15***	246.37±12.64***	91.58±6.17*

*:p<0.05; **:p<0.01; ***:p<0.001; SD: Standart error mean

Histological analysis: Hemorrhoid induced control and experimental groups exhibited severe inflammation, edema and congestion in histological sections. Histomorphometric scores of lesions excised from rats receiving *M. sylvestris* cream were evaluated histomorphometrically and the recovery of the rectoanal area was found significantly better than that of the control group (p=0.032) (Fig. 1A). The *M. sylvestris* treated group presented similar histomorphometric healing scores with base cream and Pilex cream groups. Although scores belonging to Pilex cream treated group were lower than those of the control group, the difference was not statistically significant (Fig. 2).

Myeloperoxidase (MPO) Immunohistochemical Analysis: The myeloperoxidase immunoreactivity of inflammatory cells in control and experiment groups showed no statistically significant differences in immunohistochemical analysis (Fig. 1B, Fig. 3).




Figure 1. (A) Minimum, maximum and median values referring to experiment (topical administration) and control groups are depicted as box-plot graphs. Total score of topical administered *M. sylvestris* group was significantly less than that of the control group ($*p < 0.05$). (B) Minimum, maximum and median values referring to myeloperoxidase labelling in experiment (topical administration) and control groups are depicted as box-plot graphs. Note that the scores remained similar in both groups.




Figure 2. Photomicrographs of histological sections referring to topical applications are demonstrated. Edema and congestive vessels in mucosal and submucosal connective tissue are apparently observed in control group (A) and epithelial loss is clearly seen in vehicle control (base cream) administered group (B). Note the inflammatory cell infiltration in A, B, C, and a lesser amount in D. *Ep: Epithelium, Ct: Connective tissue, V: Vessel, Ed: Edema, L: Lumen, Haematoxylin-Eosin 100x*




Figure 3. Photomicrographs of myeloperoxidase labelled sections obtained from topical administered groups are demonstrated. Inflammatory cells (arrow) showing positive immune reactivity with myeloperoxidase are marked brown. Inflammatory cells labelled with immune peroxidase are obviously seen in mucosal and submucosal connective tissue in the control (A) and the vehicle control (base cream) administered groups (B). Inflammatory cells labelled with immune peroxidase are less in C, however the appearance of the inflammatory cells is observed close to D (reference control). *L: Lumen, 200x*.

DISCUSSION AND CONCLUSION

The effects of the *M. sylvestris* methanolic extract were examined by *in vivo* studies. According to ARC scores, TNF- α and VEGF levels, SOD enzyme inhibition and histopathological-immunohistochemical results, *M. sylvestris* methanolic extract exhibited greater biological activity than the control and vehicle control groups, close to reference control group. TNF- α , which is an indicator of inflammation, was significantly lower in the Malva treatment group, which is compared to all the control groups (Yarijani et al., 2018). VEGF, which regulates vascular permeability in mucosal tissue, was detected at low levels in examined rat serum in Malva extract treated group compared to the controls including reference (Bahramsoltani et al., 2014). SOD enzyme inhibition, which is related to lowering of oxidation level in cells, closely related to many pharmacological activities, was noteworthy in Malva extract applied group compared to the control groups (Saad et al., 2016). Although it was not statistically significant, the total histomorphometric scores of the lesions were lower than the control group. Edema and congestive vessels in mucosal and submucosal connective tissue were apparently observed in control group. Unlike the results obtained with hematoxylin staining, MPO immunohistochemical analysis results have not seen a significant activity. Considering the parameters in the serum, the results of MPO immunohistochemical analysis can be neglected. The effectiveness of the traditional usage has been proven on the scientific platform. It was found that the mallow, rich in mucilage and flavonoids, which contributed to healing of hemorrhoidal edema, vasoconstriction of the vessels and inflammation. It is thought that the effective mechanism might be caused by major phytochemicals flavonoids in the Malva methanolic extract, which show effects on anti-inflammatory pathway (Gasparetto et al., 2012). Further phytochemical studies were needed to clarify the actual antihemorrhoidal mechanism/s.

The various pharmaceutical formulations containing plant extracts is an important factor considering its suitability for commercial applications in pharmacy and cosmetics. Our future goal is to prepare novel antihemorrhoidal formulations.

ACKNOWLEDGMENTS

We would like to thank Gazi University Scientific Research Projects Unit for its project (02/2016-03 code number) fund support.

REFERENCES

1. Abascal, K., & Yarnell, E. (2005). Botanical treatments for hemorrhoids. *Altern Complement Ther*, 11(6), 285-289.
2. Azeemuddin, M., Viswanatha, G. L., Rafiq, M., Thippeswamy, A. H., Baig, M. R., Kavya, K. J., & Shyam, R. (2014). An improved experimental model of hemorrhoids in rats: evaluation of antihemorrhoidal activity of an herbal formulation. *ISRN Pharmacol*, 2014.
3. Bahramsoltani, R., Farzaei, M. H., & Rahimi, R. (2014). Medicinal plants and their natural components as future drugs for the treatment of burn wounds: an integrative review. *Arch Dermatol Res*, 306(7), 601-617.
4. Dey, Y. N., Wanjari, M. M., Kumar, D., Lomash, V., & Jadhav, A. D. (2016). Curative effect of *Amorphophallus paeoniifolius* tuber on experimental hemorrhoids in rats. *J Ethnopharmacol*, 192, 183-191.
5. Evans, W.C. (2002). A taxonomic approach to the study of medicinal plants and animal derived drugs. Trease and Evans Pharmacognosy, (15th ed). WB Saunders, Edinburgh, 33-35.
6. Everhart, J. E. (2008). Burden of digestive diseases in the United States Report. Retrieved from US.
7. Gasparetto, J. C., Martins, C. A. F., Hayashi, S. S., Otuky, M. F., & Pontarolo, R. (2012). Ethnobotanical and scientific aspects of *Malva sylvestris* L.: a millennial herbal medicine. *J Pharm Pharmacol*, 64(2), 172-189.
8. Gurel, E., Ustunova, S., Ergin, B., Tan, N., Caner, M., Tortum, O., Demirci-Tansel, C. (2013). Herbal haemorrhoidal cream for haemorrhoids. *Chin J Physiol*, 56(5), 253-262.
9. Krawisz, J. E., Sharon, P., & Stenson, W. F. (1984). Quantitative assay for acute intestinal inflammation based on myeloperoxidase activity: assessment of inflammation in rat and hamster models. *Gastroenterology*, 87(6), 1344-1350.
10. Leff, E. (1987). Hemorrhoids: Current approaches to an ancient problem. *Postgrad Med*, 82(7), 95-101.
11. Lohsiriwat, V. (2012). Hemorrhoids: from basic pathophysiology to clinical management. *World J Gastroenterol*, 18(17), 2009-2017.
12. Nishiki, K., Nishinaga, K., Kudoh, D., & Iwai, K. (1988). Croton oil-induced hemorrhoid model in rat: comparison of anti-inflammatory activity of diflucortolone valerate with other glucocorticoids. *Nihon yakurigaku zasshi. Folia Pharmacologica Japonica*, 92(4), 215-225.
13. Saad, A. B., Rjeibi, I., Brahmi, D., Smida, A., Ncib, S., Zouari, N., & Zourgui, L. (2016). Malva sylvestris extract protects upon lithium carbonate-induced kidney damages in male rat. *Biomed Pharmacother*, 84, 1099-1107.
14. Sakai, Y., & Matsukuma, S. (2002). CD34+ stromal cells and hyalinized vascular changes in the anal fibroepithelial polyps. *Histopathology*, 41(3), 230-235.
15. Yarijani, Z. M., Godini, A., Madani, S. H., & Najafi, H. (2018). Reduction of cisplatin-induced renal and hepatic side effects in rat through antioxidative and anti-inflammatory properties of *Malva sylvestris* L. extract. *Biomed Pharmacother*, 106, 1767-1774.

RADICAL SCAVENGING AND ANTI-BACTERIAL ACTIVITIES OF METHANOLIC EXTRACT FROM ALGERIAN FIR (*ABIES NUMIDICA*)

Bellil Inès*, Benouchene Djamila, Khelifi Douadi

Laboratoire de Génétique Biochimie et Biotechnologies Végétales, Faculté des Sciences de la Nature et de la Vie, Université Frères Mentouri Constantine 1, Constantine, Algérie.

*Corresponding author: bines07@yahoo.fr / ies.bellil@umc.edu.dz

Abstract: The objective of this study was the extraction of secondary metabolites from algerian fir leaves (AFL) and the evaluation of the anti-oxidant and the anti-bacterial activities of the crude extract. The air-dried powdered leaves were extracted with methanol solvent using Soxhlet apparatus. The extract was evaporated under reduced pressure by rotary vacuum evaporator type Buchi R-215 at 40°C. DPPH and ABTS assays were used to evaluate the anti-oxidant activity. The results showed that methanolic extract of algerian fir leaves by DPPH method can reduce radicals ($IC_{50}=18.08\pm 0.21$) lower than ABTS ($IC_{50}=23.51\pm 0.54$). The Anti-bacterial activity was also tested using the disc diffusion method against five bacterial strains; *Escherichia coli*, *Pseudomonas aeruginosa*, *Morganella morgani*, *Staphylococcus aureus* and *Bacillus subtilus*. The methanolic extract exhibited no anti-bacterial activity against Gram negative bacteria; *E. coli*, *P. aeruginosa* as well as *M. morganii*. However, it represented a moderate activity against Gram positive; *S. aureus* and *B. subtilus*.

Keywords : Algerian fir, leaves, methanolic extract, DPPH assay, ABTS assay, anti-bacterial activity.

INTRODUCTION

Secondary metabolites or phytochemicals are aromatic, non-nutritive compounds; they are an abundant and diverse group of molecules in plant kingdom. They have an important role for plant in the defense reaction to protect against abiotic stresses and biotic factors (Mazidet *al* ; 2011), for animal because they have a great ability to scavenge free radicals (Ali khani ; 2012). There is an increasing suggestions by considerable evidence that the free radicals induce oxidative damage to biomolecules (lipids, proteins and DNA) (Leonard *et al* ; 2003, Medini *et al* ; 2014) caused a degenerative diseases as cancer (Momina *et al* ; 2017), cardiovascular diseases and alzheimer (Brieger *et al* ; 2012). Also for food spoilage caused by several bacterial strains, nowadays, there is a marked interest in the food industry toward the development and manufactures of functional products. For that, there is a huge deal of attention toward search and discover a new molecules or products from natural resources (Khoddamet *al*, 2013) that have a beneficial effect comparing with synthetic one. Algeria by its geographical position represents a high biodiversity in plant kingdom major of them are endemic and a few are known about them for exemple : algerian fir (*Abies numidica*) (Quezel ; 1964). *A. numidica* is a conifer belonging to pinaceae family, an endemic tree solely to Algeria; it covers only an area of 2300 ha, found only in Babor's mounts, in the north of Sétif. It is very branchy tree, needles arranged around twigs (Bennadja *et al* ; 2012). It used in traditional medicine as an anti-inflammatory, to treat cataplasma and respiratory system problems. (Aitkaki *et al* ; 2013). The goal of this research is to investigate the radical scavenging ability of crude methanolic extract extracted from Algerian fir needles; furthermore to evaluate its anti-bacterial activity.

MATERIAL AND METHODS

Collection and preparation of the plant: Leaves of Algerian fir were collected from Constantine, they were harvested randomly from branches during the month of September 2018, after which they were dried for 15 days in the dark at room temperature. Then the sample were powdered using manual grinder IK1a 10 type and kept in the dry, dark place until they were used.

Extraction of secondary metabolites : Secondary metabolites are extracted as the method of Ramadan *et al* ; 2003 with slight modifications, with methanol using soxhlet apparatus for 2 hours. The extract was evaporated under reduced pressure by rotary vacuum evaporator type Buchi-R15 type at 40°C.

Biological activities:

***Anti-oxidant activities**

-DPPH radical scavenging assay: The DPPH free radical scavenging assay was tested as Blois, 1958 and Tel G *et al*, 2012 methods with some modifications. The different dilutions of the methanolic extract were prepared in methanol. 40µl of various sample concentrations was added to 160 µl of methanolic DPPH solution in 96-well microplate. Methanol with DPPH reagent mixture was used as control. After 30 min of incubation in the dark at room temperature, the absorbance of each solution was determined at 517 nm using a microplate reader. BHT, BHA were used as standards. Reduction of DPPH radical in percent (R %) was calculated in following way:

$$\% \text{ Inhibition} = [A_{\text{blank}} - A_{\text{sample}} / A_{\text{blank}}] * 100$$

Where : A_{blank} : is the absorbance of control reaction.

A_{sample} : is the absorbance of test sample.

Tests were carried out in triplicates. The IC₅₀ is the inhibition percent represents the half of free radicals (50%), was calculated from the graph of DPPH radical scavenging effect percent against extract concentration.

-ABTS⁺ radical scavenging assay (2,2'-azino-bis-(3-ethylbenzothiazoline-6-sulfonic acid): This anti-radical activity was measured using a cation decolorisation assay as described by Re *et al*, 1999. Briefly, ABTS⁺ solution was prepared by mixing of ABTS⁺ solution (7 mM) with potassium persulfate solution (2.45 mM). The mixture was left in the dark at room temperature for 16 h before use. This solution was diluted and was adjusted to obtain an absorbance of 0,70 ±0,02 at 734 nm. 160µl of ABTS⁺ was added to 40µl of sample at different concentrations. The mixture was incubated in the dark for 10 min at room temperature. The blank solution was methanol, ABTS⁺ solution without methanolic extract was used as control. The antioxidant standards were BHT and BHA. The experiment was performed in triplicate. The inhibition percent was calculated the same as described in DPPH radical assay.

***Anti-bacterial activity**

The anti-bacterial assay was carried out according to the Biondi *et al*, 1993 method with minor modifications, using disc diffusion, against 5 human pathogenic bacteria strains including Gram positive ; *Bacillus subtilis*, *Staphylococcus aureus* and gram negative ; *Escherichia coli*, *Pseudomonas aeruginosa* as well as *Morganella morganii*. The bacterial strains were first cultured on nutrient agar and were incubated at 37°C for 24 hours. Bacterial suspensions were spread on the surface of Mueller-Hinton agar plates. The sterile filter papers discs (6mm in diameter) were soaked with 10µl of methanol extract for the following concentrations (mg/ml) (100, 50, 25, 12.5, 6.25, 3.125, 1.562 W/V in DMSO). The discs were placed on the inoculated agar medium, kept at 4°C for 1h, and incubated at 37°C for 24h. The anti-bacterial activity was assessed by measuring the inhibition zone surrounding the discs. Each experiment was done in triplicate.

RESULTS

Anti-oxidants activities: The anti-oxidant activities of methanolic extract were tested using DPPH and ABTS assays. The results in **Table 1** showed that the methanolic extract obtained from extraction by soxhlet apparatus has a moderate DPPH and ABTS anti-oxidant activities versa to BHA and BHT standards.

Anti-bacterial activity : Results in **Table 2** revealed that methanolic extract has a mild inhibition zones for Gram (+ve) bacterial strains however, it exhibited no activity against gram(-ve) bacteria.

DISCUSSION AND CONCLUSION

Free radicals or reactive oxygene species are the products of normal cells metaboloism, represent the essential part of aerobic life (Shettar *et al* ; 2015), but the imbalance between these molecules and body system defense, or the excess of them lead to many diseases (Sasikumar ; 2014). For that a lot of works have been done to look for new products from natural substances that have the ability to scavenge free

radicals and stop this chain reaction, in addition this group of products is well known for its ability to inhibit and suppress the growth of several microbial strains (Nascimento *et al* ; 2000).

Natural anti-oxidants have an important function in the enhancement of body system defense and help to prevent many diseases as cancer and diabetes (Kenjal *et al* : 2007). Due to the complex structure of secondary metabolites, the examination of its anti-oxidant activities cannot be tested by a single assay, herein, in this study two methods were used to evaluate the anti oxidant ability of crude methanolic extract from *A. numidica* leaves, DPPH and ABTS assays. DPPH is a stable free radical which produce a violet solution in methanol, changed to yellow color when it is reduced in the presence of an anti-oxidant molecules (Chang *et al* ; 2002). In the present study anti-oxidant activity of *A.numidica* leaves was evaluated using the crude methanol extract obtained by soxhlet extraction and was compared with standards (BHA, BHT). Results obtained from chemical assay tested showed that standards had higher scavenging activity than the extract, in comparison with results reported by Ghadbane *et al* ; 2016 revealed that DPPH anti-oxidant assay of methanol extract by maceration showed a potential radical scavenging assay. ABTS assay is a decolorization method in the presence of an anti-oxidant agents. Results obtained also showed that standards have a potential anti-oxidant ability compared with methanolic extract. These results might be due to the extraction by soxhlet that degrade and destroy molecules or phytochemicals by heat (thermal degradation) (Martino *et al* ; 2006).

Results for anti-bacterial activity showed that methanolic extract have an inhibition zone against Gram (+ve) bacterial strains however it exhibited no anti-bacterial effect against Gram (-ve) bacterial strains tested that was resistant to methanolic extract, this result is coinciding with the statement given by Ghadbane *et al* ; 2016. *E.coli* and *P. aeruginosa* well known to be multi-resistant strains. Gram (+ve) bacteria were sensitive to the spectrum-broad of phenolic compounds. This is an indication that these group of products (secondary metabolites) has different mode of action or that the metabolism of some micro-organisms are better able to stop the effect of these molecules or adapt to it. Also plant extract is less effective on gram (-ve) bacteria might due to the degradation of phenolic compounds by heat of soxhlet.

In conclusion, the crude extract showed a moderate DPPH and ABTS anti-oxidant activities due to the presence of secondary metabolites, further it exhibit an inhibition activity against Gram (+ve) bacteria: *S.aureus* and *B. subtilus* which suggested that have a potential role in food spoilage. The results should be encouraged in the future by *in vivo* studies in order to integrate this extract of plant in pharmaceutical, cosmetic and diet.

Table 1. Antioxidant activities of methanolic extract extracted from *A.numidica* leaves

Extract (0,25µg/µl)	IC ₅₀ (µg/µl) DPPH assay*	IC ₅₀ (µg/µl) ABTS assay*
Methanolic extract	18,08±0,21	23,51±0,54
BHA	15,74±0,47	7,54±0,67
BHT	6,55±0,59	1,55±0,26

*Values were expressed as means±SD. Of three parallel measurements, (p<0.05). SD: Standard deviation, BHT: butyl hydroxyltoluene, BHA: Butyl hydroxyl anisole, DPPH: 2,2-diphenyl-picrylhydrazyl, ABTS: 2,2'-azino-bis-(3-ethylbenzothiazoline-6-sulfonic acid, IC₅₀: Half inhibitory concentration.

Table 2. Anti-bacterial activity of methanolic extract of *A. numidica* leaves

Inhibition zones				
Gram (-ve) bacteria			Gram (+ive) bacteria	
<i>E. coli</i> ATCC 25922	<i>P. aeruginosa</i> ATCC 27853	<i>M. morgani</i> ATCC 25830	<i>S. aureus</i> ATCC 43300	<i>B. subtilus</i> ATCC 6633
0,0±0,0	0,0±0,0	0,0±0,0	8,33±0,57	7,66±0,57

Data are means of three replicates (n = 3) ± standard error

REFERENCES

- [1] Ali Khan,R., Rashid Khan, M., Sahreen,S., and Ahmed M.,2012. Evaluation of phenolic contents and antioxidant activity of various solvent extracts of *Sonchus asper* (L.) Hill. Chemistry Central Journal. 6(12), 1-7.
- [2] Bennadja, S., and Tlili ait kaki, Y., 2012. The Fir of Numidia: a Threatened Species. Journal of Forestry Faculty of Kastamonu University, 12 (31).
- [3] Biondi, D., Cianci, P., Gerad, C., and Rubertof, G., 1993. Antimicrobial Activity and Chemical Composition of Essential Oils from Sicilian Aromatic Plants. Flavour and Fragrance Journal, 8, 331-337.
- [4] Blois, M. S., 1958. Antioxidant determinations by the use of a stable Free Radical. Nature, 4617 (181), 1119-1200.
- [5] Briegera, K., Schiavonea, S., Francis, J. M. J., and Krause K. H., 2012. Reactive oxygen species: from health to disease. Swiss Medical Weekly, 142, 1-14.
- [6] Choi, C. W., Kim, S. C., Hwang, S. S., Choi, B. K., Ahn, H. J., Lee, M. Y., Park, S. H., and Kim, S. K., 2002. Plant science, 163, 1161-1168.
- [7] Ghadbane, M., Bounar, R., Khellaf, R., Medjekal, S., Belhadj, H., Benderradji, L., Smaili, T., and Harzallah, D., 2016. Antioxidant and antimicrobial activities of endemic tree *Abies numidica* growing in babor mountains from Algeria. Global J Res. Med. Plants & Indigen. Med, 5, 19–28.
- [8] Kanjale, R. D., Shah, R. K., and Sathay, S. S., 2007. Anti-stress and anti-oxidant effects of roots of *Chlorophytum borivilium* (Santa pau and fernandes). Indian Journal of Experimental Biology, 45 ,974-979.
- [9] Khoddami, A., Wilkes, M. A., and Roberts, T. H., 2013. Techniques for Analysis of Plant Phenolic Compounds. Journal of Molecules, 18, 2328-2375.
- [10] Leonard, S. S., Xia, C., Jiang, H. S., Tinefelt, B., Klandorf, H., Harris, G. K., and Shi, X., 2003. Resveratrol scavenges reactive oxygen species and effects radical-induced cellular responses. Biochemical and Biophysical Research Communications, 309, 1017–1026.
- [11] Martino, E., Ramaiola, I., Urbano, M., Bracco, F., and Collina, S., 2006. Microwave-assisted extraction of coumarin and related compounds from *Melilotus officinalis* (L.) Pallas as an alternative to Soxhlet and ultrasound-assisted extraction. Journal of Chromatography A, 1125, 147–151.
- [12] Mazid, M., Khan, T. A., and Mohammad, F., 2011. Role of secondary metabolites in defense mechanisms of plants. Biology and Medicine, 3 (2), 232-249.
- [13] Medini, F., Fellah, H., Ksouri, R., and Abdelly C., 2014. Total phenolic, flavonoid and tannin contents and antioxidant and antimicrobial activities of organic extracts of shoots of the plant *Limonium delicatulum*. Journal of Taibah University for Science, 216–224.
- [14] Momina, H., Abbas, M., Munir, F., Qasim Hayat, M., Keyani, R., and Amir, R., 2017. Potential of plant flavonoids in pharmaceuticals and nutraceuticals. J Biomol Biochem., 1(1), 12-17.
- [15] Nascimento, G. G. F., Locatelli, J., Freitas, P. C., Silva, G. L., 2000. antibacterial activity of plant extracts and phytochemicals on antibiotic-resistant bacteria. Brazilian Journal of Microbiology, 31, 247-256.
- [16] Quezel, P., 1964. L'endémisme dans la flore de l'Algérie, 138-149.
- [17] Ramadan, M. F., and Morsel, T. J., 2003. Analysis of glycolipids from black cumin (*Nigella sativa* L), coriander (*Coriandrum sativum* L.) and niger (*Guizotia abyssinica* cass.) oil seeds. Food Chemistry, 80, 197-204.
- [18] Re, R., Pellegrini, N., Proteggente, A., Pannala, A., Yang, M., and Rice-evans Catherine., 1998. antioxidant activity applying an improved abts radical cation decolorization assay. Free Radical Biology and Medicin, 26, 1231–1237.
- [19] Sasikumar, V., and Kalaisezhien, P., 2014. Evaluation of Free Radical Scavenging Activity of Various Leaf Extracts from *Kedrostis foetidissima* (Jacq.) Cogn. Biochemistry and Analytical Biochemistry, 3(2), 1-7.
- [20] Shettar, A. K., Kotresha, K., Kaliwal, B. B., and Vedamurthy A. B., 2015. Evaluation of in vitro antioxidant and anti-inflammatory activities of *Ximenia americana* extracts. Asian Pacific Journal of Tropical Disease, 5(11), 918-923.
- [21] Singleton, V. L., and Rossi J.A.J., 1965. Colorimetry of total phenolics with phosphomolybdic-phosphotungstic acid reagents. Amer. J. Enol. Viticult, 16, 144-58.
- [22] Tel, G., Apaydin, M., Duru, M. E., and Öztürk, M., 2012. Antioxidant and cholinesterase inhibition activities of three *Tricholoma* species with total phenolic and flavonoid contents : The edible Mushrooms from Anatolia. Food Anal Methods, 5, 495-504.
- [23] Tlili-Ait Kaki, Y., Bennadja, S., and Chefrour, A., 2013. Revalorisation d'une essence endémique: le sapin de Numidie (*Abies numidica*). Flora Mediterranea, 23, 123-129.

ETHNOMEDICINAL AND AROMATIC AND AROMATIC SALT TOLERANT PLANTS USE VALUE IN KHEWRA SALT MINE, PUNJAB, PAKISTAN

Muhammad Zafar & Moona Nazish

¹*Department of Plant Sciences, Quaid-i-Azam University, Islamabad, Pakistan*

*Corresponding author e-mail: zafar@qau.edu.pk

Abstract

It is the principal quantitative investigation of halophytic vegetation with their customary therapeutic and fragrant uses in Khewra salt mine, Punjab, Pakistan. In this investigation, the reliance of nearby individuals on halophytes gives pragmatic proof about conventional social insurance frameworks. An aggregate of 151 respondents including 13 customary healers were talked with utilizing quick evaluation strategy. The reported information was mapped and examined quantitatively and in GIS too. The mineral portrayal of the dirt was completed utilizing X-beam diffraction (XRD) strategy. Ethno-restorative information was gathered for 38 salt tolerant plant species having a place with 23 families. Results showed the Fabaceae as much of the time referred to plant family (5 species) and *Acacia nilotica* (L.) Delile as most referred to plant species. Herbs (half) were the prevailing living thing and leaf (37%) was the generally utilized plant section (24 reports). The general significance of the halophytes and homogeneity of the respondents' learning was dictated by utilizing source agreement factor (ICF) and use esteem (UV). The most noteworthy ICF (0.62) was found for gastrointestinal illness class. Relative recurrence of reference and UV ranges from 0.08 to 0.3 and 0.22 to 0.48 separately. The XRD examination demonstrated the copper sulfide nature of soil which is the pointer of halophytes dispersion in the investigation territory. Lion's share of the halophytes has been seen to have solid pharmacological and sweet-smelling proof. It is the main recorded provide details regarding halophytes utilized for therapeutic purposes in this locale. The discoveries of this study will be valuable for the safeguarding of Ethnomedicinal and sweet-smelling legacy and for future pharmacological, sweet-smelling and phytochemical screening of salt tolerant plants to discover their potential for normal medication improvement.

Keywords: Halophytes, Medicinal, aromatic, GIS, X-ray diffraction, Khewra Salt Mines, Phytochemicals, Pakistan.

Acknowledgements: Authors acknowledge Pakistan Science Foundation (PSF) Pakistan under letter No. PSF/P&D/TG-ND (384)19. by providing travel grant for oral presentation and to attend Symposium (MESMAP-2019).

INTRODUCTION

Ethnobotanical appraisals of therapeutic plants give information about their customary uses, protection, indigenous networks, and vital for the natural medications improvement (Heinrich and Gibbons, 2001; Vitalini et al., 2013). These reviews are pivotal to illuminate the essential plant species, principally to grow new rough medications (Cox, 2000; Leonti et al., 2002). Indigenous learning has been hold by the customary social orders taken as principal to their wellbeing and social premiums (Cotton, 1996; Getahun, 1976). In addition, Ethnomedicinal and fragrant examinations bolster the network in sociocultural and financial setting and vital for the preservation of worldwide legacy with respect to plants (Sanz-Biset et al., 2009).

The therapeutic plants are as yet perceived as fundamentally medicinal services in immature networks in view of their more viability and social inclinations while in created nations it give elective for human services (Svarstad and Dhillon, 2000). The evaluated report demonstrated the unwavering quality of 70-80% individuals on the ethnobotanical and ethno-restorative employments of therapeutic plants world generally (Muthu et al., 2006). A harsh gauge revealed 35-000 to 75,000 restorative plants giving the premise to social insurance frameworks around the world (Farnsworth and Soejarto, 1991). Natural pharmaceuticals assume huge job in fix of number of human afflictions (Rehecho et al., 2011). The ethnobotanical examines have increased huge consideration among scientific networks concerning therapeutic plants use (Tripathi et al., 2017). Various Ethnomedicinal and sweet-smelling reviews have

been changed over into preservation and medicinal services programs (Quamar and Bera, 2014). The methods for plants use are differing in different networks far and wide (Abebe et al., 2003; Kassaye et al., 2006).

Numerous medications accessible in market are engineered analogs of disconnected mixes from plants (Amjad et al., 2017). Customary phytotherapies are still prevalently polished in numerous country networks passed on from one age to next inspite the assortment of clinical operators created by the pharmaceutical businesses (Abbasi et al., 2010). The broad social esteem considers and their methodical documentation have been put on high need to feature the pharmaceutical capability of plants and to investigate their new properties in the previous decades. The customary investigations of plants assume vital job in cultivating, nutraceutical and pharmaceutical enterprises (Cox, 2000; Idrisi et al., 2010). The cooperations of number of synthetic constituents in restorative plants cause amicability and collaboration in human body and recuperate the body tissues logically. The medication ventures have created prescriptions utilizing confined substance mixes from restorative plants which deliver physiological reaction (Shinwari and Gilani, 2003). So as to find out the remedial employments of plant based medications, it is important to lead the pharmacological and pharmacognostic examines and to investigate their utilizations (Ahmad et al., 2009)

Pakistan has assortment of atmosphere zones and topographic locales and is supplied with decent variety of therapeutic plants (Yaseen et al., 2015). The parochial information and home grown use mirror the diverse common assets present in various parts of the nation. In Pakistan, around 400-600 therapeutic plants has been accounted for having use in customary medicinal services rehearses (Ahmad et al., 2014; Shinwari and Qaisar, 2011). Qureshi et al., (2007) uncovered that 84% masses of Pakistan depends upon standard drugs for the fix of various issue. Past review portrayed that around 60,000 customary healers in remote territories use restorative plants against a few infirmities (Rehman et al., 2017). The indigenous home grown meds depend on the properties of restorative plants (Bartam, 1995). Conventional Yunani restorative framework is drilled famously among the provincial populace. Yunani therapeutic framework was embraced and archived by Muslim researchers in the glorious time of Islamic human advancement and they rehearsed it for a long time (Khan et al., 2012). In Pakistan, indigenous information is diminishing step by step because of quick land corruption, present day culture presentation, urban improvement, and war impacts yet in rustic networks the customary learning about restorative plants makes due because of its change from one age to people to come (Figueiredo et al., 1993; Kayani et al., 2015; Manandhar, 1995; Tabuti et al., 2003). Different Ethnomedicinal and sweet-smelling contemplates led in Pakistan revealed the long time routine with regards to therapeutic plants a few infections including extraordinary number of halophytes against (Adnan et al., 2010; Akhtar et al., 2013; Khan et al., 2013; Qasim et al., 2010; Shah and Hussain, 2012; Shinwari, 2010).

In excess of 400 types of halophytes are accounted for from Pakistan (Khan and Qaiser, 2006). The saltiness of soil is taken as limitation for the development of plants as about 6% of the world aggregate land region is salt affected (Khan et al., 2011). The 6.30 million hectares of Pakistan's aggregate land is salt influenced and 0.45 hectares of this aggregate saline land is available in Punjab. The broad review gave the wide record on the monetary uses and circulation of halophytes in Pakistan with rich decent variety of halophytes from Punjab because of differing biological living spaces (Khan and Qaiser, 2006). Ethnobotanical contemplates have been done wherever scale in the Punjab domain (Arshad et al., 2011; Ikram et al., 2014; Mahmood et al., 2013; Parvaiz, 2014; Qureshi et al., 2009; Sardar et al., 2015) which furthermore demonstrates the significant number of halophytes use in indigenous social orders. Halophytes are accounted for as underutilized source (Lieth and Hamdy, 1999) which verify their centrality for subsistence and their job in nearby social legacy. On the opposite side, the rising enthusiasm for halophytes demonstrates their gigantic potential (Khan et al., 2009).

Sadly, the halophytic greenery of a few sections of Punjab territory, for example, Khewra is unexplored with respect to the ethnobotanical investigations of restorative plant species. Khewra salt mine is situated in the Khewra city and is among the 5 salt mines of Pakistan and it is the Pakistan's old and biggest salt mine. In this angle, the present overview can be taken as the primary review in the zone as before the Khewra salt mine has not been considered for ethno-therapeutic examination. The investigation region has particular geology with incredible assorted variety of halophytes. The neighborhood individuals of

Khewra have been utilizing salt tolerant plants with long time experienced strategies and amount for treating different sicknesses since ages. The elderly individuals want to utilize restorative plants against a few illnesses and have adequate learning with respect to Ethnomedicinal and sweet-smelling rehearses. There is absence of essential conveniences, instruction and human services offices for the local individuals; in this way, they depend on the use of restorative halophytic vegetation in customary route as an essential wellspring of drug and as their domesticated animals.

The GIS and vegetation system together with PC based robotized field mapping methodology have been key gadgets for vegetation mapping (Ismail, 2010). On the opposite side, X-beam diffraction (XRD) is a valuable nondestructive strategy for the portrayal of crystalline materials (Bunaciu et al., 2015) which is utilized to abuse minerals recognizable proof in the dirt of study region. Halophytes got little consideration notwithstanding of having gigantic potential as they have capacity to blossom with poor soil conditions. The re-development of worry in the utilization of halophytes as restorative plants in social insurance rehearses has been energized because of the symptoms and increasing expense of engineered drugs. Besides, the conventional learning in the investigation region is draining because of deforestation and numbness with respect to ethnobotanical documentation before. Remembering every one of these issues, it was felt important to record the indigenous information with respect to salt tolerant plant species. The reason for the present study is the request and documentation of the assorted variety and ethno-therapeutic information of halophytes to depict the status of current ethnobotanical data among associates of novel ages, to get to the component of traditional getting the hang of using quantitative records, to find the association between soil minerals and halophytic vegetation by performing XRD, and to make the vegetation layout GIS to portray the varying scattering of halophytes in the examination locale.

MATERIALS AND METHODS

Geo-ethnographical diagram: In Pakistan, Khewra salt mine is the subdivision of Jhelum locale in region Punjab. The study was conducted from March to October 2018 regarding plant collection and data compilation. It lies 102 km far from the Jhelum. It is situated at 32° 38' N and at 73° 00' E. The underground piece of salt mine spread on the region of 110 km². It is the Pakistan's first and world's second biggest salt mine. It lies in the mountains of salt scope of Punjab. The half of salt is separated utilizing old Room and Pillar mining techniques while half of salt is left as columns to hold mountains. There is a distinction in temperature and mugginess because of the blasts and ignition motors. The nearby individuals use Punjabi dialect chiefly in discourse. The investigation zone incorporates sub-muggy zone, uneven and subtropical semi-bone-dry zone.

The region of Khewra salt mine has expansive number of minerals and have rich assorted variety of salt tolerant plants. Khewra salt dig is the site for travelers. Out of 19 stories, 6th one is intended for sightseers in whom salt made mosque, saline solution lakes, and Minar-e-Pakistan is available. The nearby individuals have adequate wellbeing offices as number of private facilities and single government healing facility however the vast majority of the general population still depend on standard medications due to ordinary tradition that the development people support to use indigenous standard techniques for the treatment of various issue instead of later remedial system. Agribusiness is the essential wellspring of benefit for inhabitants as half of the total tenants of the examination region are associated with agriculture. The money related conditions of the examination locale can end up being better unequivocally by engaging the advancement of salt tolerant therapeutic plants.

Ethnobotanical examination: The field work was directed between February 2017 and October 2018 after conventional methodologies. Ethno-therapeutic information was gathered utilizing distinctive methods like semi-organized discourses, open-finished meetings and field understandings (Martin, 1995). A fast evaluation approach and survey was utilized to record the therapeutic employments of halophytes. This strategy is viewed as successful in ethnobotanical examinations since it go about as an extension between formal overviews and network support. The survey contained data on the demography of nearby respondents and in addition on their sexual orientation, instruction, experience and age. In addition, it incorporated the plant nearby name, society formulas, readiness techniques, parts utilized, usage mode, and the illnesses treated with halophytic therapeutic plants.

One fifty-one individuals from network including customary healers, agriculturists and ladies took an interest in this review. These respondents included 13 conventional wellbeing experts, 83 men, and 68 ladies. A Punjabi talking individual was locked in as interpreter to record data precisely that would be lost amid meetings and translation. The approval of salt tolerant plant for restorative use was made just when at least two sources concurred to a similar use.

Recognizable proof and safeguarding of Halophytes: Five examples of each salt tolerant plant species was gathered from study zone. The open writing and Flora of Pakistan (<http://www.efloras.org>) was utilized to recognize the gathered halophytes. The halophytes were additionally contrasted and herbarium examples in Herbarium of Pakistan (ISL), Quaid-I-Azam University Islamabad for their recognizable proof. The Plant List (www.theplantlist.org) and Medicinal plant names administrations (www.kew.org/mpns) were counseled for ordered and organic names verification. The salt tolerant therapeutic plants were mounted on standard herbarium sheets subsequent to squeezing, drying, and treatment with mercuric chloride and ethanol. The voucher numbers were appointed to gathered plant examples and put in the Pakistan Herbarium (ISL), Quaid-I-Azam University Islamabad.

Information examination: Various quantitative files, for example, relative recurrence of reference (RFC), use esteem (UV) and witness agreement factor (ICF) was connected on ethno-restorative information to break down it quantitatively.

Source Consensus Factor (ICF): The ICF was connected on ethnobotanical information utilizing given recipe (Logan, 1986; Tabuti et al., 2003; Teklehaymanot, 2009):

$$ICF = (Nur - Nt) / (Nur - 1)$$

Where Nur demonstrates the utilization reports in number for a distinct disease class and Nt shows the taxa in number for the infirmity classification. ICF depict the agreement of data on therapeutic plant species use for an explicit issue (Canales et al., 2005). The estimation of ICF esteem fluctuates from 0 to 1. The most noteworthy ICF evaluation demonstrates the fix of number of different illnesses utilizing taxa educated by vast number of respondents. Then again, low ICF appraisal demonstrates the fix of couple of various illnesses utilizing taxa arbitrarily gathered from respondents having absence of information about plant species use (Abu-Irmaileh and Afifi, 2003; Gazzaneo et al., 2005; Kloutsos et al., 2001; Teklehaymanot, 2009).

Use esteem (UV): The UV was connected in ethnobotanical information as appeared in writing utilizing standard strategy (Phillips et al., 1994; Šavikin et al., 2013):

$$UV = U/n$$

Where U demonstrates the utilization reports with aggregate number for a notice plant animal varieties while 'n' demonstrates the aggregate number of respondents addressed for a notice plant animal varieties. In the event that UV is near 1, the esteem will be high with the sign of many use reports and plant species essentialness among respondents. In the event that UV is near zero, the esteem will be low with the sign of notice plant species with few use reports.

Relative Frequency Citation (RFC): The RFC was connected on ethno-restorative information which shows the nearby essentialness of therapeutic halophytes utilizing following equation (Tardío and Pardo-de-Santayana, 2008; Vitalini et al., 2013):

$$RFC = FC/N \quad (0 < RFC < 1)$$

Where FC shows number of sources who reported the use of species and N is the total number of observers who appreciated the examination. High RFC regard shows the obvious nature of a plant creature assortments among the respondents. In case respondents see the plant species as supportive, the

RFC regard will be 1 for notice plant species while if no respondent give information about the plant species use, the RFC will be zero for notice plant species (Medeiros et al., 2011).

Geographic data framework mapping: In geographic examination, the method for showing information on the guide is a pivotal advance. A GIS is an electronic database intended for spatial information the executives. The halophytes were gathered amid field review utilizing GPS. The information inspecting and mapping is made through GIS. The choice in the GIS database for the structure of fundamental organizer and subfolder was chosen. The recorded halophytes later on were moved into Arc GIS programming 10.5 for creating halophytes mapping framework.

X-Ray Diffraction: X-beam diffraction is utilized for the distinguishing proof of crystalline strong and to decide precious stone introduction and other basic parameters (Bish et al., 2014). It gives data as pinnacles delivered by productive impedence from a cross section planes in test utilizing monochromatic light emission beams dissipated at various edges. So as to perform XRD, salt example from study region is crush utilizing mechanical pounding to go through 45 µm strainer. The example in powder shape is cautiously arranged to stay away from spottiness, changes in force, and to acquire molecule size and introduction. The unreasonable pounding results in grid mutilation. The got outcomes were broke down utilizing Origin and High Score software's.

Moral endorsement/Intellectual property assentment: The present investigation was directed with the full assent of members. Formed Prior taught consent' (PIC) was gotten from the close-by systems which consolidate the benefit of an indigenous system to give or hold its consent to the proposed research adventure that may impact the grounds, resources, ordinary learning and conventions that they by and large have, include or for the most part use. The verbal understanding was additionally made that this review will be utilized just to illuminate the network and Khewra, Punjab on the salt tolerant plants and their restorative uses and recorded information won't be utilized for business purposes. The morals board of Quaid-I-Azam University, Islamabad allowed moral endorsement for the examination.

RESULTS AND DISCUSSION

Socio-statistic data: One fifty-one respondents including laypeople, cultivators and customary healers between the 18-80 years of ages took part in the study. Regarding sexual orientation, 55% of the respondents were men and 45.03% were ladies. The more respondents were guys because of the impediments on females in the investigation territory. The ladies were not taking into account discussion with obscure guys or outsiders. The female respondents have been seen to have critical learning about the home grown arrangements utilizing halophytes. The provincial ladies have noteworthy job in the conveyance of customary social insurance framework because of her culinary exercises and home garden development additionally (D'Avigdor et al., 2014).

The inconstancy in respondent's age has critical effect on the customary restorative learning of salt tolerant plants in Khewra. Based on age, respondents were partitioned into 6 noteworthy companions. The 73% of the witnesses were over 40 years, while 27% were beneath the 40 years. This suggests the customary therapeutic learning of salt tolerant plants is declining step by step among the youthful age in the investigation region. A few different examinations have detailed that the decrease in customary information among youthful age isn't restricted to think about zone or halophytes, yet in different indigenous populaces or provincial networks the conventional learning is declining step by step among young people (Giday et al., 2009; Hong et al., 2015; Rehman et al., 2017). Most of the respondents (28.47%) were uneducated. The unskilled sources shared increasingly conventional information of therapeutic halophytes when contrasted with instructed witnesses (24.50%). This may be because of the urbanization and utilization of allopathic meds. Comparable discoveries were additionally announced by Umair et al., (2017), Kadir et al., (2012) and Hayta et al., (2014). The conventional healers are the caretakers of the therapeutic information in the investigation region. This demonstrates laypeople can treat minor sicknesses, for example, skin issue, fever, and migraine while treatment of significant illnesses, for example, Hepatitis, fruitlessness, and diabetes are as yet limited to cultivator and customary healers. The dominant part of customary healers was noted to have 5-10 years' training (2.64%). [Table 1]

Diversity of halophytes and their Life form: This investigation establishes thirty-eight salt tolerant plant species dispersed among twenty-three families which is the sign of rich assorted variety of halophytes. The herbal names of halophytes with their family name, neighborhood names, part(s) utilized, society formulas, strategy for usage, living thing, development shape and remedial uses alongside quantitative information examination are displayed in Table 2. Fabaceae contributed the most taxa with five species while different families have species under five in number (Figure 2). Fabaceae is prevailing family in light of the fact that a large portion of the individuals from this family are developed for sustenance and the nearby individuals utilized them for the fix of different diseases. Fabaceae is among the transcendent families on the planet (Ali, 2008). Different examinations additionally demonstrate the comparative outcomes in term of overwhelming family (Cheikhoussef et al., 2011; Mahwasane et al., 2013; Qayum et al., 2016; Wagh and Jain, 2015). Most of the prevailing halophytes were gathered from wild (80%) trailed by developed (20%) (Figure 3). A few ethnobotanical examinations additionally revealed the restorative proficiency of wild plant species (Hong et al., 2015; Soukand et al., 2017). The WHO announced that 80% of Asian populace relies upon wild therapeutic plant species because of their simple access, adequacy and social recognition. The examination dependent on remedial utilizations demonstrated numerous restorative employments of single species. The developing worry in the present period about restoratively critical plant species is their over-misuse and intemperate utilization (Anon, 2009).

The prevailing halophytes were enduring plants (65%) trailed by yearly (31%) and biennial plants (2%). The explanation for the most perpetual plants is the flighty and uncommon precipitation in the dry and semi-bone-dry locales (Qasim et al., 2010). The recorded overwhelming living thing was herbs (half) (Figure 4). The purposes behind herbs high rate are their simple openness in assortment of living spaces as in fields, woodland territories, rocks and prairies (Ayyanar and Ignacimuthu, 2005; Sanz-Biset et al., 2009; Uniyal et al., 2006) and their adequacy in the fix of a few sicknesses because of the nearness of dynamic pharmacological constituents and straightforwardness in gathering accumulation (Adnan et al., 2012). Another reason is the straightforwardness in home grown arrangements (Arshad et al., 2011; Lulekal et al., 2013) and have a noteworthy part in keeping up processing and digestion (George and Nimmi, 2011). Herbaceous living thing is the regular broad environmental wonders around the world (Jan et al., 2011). Jan et al., (2017) affirm the discoveries of this examination. In the examination of the organic range of halophytic vegetation following Raunkiaer (Akman and Ketenoglu, 1992), it was discovered that 47% of the halophytes were phanerophytes. The strength of phanerophytes over other living things gives the thought regarding their better reaction to the topographic difference.

[Table 2]

[Figure 2]

[Figure 3]

[Figure 4]

The present examination moreover selected the most-a significant part of the time referred to halophytes with something close to 55 references or progressively (Table 2). The regularly alluded to halophytes are utilized by the broad number of adjacent individuals especially development people because of their simple openness and normal presence. The expansive number of respondents was likewise found to have belief in the viability of usage of halophytes customarily in light of the fact that they support to use the home grown arrangements over manufactured medications in view of different reactions and their surprising expense.

Plant parts used and methods of utilization: In this examination, nearby individuals use a few sections of therapeutic halophytes in rough home grown medication arrangements; consequently, different society formulas were recorded. For the most part leaves (37%) was the real plant part utilized by neighborhood networks for a few infections because of simplicity in culling (Kadir et al., 2012), their most elevated corrective powder and high auxiliary metabolites (Verpoorte, 1998; Verpoorte and Memelink, 2002) (Figure 5). In various examinations at various spots, ethnic peoples were found to exhibit practically identical results if there ought to emerge an event of plant parts used for the arranging

of home developed medications as in the present examination (Ahmed et al., 2015; Dolatkhahi et al., 2014; Giday et al., 2009; González et al., 2010; Mahishi et al., 2005; Mowobi et al., 2016; Parvaiz, 2014; Srithi et al., 2009). The products of the soil of the plants after leaves are fundamental for the treatment of different sicknesses since they contain high dynamic substance of bioactive mixes (Moore and Thomas, 1994). Other plant parts detailed for treatment of a few illnesses are given in Table 2. [Figure 5]

Mode of administration and utilization: The strategy utilized for medication planning and organization differ as indicated by the illness type. In this review, 80 home grown arrangements are recorded. Greater part of the plans (59%) were conveyed orally in ailments classifications. In couple of diseases, the details utilizing salt tolerant therapeutic plants (41%) were topically directed (Figure 6). Liquor, water and couple of added substances like salt, oil and nectar were utilized regularly in the cures readiness. The vast majority of the cures arrangements included blending of various plant parts or distinctive plant species while the utilization of single plant or a solitary piece of the plant was less experienced in the enquired territory. The oral course was likewise recorded as the significant method of organization in prior examinations (Brandão et al., 2012; Chekole et al., 2015; Kadir et al., 2012; Mood, 2008; Nadembega et al., 2011; Poonam and Singh, 2009). The oral organization of prescription is constantly powerful for inside medical issues (Shinwari et al., 2010).

The most utilized arrangement technique was glue (25%) trailed by decoction (24%) (Figure 7). These discoveries demonstrate that for the most part individuals in the examination territory want to utilize glue of different plant parts either through oral or topical course of organization to mend different afflictions. Customary healers of the investigation region educated about the numerous arrangements with different plants parts because of high viability for relieving infections. Decoction is set up by bubbling diverse parts of the plants until the required volume of water is gotten. Warming or bubbling velocity up different natural responses which deliver a few valuable and dynamic mixes (Chen et al., 2008). Indigenous individuals favored to use crisp plant parts particularly to get their concentrate and squeeze by decoction or imbue (extraction of flavors or synthetic mixes by suspending the plant parts in a dissolvable, for example, water or liquor after some time) because of restorative adequacy and don't store and dry diverse parts of the plant. Sweet fixings, for example, sugar and nectar were included the home grown medications to lessen their severe taste particularly for kids (Balangcod and Balangcod, 2011). It was watched additionally that the respondents utilize different portion of medications relying upon the malady and age.

[Figure 6]

[Figure 7]

Quantitative analysis

Relative frequency of citation (RFC): The estimation of RFC portrays the successive utilization of halophytes with therapeutic esteem and the essentialness of that plant species with reference to respondents. In the present investigation, the most astounding RFC esteem alludes to the need and use of plant species announced by respondents. The estimation of RFC fluctuates from 0.22 to 0.48. The therapeutic halophytes having most noteworthy RFC were *Acacia nilotica* (L.) Delile and *Abelmoschus moschatus* Medik. *Acacia nilotica* (L.) has the most elevated RFC esteem as a result of its therapeutic regard yet it was also observed that people moreover use it for fuel, timber and advancement purposes; along these lines, it is an exceptional plant of the district. Most referred to plant species are most normal and valuable with respect to restorative purposes among the masses (Samoisy, et al., 2015). Other imperative halophyte was *Abelmoschus moschatus* Medik. with high RFC esteem because of its simple openness to respondents and straightforwardness in development in fields and home patio nurseries. The halophytes with minimum RFC values were *Podocarpus latifolius* (Thunb.) R.Br. ex Mirb. Furthermore, *Verbascum thapsus* L. The ethno-restorative investigations directed in other place additionally announced some plant species with most astounding RFC values like the present examination (Kankara et al., 2015). The plant species with high RFC esteem ought to be exposed to organic, phramcological and phytochemical thinks about (Mukherjee et al., 2012).

Use value (UV): UV list has been utilized for the quantitative examination of information and to affirm the general criticalness of species or family implied for populace (Vendruscolo and Mentz, 2006). The overall significance of salt tolerant plant species utilized by neighborhood individuals for drugs was validated by assessing the information utilizing use esteem. In this investigation, UV esteem ranges from 0.08 to 0.3. High use esteems demonstrate the huge number of reports for plants by respondents while plants with less use esteem show the less use reports. The halophytes with most astounding UV were *Podocarpus latifolius* (Thunb.) R.Br. ex Mirb. (0.3) and *Pentatropis nivalis* (J.F.Gmel.) D.V. Field and J.R.I. Wood (0.28). The most elevated UV of these halophytes demonstrates their basic presence and the customary restorative essentialness of these halophytes by respondents against a few issue. The over and again revealed plant species are seen as more naturally dynamic than other plant species (Logan, 1986). The halophytes having minimum UV are given in Table 2. The previous reports show that the plant species in the present investigation with minimum UV are likewise compelling as these halophytes can be utilized for the human pharmaceuticals advancement (Holling et al., 2012; Macuja et al., 2015; Oliveira et al., 2015). Plant species with high use esteem should prepared further to acquire significant mixes for the advancement of characteristic medications (Mahmood et al., 2012).

Informant Consensus Factor (ICF): The ICF is a fair execution for the elaboration of utilization repeat of therapeutic halophytes for various suffering arrangements. The present examination delineated 12 essential disturbance classes reliant on estimations of ICF. The ICF estimations of halophytes move from 0 to 0.62 (Table 3). The most astonishing ICF regard (0.62) was represented gastrointestinal dissipates. The more inescapable disease class was GIT issue in the examination domain due to the low-quality sustenance permit by teenagers and adolescents who need to eat drive-through nourishment than cautiously collected dishes in home. Equivalent revelations were moreover filed by a couple of various examinations coordinated in Pakistan and in various countries (Logan, 1986; Rokaya et al., 2010; Teklehaymanot, 2009). The second most shocking ICF regard (0.38) was represented on edge disarranges in the examination zone. These disclosures reveal the manner in which that the respondents living in salt mine work barely and exhaustingly accumulate things from the forested areas for cooking as they live in rough zone or with the agriculture occupation, lock in fields as they have poor money related conditions. The smallest ICF regard (0.00) was found for immunizing agent and urinary issue. The need consistency about the usage of salt tolerant therapeutic plants was seen among respondents and spoke to the base ICF values. Scarcest ICF regard exhibits that respondents have no simultaneousness on the use of assembled and referenced therapeutic halophytes. Assortments in relationship of helpful plants for easing sicknesses show the social example in the examination zone.

[Table 3]

GIS mapping of halophytic vegetation: The database configuration has a noteworthy effect to plant mapping as it decides the highlights and information association. The GIS essential introduce is the relationship between's highlights traits and spatial information. The computerized guide delineating the halophytic vegetation of Khewra salt mine was delivered utilizing GIS (Figure 8). The vegetation mapping has been utilized in the present investigation to discover the conveyance of halophytes. The vegetation mapping utilizing GIS these days is turning into a critical instrument so as to make a database including a spatial area and property table data of the plants utilizing both subjective and quantitative accumulation procedures. The utilization of GIS in vegetation mapping was likewise introduced different examinations (Monto et al., 2005; Guisan and Thuiller, 2005).

[Figure 8]

X-Ray Diffraction: The XRD technique is extensively used in characterization of rock-based minerals (Wu *et al.*, 2017). In the current study, this technique was used to determine the nature of salt sample and to understand the relationship between salt type and distribution of halophytes in study area. The XRD analysis of salt sample using High Score software demonstrates that the salt nature of study area is copper sulfide. It contains copper sulfide mineral Cu₂S which is chalcocite. This technique was performed for structural studies using salt sample in powder form. The XRD pattern shows that salt of study area is crystalline in nature and reported as cubic crystal Cu₂S (a=5.6286 Å, b=5.6286 Å, c=5.6286 Å).

A), with spacer group Fm3m and its density was calculated as 5.93 g/cm³. Copper sulfide nature of soil in relation with plants attained interest in recent years for many reasons. Copper sulfide is chemically composed of copper and sulfide. The 'CU' and 'S' both are important element for plant growth, plant sterility and photosynthesis and they also help in maintaining soil PH.

On the other hand, sulfide inhibits the rate of plant growth in several ways as it is toxic for plant roots, cause deficiency of essential elements in plant roots and also results in sulfur deficiency in plants (Wu *et al.*, 2017). The copper sulfide based salty soil as compare to loamy soil has small particle size than loamy soil which affects the PH of soil and flow of water and minerals in soil. The halophytes distribution in the study area revealed that they have the ability to adopt such type of soil conditions. The local floras composition could be altered by global environmental changes therefore it should revise regularly and knowledge about indigenous use of medicinal plants, soil and environ The XRD method is widely utilized in portrayal of shake based minerals (Wu *et al.*, 2017). In the present investigation, this procedure was utilized to decide the idea of salt example and to comprehend the connection between salt kind and appropriation of halophytes in study zone. The XRD examination of salt example utilizing High Score programming exhibits that the salt idea of study region is copper sulfide. It contains copper sulfide mineral Cu₂S which is chalcocite. This method was performed for basic examinations utilizing salt example in powder frame. The XRD design demonstrates that salt of study region is crystalline in nature and announced as cubic gem Cu₂S (a=5.6286 Å, b=5.6286 Å, c=5.6286 Å), with spacer aggregate Fm3m and its thickness was determined as 5.93 g/cm³. Copper sulfide nature of soil in connection with plants accomplished enthusiasm for ongoing years for some reasons. Copper sulfide is synthetically made out of copper and sulfide. The 'CU' and 'S' both are vital component for plant development, plant sterility and photosynthesis and they additionally help in keeping up soil PH.

Then again, sulfide represses the rate of plant development in a few different ways as it is poisonous for plant roots, cause inadequacy of basic components in plant establishes and furthermore results in sulfur lack in plants (Wu *et al.*, 2017). The copper sulfide based salty soil as contrast with loamy soil has little molecule estimate than loamy soil which influences the PH of soil and stream of water and minerals in soil. The halophytes dispersion in the investigation territory uncovered that they can embrace such sort of soil conditions. The nearby verdures structure could be modified by worldwide natural changes along these lines it should overhaul routinely and information about indigenous utilization of therapeutic plants, soil and condition ought to be reported legitimately. ment should be documented properly.

[Figure 9]

Phytochemical constituents of Medicinal halophytes: The pharmacological confirmation of any plant species dependent on ethno-therapeutic information assume critical job for the fuse of customary natural meds into current human services framework. In the present examination, the vast majority of the salt tolerant plant species have been seen to have a solid pharmacological movement while couple of halophytes demonstrate some pharmacological action. The reported restorative halophytes with their dynamic phytochemicals have been given in Table 2. These phytochemical mixes are connected with a few pharmacological exercises utilized for the treatment of different ailments in everywhere throughout the world (Verpoorte *et al.*, 2002). The nearness of glycosides, phytohormones, saponins, alkaloids and flavonoids in plant species are in charge of their pharmacological exercises (Khan *et al.*, 2012). *Podocarpus latifolius* is an essential salt tolerant plant species which have solid ethno-therapeutic confirmations however have no pharmacological record. This halophyte is suggested for cutting edge exploratory pharmacological strategies for future investigations. Phytochemical thinks about give pharmacological proof to the confirmation of Ethnomedicinal and fragrant employments of plants species. The past writing announced the recognized phytochemical constituents of revealed salt tolerant plant species. It is broke down that these pharmacological exercises can be favored in future because of co-dismal circumstance and moral pondering and can handled further for more subtleties. In the ebb and flow examination, the salt tolerant restorative plants not revealed in past research ought to be surveyed for pharmacological investigation which can help in the revelation of new medications.

CONCLUSIONS

The present investigation reported the conventional restorative practices of halophytes, mapped the halophytic vegetation utilizing GIS and portrayed the dirt minerals of Khewra salt mine utilizing X-beam diffraction examination. This investigation is embraced to record the indigenous restorative information about halophytes for future confirmation utilizing phytochemicals and phramcological exercises. Thirty-eight salt tolerant restorative plants having a place with twenty-three families were reported. Fabaceae was the much of the time refered to plant family. Lion's share of the salt tolerant plants were herbs and leaf was the most well-known utilized plant part. This review uncovers the novel data that the investigation region is enriched with rich decent variety of halophytes which are utilized as nourishment and additionally for home grown details for the treatment of different afflictions. The discoveries show the insufficient wellbeing administrations for provincial populace by the legislature of Pakistan, along these lines, lion's share of the neighborhood individuals utilize conventional restorative information which additionally include the huge number of halophytes. The outcomes help the phytochemistry analysts for the screening of therapeutic salt tolerant plants for medication disclosure in future with progressively worldwide intrigue. The reconciliation of characteristic prescriptions in the essential human services for neighborhood individuals is approved by the logical bio-movement validation of the therapeutic plants. This exhaustive investigation is of exceedingly importance since it is first archived ethno-therapeutic write about salt tolerant plants of Khewra salt mine. The further research on preservation systems ought to be directed for the natural medications reasonable advancement since it will give the better financial conditions in the investigation region.

Acknowledgements: Authors acknowledge Pakistan Science Foundation (PSF) Pakistan under letter No. PSF/P&D/TG-ND (384)19. by providing travel grant for oral presentation and to attend Symposium (MESMAP-2019).

Table 1. Demographics of informants in Khewra (Punjab)

S. NO.	Variable	No. of persons	Percentage
1.	Informant category -THPs	13	8.60
	Local People	138	91.39
2.	Gender : F/M	68	45.03
		83	54.96
3.	Age	06	3.97
		11	7.28
		25	16.55
		27	17.88
		49	32.45
		33	21.85
4.	Educational background	43	28.47
		12	7.94
		04	2.64
		29	19.20
		15	9.93
		11	7.28
		37	24.50
5.	Experience of the traditional health practitioners	02	1.32
		02	1.32
		04	2.64
		03	1.98
		02	1.32

Table 3. ICF value of halophytic medicinal plants used against various diseases

GIT diseases	72	189.47	28	73.68	0.62
Respiratory diseases	29	76.32	24	63.16	0.18
Sexual disorders	19	50	14	36.84	0.28
Urinary disorders	17	44.74	17	44.74	0.00
Muscle and skeletal disorders	40	105.26	26	68.42	0.36
Nervous disorders	41	107.89	26	68.42	0.38
Glandular disorders	31	81.58	25	65.79	0.2
Cardiovascular disorders	35	92.11	27	71.05	0.24
Ear, nose and eye disorders	17	44.74	13	34.21	0.25
Nail, skin and hair disorders	18	47.37	23	60.53	0.29
Body energizers	6	15.79	5	13.16	0.2
Antidote	10	26.32	10	26.32	0.00

*ICF = Informant Consensus Factor

Table 2. List of halophytes and their medicinal used by the local people of Khewra salt mine, Punjab, Pakistan

Family Plant Names/ Voucher Specimen No	Common name	Flowering season	Habit	Status	Life form	Growth form	Part used	Mode of preparation	Mode of administra tion	Phytochemicals	Ethno-medicinal uses	FC *	UV*	RFC*	Herbal Recipes
<i>Aizoaceae</i> <i>Trianthema</i> <i>portulacastrum</i> L. ISL- 97	Itsit	Sep-Oct	Herb	Wild	Therophyte	Annual	Whole Plant	Decoction , Paste	Oral	Ecdysterone, Alkaloids (Trianthemine, Punarnavine), Saponins, Tannins, Trianthenol, Steroids, Flavonoids, Anthraquinones, Tri-Terpenoids, Leptorumol, Nicotinic acid, Ascorbic acid, Beta cyanin, Cardiac Glycosides ^{72,73}	**Respiratory disorders , Headache , Liver disorders , Arthritis , Heart problems , Tuberculosis , Obesity , Hypersomnia	57	0.14	0.38	3 spoon paste of whole plant is mixed with 1 spoon of <i>Piper nigrum</i> powder; take it twice in day for respiratory diseases Decoction of plant is used to treat arthritis
<i>Amaranthaceae</i> <i>Achyranthes aspera</i> L. ISL- 175	Putkanda	Sep-Aug	Herb	Wild	Chamaephyte	Annual	Root, Leaf	Infusion, Paste	Oral, Topical	Alkaloids (betaine), Carbohydrates, Tannins, Saponin, Flavonoids, Proteins, Amino acids, Tannins, Phenols, Steroids, Glycosides, Ecdysterone, β -sitosterol, Hydroxyecdysone, Pentatriaontane, Hexatriacontane, Tritriacontane, Quercetin ^{74,75}	Abdominal pain , Scorpion sting , Headache , Heart problems , Pneumonia , <i>Tuberculosis</i> , <i>Kidney stones</i> , <i>Obesity</i> , <i>Skin diseases</i> , <i>Constipation</i> (DS/ I), <i>Fever</i>	59	0.19	0.39	Roots are placed in hot boiled water overnight then used after straining for abdominal pain The fresh leaves paste is applied on scorpion bite
<i>Amaranthus viridis</i> L. ISL- 586	Cholai	Aug-Sep	Herb	Wild	Therophyte	Annual	Leaf	Decoction, Paste	Oral	Rutin, Quercetin, 24-ethyl-22-dehydrolathosterol, Dehydrolathosterol, Amasterol (24-methylene-20-hydroxycholesta-5,7-dien-3 β -ol) ⁷⁶	Diuretic , Diabetes , , Depression , Influenza , Skin diseases , Snake bite (CS/E)	63	0.11	0.42	1 teaspoon of leaf paste is taken with milk once in day which is diuretic The leaves decoction is given twice a day to treat influenza
<i>Spinacia oleracea</i> L. ISL- 111	Palak	Jun-Oct	Herb	Cultivat ed	Therophyte	Annual	Leaf	Cooked, Raw	Oral	Coumerin, Anthocyanins, Emodins, Phlobatannins, Spinacetin, Jaceidin, Violaxanthin, β -carotene, Folic acid, Oxalic acid, Diacylglycerol, Sulfoquinovosyl diacylglycerol, beta-basrubrins, 4'pentahydroxi-6-methoxyflavone ^{77,78,79}	Digestive disorders , , Skin diseases , Influenza , Diabetes , Liver inflammation , Urethral ulcer , Obesity , Diuretic	65	0.14	0.43	Leaves are eaten in raw form to treat liver inflammation Leaves are cut into pieces and boiled with <i>Trigonella foenum-graecum</i> , <i>Menthapiperita</i> and <i>Coriandrumsativum</i> leaves and then add 1 spoon salt and fruit paste of <i>Capsicum annum</i> ; take half plate

MESMAP-5 PROCEEDINGS BOOK
April 24th - 26th, 2019 / Cappadocia – Turkey

<i>Suaeda vermiculata</i> Forssk. Ex J. F. Gmel. ISL- 399	Laani	Mar-Sep	Shrub	Wild	Nanophanerophyte	Perennial	Leaf	Powder, Paste	Oral, Topical	Alkaloid, Coumarins, Flavonoids (Genistein), Saponins, Sterols, Terpenes, Tannins, Triterpenoids, Polyphenols ^{80,81,82}	Constipation, Skin diseases , Stomach ache, Dysentery, Hepatitis, Nausea, Insomnia, Cholera	49	0.16	0.32	in twice a day to treat obesity Dried leaves powder is mixed with sugar and 1 spoon is taken thrice in day Leaves paste is applied on skin topically to treat skin rashes and wounds Leaf paste is applied on wounds for healing Leaf juice with rose water is applied topically for skin diseases
<i>Apocynaceae</i> <i>Nerium oleander</i> L. ISL- 155	Kaneri	Mar-Sep	Shrub	Wild	Phanerophyte	Perennial	Leaf	Paste, Juice	Topical	Rhamnose, Arabinose, Galactose, Cardenolides, Neriucoumaric, Isoneriucoumaric acids, Kaneroside, Neriumoside, Proceragenin, Neridienone A, Beta-neriursate, D-sarmentose, D-diginose, Neridiginoside, Nerizoside, Neritaloside, Odoroside, Isoricinoleic acid, Folinerin ^{83,84}	Arthritis, Wound healing (MS/E), Skin diseases , Tuberculosis, Malaria, Insomnia, Diabetes, Meningitis, Blood purifier, Scorpion sting, Body swellings (MS/E)	53	0.21	0.35	Leaf juice with rose water is applied topically for skin diseases
<i>Arecaceae</i> <i>Phoenix sylvestris</i> L. ISL- 185	Khajoor	Mar-Apr	Tree	Cultivated	Phanerophyte	Perennial	Fruit	Paste	Oral	β -sitosterol, β -amyrin, Quercetin, Quercitrin, Alkaloids, Flavonoids, Phenols, Tannins, Saponins ^{85,86}	Obesity, Fever , Brain tonic , Heart tonic, Sexual problems (RS/I), Epilepsy, Cold, Dysentery, Aphrodisiac (RS/I)	71	0.13	0.47	1 teaspoon of fruit paste is given with ½ teaspoon of vinegar to cure fever Fruit paste is given with warm milk which is brain tonic
<i>Asclepiadaceae</i> <i>Calotropis procera</i> (Aiton) W.T.Aiton ISL- 58	Aak	Throughout year	Shrub	Wild	Nanophanerophyte	Perennial	Latex, Flower, Leaf	Raw	Oral, Topical	Alkaloids, Terpenoids, Flavonoids, Tannins, Saponins, Sterols (β -sitosterol, Stigmasterol, Multiflorenol) Cardiac glycosides, Terpinoids, Cardenolides ^{87,88}	Piles , Diabetes, Obesity, Jaundice, Malaria, Tuberculosis, Toothache (DS/E), Hysteria, Snake bite , Wound healing (MS/E), Teeth problems (DS/E), Stomach disorders	65	0.18	0.43	Latex is applied on the wounds of piles to get relieve. Fresh flowers and leaves are eaten in raw form for snakebite
<i>Pentstemon nivalis</i> (J.F.Gmel.) D.V.Field&J.R.I.Wood ISL- 177	-	June-Oct	Climber	Wild	Phanerophyte	Perennial	Root	Paste, Juice	Oral	Glycoside, Steroids, Terpenoids, Phenolic, Saponins, Flavonol glycosides, Flavonol sulphates, Flavonol disulphates ^{89,90}	Dysentery, Hepatitis, Piles , Wound healing (MS/E), Obesity , Lung disorders, Eczema (MS/E), Skin allergy, Earache (SO/E), Leucorrhoea (RS/I), Nail disorders (MS/E)	39	0.28	0.26	One teaspoon of roots paste is used to treat piles Root juice is taken in the night before sleeping to treat obesity

MESMAP-5 PROCEEDINGS BOOK
April 24th - 26th, 2019 / Cappadocia – Turkey

Asteraceae <i>Carthamus oxyacantha</i> M. Bieb. ISL- 137	Pohli	Apr-July	Herb	Wild	Therophyte	Annual	Seed	Powder, Oil	Oral, Topical	Flavonoids, Cardiac glycosides, Triterpenoids, Tannins, Carbohydrates, Phenols, Palmitic acid, Stearic acid, Oleic acid, Linoleic acid, Carthamidin, Serotonin ^{91,92,93}	Fever , Skin allergy , Influenza , Impotence (RS/I), Arthritis , Hepatitis , Dog bite (CS/E), Itching , Skin rashes , Conjunctivitis (SO/I), Heart disorders (MS/E)	41	0.27	0.27	Dried seeds are ground to make powder which is taken after boiling in water once in day to treat fever
Brassicaceae <i>Malcolmia africana</i> (L.) R.Br. ISL- 359	Kambar	Mar-July	Herb	Wild	Therophyte	Annual	Leaf, Flower	Decoction, Extract	Topical	Oleic acid, Linoleic acid, Linolenic acid, Erucic acid, Flavonoids, Phenols ⁹⁴	, Skin allergy , Lung cancer , Dog bite	49	0.08	0.32	Seeds oil is applied topically on skin to treat itching and rashes Leaves are boiled to obtain juice of leaves which is applied thrice in day to treat dog bite
Capparaceae <i>Capparis decidua</i> (Forssk.) Edgew. ISL- 470	Karineh	Mar-Sep	Shrub	Wild	Phanerophyte	Perennial	Fruit, Stem	Raw, Paste	Oral, Topical	Alkaloids (Capparine, Cappariline), Capparinine), Terpenoids (Lupine), Glycosides, Sterol, Methyl isothiocyanate, β-Carotene, Ascorbic acid, Phthalic acid ⁹⁵	Digestive disorders , Blood purifier , Arthritis , Aphrodisiac (RS/I), Impotence (RS/I), Tuberculosis , Stomach ulcer , Teeth disorders (DS/E), Arthritis , Alzheimer's disease	53	0.19	0.35	Crushed flowers are placed in hot water to obtain extract which is used topically to treat skin allergy Fruit is eaten in raw form for digestive disorders and blood purification The twig paste is applied on teeth for shining
Cucurbitaceae <i>Citrullus colocynthis</i> (L.) Schrad. ISL- 351	Tumma	June-Sep	Herb	Wild	Therophyte	Annual	Fruit, Root	Decoction	Oral	Colocynthin, Colocynthetin, Albuminoids, Starch, Tannins, Saponins, Proteins, Reducing sugars, Alkaloids, Glycosides, Diterpenes, Carotenoids, Anthraquinones, Phlobatannins, Flavanoids ^{96,97}	Constipation (DS/I), Earache (SO/I), Haemorrhoids , Dysentery , Liver diseases	55	0.11	0.36	Fresh fruit pieces are boiled in water with sugar to make paste which is used for constipation Root decoction is used to get relieve from earache
Cyperaceae <i>Cyperus rotundus</i> L. ISL- 89	Delluca	May-Oct	Shrub	Wild	Nanophanerophyte	Perennial	Root	Powder, Paste	Oral	Flavonoids, Coumarin, Tannins, Polyphenols, Sterols, Carbohydrates, Glycosides, Anthraquinone, Anthocyanin, Saponins, Alkaloids ^{98,99}	Diuretic , Diabetes , Digestive disorders , Skin diseases , Nausea , Cold , Cough , Dysentery , Stomach ache (DS/ I), Oedema , Headache	46	0.24	0.3	Root powder is taken with milk once in day which is diuretic 1 teaspoon of root paste is used with rose water to cure digestive disorders

MESMAP-5 PROCEEDINGS BOOK
April 24th - 26th, 2019 / Cappadocia – Turkey

<i>Euphorbiaceae</i> <i>Ricinus communis</i> L. ISL- 193	Harnoli	Mar-Sep	Herb	Wild	Phanerophyte	Perennial	Leaf, Root	Paste	Topical	Chlorogenic acid, Decanamine, Ellagic acid, Linoleic acid, Oleic acid, Palmitic acid, Palmitoleic acid, Stigmasterol, Tartaric acid, Tannins, Arachidic acid, Glycine, Ricinine, Tryptophan ¹⁰⁰	Wound healing (MS/E), Snake bite (CS/E), Arthritis , Constipation (DS/ I), Skin diseases, Fever, Diuretic, Digestive disorders, Liver diseases <i>.Haemorrhoids</i>	67	0.15	0.44	Leaves are crushed and heated to make poultice for healing wounds Young root paste is applied on joints to treat arthritis
Fabaceae <i>Acacia modesta</i> Wall. ISL-103	Phulai	Nov- Mar	Tree	Wild	Phanerophyte	Perennial	Bark, Stem	Decoction, Powder, Raw	Oral, Topical	Flavonoids (Quercetin, Kaempferol) Alkaloids, Terpenoids (Lupeol, Betulin, α -amyrin), Tannins, Linoleic acid, Oleic acid, Cyclitols (Pinitol) ¹⁰¹	**Digestive disorders , Obesity, Teeth disorders (DS/E), Kidney pain, Gastric disorders, Backache, Epilepsy, Sexual infertility (RS/I), Jaundice, Skin dryness, Skin allergy (MS/ E), Skin boils(MS/ E)	67	0.18	0.44	Bark is grinded to make powder which is used for digestive disorders Bark is boiled in water to make decoction which is used to reduce obesity Fresh stem is used for cleansing teeth
*Acacia nilotica (L.) Delile ISL- 241	Kikar	May-Oct	Tree	Wild	Mesophanerophyte	Perennial	Bark, Leaf	Decoction, Infusion	Oral, Topical	Alkaloids, Saponin, Proteins, Amino Acids, Anthraquinone, Tannin, Flavonoids, Fixed oils, Fats, Cardiac Glycosides, Leucocyanidin, Polyphenols (dicatchin, Quercetin, Gallic acid), Amino acids (Histidine, Methionine, Valine, Cystine) ^{102,103}	Teeth and gum disorders (DS/E), Blood purifier, Dysentery , Diabetes, Gonorrhoea (RS/I), Diabetes, Asthma, Aphrodisiac (RS/I), Liver diseases, Arthritis, Eye diseases (SO/I)	73	0.16	0.48	Bark is soaked in tepid water for few hours; use this water for mouth sores and teeth and gum disorder Fresh leaves are boiled in water which is orally used for straining twice in day for dysentery
<i>Medicago polymorpha</i> L. ISL- 146	Maina	Mar-May	Herb	Wild	Therophyte	Perennial	Leaf	Juice, Raw	Oral	Triterpene saponins, echinocystic acid, Isoflavones ^{104,105}	Skin dryness, Diuretic, Stomach ache (DS/ I), Alzheimer's disease , Skin boils	57	0.09	0.38	One spoon of leaf juice is taken for stomach ache Leaves are eaten in raw form to treat Alzheimer's disease
<i>Melilotusindicus</i> (L.) All. ISL- 484	Sainji	Mar-Aug	Herb	Wild	Therophyte	Annual	Leaf, Seed	Decoction, Raw	Oral	Flavonoids (Catechin, Epicatechin, Taxifolin), Tannins, Xanthoprotein, Starch, Cystine, Sterols, Triterpenoids, Reducing sugars, Saponins, Alkaloids ^{106,107}	, Abdominal pain , Arthritis, Pneumonia, Skin diseases, Dysentery , Obesity, Skin cancer	44	0.18	0.29	Leaves are boiled in water which is used twice in a day after straining for and abdominal pain Seeds are taken in raw form with <i>Piper nigrum</i> to treat dysentery
<i>Prosopis juliflora</i> (Sw.) DC. ISL- 76	Pahari kikar	Mar-June	Shrub	Wild	Mesophanerophyte	Perennial	Bark, Fruit	Decoction	Oral	Flavonoids, Alkaloids, Saponins, Phenols, Tannins, Phlobatanins, Triterpenes, Cteroids, cardiac glycoside ¹⁰⁸	Asthma , Dysentery, Pneumonia, Influenza (RS/I), Improves memory, Fever, Bladder tonic, Paralysis, Testis swellings (RS/I)	53	0.19	0.35	Bark is boiled in water for 15 minutes which is used twice in a day to treat asthma

MESMAP-5 PROCEEDINGS BOOK
April 24th - 26th, 2019 / Cappadocia – Turkey

Malvaceae <i>Abelmoschus moschatus</i> Medik. ISL- 272	Junglibhindi	Apr-Sep	Herb	Wild	Therophyte	Perennial	Leaf, Seed	Paste, Oil	Topical	Resins, Farnesol, Dodecyl acetate, beta-Sitosterol, Anthocyanins (cyanidin - 3 - sambubioside ,cyanidin - 3 – glucoside), Linoleic acid, Carbohydrates, Phenols, Flavonoids, Terpenoids, Tannins, Saponins ^{109,110}	Headache , Wound healing (MS/E), Hypertension , Skin diseases , Obesity , Mouth sores , , Hepatitis , Piles	72	0.13	0.48	A tea is prepared from the dried pods which is useful for Leaf paste is applied on wounds for healing Seeds oil is applied on forehead to get relieve from headache
Meliaceae <i>Melia azedarach</i> L. ISL- 289	Dharek	Apr-June	Tree	Wild	Phanerophyte	Annual	Leaf, Fruit	Infusion, Powder	Oral, Topical	Terpenoids (β-pinene, a-terpinene), Limonoid (Kulnionone, Fraxinellone), Steroid , Flavonoids (Quercetin), Saponin, Anthraquinones, Alkaloids, Phytosterols (Stigmastrol, Campesterol), β-carotene, Tocopherol, Organic acids (Linoleic acid, Linolenic acid, Oleic acid, Benzoic acid, Vanillic acid) ¹¹¹	Itching , Piles , Migraine , Diabetes , Malaria , <i>Tuberculosis</i> , <i>Earache</i> (SO/I), <i>Skin diseases</i> , <i>Blood purifier</i> , <i>Wound healing</i> (MS/E), <i>Depression</i> , <i>Constipation</i> (DS/ I), Sciatica	70	0.19	0.46	Leaves are crushed and placed in tepid water; apply this water topically to get relieve from itching Dried fruits are crushed to make powder and taken once in day for piles
Myrtaceae <i>Eucalyptus camaldulensis</i> Dehnh. ISL- 427	Lhacesufaida	Sep-Nov	Tree	Cultivated	Macrophanerophyte	Perennial	Leaf	Decoction, Oil	Oral	Tannins, Sterols, Triterpenes, Saponins, Flavonoids, Phenolic compounds, Anthraquinones, Alkaloids ^{112,113}	Influenza , Pneumonia , <i>Liver diseases</i> , <i>Brain tonic</i> , <i>Cough</i> , <i>Sedative</i> , <i>Sexual infertility</i> (RS/I), <i>Epilepsy</i>	63	0.13	0.42	Five leaves are boiled in water which is taken after staining for influenza twice a day One drop of leaves oil is taken twice in a day to treat pneumonia
<i>Eucalyptus globulus</i> Labill. ISL- 367	Sufaيدا	Mar-Aug	Tree	Cultivated	Phanerophyte	Perennial	Leaf	Decoction, Paste	Oral	Tannins, Saponins, Alkaloids, Phenols, Glycosides, Steroids, Terpenes, Camphene, Caproic acid, Citral, Eudesmol, Glutamic acid, Dextrin, Homoserine, Chrysanthemine, Chrysin, Cyanin, Cynaidin, Oleanolic acid, Masinic acid ^{114,115}	Digestive disorders , Diabetes , Cough , Cold , Tuberculosis , Constipation (DS/ I), Brain tonic , Jaundice , Kidney stones , Piles , Backache , Paralysis , Ringworm	63	0.21	0.42	Leaves decoction is used for cough Leaves paste is placed in hot water overnight to obtain extract; used it after straining twice in a day for diabetes

MESMAP-5 PROCEEDINGS BOOK
April 24th - 26th, 2019 / Cappadocia – Turkey

<i>Nitrariaceae</i> <i>Peganum harmala</i> L. ISL- 141	Hermal	Apr-Oct	Herb	Wild	Chamaephyte	Annual	Seed, Leaf	Raw, Smoke	Oral, Inhale	Flavonoids, Alkaloids (Harmine, Harmaline, Vasicine, Vasicinone, Tetrahydroharmine), Saponins, Tannins, Terpenoids , Steroids , Anthraquinons ^{116,117}	Abdominal pain , Liver diseases , Malaria , Cataract (SO/E), Digestive disorders , Hypertension , Eye weakness (SO/I), Narcotic , Stomach ulcer , Sciatica	58	0.17	0.38	Seeds are taken with water for abdominal pain Burnt leaves are smoke for reducing hypertension
<i>Poaceae</i> <i>Arundo donax</i> L. ISL- 445	Nari	June-Oct	Herb	Wild	Microphanerophyte	Perennial	Root, Leaf	Decoction, Paste	Oral , Topical	Carbohydrates (Rhamnose, Mannose, Arabinose, Xylose, Galactose), Lignin, Alkaloids (Tryptamine, Bufotenidine, Gramine, Arundamine), n-Eicosanoic acid, B-Sitosterol, α -Tocopherol ¹¹⁸	Haematuria , Constipation (DS/ I), Migraine , Eye sight (SO/E), Blood cancer , Epilepsy , Female infertility (RS/I), Liver problems	52	0.15	0.34	Root paste is applied on forehead to treat migraine Leaves decoction is take once in a day to treat constipation
<i>Cynodon dactylon</i> (L.) Pers. ISL- 101	Khabbalghass	Throughout year	Herb	Wild/ Cultivated	Sub-FruticoseChamaephyte	Perennial	Leaf, Stem	Decoction, Paste, Raw	Topical	Terpenoids, , Alkaloids, Flavonoids (Apigenin, Luteolin, Orientin, Vitexin), Carotenoids (Neoxanthin, Violaxanthin), Phenolic phytotoxins (Ferulic, Syringic, Paracoumaric, Vanillic), Phytosterols, Glycosides, Saponins, Palmitic acid, Linoleic acid ^{119,120}	Wound healing (MS/E), Eczema (MS/E), Skin diseases , Depression , Earache (SO/I), Headache , Arthritis , Urethra inflammation (US/ I), Constipation (DS/ I), Diabetes , Stomach ache (DS/ I), Eye infections (SO/E)	69	0.17	0.46	Leaf paste is applied on wounds for healing Aerial parts are boiled in water which is applied on skin to treat eczema The walk with barren feet on this plant give relieve from depression
<i>Hordeum vulgare</i> L. ISL- 309	Joo	Sep-Feb	Herb	Cultivated	Therophyte	Annual	Seed	Decoction, Raw	Oral	Phenolic acids (Ferulic acid, Vanillic acid, Syringic acid, p-coumaric acid), Flavonoids (Flavanols, Anthocyanins, Proanthocyanidins) Lignans (Pinoselinol , Syringaresinol, Laricresinol), Tocopherols, Phytosterol (Sistostanol , Campestanol), Foliates, Ubiquinones ^{121,122}	Kidney disorders , Diabetes , Malaria , Blood purifier	49	0.08	0.32	Seeds are boiled in water and taken for kidney pain Roasted seeds are taken with milk twice in a day to treat diabetes

MESMAP-5 PROCEEDINGS BOOK
April 24th - 26th, 2019 / Cappadocia – Turkey

Podocarpaceae <i>Podocarpus latifolius</i> (Thunb.) R.Br. ex Mirb. ISL- 38	-	Mar-May	Tree	Wild	Mesophanerophyte	Perennial	Bark, Leaf	Decoction, Paste	Oral, Topical	Not known	Backache , Constipation (DS/ I), Asthma , High Blood pressure , Blood cancer , Bone fracture , Cough , Wound healing (MS/E), Skin boils (MS/ E), Ringworm	33	0.3	0.22	1 spoon of bark decoction is taken twice in a day to treat constipation Leaves paste is applied on wounds for healing
Rhamnaceae <i>Ziziphus mauritiana</i> Lam. ISL- 158	Bairi	Sep	Tree	Wild/ Cultivated	Phanerophyte	Perennial	Bark, Fruit, Leaf	Decoction, Raw, Powder	Oral	Alkaloids (Berberine, Quercetin, Kaempferol, Sitosterol), Flavonoids, Glycosides, Phenols, Lignins, Tannins, Sterols, Saponins, Ceryl alcohol, Holosides, Mucilages, , Triterpenoids, Carotene, Thiamine, Riboflavin, Niacin, Citric Acid, Ascorbic Acid, Fluoride, Pectin, Malic acid, Oxalic acid ^{123,124}	71	0.13	0.47	Fresh leaves are boiled in water to make decoction; add honey in it. Take two times in day for blood purification Fruit is eaten in raw form to reduce obesity Bark powder is used with water once in a day for dysentery	
<i>Ziziphusnummularia</i> (Burm.f.) Wight & Arn. ISL- 279	Jungli Bair	Mar-Aug	Shrub	Wild	Phanerophyte	Perennial	Leaf	Paste, Infusion	Oral, Topical	Alkaloids, Glycosides (Nummularogenin), Saponin (Dammarane), Steroids Triterpenoids, Flavonoids, Oil Protein, Tannins, Phenolic compounds ^{125,126,127}	67	0.16	0.44	Shining Hairs (MS/E), Dog bite (CS/E), Fever , Female infertility (RS/I), Nausea , Epilepsy , Skin diseases , Toothache (DS/E), Flatulence , Leucorrhoea (RS/I), Amenorrhoea (RS /I)	Fresh leaves are crushed and place in tepid water, after straining used it for hair washing for shining hairs Leaf paste is placed in hot water for 1 hour then take orally for skin diseases

MESMAP-5 PROCEEDINGS BOOK
April 24th - 26th, 2019 / Cappadocia – Turkey

<i>Salvadoraceae</i> <i>Salvadora oleoides</i> Decne. ISL- 60	Peelu	Mar-June	Tree	Wild	Microphanerophyte	Perennial	Fruit, Root	Paste, Raw	Oral, Topical	Carbohydrates, Glycosides, Sterols (Beta-sitosterol), Steroids, Triterpenes, Proteins, Amino acids, Mucilage, Tannins, Flavonoids, Dihydroisocoumarin, n-hexadecanoic acid, Octacosane, Nonacosane, Octadecene, Heptacosane, Pentacosane, Squalene ^{128,129}	Digestive disorders , Blood purifier , Periodontitis (DS/E), Asthma , Arthritis , Dengue fever , Obesity , Abortive (RS/I), Night blindness (SO/I), Eczema (MS/E)	71	0.14	0.47	Fruit paste is used to treat digestive disorder Root is chewed to treat periodontitis
<i>Scrophulariaceae</i> <i>Verbascum thapsus</i> L. ISL- 176	Geddarta mbakoo	May-Sep	Herb	Wild	Therophyte	Biennial	Leaf, Flower	Juice, Powder	Oral, Inhale	Iridoid glycosides, Phenylethyl glycoside, Sesquiterpenes, Diterpene, Biflavonoid, Triterpene saponins, Polysaccharides, Phenolic acids, Phytosterols (β-sitosterol, Ergosterol peroxide) ^{130,131}	Liver diseases , Tuberculosis , Scorpion sting (CS/E), Diarrhoea , , Diuretic , Asthma	35	0.2	0.23	Flowers juice is taken with tuberculosis Leaves powder is smoke to treat asthma
<i>Solanaceae</i> <i>Solanum incanum</i> L. ISL- 402	Peele mokre	Mar-May	Shrub	Wild	Nanophanerophyte	Perennial	Leaf, Seed	Decoction, Powder	Oral, Inhale	Alkaloid, Steroid, Resin, Glycosides, Flavonoid, Saponin, Tannins, Triterpens, Cardiacglycosides ^{132,133}	Heart diseases , Fever , Narcotic , Blood purifier , Digestive problems , Skin diseases , Cold , Rheumatism , Depression , Sexual tonic (RS/I), Kidney stones , Abdominal worms	45	0.27	0.3	Leaves are boiled in water which is used after straining twice a day for heart diseases Seeds powder with tobacco is smoke which is narcotic
<i>Solanum virginianum</i> L. ISL- 262	Kindair	Throughout year	Herb	Wild	Chamaephyte	Perennial	Fruit, Seed	Powder, Raw, Smoke	Oral, Topical	Alkaloid (Solasodine), Solasonine, Solamargine, Solanocarpine, beta – Solamargine, Solanocarpidine, Isochlorogenic, Neochronogenic, Chronogenic, Caffeic acids, Sterol (Carpesterol) ¹³⁴	, Malaria , Toothache (DS/E) , Asthma , Skin diseases , Liver diseases , Wound healing (MS/E), Alzheimer's disease , Leprosy , Influenza , Fever , Nausea , Rheumatism	55	0.24	0.36	Dried fruit is crushed to make powder which is taken twice in a day for gastric problem Seeds are burnt and smoke is topically used to treat leprosy
<i>Withania coagulans</i> (Stocks) Dunal ISL- 183	Khamjira	Jan-Apr	Herb	Wild	Phanerophyte	Perennial	Seed, Fruit	Decoction, Paste	Oral, Topical	Alkaloids, Steroids (Dihydrostigmasterol), Phenolic compounds, Tannins, Saponins, Carbohydrates (D-galactose, D-arabinose), Proteins, Amino acids, Organic acids,	Diabetes , Diuretic , Improves memory , Hypertension , Hypersomnia , Heart diseases , Wound healing (MS/E)	55	0.13	0.36	Fruit is chewed to treat toothache Seeds are boiled in hot water which is taken with water after straining to treat diabetes Fruit paste is applied topically on wound for healing

MESMAP-5 PROCEEDINGS BOOK
 April 24th - 26th, 2019 / Cappadocia – Turkey

<i>Withaniasomnifera</i> (L.) Dunal ISL- 85	Aksun	Throughout year	Shrub	Wild/ Cultivated	Phanerophyte	Perennial	Leaf, Root	Decoction, Powder	Oral	Withanolides, Linoleic acid ^{135,136} Withanolides, Alkaloids (Withanine, Withasomnine, Somniferinine, Pseudotropine, Cuscohygrine, Scopoletin, Tropanol), Alcohols (Withaniol, Sommirol, Sommitol), Saponins, Cysteine, Chlorogenic acid ^{137,138,139}	Rheumatism , Aphrodisiac (RS/I), Diabetes , Kidney stones , Constipation , Improves memory , Scorpion sting , Hypertension , Fever	59	0.14	0.39	Fresh leaves are boiled in water; used this one cup of decoction once in a day for rheumatism Root powder is given with milk to remove kidney stones
<i>Zygophyllaceae</i> <i>Tribulus terrestris</i> L. ISL- 655	Bhaakra	July-Sep	Herb	Wild	Chamaephyte	Annual	Root, Fruit	Powder, Paste	Oral	Saponins (Furostanol, Spirostanol) Flavonoids , Glycosides, Alkaloid (Harmane, Norharmane, Tribulusterine), Tannins, Terrestriamide, Hecogenin, Aurantiamide acetate, Xanthosine, Fatty acid ester, Ferulic acid, Vanillin, Sterols (β-sitosterol, Stigmasterols) ^{140,141}	Rheumatism , Diuretic , Aphrodisiac (RS/I), Impotence (RS/I), Tuberculosis , Blood purifier , Headache , Piles , Migraine , Depression	66	0.15	0.44	Dried plant without root is crushed to make powder which is taken for rheumatism for ten days Fruit paste is taken with milk which is diuretic

MESMAP-5 Draft Proceedings

List of Author

Participant	Country
Adem Gharsallaoui	France
Adil Farooq Wali	UEA
Ahmad Ali	India
Ahmet Altay	Turkey
Ahmet Özdemir	Turkey
Ahmet Yenikalayci	Turkey
Albert Katchborian Neto	Brazil
Alexander Tashev	Bulgaria
Alper Çimik	Turkey
Alvaro Viljoen	South Africa
Amantay Alzhan	Kazakhstan
Ana Flavia Burlec	Romania
Anake Kijjoa	Portugal
Antoaneta Trendafilova	Bulgaria
Archana Thite	India
Argentieri Maria Pia	Italy
Asih Triastuti	Indonesia
Ataman Bilge Sari	Turkey
Aurel VasIU	Romania
Awatif Boumaza	Algeria
Ayşe Esra Karadağ	Turkey
Ayşegül ÇaşkurIU	Turkey
Badra Boubakeur	Algeria
Belgin Sever	Turkey
Berna Goker	Turkey
Biljana Bauer	Macedonia
Bouhenni Hasna	Algeria
Büşra Karpuz	Turkey
Carmen Dana Şandru	Romania
Ceren Anlas	Turkey
Ceylan Dönmez	Turkey
Charafeddine Jama	France
Ciprian Ilaşcă	Romania
Cornelia Mircea	Romania
Çimen Karasu	Turkey
Decha Kumla	Portugal
Diana Mihaela Dumitrascu	Romania
Didem Koçum	Turkey

Participant	Country
Ebru Yilmaz	Turkey
Ece Sözer	Turkey
Eda Şensu	Turkey
Ela Nur Şimşek Sezer	Turkey
Elif Dilmaç	Turkey
Elif Güzel	Turkey
Elsadig A. Eltayeb	Oman
Emine Saliha Toprak	Turkey
Emrah Şirin	Turkey
Enkelejda Goci	Albania
Erden Banoglu	Turkey
Esmayagdi Efe	Turkey
Ezgi Şenol	Turkey
F. Sezer Şenol Deniz	Turkey
Faruk Karahan	Turkey
Fatma Ayaz	Turkey
Fatma Güneş	Turkey
Filiz Meriçli	Turkey
Grozi Delchev	Bulgaria
Gülaçtı Topçu	Turkey
Gülmira Özek	Turkey
Gülsüm Yaldiz	Turkey
Gülsüm Yıldız	Turkey
Habiba Daroui-Mokaddem	Algeria
Hafida Bouamama	Morocco
Halil I Ciftci	Japan
Hallal Nouria	Algeria
Hasna Bouhenni	Algeria
Havva Atar	Turkey
Helene De Wet	South Africa
Hilal Top	Turkey
Hilal Zaid	Palestine
Inès Bellil	Algeria
Iram Nizam	India
Irmak Alev	Turkey
Ivan Salamon	Slovakia
İlkay Erdoğan Orhan	Turkey
İrem Ayran	Turkey
İrem Şengür	Turkey
Janar Jenis	Kazakhstan

Participant	Country
Jintanaporn Wattanathorn	Thailand
Juthamart Maneenet	Thailand
Kabouche Ahmad	Algeria
Kanyarat Holasut	Thailand
Kawintra Tamprasit	Thailand
Kevser Kusat Ol	Turkey
Khaled El Omari	Lebanon
Koula Doukani	Algeria
Lahouel Zakia	Algeria
Lamia Hamrouni	Tunisia
Lütfi Nohutçu	Turkey
Madalena M. M. Pinto	Portugal
Mahmoudi Kheira	Algeria
Marina Spînu	Romania
Marwan Azubaidi	Malaysia
Masami Otsuka	Japan
Mebarka Bouziane	Algeria
Medjeber Nacera	Algeria
Mehlika Dilek Altintop	Turkey
Mehmet Engin Celep	Turkey
Mehtap Kılıç	Turkey
Meliani Samiha	Algeria
Meryem Bozkurt	Turkey
Mihaela Niculae	Romania
Mikako Fujita	Japan
Mohamed O. Radwan	Japan
Mohammad Rais Mustafa	Malaysia
Monica Hancianu	Romania
Muhammad Ajmal Shah	Pakistan
Muhammad Zafar	Pakistan
Murat Tunçtürk	Turkey
N. Gökçe Ulusoy	Turkey
Nabila Slougui	Algeria
Nacira Benahmed	Algeria
Nadjia Hamlat	Algeria
Naima Hellali	Algeria
Najjaa Hanen	Tunisia
Nalan Yilmaz	Turkey
Narin Sadıkoğlu	Turkey
Natthida Weerapreeyakul	Thailand

Participant	Country
Nazım Şekeroğlu	Turkey
Nidhi Pathak	India
Nour-Eddine Chihib	France
Noureddine Djebli	Algeria
Nut Palachai	Thailand
Nuttinee Teerakulkittipong	Thailand
Orawan Monthakantirat	Thailand
Orçun Çinar	Turkey
Pamela Jha	India
Paweena Pradniwat	Thailand
Pratthana Srichomphu	Thailand
Priya Sundarrajan	India
Rachadaporn Benchawattananon	Thailand
Rafik Bensegueni	Algeria
Raksaworn Jaisaard	Thailand
Ramanpreet Walia	India
Ramdane Farah	Algeria
Rana Samara	Palestine
Reecha Madaan	India
Reem Hassan Ahmed	Sudan
S. Ayşe Çelik	Turkey
Safiye Emirdağ-Öztürk	Turkey
Sagarika Damle	India
Saiqa Ishtiaq	Pakistan
Salima Douichene	Algeria
Sana Janakat	Jordan
Sandy Van Vuurena	South Africa
Satish Sardana	India
Serpil Demirci	Turkey
Sevgi Gezici	Turkey
Sinan Akgöl	Turkey
Siriporn Chuchawankul	Thailand
Sokar Zahra	Morocco
Stan Cataliana	Romania
Stan Cristi	Romania
Štefica Findri Guštek	Croatia
Sultan Pekacar	Turkey
Sumon Sakolchai	Thailand
Supaporn Muchimapura	Thailand
Supawadee Daodeec	Thailand

Participant	Country
Suresh Kumar	India
Susmita Gudulkar	India
Suwakon Wongjaikam	India
Svetla Angelova	Bulgaria
Svetlana Kulevanova	Bulgaria
Tabassum Khan	India
Tala Reihanifar	Turkey
Taner Özcan	Turkey
Tansu Uskutoğlu	Turkey
Tarapong Srisongkram	Thailand
Tatjana Kadifkova Panovska	Macedonia
Tavani Rocha Camargo	Brazil
Tejashree Shanbhag	India
Temel Özek	Turkey
Thananya Nantarat	Thailand
Theera Rittirod	Thailand
Thomas Jakschitz	Austria
Tofiq Mammadov	Azerbaijan
Tuba Kuşman	Turkey
Tuğba Günbatan	Turkey
Tuna Uysal	Turkey
Ufuk Koca Çalışkan	Turkey
Usha Mukundan	India
Ünal Karık	Turkey
Ünsal Açık	Turkey
Vikneswaran Murugaiyah	Malaysia
Višnja Oreščanin	Croatia
Wanna Sirisangtakul	Thailand
Warin Ohn-on	Thailand
Worrapan Poomanee	Thailand
Yaowared Chulikhit	Thailand
Yeşim Sağ Açık	Turkey
Yigit Gunes	Turkey
Zahia Kabouche	Algeria
Zamri Chik	Malaysia
Zehra Aydin Bek	Turkey
Zglinska Klara	Poland
Zine El Abidine Ababsa	Algeria
Zubeyir Elmazoglu	Turkey
Zvijezdana Štaba	Croatia

UPCOMING SYMPOSIUMS

MESMAP-6

&

ISPBS-6

April, 2020

Call for Papers

Current Perspectives on Medicinal and Aromatic Plants (CUPMAP)

ISSN: 2619-9645 (Print) / e-ISSN: 2667-5722 (On-line)

Current Perspectives on Medicinal and Aromatic Plants (CUPMAP) is an open access, peer-reviewed and refereed international journal published by MESMAP scientific group. The main objective of the CUPMAP is to provide an intellectual outlook on the scientific researches on Medicinal and Aromatic Plants. CUPMAP have distinguished goals to promote interdisciplinary scientific studies in which results could easily be used in industrial production on MAPs.

This international scientific journal publishes research papers related to Medicinal and Aromatic Plants in the fields of science and technology such as Biology, Molecular Biology and Genetics, Chemistry, Agriculture, Biochemistry, Botany, Ethnobotany, Environmental Science, Forestry, Horticulture, Health Care & Public Health, Nutrition and Food Science, Pharmaceutical Sciences, and so on.



<http://dergipark.gov.tr/cupmap> / <http://www.cupmap.org>

The journal is published in both print and on-line versions

CUPMAP publishes original research papers, applied studies, and review articles in MAPs science and technology. Special Issues devoted to important topics in the MAPs science and technology could also be published.

Having well known board members distinguished scientists from different disciplines with huge experiences on MAPs all over the world, CUPMAP will be indexed in many databases after first issue. The goal of the journal is to be indexed in Thomson Reuters in a short time.

CUPMAP is inviting papers for Vol. 2 No. 1 which is scheduled to be published on July 31, 2019. Last date of submission: May 31, 2019. However, an early submission will get preference in case of review and publication process.

Send your manuscript to the editor editor@cupmap.org

For more information, visit the official website of the journal www.cupmap.org

With regards,

Prof. Dr. Nazım ŞEKEROĞLU

Chief Editor, Current Perspectives on Medicinal and Aromatic Plants (CUPMAP)

Contact: editor@cupmap.org / sekeroglunazim@gmail.com
