




**BRIEF REPORT**

# Homozygous loss-of-function variants of *TASP1*, a gene encoding an activator of the histone methyltransferases *KMT2A* and *KMT2D*, cause a syndrome of developmental delay, happy demeanor, distinctive facial features, and congenital anomalies

Jehan Suleiman<sup>1,2\*</sup> | Korbinian M. Riedhammer<sup>3,4,5\*</sup>  | Timothy Jicinsky<sup>6</sup> |  
 Melinda Mundt<sup>6</sup> | Laurie Werner<sup>6</sup> | Mirjana Gusic<sup>3,4</sup> | Anna L. Burgemeister<sup>7</sup> |  
 Hessa S. Alsaif<sup>8</sup> | Maha Abdulrahim<sup>9</sup> | Nabil N Moghrabi<sup>10</sup> | Manal Nicolas-Jilwan<sup>11</sup> |  
 Moeenaldeen AlSayed<sup>9,12</sup> | Weimin Bi<sup>13,14</sup> | Srirangan Sampath<sup>6</sup> |  
 Fowzan S. Alkuraya<sup>8,12,15</sup>  | Ayman W. El-Hattab<sup>16,17</sup> 

<sup>1</sup>Division of Neurology, Department of Pediatrics, Tawam Hospital, Al Ain, United Arab Emirates

<sup>2</sup>Department of Pediatrics, College of Medicine and Health Sciences, United Arab Emirates University, Al Ain, United Arab Emirates

<sup>3</sup>Institute of Human Genetics, Klinikum Rechts der Isar, Technical University of Munich, Munich, Germany

<sup>4</sup>Institute of Human Genetics, Helmholtz Zentrum Munich, Neuherberg, Germany

<sup>5</sup>Department of Nephrology, Klinikum Rechts der Isar, Technical University of Munich, Munich, Germany

<sup>6</sup>PreventionGenetics, LLC, Marshfield, Wisconsin

<sup>7</sup>Genetikum, Genetic Counseling and Diagnostics, Stuttgart, Germany

<sup>8</sup>Department of Genetics, King Faisal Specialist Hospital and Research Center, Riyadh, Saudi Arabia

<sup>9</sup>Department of Medical Genetics, King Faisal Specialist Hospital and Research Center, Riyadh, Saudi Arabia

<sup>10</sup>Molecular Diagnostic Laboratory, Department of Genetics, King Faisal Specialist Hospital and Research Center, Riyadh, Saudi Arabia

<sup>11</sup>Division of Neuroradiology, Department of Radiology, King Faisal Specialist Hospital and Research Center, Riyadh, Saudi Arabia

<sup>12</sup>Department of Anatomy and Cell Biology, College of Medicine, Alfaisal University, Riyadh, Saudi Arabia

<sup>13</sup>Department of Molecular and Human Genetics, Baylor College of Medicine, Houston, Texas

<sup>14</sup>Baylor Genetics, Houston, Texas

<sup>15</sup>Saudi Human Genome Program, King Abdulaziz City for Science and Technology, Riyadh, Saudi Arabia

<sup>16</sup>Department of Clinical Sciences, College of Medicine, University of Sharjah, Sharjah, United Arab Emirates

<sup>17</sup>Genetics Clinics, KidsHeart Medical Center, Abu Dhabi, United Arab Emirates

**Correspondence**

Ayman W. El-Hattab, Department of Clinical Sciences, College of Medicine, University of Sharjah, Sharjah, United Arab Emirates.  
 Email: elhattabaw@yahoo.com

**Abstract**

We report four unrelated children with homozygous loss-of-function variants in *TASP1* and an overlapping phenotype comprising developmental delay with hypotonia and microcephaly, feeding difficulties with failure-to-thrive, recurrent respiratory infections, cardiovascular malformations, cryptorchidism, happy demeanor, and distinctive facial features. Two children had a homozygous founder deletion encompassing exons 5–11 of *TASP1*, the third had a homozygous missense variant, c.701 C>T (p.Thr234Met), affecting the active site of the encoded enzyme, and the

\*Jehan Suleiman and Korbinian M. Riedhammer contributed equally to this work.

fourth had a homozygous nonsense variant, c.199 C>T (p.Arg67\*). *TASP1* encodes taspase 1 (TASP1), which is responsible for cleaving, thus activating, the lysine methyltransferases KMT2A and KMT2D, which are essential for histone methylation and transcription regulation. The consistency of the phenotype, the critical biological function of TASP1, the deleterious nature of the *TASP1* variants, and the overlapping features with Wiedemann–Steiner and Kabuki syndromes respectively caused by pathogenic variants in *KMT2A* and *KMT2D* all support that *TASP1* is a disease-related gene.

**KEYWORDS**

chromosomal microarray, histone methylation, novel gene, novel syndrome, TASP1, whole exome sequencing

The marked contrast between the static genome and the highly variable transcriptome and proteome of a given organism is a testimony to the complexity of gene expression control. Key to the dynamic spatial and temporal control of gene expression is the chromatin state (packaging of DNA around histone proteins), which in turn dictates DNA accessibility to the transcriptional machinery (Greer & Shi, 2012). There are numerous posttranslational modifications of histones for example, phosphorylation, acetylation, ubiquitylation, and methylation that net effect of which determines the level of gene expression (Tan et al., 2011). One of the best known forms of histone methylation involves histone 3 (H3) lysine residues, which can be monomethylated, dimethylated, or trimethylated (Greer & Shi, 2012). The state of methylation of a number of key lysine residues represents an important component of the epigenetic mark which correlates with gene expression states, and its dynamicity is achieved by the opposing action of methyltransferases and demethylases (Young, Dimaggio, & Garcia, 2010).

While histone methylation has been extensively studied in the fields of cancer, there is a growing appreciation that abnormal histone methylation plays an important etiological role in intellectual disability whether in isolation or as part of multiple congenital anomalies syndromes (Mastrototaro, Zaghi, & Sessa, 2017). Some of these monogenic forms of intellectual disability are well established in the literature such as Sotos syndromes, which is caused by mutations in *NSD1* encoding a histone methyltransferase (Kurotaki et al., 2002). The list of intellectual disability syndromes caused by mutations in genes encoding various players in histone methylation has been expanding quickly due to the growing use of genome sequencing. A few notable examples include Wiedemann–Steiner syndrome and Kabuki syndrome 1, which are caused by mutations in *KMT2A* (lysine methyltransferase 2A) and *KMT2D* (lysine methyltransferase 2D), respectively (Jones et al., 2012; Ng et al., 2010).

Factors that control the activity of methyltransferases and demethylases represent an additional layer of complexity in the dynamic fine-tuning of the epigenetic mark. In view of the established role of histone methylation in intellectual disability syndromes, genes which encode these factors are attractive candidates in the search for novel candidate genes (Larizza & Finelli, 2019). In this report, we

suggest one such gene is *TASP1* (MIM# 608270). This gene encodes the proenzyme taspase 1 (TASP1) which is autocatalytically cleaved into  $\alpha$  and  $\beta$  subunits resulting in a mature heterodimeric enzyme. This enzyme is an endopeptidase that utilizes the N-terminal threonine of its mature  $\beta$  subunit as the active-site nucleophile to proteolyze polypeptide substrates following aspartate, hence called threonine aspartase (taspase). The active TASP1 is responsible for cleaving, thus activating, a number of broad-acting nuclear factors which play important roles in transcription regulation (Hsieh, Cheng, & Korsmeyer, 2003; Niizuma, Cheng, & Hsieh, 2015). TASP1 activates the lysine methyltransferase KMT2A and KMT2D which subsequently induce *HOX* and cyclin genes that are crucial for cell fate determination and cell cycle progression, respectively (Hsieh et al., 2003; Takeda et al., 2006).

We present here four unrelated children with an autosomal recessive syndrome caused by homozygous loss-of-function variants in *TASP1*. The first two were diagnosed in a single institution in the Arabian Peninsula. The third was diagnosed in another institution in the Arabian Peninsula. The fourth was diagnosed in Germany and recruited through GeneMatcher (Sobreira, Schiettecatte, Valle, & Hamosh, 2015). The common clinical features among these unrelated children and the deleterious nature of the variants suggest that *TASP1* is a disease-associated gene responsible for a consistent, previously unrecognized syndrome. The study was approved by Al-Ain Medical District Human Research Ethics Committee, Ethics Committee of the Technical University of Munich, and the Ethics Committee of the King Faisal Specialist Hospital and Research Center.

The first child was a 4-year-old boy with global developmental delay, hypotonia, microcephaly, feeding difficulties, drooling, failure-to-thrive, recurrent respiratory infections, cryptorchidism, cardiovascular malformations (atrial septal defect [ASD] and ventricular septal defect [VSD]), hearing impairment in the right ear, clinodactyly, left single palmer crease, brachydactyly, excessive hair on trunk and extremities, a happy demeanor with frequent laughing and clapping, and distinctive facial features (Figure 1a; Supporting Information Clinical Data). Chromosomal microarray (CMA), performed at Baylor Genetics Laboratory, Houston, TX as previously

described (Gambin et al., 2017), revealed a homozygous deletion at 20p12.1 corresponding to a minimum deletion boundary of chr20:13,463,860-13,532,560 (hg19) (Figure S1A). The minimum deleted region (69 kb) only included exons 9–11 of *TASP1* (NM\_017714.2). Whole exome sequencing (WES) did not reveal any relevant pathogenic variants in known disease-causing genes potentially responsible for the phenotype. Furthermore, no sequence coverage was obtained by WES for part of the *TASP1* gene, which is consistent with the homozygous deletion affecting this gene.

The second child was a 1-year-old girl with global developmental delay, hypotonia, microcephaly, feeding difficulties, failure-to-thrive, recurrent respiratory infections, cardiovascular malformation (patent foramen ovale (PFO) and VSD), left hydronephrosis, left preauricular skin tag, right hand preaxial polydactyly, excessive hair on trunk and extremities, bilateral single palmar creases, congenital dermal melanocytosis and a tuft of hair in the sacral region, bilateral hyperopia, happy demeanor with frequent laughing, and distinctive facial features (Figure 1b; Supporting Information Clinical Data).




FIGURE 1 Continued.

CMA performed at PreventionGenetics, Marshfield, WI, as previously described (Alabdullatif, Al Dhaibani, Khassawneh, & El-Hattab, 2017) revealed a homozygous deletion at 20p12.1, corresponding to a minimum deletion boundary of chr20:13,466,774-13,593,390 (hg19) (Figure S1B). The minimum deleted region (126 kb) only includes exons 5–10 of *TASP1* (NM\_017714.2). Both parents were found to be heterozygous for this deletion and none of the five healthy siblings was found to have a homozygous deletion in this region (Figure 1b). WES did not reveal any relevant pathogenic variants. Furthermore, no coverage was obtained by WES for a part of *TASP1*, which is consistent with the homozygous deletion affecting this gene. This child was previously reported at an earlier age (Suleiman, Mundt, Sampath, & El-Hattab, 2018).

Breakpoint mapping for the first and second child was performed using polymerase chain reaction strategy where custom primers were designed to sequence across the deletion breakpoints. Breakpoint mapping revealed that both children had the exact same deletion which was larger than the minimum deletions predicted by CMA. The *TASP1* deletion has a size of 149.4 kb spanning chr20:13,448,380-13,597,783 (hg19) and included the entirety of exons 5–11 (Figure 1e; Figure S2). This deletion includes the active site of *TASP1* and is predicted to lead to a frameshift (NC\_000020.10:g.13448380\_13597783del [p.Ser329Valfs\*15]), supporting the loss-of-function nature of this variant. Breakpoint mapping also provided an insight into the possible mechanism of this deletion. Reviewing the sequence around breakpoints revealed regions of microhomology nucleotides TG (Figure S2). These TG

nucleotides at both 5' and 3' breakpoints possibly cause the microhomology-mediated deletion (Hastings, Ira, & Lupski, 2009). The two children were from different countries of the Arabian Peninsula. To investigate whether this deletion is a founder or recurrent, we performed autozygote analysis for their families as previously described (Alkuraya, 2016). They were found to have the same haplotype indicating that this deletion is a founder Arab variant (Figure S3).

The third child was a 3-year-old boy with global developmental delay, hypotonia, microcephaly, drooling, feeding difficulties, failure-to-thrive, short stature, cryptorchidism, an episode of generalized tonic clonic seizure, happy demeanor, and distinctive facial features (Figure 1c; Supporting Information Clinical Data). CMA did not reveal any pathogenic copy number variants (CNVs). WES, performed at Medical Diagnostic Laboratory, Riyadh, Saudi Arabia as previously described (Monies et al., 2017), revealed a homozygous missense variant in *TASP1* (NM\_017714.2): c.701C>T (p.Thr234Met; NC\_000020.10:g.13514763G>A). Both parents were heterozygous carriers and the variant was confirmed by Sanger sequencing (Figure 1c,e; Figure S1C). This variant was not found in over 140,000 individuals of the Genome Aggregation Database (gnomAD, <http://gnomad.broadinstitute.org/>). The amino acid residue p.Thr234 of *TASP1* is highly conserved during evolution and the p.Thr234Met missense variant was found consistently to be disease-causing or damaging by multiple in silico prediction tools. Furthermore, the p.Thr234 residue is the active-site nucleophile to proteolyze polypeptide substrates (Takeda et al., 2015). In addition, the

**FIGURE 1** (a) Pedigrees and photos of the first child showing the following distinctive facial features: prominent glabella, excessive forehead hair, thick and arched eyebrows with synophrys, epicanthus, downslanted palpebral fissures, hypertelorism, periorbital fullness with thick eyelids, broad nasal bridge, long and smooth philtrum, wide mouth with thin upper lip and thick lower lip, microretrognathia, prominent low-set ears, and a webbed neck. (b) Pedigrees and photos of the second child showing the following distinctive facial features: excessive forehead hair, arched and thick eyebrows with synophrys, epicanthus, downslanted palpebral fissures, hypertelorism, periorbital fullness with thick eyelids, broad nasal bridge, long and smooth philtrum, thin upper lip, downturned corners of mouth, microretrognathia, prominent low-set ears with overfolded right ear helix, and a webbed neck. (c) Pedigrees and photos of the third child showing the following distinctive facial features: prominent glabella, thick and arched eyebrows with synophrys, epicanthus, hypertelorism, thick eyelids with periorbital fullness, broad nasal bridge, long and smooth philtrum, wide mouth with thin upper lip and thick lower lip, and low-set prominent ears. (d) Pedigree of the fourth child. Parents did not give permission to publish photographs of this child. The child had the following distinctive facial features: excessive forehead hair, thick, and arched eyebrows with synophrys, epicanthus, hypertelorism, periorbital fullness, broad nasal bridge, microretrognathia, high-arched palate, and low-set and prominent ears with overfolded ear helix. (e) Diagram presenting the *TASP1* gene, *TASP1* protein, and the reported variants. The 14 exons of *TASP1* (NM\_017714.2) are presented as black rectangles that are numbered from 1 to 14. Note that exon 1 of *TASP1* is noncoding and the start Met is in exon 2. Corresponding chromosomal position of *TASP1*: chr20: 13,368,036 - 13,621,583 (hg19). *TASP1* protein (NP\_060184.2) is presented as blue and green rectangle. The blue/light part represent the alpha subunit (amino acids 1–233) and the green/dark part is the beta subunit (amino acid 234–420). Note that the p.Thr234 is the first amino acid of the beta subunit and the active-site nucleophile to proteolyze polypeptide substrates (Hsieh et al., 2003; Khan, Dunn, & Tong, 2005). The red/dark rectangle on the top represent the deletion of exons 5–11; chromosomal position: chr20: 13,448,380 - 13,597,783 (hg19). The location of nonsense variant c.199C>T (p.Arg67\*) and the missense variant c.701C>T (p.Thr234Met) are marked at the bottom of the diagram. Note that the amino acid residue p.Thr234 of *TASP1* is highly conserved during evolution with a 5.7899 score in GERP (<http://mendel.stanford.edu/SidowLab/downloads/gerp/>), a 0.953 mammalian score in phyloP20way and a 9.362 vertebrate score in phyloP100way (<https://omictools.com/phyloP-tool/>), a 0.984 mammalian score in phastCons20way and 1 vertebrate score in phastCons100way (<http://compugen.cshl.edu/phast/>), and a 4.4549 score in MutationAssessor (<http://mutationassessor.org/r3/>). Furthermore, the p.Thr234Met missense variant was found consistently to be disease causing or damaging by multiple in silico prediction tools including: SIFT (<http://sift.bii.a-star.edu.sg/>) (score 0; converted rankscore 0.9122), MutationTaster (<http://www.mutationtaster.org/>) (accuracy 1; converted rankscore 0.8103), DANN ([https://cbcl.ics.uci.edu/public\\_data/DANN/](https://cbcl.ics.uci.edu/public_data/DANN/)) (score 0.9992), FATHMM (<http://fathmm.biocompute.org.uk/>) (score -3.5; converted rankscore 0.9466), MetaSVM (<https://omictools.com/meta-svm-tool/>) (score 1.0907; rankscore 0.9927), and Provean (<http://provean.jcvi.org/index.php>) (score -5.49; converted rankscore 0.8573). Variants have been submitted to Global Variome shared LOVD (<http://www.lovd.nl>) and ClinVar (<https://www.ncbi.nlm.nih.gov/clinvar/>)

**TABLE 1** Clinical features of the reported children

Age* and gender <i>TASP1</i>	1 4-year boy Hom ex 5–11 del	2 1-year girl Hom ex 5–11 del	3 3-year boy Hom c.701 C>T (p.T234M)	4 4-year boy Hom c.199 C>T (p.R67*)	
Neurodevelopment					
Developmental delay	+	+	+	+	4/4
Microcephaly	+	+	+	+	4/4
Hypotonia	+	+	+	+	4/4
Feeding and growth					
Feeding difficulties	+	+	+	+	4/4
Failure to thrive	+	+	+	-	3/4
Happy demeanor	+	+	+	-	3/4
Recurrent respiratory infections	+	+	-	+	3/4
Cardiovascular malformation	VSD/ASD	PFO/VSD	-	VSD	3/4
Cryptorchidism	+	NA	+	+	3/3
Single palmar crease	+	+	-	+	3/4
Distinctive facial features					
Prominent glabella	+	-	+	-	2/4
Excessive forehead hair	+	+	-	+	3/4
Thick eyebrows	+	+	+	+	4/4
Arched eyebrows	+	+	+	+	4/4
Synophrys	+	+	+	+	4/4
Epicanthus	+	+	+	+	4/4
Downslanted palpebral fissures	+	+	-	-	2/4
Hypertelorism	+	+	+	+	4/4
Thick eyelids	+	+	+	-	3/4
Periorbital fullness	+	+	+	+	4/4
Broad nasal bridge	+	+	+	+	4/4
Long and smooth philtrum	+	+	+	-	3/4
Wide mouth	+	-	+	-	2/4
Thin upper lip	+	+	+	-	3/4
Thick lower lip	+	-	+	-	2/4
Downturned corners of mouth	-	+	-	-	1/4
Microretrognathia	+	+	-	-	2/4
High arched palate	-	-	-	+	1/4
Low-set ears	+	+	+	+	4/4
Prominent ears	+	+	+	+	4/4
Overfolded ear helix	-	+	-	+	2/4
Webbed neck	+	+	-	-	2/4
Less common manifestations					
Dermatological					
Hirsutism	+	+	-	-	2/4
Preauricular skin tag	-	+	-	+	2/4
Congenital dermal melanocytosis	-	+	-	-	1/4
Skeletal					
Polydactyly	-	+	-	-	1/4
Brachydactyly	+	-	-	-	1/4

(Continues)

**TABLE 1** (Continued)

Age* and gender	1	2	3	4	
TASP1	4-year boy	1-year girl	3-year boy	4-year boy	
	Hom ex 5–11 del	Hom ex 5–11 del	Hom c.701 C>T (p.T234M)	Hom c.199 C>T (p.R67*)	
Clinodactyly	+	–	–	–	1/4
Short stature	–	–	+	–	1/4
Abdominal/gastrointestinal					
Inguinal hernia	–	–	–	+	1/4
Drooling	+	–	+	–	2/4
Renal					
Hydronephrosis	–	+	–	–	1/4
Neurological					
Seizures	–	–	+	+	2/4
Hearing impairment	+	–	–	–	1/4
Ophthalmologic					
Hyperopia	–	+	–	–	1/4
Strabismus	–	–	–	+	1/4
Amblyopia	–	–	–	+	1/4
Pale optic disc	–	–	–	+	1/4

\*Age at last examination.

Abbreviations: ASD, atrial septal defect; NA, not applicable; PFO, patent foramen ovale; VSD, ventricular septal defect.

substitution of p.Thr234 with Alanine has already been shown in vitro to render TASP1 inactive (Hsieh et al., 2003). Given the novelty, the conservation, the location at the active site of TASP1, the consistent in silico prediction of the variant, and the strikingly overlapping phenotype to other affected children, the missense variant c.701 C>T (p.Thr234Met) is very likely to be disease-related.

The fourth child was a 4-year-old boy with global developmental delay, hypotonia, microcephaly, seizures, feeding difficulties, recurrent respiratory infections, VSD, cryptorchidism, inguinal hernias, pale optic disks, convergent strabismus, amblyopia of left eye, periauricular skin tag on the left tragus, bilateral single palmar crease, and distinctive facial features (Figure 1d; Supporting Information Clinical Data). CMA did not reveal any pathogenic CNVs. WES, performed at the Institute of Human Genetics, Klinikum rechts der Isar, Technical University of Munich, Munich, Germany as previously described (Wortmann et al., 2015), revealed a homozygous nonsense variant in TASP1 (NM\_017714.2): c.199 C>T (p.Arg67\*) (NC\_000020.10:g.13605846 G>A). This variant was confirmed by Sanger sequencing and both parents were confirmed to be heterozygous carriers (Figure 1d,e; S1D). This variant was reported once in a heterozygous state in gnomAD with an allele frequency of  $4 \times 10^{-6}$  indicating that it is an extremely rare variant. The pathogenicity of this nonsense variant is supported by the expected outcome which is a severely truncated protein or nonsense-mediated decay most likely leading to a complete loss of function.

The four unrelated children presented here shared common features including developmental delay with hypotonia and microcephaly, feeding difficulties with failure-to-thrive, recurrent

respiratory infections, cardiovascular malformations, cryptorchidism, happy demeanor, and the following distinctive facial features: excessive forehead hair, arched and thick eyebrows with synophrys, epicanthus, hypertelorism, thick eyelids with periorbital fullness, broad nasal bridge, long and smooth philtrum, thin upper lip, and low-set prominent ears (Table 1). All had homozygous loss-of-function variants in TASP1 (Figure 1e). The consistency of the phenotype observed in multiple unrelated individuals, the family co-segregation studies, critical biological function of TASP1, and the deleterious nature of the TASP1 variants support that TASP1 is a disease-related gene and TASP1 deficiency leads to a distinctive syndrome characterized by developmental delay, respiratory infections, cryptorchidism, cardiac anomalies, happy demeanor, and distinctive facial features. Interestingly, the reported children shared unique neurobehavioral features characterized by severe expressive language impairment with better receptive skills and happy affect with frequent laughing that is reminiscent of Angelman syndrome. Such characteristic neurobehavioral features along with the common distinctive facial appearance can allow the clinical recognition of this syndrome.

Of note, the phenotype observed in the children reported here overlaps with Wiedemann–Steiner and Kabuki syndromes, which are caused by loss-of-function monoallelic pathogenic variants in KMT2A and KMT2D, respectively. Wiedemann–Steiner syndrome is characterized by developmental delay, distinctive facial features (long eyelashes, thick or arched eyebrows, synophrys, broad nasal bridge, downslanted and vertically narrow palpebral fissures, hypertelorism, epicanthus, low-set ears, thin upper lip, and exaggerated Cupid's



bow), hypertrichosis cubiti, feeding difficulties, and growth failure (short stature, microcephaly, and failure-to-thrive), skeletal malformations (hands and feet anomalies, scoliosis, rib anomalies, and sacral dimple), and cardiac malformations (ASD and PDA; Jones et al., 2012). Kabuki syndrome is characterized developmental delay, typical facial features (long palpebral fissures, thick and arched eyebrows, short columella with depressed nasal tip, and large prominent ears), skeletal anomalies (joint hyperlaxity, kyphoscoliosis, brachydactyly, and clinodactyly), persistence of fetal fingertip pads, recurrent infections, feeding problems, growth failure, seizures, congenital heart defects, genitourinary and gastrointestinal anomalies, cleft lip and palate, ptosis, strabismus, hirsutism, hypodontia, and hearing loss (Adam, Hudgins, & Hannibal, 1993). The homozygous *TASP1* loss-of-function variants in the reported children here are expected to result in impaired activation of the *TASP1* downstream targets *KMT2A* and *KMT2D* explaining the overlap between the features of Wiedemann-Steiner and Kabuki syndromes and the phenotype of these children. This observation further supports that *TASP1* is a novel disease-related gene which is associated with a syndrome overlapping with Wiedemann-Steiner and Kabuki syndromes as these syndromes are related to abnormal histone modification.

Furthermore, *taspase 1*-null mice have been generated and found to share some features with the reported children with the identified homozygous *TASP1* loss-of-function variants. *Taspase 1*<sup>-/-</sup> mice were found to be smaller in size compared with wild-type and heterozygous knock-out mice. This phenotype commences *in utero* with embryos being significantly smaller and the animals who survived the newborn period were smaller through adulthood. *Taspase 1*<sup>-/-</sup> mouse embryonic fibroblasts exhibit impaired proliferation with downregulation of the cyclin genes supporting that the small size is due to deregulation of signaling pathways resulting in impaired cell proliferation (Takeda et al., 2006). Similar to *taspase1*-null mice, children with *TASP1* homozygous loss-of-function variants demonstrated growth impairment with microcephaly and failure-to-thrive. *Taspase 1* deficient mice also demonstrated skeletal abnormalities of vertebra, ribs, and sternum (Takeda et al., 2006). Skeletal abnormalities were not major features in the reported children, however, digital deformities including polydactyly, brachydactyly, and clinodactyly have been observed each in one child. As *taspase 1* is essential for the cleavage and activation of *KMT2A* and *KMT2D*, *taspase 1* deficient cells demonstrated markedly decreased histone H3 methylation (Takeda et al., 2006). Defects of *TASP1* downstream transcription factors can potentially contribute to the observed phenotype. During mouse head formation, *TASP1*-mediated cleavage of the general transcription factor *TFIIA* (*GTFIIA*) regulates cell proliferation and morphogenesis by maintaining limited transcription of the negative cell cycle regulators. *TASP1* deficiency in mice has been shown to lead to multiple craniofacial malformations associated with inadequate cell proliferation. Evaluation of mice expressing noncleavable *TASP1* targets revealed that *TFIIA* is the principal *TASP1* substrate that orchestrates craniofacial morphogenesis (Niizuma et al., 2015;

Takeda et al., 2015). Therefore, defective *GTFIIA* may play a role in the observed distinctive facial features in individuals with *TASP1* deficiency. *KMT2A*, which is important in neuronal differentiation (Lim et al., 2009), may play a role in the neurodevelopmental phenotype.

In conclusion, we show that the presence of homozygous loss-of-function variants in *TASP1* is associated with a recognizable syndrome. The phenotypic overlap with Wiedemann-Steiner and Kabuki syndromes and the established role of *TASP1* in the activation of *KMT2A* and *KMT2D* strongly hint at the potential pathogenesis which is likely related to abnormal histone modification. The identification of more cases is needed to further delineate the phenotypic spectrum of this newly described syndrome.

## CONFLICT OF INTERESTS

S. S., T. J., M. M., and L. W. are employees of PreventionGenetics, LLC. B. W. is an employee of Baylor Genetics.

## ORCID

Korbinian M. Riedhammer  <http://orcid.org/0000-0002-7503-5801>

Fowzan S. Alkuraya  <http://orcid.org/0000-0003-4158-341X>

Ayman W. El-Hattab  <http://orcid.org/0000-0002-5737-5271>

## REFERENCES

- Adam, M. P., Hudgins, L., & Hannibal, M. (1993). In Adam, MP, Ardinger, HH, Pagon, RA, Wallace, SE, Bean, LJ, Stephens, K, & Amemiya, A (Eds.), *GeneReviews*®. Seattle, WA: University of Washington. <http://www.ncbi.nlm.nih.gov/books/NBK62111/>
- Alabdullatif, M. A., Al Dhaibani, M. A., Khassawneh, M. Y., & El-Hattab, A. W. (2017). Chromosomal microarray in a highly consanguineous population: Diagnostic yield, utility of regions of homozygosity, and novel mutations. *Clinical Genetics*, 91(4), 616–622. <https://doi.org/10.1111/cge.12872>
- Alkuraya, F. S. (2016). Discovery of mutations for Mendelian disorders. *Human Genetics*, 135(6), 615–623. <https://doi.org/10.1007/s00439-016-1664-8>
- Gambin, T., Yuan, B., Bi, W., Liu, P., Rosenfeld, J. A., Coban-Akdemir, Z., & Stankiewicz, P. (2017). Identification of novel candidate disease genes from de novo exonic copy number variants. *Genome Medicine*, 9(1), 83. <https://doi.org/10.1186/s13073-017-0472-7>
- Greer, E. L., & Shi, Y. (2012). Histone methylation: A dynamic mark in health, disease and inheritance. *Nature Reviews Genetics*, 13(5), 343–357. <https://doi.org/10.1038/nrg3173>
- Hastings, P. J., Ira, G., & Lupski, J. R. (2009). A microhomology-mediated break-induced replication model for the origin of human copy number variation. *PLOS Genetics*, 5(1), e1000327. <https://doi.org/10.1371/journal.pgen.1000327>
- Hsieh, J. J. -D., Cheng, E. H. -Y., & Korsmeyer, S. J. (2003). *Taspase1*: A threonine aspartase required for cleavage of MLL and proper HOX gene expression. *Cell*, 115(3), 293–303.
- Jones, W. D., Dafou, D., McEntagart, M., Woollard, W. J., Elmslie, F. V., Holder-Espinasse, M., & Simpson, M. A. (2012). De novo mutations in MLL cause Wiedemann-Steiner syndrome. *American Journal of Human Genetics*, 91(2), 358–364. <https://doi.org/10.1016/j.ajhg.2012.06.008>
- Khan, J. A., Dunn, B. M., & Tong, L. (2005). Crystal structure of human *Taspase1*, a crucial protease regulating the function of MLL. *Structure*

- (London, England: 1993), 13(10), 1443–1452. <https://doi.org/10.1016/j.str.2005.07.006>
- Kurotaki, N., Imaizumi, K., Harada, N., Masuno, M., Kondoh, T., Nagai, T., & Matsumoto, N. (2002). Haploinsufficiency of NSD1 causes Sotos syndrome. *Nature Genetics*, 30(4), 365–366. <https://doi.org/10.1038/ng863>
- Larizza, L., & Finelli, P. (2019). Developmental disorders with intellectual disability driven by chromatin dysregulation: Clinical overlaps and molecular mechanisms. *Clinical Genetics*, 95(2), 231–240. <https://doi.org/10.1111/cge.13365>
- Lim, D. A., Huang, Y. -C., Swigut, T., Mirick, A. L., Garcia-Verdugo, J. M., Wysocka, J., & Alvarez-Buylla, A. (2009). Chromatin remodelling factor Mll1 is essential for neurogenesis from postnatal neural stem cells. *Nature*, 458(7237), 529–533. <https://doi.org/10.1038/nature07726>
- Mastrototaro, G., Zaghi, M., & Sessa, A. (2017). Epigenetic mistakes in neurodevelopmental disorders. *Journal of Molecular Neuroscience: MN*, 61(4), 590–602. <https://doi.org/10.1007/s12031-017-0900-6>
- Monies, D., Abouelhoda, M., AlSayed, M., Alhassnan, Z., Alotaibi, M., Kayyali, H., & Alkuraya, F. S. (2017). The landscape of genetic diseases in Saudi Arabia based on the first 1000 diagnostic panels and exomes. *Human Genetics*, 136(8), 921–939. <https://doi.org/10.1007/s00439-017-1821-8>
- Ng, S. B., Bigham, A. W., Buckingham, K. J., Hannibal, M. C., McMillin, M. J., Gildersleeve, H. I., & Shendure, J. (2010). Exome sequencing identifies MLL2 mutations as a cause of Kabuki syndrome. *Nature Genetics*, 42(9), 790–793. <https://doi.org/10.1038/ng.646>
- Niizuma, H., Cheng, E. H., & Hsieh, J. J. (2015). Taspase 1: A protease with many biological surprises. *Molecular & Cellular Oncology*, 2(4), e999513. <https://doi.org/10.1080/23723556.2014.999513>
- Sobreira, N., Schiettecatte, F., Valle, D., & Hamosh, A. (2015). GeneMatcher: A matching tool for connecting investigators with an interest in the same gene. *Human Mutation*, 36(10), 928–930. <https://doi.org/10.1002/humu.22844>
- Suleiman, J., Mundt, M., Sampath, S., & El-Hattab, A. W. (2018). TASP1 is deleted in an infant with developmental delay, microcephaly, distinctive facial features, and multiple congenital anomalies. *Clinical Genetics*, 94(1), 170–173. <https://doi.org/10.1111/cge.13258>
- Takeda, S., Chen, D. Y., Westergard, T. D., Fisher, J. K., Rubens, J. A., Sasagawa, S., & Hsieh, J. J. -D. (2006). Proteolysis of MLL family proteins is essential for taspase1-orchestrated cell cycle progression. *Genes & Development*, 20(17), 2397–2409. <https://doi.org/10.1101/gad.1449406>
- Takeda, S., Sasagawa, S., Oyama, T., Searleman, A. C., Westergard, T. D., Cheng, E. H., & Hsieh, J. J. (2015). Taspase1-dependent TFIIA cleavage coordinates head morphogenesis by limiting Cdkn2a locus transcription. *The Journal of Clinical Investigation*, 125(3), 1203–1214. <https://doi.org/10.1172/JCI77075>
- Tan, M., Luo, H., Lee, S., Jin, F., Yang, J. S., Montellier, E., & Zhao, Y. (2011). Identification of 67 histone marks and histone lysine crotonylation as a new type of histone modification. *Cell*, 146(6), 1016–1028. <https://doi.org/10.1016/j.cell.2011.08.008>
- Wortmann, S. B., Ziętkiewicz, S., Kousi, M., Szklarczyk, R., Haack, T. B., Gersting, S. W., & Wevers, R. A. (2015). CLPB mutations cause 3-methylglutaconic aciduria, progressive brain atrophy, intellectual disability, congenital neutropenia, cataracts, movement disorder. *American Journal of Human Genetics*, 96(2), 245–257. <https://doi.org/10.1016/j.ajhg.2014.12.013>
- Young, N. L., Dimaggio, P. A., & Garcia, B. A. (2010). The significance, development and progress of high-throughput combinatorial histone code analysis. *Cellular and Molecular Life Sciences: CMLS*, 67(23), 3983–4000. <https://doi.org/10.1007/s00018-010-0475-7>

## SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

**How to cite this article:** Suleiman J, Riedhammer KM, Jicinsky T, et al. Homozygous loss-of-function variants of *TASP1*, a gene encoding an activator of the histone methyltransferases KMT2A and KMT2D, cause a syndrome of developmental delay, happy demeanor, distinctive facial features, and congenital anomalies. *Human Mutation*. 2019;1–8. <https://doi.org/10.1002/humu.23844>