

# Research Article

## AUTONOMIC CARDIAC CONTROL IN DEPRESSED ADOLESCENTS

Michelle L. Byrne, B.A.,<sup>1,2</sup> Lisa Sheeber, Ph.D.,<sup>3</sup> Julian G. Simmons, B.Sc.,<sup>1,2</sup> Betsy Davis, Ph.D.,<sup>3</sup> Joann Wu Shortt, Ph.D.,<sup>4</sup> Lynn Fainsilber Katz, Ph.D.,<sup>5</sup> and Nicholas B. Allen, Ph.D.<sup>1,2\*</sup>

**Background:** *The aim of this study was to identify the aspects of cardiac physiology associated with depressive disorder early in life by examining measures of autonomic cardiac control in a community-based sample of depressed adolescents at an early phase of illness, and matched on a number of demographic factors with a nondepressed comparison group. Methods:* Participants were 127 adolescents (44 boys), ages 14–18, who formed two demographically matched groups of clinically depressed and nondepressed participants. Adolescents were excluded if they evidenced comorbid externalizing or substance-dependence disorders, were taking medications with known cardiac effects, or reported regular nicotine use. Resting measures of heart rate, respiratory sinus arrhythmia, skin conductance level, blood pressure, and pre-ejection period were collected. **Results:** Depressed adolescents had resting heart rates significantly higher than those of healthy adolescents. No other measure of autonomic functioning differentiated the groups. Post hoc analyses were conducted to examine the influence of illness chronicity, severity, comorbidity, and sex on cardiac psychophysiology. These variables did not appear to exert a significant influence on the findings. **Conclusions:** Our findings suggest that neither autonomic cardiac control, illness chronicity, or severity, nor medication effects fully explain resting heart rate differences between depressed and nondepressed adolescents. Future research on depression and heart rate should consider mechanisms other than sympathetic or parasympathetic control as potential explanations of heart rate differences, including blood-clotting mechanisms, vascular and endothelial dysfunction of the coronary arteries, and inflammatory immune system response. *Depression and Anxiety 27:1050–1056, 2010.* © 2010 Wiley-Liss, Inc.

**Key words:** heart rate; autonomic nervous system; major depressive disorder; cardiovascular diseases

### INTRODUCTION

Depression is a strong independent predictor of coronary heart disease (CHD).<sup>[1]</sup> Strikingly, the magnitude of the association has been found to be on par with that of known behavioral and biological risk

*The authors disclose the following financial relationships within the past 3 years: Contract grant sponsor: National Institute of Mental Health; Contract grant number: R01 MH65340.*

\*Correspondence to: Nicholas B. Allen, Psychological Sciences, University of Melbourne, Parkville, Vic. 3052, Australia. E-mail: nba@unimelb.edu.au

Work performed at Oregon Research Institute, Eugene, Oregon.

Received for publication 7 March 2010; Revised 9 May 2010; Accepted 10 May 2010

DOI 10.1002/da.20717

Published online 11 June 2010 in Wiley Online Library (wileyonlinelibrary.com).

<sup>1</sup>ORYGEN Youth Health Research Centre, University of Melbourne, Melbourne, Vic., Australia

<sup>2</sup>Department of Psychological Sciences, University of Melbourne, Melbourne, Vic., Australia

<sup>3</sup>Oregon Research Institute, Eugene, Oregon

<sup>4</sup>Oregon Social Learning Center, Eugene, Oregon

<sup>5</sup>University of Washington, Seattle, Washington

factors such as smoking or high cholesterol.<sup>[2]</sup> Despite the clear associations between depressive and cardiac conditions, the mechanisms underlying the association are not yet well understood.<sup>[3]</sup>

To date, most of the research on the association between depressive disorder and cardiac functioning has been conducted in adults, presumably because cardiac disease is typically of adult onset. However, depression is very frequently an adolescent-onset disorder.<sup>[4]</sup> The examination of cardiac functioning in depressed adolescents may be particularly beneficial in that, relative to adults, there is less heterogeneity with regard to disease and treatment history—factors which have been hypothesized to obscure previous research findings.<sup>[5,6]</sup> If cardiac irregularities known to predict subsequent cardiac disease are experienced disproportionately by depressed adolescents, then this knowledge would have significant public health implications. First, it would suggest that depressed youth may be a population at risk for subsequent cardiac disease and hence, an important population target for prevention efforts. This approach would build on preliminary evidence regarding the efficacy of preventive efforts aimed at unselected samples of children and adolescents as well as at those selected on the basis of biological risk factors.<sup>[7,8]</sup> Second, it would suggest that the examination of potential mechanisms to explain the comorbidity, including the identification of shared risk factors, should focus on those that are present early in the life span, well before cardiac illnesses typically emerge.

One plausible mechanism for the association between depression and CHD is dysregulation of the autonomic nervous system, characterized by greater sympathetic and/or weaker parasympathetic activity. Consistent with this model is evidence that depressed persons without known cardiac disease demonstrate higher resting heart rates than do nondepressed persons.<sup>[5,9–12]</sup> Given that elevated heart rate is a predictor of subsequent cardiac events,<sup>[13,14]</sup> elevated heart rate in depressed persons may be relevant to understanding the association between cardiac and depressive disorders.

Heart rate, however, is a rather blunt index of autonomic activity as it is influenced by the heart's autonomous rate as well as by innervations from both the parasympathetic nervous system (PNS) and the sympathetic nervous system (SNS). Hence, increasing attention has been directed to comparing more specific indices of autonomic cardiac control in depressed patients. Most research has focused on respiratory sinus arrhythmia (RSA) as an index of vagal tone, and therefore of parasympathetic functioning.<sup>[15,16]</sup> Because vagal tone has been associated with a wide array of physical and mental health outcomes, including both cardiac disease and difficulties in emotion regulation,<sup>[6]</sup> it has been considered a promising candidate for investigation.

Overall, this body of research has yielded mixed results. Some research has reported lower vagal tone in

people suffering from Major Depressive Disorder,<sup>[17]</sup> whereas other studies have failed to find such differences.<sup>[5,11,18]</sup> In a recent meta-analysis, Rottenberg<sup>[6]</sup> concluded that the association between depression and cardiac vagal control was modest, noting that heterogeneity in symptoms, comorbidity, and severity could obscure relationships.

A smaller body of research has examined indices of sympathetic activity. For example, plasma norepinephrine has been found to be elevated in patients with major depression, suggesting increased sympathetic tone.<sup>[12]</sup> Again, however, findings have not been consistent. Studies examining other indices of sympathetic activation, such as systolic blood pressure, have not found between-group differences.<sup>[5]</sup>

As suggested above, one notable limitation of the research to date has been the use of adult patient samples, especially inpatient samples, whose depressive conditions are likely of unusual severity and chronicity. Thus, even in regard to the well-replicated effect of elevated heart rate, it is difficult to discern the extent to which the effect is inherent to depressive disorder versus a consequence of either lifestyle factors associated with chronic disorder or treatment effects.<sup>[5]</sup> Indeed, there are many potentially confounding variables affecting autonomic tone in studies of depressed adult patients. In particular, it appears that medication effects may account for observed associations between autonomic function and depression,<sup>[11,18,19]</sup> as controlling for antidepressant use appears to significantly attenuate associations, especially in regards to heart rate variability. Notably, medicated depressed persons have been found to differ from both nonmedicated depressed persons and nondepressed controls.<sup>[11,19]</sup>

To our knowledge, no studies have examined whether there are disturbances in cardiac physiology associated with depressive disorder during adolescence. However, a few studies have revealed associations between cardiac physiology and anxiety disorders and internalizing symptoms in children and adolescents.<sup>[20,21]</sup> For example, findings from a small sample of youth presenting for treatment of anxiety disorders indicated that they had higher heart rates and lower heart rate variability relative to nonanxious youth.<sup>[22]</sup> As well, self-reports of depressive symptoms in community samples of children have been associated with resting RSA,<sup>[23]</sup> although this effect has not been obtained consistently,<sup>[24]</sup> even with other autonomic measures such as heart rate variability and baroreflex sensitivity.<sup>[25]</sup> Though these findings suggest that aspects of cardiac physiology associated with internalizing symptoms and disorder are identifiable in youth, decades before the comorbidity between depression and cardiac illness or events would be evident, the associations have not been examined in samples of youth with case-level depressive disorder.

In this investigation, we examined aspects of cardiac physiology hypothesized to be associated with depressive disorder in a community-based sample of adolescents. We strove to provide a precise characterization

of the nature of potential autonomic dysregulation by examining a wide range of measures, including those that provide presumptively pure indices of sympathetic and parasympathetic activation. In this way, we sought to both examine whether associations between cardiac functioning and depression were evident in youth, and to provide preliminary data regarding the extent to which one potential explanation for comorbidity, excessive sympathetic as compared to parasympathetic activation, might be responsible for these effects.

## MATERIALS AND METHODS

### PARTICIPANTS

Participants were 127 adolescents (44 boys) between the ages of 14 and 18 participating in a larger study of emotion processes associated with unipolar depression. Adolescents had to meet research criteria for placement in one of two groups (Depressed,  $n = 54$  or Healthy,  $n = 73$ ). Depressed adolescents evidenced elevated scores on the Center for Epidemiological Studies—Depression Scale (CES-D,<sup>[26]</sup>  $\geq 31$  for males and  $\geq 38$  for females) during a school-based screening and subsequently met DSM IV<sup>[27]</sup> diagnostic criteria for a current unipolar depressive disorder during a diagnostic interview. Consistent with guidelines for establishing the offset of depressive episodes, a diagnosis was considered current if it was ongoing or had an offset within 2 months preceding the diagnostic interview.<sup>[27]</sup> Twelve of the depressed adolescents had one or more comorbid psychiatric disorders, including four participants with comorbid anxiety disorders. Healthy adolescents scored below an adolescent-appropriate cutoff on the CES-D ( $< 21$  for males and  $< 24$  for females), had no current or lifetime history of psychopathology, and no history of mental health treatment.

**Exclusion criteria.** Adolescents were excluded if they evidenced comorbid externalizing disorders, were regularly taking medications with known cardiac effects (e.g. antidepressants, oral contraceptives, benzodiazepines), reported regular nicotine use, or reported marijuana use on the day of testing. Allowable medications were: asthma medication (six participants), antibiotics (one participant), acne medication (one participant), and antihistamines (one participant). Three potential participants whose physiological data were missing or unusable were also excluded. Demographic data are presented in Table 1.

**TABLE 1. Demographic data**

Demographic category	Depressed ( $N = 54$ )	Healthy ( $N = 73$ )	Test statistic
Sex			$\chi^2(1) = 1.04, P = .31$
Male	16	28	
Female	38	45	
Mean age ( $SD$ )	16.06 (1.08)	16.17 (1.06)	$t(125) = -0.51, P = .61$
Md income	\$45K	\$53K	$\chi^2(1) = 4.06, P < .05$
Race and ethnicity			$\chi^2(1) = 0.01, n.s.^a$
Caucasian	37	53	
African American	2	2	
Asian	0	1	
Native American	1	0	
More than one race	10	15	
Unknown	4	2	

<sup>a</sup> $\chi^2$  is based on analysis of Caucasian–non-Caucasian dichotomy due to limited frequencies in specific non-Caucasian racial groups.

### RECRUITMENT AND ASSESSMENT PROCEDURES

The research was approved by the Institutional Review Board of the Oregon Research Institute. Informed consent was obtained from the participants and guardians after each procedure was explained. Adolescents were recruited and selected using a two-gate procedure consisting of an in-school screening and an in-home diagnostic interview.

**School screening.** Students ( $N = 4,182$ ) from 10 area high schools participated in the school screening which was conducted during class time. Approximately 70% of eligible students participated, 12% declined or had parents who declined their participation and 18% were absent on the day of the assessment. Participating students completed the CES-D, a demographic information form, and a contact form.

**Diagnostic assessment.** Interviewers conducted the Schedule of Affective Disorders and Schizophrenia—Children's Version (K-SADS,<sup>[28]</sup>) with adolescents who demonstrated elevated CES-D scores. Subsequent to the interviews, adolescents who met diagnostic criteria for a unipolar depressive disorder were invited to participate in a lab-based assessment. A healthy comparison participant, demographically matched to the depressed student, was recruited from the pool of students who scored within the normal range on the CES-D and invited to participate. Of adolescents invited to participate in the diagnostic assessment, approximately 26% declined or had parents decline their participation. Rates of decline did not vary as a function of pre-interview group status (i.e. elevated or healthy CES-D score), age, or race. Rates of decline were higher for males than females (31.6 versus 23%),  $\chi^2(1, n = 498) = 4.57, P < .05$ .

**Lab assessment.** Approximately 4% of adolescents and parents invited to participate in the lab assessment declined. The decline rate did not vary as a function of group status, age, race, or gender. The assessment consisted of a battery of questionnaire, family interaction, and interview tasks designed to obtain measures of emotional functioning, including indices of cardiac and respiratory physiology. Data for the current report were obtained during a 3-minute quiet baseline, which occurred after the questionnaires and interviews and before the family interaction tasks. During the baseline assessment, participants were seated and instructed to remain still and quiet. Participants were asked to abstain from coffee, tea, alcohol, and illicit drugs for at least 2 h before the assessment. Compliance with this instruction was confirmed on the day of the assessment via self report.

### MEASURES

**Depression screener.** The CES-D is a self-report measure of depressive symptomatology that has acceptable psychometric properties for use with adolescents.<sup>[29,30]</sup> It has a well-established

record of use as a screener for depressive symptomatology in adolescent samples.<sup>[31–33]</sup>

**Diagnostic interview.** The K-SADS interview was conducted with the adolescents only to obtain current and lifetime diagnoses. Adolescent-only diagnostic interviews have been used successfully in past research.<sup>[34,35]</sup> Interviewers, who were bachelor and masters level research staff, participated in a rigorous training program and demonstrated agreement with a senior interviewer ( $\kappa \geq .80$ ) on at least two interviews before conducting independent interviews. All interview-derived diagnoses were confirmed by supervisors who reviewed both item-endorsement and interviewers' notes. Reliability ratings were obtained on approximately 20% of the interviews, chosen at random. The average agreement was  $\kappa = .94$ .

**Brief health status assessment.** Height and weight measurements were completed by research staff at the time of the lab appointment.<sup>1</sup> Adolescents reported on their intake of caffeinated beverages, nicotine, medications, alcohol, and illicit substances on the day of the assessment as well as of their average use of caffeinated beverages and nicotine during the last 12 months.

**Physiological measures and procedures.** All data were acquired using software and equipment from the James Long Company (www.jameslong.net) except where otherwise noted. Electrocardiograph (ECG), skin conductance, and respiratory signals were input to an isolated bioelectric amplifier custom built for research ("Bioamp"). Impedance cardiogram (ICG) signals were amplified and processed by a Hutchinson Impedance Cardiogram model HIC-2000 produced by Bio-Impedance Technology Inc. (Chapel Hill, NC). Blood pressure was monitored via a Portapres portable continuous blood pressure monitor produced by Finapres Medical Systems (Amsterdam, The Netherlands).

The ECG and ICG signals were recorded using Ag-AgCl electrodes. To record the ECG signal, we used a three-lead system to maximize the *r*-wave amplitude and minimize movement artifact and *t*-wave amplitude. The ECG signals were amplified with the Bioamp, with a gain of 250 and bandpass of frequencies between 0.1–1,000 Hz. ICG signals were produced using two current electrodes placed on the back at thoracic vertebra T9 and on the neck at cervical vertebra C4<sup>25</sup> through which a 2 mA RMS current was passed. The basal thorax impedance ( $Z_0$ ) was measured (in ohms) by two electrodes placed between the current electrodes between the shoulder blades and in the mid back, and the rate of change in impedance waveform ( $dZ/dt$ ) was calculated.

The Portapres blood pressure monitor measured systolic and diastolic blood pressures, and over time calculated the mean blood pressure from the aggregate blood pressure waveform.

Respiration signal was detected by a pneumo-bellows chestband and recorded as input into the Bioamp. Skin conductance level (SCL) was measured with a 10 mA constant AC current passed through two Ag-AgCl electrodes attached to the palmar surfaces of the medial phalanges of first and second fingers of the participant's left hand. The SCL signal was also amplified by the Bioamp.

<sup>1</sup>Data from 40 participants were collected before we began taking height and weight measurements. All participants provided self-reports of height and weight. Self- and lab-assessed height and weight were highly correlated for participants on which we had both ( $r_s = .97$  and  $.97$ ). There were no differences between self-report and lab-assessed height. On average, self-reported weight was 2.75 and 6.5 lb less than lab-assessed weight for boys and girls, respectively. These amounts did not differ as a function of group. Hence, in participants missing lab-assessed data, we used self-reported height data and self-reported weight plus 2.75 or 6.5 lb, for boys and girls, respectively.

The ECGRWAVE program from the James Long Company identified *r*-waves from the ECG signal with an automated, multiple-pass, self-scaling algorithm. These signals were then visually inspected to see if the program identified the morphology of the *r*-wave correctly and manually corrected for missed or misrepresented *r*-waves. Sections of movement, noise artifact or flat line artifact were removed. Overall, this accounted for only 0.5% (in seconds) of the total data that had to be marked and removed as artifact. Other physiological data such as skin conductance, respiration, and blood pressure were visually inspected to ensure signal quality, and quantitative data were examined to ensure that values fell within a biologically plausible range (e.g. values of 0 in any signal were removed as artifact).

RSA reflects the variation in heart rate due to changes in respiration. We calculated a "time-domain" RSA variable by measuring the difference in milliseconds between the maximum inter-beat interval (IBI, or *r-r* interval) during expiration and the minimum IBI during inspiration (peak-to-trough method<sup>2</sup>). Because the RSA variable had a non-normal distribution, we transformed it to  $\log(\text{RSA})$ .  $\log(\text{RSA})$  has been used as a time-domain measure of RSA in previous research.<sup>[5,11]</sup>

Pre-ejection period (PEP) estimates the period of time commencing with onset of ventricular depolarization as represented by the ECG Q wave and ending with the onset of left ventricular ejection as indicated by the B point of the  $dZ/dt$  signal.<sup>[36]</sup> The positions in time of the Q peak in the ECG and the B point in the  $dZ/dt$  signal were detected automatically and were subsequently checked visually and edited where the detection was incorrect.

We considered RSA to be a measure of vagal tone, and therefore of PNS functioning. We used blood pressure, SCL, and PEP as indicators of SNS functioning.<sup>[36]</sup>

## RESULTS

### PRELIMINARY ANALYSES

We conducted preliminary analyses to assess the equivalence of the depressed and healthy groups. Participants in the depressed and healthy groups were similar on demographic variables, with the exception of income, which was somewhat higher in families of healthy adolescents (Table 1). They were also equivalent with regard to body mass index (BMI), and caffeine use (Table 2). Finally, as assessments were done at all times of the day, and some autonomic processes can be more active in the morning,<sup>[37]</sup> we conducted a Pearson  $\chi^2$  test that showed no between group difference in the time of day that the laboratory protocol was conducted (Table 2).

### BETWEEN GROUP DIFFERENCES IN CARDIAC CONTROL

Between-group tests on measures of heart rate and autonomic nervous system activity are presented

<sup>2</sup>We used the peak-to-trough method (time-domain) of measuring RSA instead of the spectral analysis method (frequency-domain) because we only had time-domain RSA separately for each task. However, we also had a frequency-domain RSA value for the entire 2-hour session. When we ran *t* tests by diagnostic group to look for differences in frequency-domain RSA across the entire session, no group differences emerged [ $t(127) = -0.513$ ,  $P = .609$ ]. Therefore, the time-domain RSA reported for the quiet baseline would be similar to frequency-domain RSA for the quiet baseline.

TABLE 2. Assessment and subject differences

	Depressed (N = 54)	Healthy (N = 73)	Test statistic
<i>Assessment differences</i>			
Time of day			$\chi^2(2) = 0.19, P = .91$
Morning	21	28	
Afternoon	20	26	
Evening	12	19	
<i>Subject differences</i>			
Mean BMI (SD)	23.29 (4.51)	23.02 (4.15)	$t(125) = 0.36, P = .72$
Mean typical coffee use (SD) (scale 0–9)	3.39 (2.90)	2.88 (2.92)	$t(125) = 0.98, P = .33$
Mean typical nicotine use (SD) (scale 0–9)	0.07 (0.33)	0.01 (0.12)	$t(125) = 1.45, P = .15$
Mean typical tea use (SD) (scale 0–9)	2.50 (2.48)	2.23 (2.57)	$t(125) = 0.59, P = .56$
Mean typical caffeine soda use (SD) (scale 0–9)	6.15 (1.97)	5.55 (2.48)	$t(125) = 1.47, P = .15$

TABLE 3. Physiological differences

	Depressed (N = 54)	Healthy (N = 73)	Test statistic
Mean heart rate (beats/min) (SD)	75.14 (10.38)	70.75 (9.93)	$t(125) = 2.41, P = .02$
Mean log <sub>10</sub> (RSA) (SD)	−1.05 (0.26)	−1.01 (0.23)	$t(125) = −0.82, P = .40$
Mean skin conductance level (micromhos) (SD)	10.05 (3.59)	11.17 (4.87)	$t(125) = −1.43, P = .16$
Mean systolic blood pressure (SD)	112.62 (29.89)	116.24 (22.60)	$t(125) = −0.78, P = .44$
Mean diastolic blood pressure (SD)	62.23 (17.52)	61.63 (14.69)	$t(125) = 0.21, P = .83$
Mean pre-ejection period (s) (SD)	0.13 (0.02)	0.13 (0.02)	$t(125) = 0.82, P = .41$

in Table 3. Consistent with hypotheses, depressed adolescents had resting heart rates significantly higher than those of healthy adolescents. Contrary to expectations, no other measure of autonomic functioning differentiated depressed from healthy adolescents.

Because as described earlier, clinical characteristics of disorder may influence findings,<sup>[5]</sup> post hoc analyses were conducted to examine the influence of illness chronicity, severity, and comorbidity on cardiac psychophysiology. Overall, these variables did not appear to exert a significant influence on the findings. Analyses conducted within the depressed group indicated that there were no significant correlations between either the total duration of MDD episodes (past and current) or the age of the onset of the first MDD episode and any autonomic measure ( $r_s = -.06$  to  $.12, P = .40$  to  $.91$ ). Similarly, analyses comparing adolescents experiencing mild episodes/episodes in partial remission to those with moderate to severe episodes indicated no group differences for any autonomic measure ( $t = -1.88$  to  $1.32, P = .07$  to  $.83$ ). Our depressed sample included only four participants who had current comorbid anxiety disorders. This number was too low to run any statistical analyses to control for anxiety. However, we re-ran the  $t$  tests without these four participants and the pattern of findings was unchanged. Therefore, we conclude that the presence of comorbid anxiety disorders did not account for the significant difference in heart rate nor the absence of significant differences in other autonomic measures between the depressed and healthy groups.

Finally, given evidence that sex can moderate the relations between depression and cardiac control,<sup>[38]</sup> we examined group by sex interactions for all dependent variables. No significant interactions emerged ( $F_s = 0.15$  to  $3.28, P = .07$  to  $.70$ ).

## DISCUSSION

Our findings replicate, in an adolescent community sample, the association between higher heart rate and depression that has been well-established in adult samples. Contrary to our hypothesis, however, no differences in measures of sympathetic or parasympathetic activation during rest emerged between the depressed and healthy groups, despite the higher heart rate in the depressed group. Furthermore, the association between depressive status and higher heart rate was not significantly moderated by sex, depressive severity, or illness duration as assessed by length of current episode or age of first onset. Given the importance of these null findings it is worth noting that the current study has adequate power ( $>.80$ ) to detect medium effect sizes (i.e. Cohen's  $= .45$ , critical  $t = 1.66$ ) at the conventional significance level. Further, the observed effect sizes of the null findings for these variables were generally very small, suggesting that additional statistical power would be unlikely to reveal non-null effects. The significance of these findings are twofold and lie primarily in demonstrating that: (1) community-based depressed adolescents who are young, physically healthy, and free from the effects of

cardioactive substances had higher resting heart rates than demographically matched, nondepressed controls; and (2) this difference in heart rates does not appear to be associated with concomitant differences in sympathetic or parasympathetic cardiac control.

The lack of difference in RSA between groups is consistent with other research.<sup>[11]</sup> For example, Licht et al.<sup>[11]</sup> concluded that associations between HRV and depression were largely attributable to medication effects.

Previous research has suggested that if there are no differences in parasympathetic activation, then the higher heart rate in depressed patients may be due to differences in sympathetic activation.<sup>[11]</sup> However, our findings show no difference between depressed and healthy participants in sympathetic activation as measured by baseline PEP, BP, and SCL, and as such is consistent with the negative findings of Moser et al.<sup>[5]</sup> regarding sympathetic arousal and depression. Some research has noted that higher resting heart rate in depressed patients could be attributable to medication effects<sup>[18]</sup> or lifestyle changes.<sup>[5]</sup> Data from participants who were taking medication or substances that could affect autonomic parameters were removed, as was that of participants who reported regular nicotine use. Given the young age of our sample, any lifestyle changes due to depressive symptoms would likely be less pervasive and long standing than those observed in adults, especially patient- as opposed to community-based samples. As noted earlier, preliminary analyses to compare possible fitness and lifestyle differences, indicated no differences in BMI, or in caffeine use between the depressed and healthy groups. Moreover, indices of illness severity or chronicity did not appear to influence findings.

## LIMITATIONS

It is important to note that we have only examined between group differences in the tonic levels of these measures, which does not rule out the possibility that there are between group differences in their reactivity. Some research has shown that differences in parasympathetic functioning between depressed and nondepressed groups may occur in the presence of stressors.<sup>[6,39,40]</sup> It is conceivable that depressed adolescents display autonomic dysregulation only in response to stress, and that baseline autonomic functioning is similar to controls.

However, given that a robust difference in resting heart rate was observed, our findings suggest that differences in resting parasympathetic and sympathetic innervation of the heart cannot explain the observed heart rate difference between depressed and nondepressed participants.

Another limitation of this study is that it is not based on a clinical sample, and data is not included from patients taking antidepressant medication (which may have influenced cardiovascular factors). Therefore, our sample may be biased in that it only includes adolescents with depression not severe enough to

require pharmacological treatment or is not generally representative of adolescents requiring clinical treatment. Finally, the study design does not allow us to tease apart the specificity of the observed effects of depression from those associated with other symptom dimension that are correlated or comorbid with depression. Future studies would be enhanced by the inclusion of nondepression psychiatric control groups, especially anxiety and conduct disorders groups, in order to address this issue.

## CONCLUSION

Consistent with some previous research,<sup>[5]</sup> our findings suggest that autonomic cardiac control cannot explain resting heart rate differences between depressed and nondepressed groups and that perhaps the autonomous heart rate (or “intrinsic” heart rate) of depressed patients is inherently higher. Bernardi et al.<sup>[41]</sup> suggested that the heart is capable of maintaining a heart rate independent of autonomic functioning based on a study of denervated transplanted human hearts. Given the strong test of whether higher heart rate is characteristic of depression provided by the current study (i.e. by controlling for potentially confounding factors) it seems that future research on depression and heart rate should consider mechanisms other than sympathetic or parasympathetic control as potential explanations of heart rate differences. A range of biological mechanisms have been hypothesized to explain the association between depression and heart disease, including disturbances in blood clotting mechanisms, vascular and endothelial dysfunction of the coronary arteries, and inflammatory immune system response.<sup>[2]</sup> Investigating the links between these mechanisms and higher heart rate in depression may reveal critical aspects of the pathophysiology of depression, potentially including those that confer increased risk for cardiovascular disease.

## REFERENCES

1. Nemeroff CB, Musselman DL, Evans DL. Depression and cardiac disease. *Depress Anxiety* 1998;8:71–79.
2. Goldston K, Baillie AJ. Depression and coronary heart disease: a review of the epidemiological evidence, explanatory mechanisms and management approaches. *Clin Psychol Rev* 2008;28:288–306.
3. Carney RM, Freedland KE, Veith RC. Depression, the autonomic nervous system, and coronary heart disease. *Psychosom Med* 2005;67:S29–S33.
4. Seeley JR, Lewinsohn PM. Epidemiology of mood disorders during adolescence: implications for lifetime risk. In: Allen NB, Sheeber LB, editors. *Adolescent Emotional Development and the Emergence of Depressive Disorders*. Cambridge: Cambridge University Press; 2008:33–55.
5. Moser M, Lehofer M, Hoehn-Saric R, et al. Increased heart rate in depressed subjects in spite of unchanged autonomic balance? *J Affect Disord* 1998;48:115–124.
6. Rottenberg J, Clift A, Bolden S, Salomon K. RSA fluctuation in major depressive disorder. *Psychophysiology* 2007;44:450–458.

7. Hayman LL, Williams CL, Daniels SR, et al. Cardiovascular health promotion in the schools: a statement for health and education professionals and child health advocates from the Committee on Atherosclerosis, Hypertension, and Obesity in Youth (AHOY) of the Council on Cardiovascular Disease in the Young, American Heart Association. *Circulation* 2004;110:2266–2275.
8. von Schacky C. Primary prevention of cardiovascular disease—how to promote healthy eating habits in populations? *J Public Health* 2008;16:13–20.
9. Lahmeyer HW, Bellur SN. Cardiac regulation and depression. *J Psychiatr Res* 1987;21:1–6.
10. Lechin F, van der Dijs B, Orozco B, et al. Plasma neurotransmitters, blood pressure, and heart rate during supine-resting, orthostasis, and moderate exercise conditions in major depressed patients. *Biol Psychiatry* 1995;38:166–173.
11. Lehofer M, Moser M, Hoehn-Saric R, et al. Major depression and cardiac autonomic control. *Biol Psychiatry* 1997;42:914–919.
12. Veith RC, Lewis N, Linares OA, et al. Sympathetic nervous system activity in major depression: basal and desipramine-induced alterations in plasma norepinephrine kinetics. *Arch Gen Psychiatry* 1994;51:411–422.
13. Kannel WB, Kannel C, Paffenbarger RS, Cupples LA. Heart rate and cardiovascular mortality: the Framingham study. *Am Heart J* 1987;113:1489–1494.
14. Seccareccia F, Pannoizzo F, Dima F, et al. Heart rate as a predictor of mortality: the MATISS Project. *Am J Public Health* 2001;91:1258–1263.
15. Berntson GG, Cacioppo JT, Quigley K. Respiratory sinus arrhythmia: autonomic origins, physiological mechanisms, and psychophysiological implication. *Psychophysiology* 1993;30:183–196.
16. Porges SW. Orienting in a defensive world: mammalian modifications of our evolutionary heritage. A polyvagal Theory. *Psychophysiology* 1995;32:301–318.
17. Rechlin T, Weis M, Spitzer A, Kaschka WP. Are affective disorders associated with alterations of heart rate variability? *J Affect Disord* 1994;32:271–275.
18. Bär K-J, Greiner W, Jochum T, et al. The influence of major depression and its treatment on heart rate variability and pupillary light reflex parameters. *J Affect Disord* 2004;82:245–252.
19. Licht CMM, de Geus EJC, Zitman FG, et al. Association between major depressive disorder and heart rate variability in the Netherlands Study of Depression and Anxiety (NESDA). *Arch Gen Psychiatry* 2008;65:1358–1367.
20. Mezzacappa E, Tremblay RE, Kindlon D, et al. Anxiety, antisocial behavior, and heart rate regulation in adolescent males. *J Child Psychol Psychiatry* 1997;38:457–469.
21. Pine DS, Wasserman GA, Miller L, et al. Heart period variability and psychopathology in urban boys at risk for delinquency. *Psychophysiology* 1998;35:521–529.
22. Monk C, Kovelenco P, Ellman LM, et al. Enhanced stress reactivity in paediatric anxiety disorders: implications for future cardiovascular health. *Int J Neuropsychopharmacol* 2001;4:199–206.
23. Shannon KE, Beauchaine TP, Brenner SL, et al. Familial and temperamental predictors of resilience in children at risk for conduct disorder and depression. *Dev Psychopathol* 2007;19:701–727.
24. Bosch NM, Riese H, Ormel J, et al. Stressful life events and depressive symptoms in young adolescents: modulation by respiratory sinus arrhythmia? The TRAILS study. *Biol Psychiatry* 2009;81:40–47.
25. Bosch NM. Preadolescents' somatic and cognitive-affective depressive symptoms are differentially related to cardiac autonomic function and cortisol: the TRAILS Study. *Psychosom Med* 2009;71:944–950.
26. Radloff LS. A CES-D scale: a self-report depression scale for research in the general population. *Appl Psychol Measurement* 1977;1:385–401.
27. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. 4th edition. Washington, DC: American Psychiatric Press; 1994.
28. Orvaschel H, Puig-Antich J. *Schedule for Affective Disorder and Schizophrenia for School-age Children, Epidemiologic Version*. Ft. Lauderdale, FL: Nova University, Center for Psychological Study; 1994.
29. Radloff LS. The use of the Center of Epidemiologic Studies Depression Scale in adolescents and young adults. *J Youth Adolesc* 1991;20:149–166.
30. Roberts RE, Andrews JA, Lewinsohn PM, Hops H. Assessment of depression in adolescents using the Center for Epidemiologic Studies Depression Scale. *Psychol Assess* 1990;2:122–128.
31. Asarnow JR, Jaycox LH, Duan N, et al. Effectiveness of a quality improvement intervention for adolescent depression in primary care clinics: a randomized control trial. *J Am Med Assoc* 2005;293:311–319.
32. Dierker LC, Albano AM, Clarke GN, et al. Screening for anxiety and depression in early adolescence. *J Am Acad Child Adolesc Psychiatry* 2001;40:929–936.
33. Roberts RE, Lewinsohn PM, Seeley JR. Screening for adolescent depression: a comparison of depression scales. *J Am Acad Child Adolesc Psychiatry* 1991;30:58–66.
34. Lewinsohn PM, Hops H, Roberts RE, et al. Adolescent psychopathology: I. Prevalence and incidence of depression and other DSM-III-R disorders in high school students. *J Abnorm Psychol* 1993;102:133–144.
35. Sheeber LB, Davis B, Leve C, et al. Adolescents' relationships with their mothers and fathers: associations with depressive disorder and subdiagnostic symptomatology. *J Abnorm Psychol* 2007;116:144–154.
36. Cacioppo JT, Uchino BN, Berntson GG. Individual differences in the autonomic origins of heart rate reactivity: the psychometrics of respiratory sinus arrhythmia and preejection period. *Psychophysiology* 1994;31:412–419.
37. Muller JE, Tofler GH, Stone PH. Circadian variation and triggers of onset of acute cardiovascular disease. *Circulation* 1989;79:733–743.
38. Thayer JF, Smith M, Rossy LA, et al. Heart period variability and depressive symptoms: gender differences. *Biol Psychiatry* 1998;44:304–306.
39. Guinjoan SM, Bernabó JL, Cardinali DP. Cardiovascular tests of autonomic function and sympathetic skin responses in patients with major depression. *J Neurol Neurosurg Psychiatry* 1995;58:299–302.
40. Hughes JW, Stoney CM. Depressed mood is related to high-frequency heart rate variability during stressors. *Psychosom Med* 2000;62:796–803.
41. Bernardi L, Keller F, Sanders M, et al. Respiratory sinus arrhythmia in the denervated human heart. *J Appl Physiol* 1989;67:1447–1455.

Copyright of Depression & Anxiety (1091-4269) is the property of John Wiley & Sons, Inc. and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.