

Chalcone-Flavanone Equilibrium in the Isomeric Mixture: 2',3,4,4',6'-Pentahydroxychalcone and Eriodictyol from *Stevia lucida*

Pablo A. Chacon-Morales¹, Carolina Santiago-Dugarte^{1,2} Juan M. Amaro-Luis^{1*}

¹) Natural Products Laboratory, Department of Chemistry, Faculty of Science, University of Los Andes, Mérida 5101, Venezuela

²) Biotechnology Engineering, Universidad Politécnica Salesiana, Km 19.5 Vía a la Costa, Guayaquil 090901, Ecuador

(*) jmamaroluis@gmail.com

Recibido: 29/09/2020

Revisado: 15/12/2020

Aceptado: 28/12/2020

Abstract

From the leaves and stems of *Stevia lucida* Lagasca (Asteraceae), an equilibrium chalcone-flavanone isomeric mixture, composed of 2',3,4,4',6'-pentahydroxychalcone (**8**) (eriodictyol-chalcone) and eriodictyol (**9**) [ratio **8/9** 5:3], was isolated as an isomorphous crystal. The mixture was structurally characterized by spectroscopic methods, including 1D- and 2D-NMR experiments. The presence of chalcones in the genus *Stevia* is reported here for the second time.

Keywords: Chalcones; Flavanones; Equilibrium Mixture; 2',3,4,4',6'-Pentahydroxychalcone; Eriodictyol; *Stevia*

Resumen

A partir de las hojas y tallos de *Stevia lucida* Lagasca (Asteraceae), se aisló una mezcla isomérica chalcona-flavanona en equilibrio, compuesta de 2',3,4,4',6'-pentahidroxichalcona (**8**) (eriodictyol-chalcona) y eriodictyol (**9**), como un cristal isomorfo en una proporción (**8/9**) 5:3. La mezcla se caracterizó estructuralmente mediante métodos espectroscópicos, incluidos experimentos de RMN 1D y 2D. La presencia de chalconas en el género *Stevia* se reporta aquí por segunda vez.

Palabras claves: chalconas; flavanonas; mezcla en equilibrio; 2',3,4,4',6' pentahidroxichalcona; eriodictyol; *Stevia*

Introduction

Flavonoids constitute one of the most important classes of naturally occurring phenols with interesting biological properties that demonstrate their great potential as treatment options to several human pathologies¹⁻⁴. The scientific literature on the wide range of biological and pharmacological activities of flavonoids is extensive and review articles abound⁵⁻¹². The so-called "minor flavonoids", among which stand out chalcones and flavanones, have received considerable attention in recent years, in the field of their biological activities¹³⁻¹⁸.

It is currently well established that, from a biosynthetic perspective, flavonoids originate through the "general phenylpropanoid pathway", starting with *L*-phenylalanine (**1**) and then transforming it into *trans*-cinnamic acid (**2**) which it is hydroxylated to *p*-coumaric acid (**3**) and this evolves by esterification to *p*-coumaroyl-CoA (**4**). On some plants, the latter (**4**) may be hydroxylated in position 3 to give caffeoyl-CoA (**5**), which can also serve as substrate in the next biosynthetic stage¹⁹⁻²¹ (Fig. 1). The known enzymes related to this pathway are phenylalanine ammonia-

lyase (PAL), cinnamic acid 4-hydroxylase (C4H), *p*-coumarate:CoA ligase (4CL), and 4-coumaroyl:CoA 3-hydroxylase (CC3H)²². Since the mid-1970s, substantial progress has been made on plant cell suspension cultures and these were a valuable source for the isolation and characterization of flavonoid enzymes^{23,24}. The next biosynthetic stage carried out by the enzyme chalcone synthase (CHS)²⁵ focuses on the condensation of *p*-coumaroyl-coenzyme-CoA (**4**) with three molecules of malonyl-CoA (**5**) [generated by the reaction of acetyl-CoA and CO₂ catalyzed by acetyl: CoA carboxylase (ACC)] giving the C-15 intermediate 2',4,4',6'-tetrahydroxychalcone (**6**). The most important stage in flavonoid biosynthesis is that corresponding to the conversion of chalcone (**6**) into the flavanone naringenin (**7**) since all the other flavonoids are generated from the latter. This stage is catalyzed by the chalcone isomerase (CHI) enzyme, which generates a chalcone-flavanone equilibrium²⁶.

Stevia lucida Lagasca (Asteraceae) is a widely distributed species in the Venezuelan Andean Region²⁷. From a phytochemical point of view, we have studied this species for

more than three decades and have achieved to isolate and identify several steroids²⁸, flavonoids^{28,29}, diterpenes of the labdane³⁰ and *ent*-kaurane series including an interesting new *ent*-kaurenolido³¹, longipinane sesquiterpenes³², some eudesmanolides³³, and (+)-mellein²⁹, the first isocoumarin reported for the genus *Stevia*. 7 β ,9 α -dihydroxyongipin-2-en-1-one, the most abundant sesquiterpene in *S. lucida*, has been an excellent substrate for chemical reactions^{34,35}, particularly Wagner-Meerwein rearrangements³⁶⁻³⁸, which lead to new regrouped sesquiterpene skeletons. Other researchers have also investigated *S. lucida*, describing as components of this species a new labdane diterpenic acid³⁹ and several longipinene diesters⁴⁰.

Recently, we reported the isolation and characterization of the first three chalcones present in the genus *Stevia*⁴¹. Now, we describe a fourth chalcone, the 2',3,4,4',6'-pentahydroxy-chalcone (also called eriodictyol chalcone) (**8**) obtained from *Stevia lucida*, with the particularity that it is in its natural state in equilibrium with the corresponding flavanone eriodictyol (**9**). The mixture could not be separated by conventional chromatographic methods and it was characterized as such, through of a detailed spectral study that includes 1D and 2D-NMR techniques.

Experimental

General experimental procedures

Melting point was determined using a Fisher-Johns apparatus and it is uncorrected. UV spectra were obtained in a Perkin-

Elmer spectrophotometer, Lambda 3B, using quartz cells with 1 cm thick and methanol (Merck-Uvasol) as solvent. IR measurement was obtained on a Perkin-Elmer FT-1725X spectrophotometer as KBr pellets. 1D and 2D NMR spectra in DMSO-*d*₆ were acquired using a Bruker-Avance DRX-400 instrument, operating at 400 MHz for ¹H and 100 MHz for ¹³C. Mass spectra were recorded on a Hewlett-Packard mass spectrometer, model 5890 (70 eV). TLC were developed on 0.25 mm layers of silica gel PF 254 (Merck); spots were visualized using UV light (254 and 365 nm) and subsequently by spraying with a mixture *v/v* CH₃COOH-H₂O-H₂SO₄ (20:4:1) and then heating with air-flow at 100 °C for few minutes. VCC was performed with silica gel Merck 60 (63-200 μ m, 70-230 mesh).

Plant material

Plant material (leaves and stems) was collected at "Páramo de la Negra, Municipio Rivas Dávila, Estado Mérida, Venezuela". Species was identified as *Stevia lucida* Lagasca by Eng. Juan Carmona Arzola, Department of Pharmacognosy and Organic Medicaments, Faculty of Pharmacy and Bioanalysis, University of Los Andes (ULA); a *voucher specimen* (JM Amaro-Luis & P. Chacón, No. 2332) was deposited at the Herbario MERF of this faculty.

Extraction and purification of the equilibrium mixture of **8** and **9**

The dry uncrushed leaves and stems (\approx 4.0 kg) were exhaustively extracted with ethanol in a soxhlet. The obtained solu-




Fig. 1: Biosynthesis of Flavonoids.




Fig. 2: ¹H NMR and ¹³C NMR spectra of 2'equilibrium 2',3,4,4',6' Pentahydroxychalcone (Eriodictyol-Chalcone) (**8**) and Eriodictyol (**9**).

tion was filtered and concentrated “*in vacuo*” in a rotary evaporator at temperature not exceeding 40 °C, to produce a crude extract (970 g), which was preadsorbed on silica gel and extracted successively with petroleum ether, acetone, and methanol, exhaustively in each case. Acetone-solution was concentrated under reduced pressure to dryness and a brown residue was obtained (\cong 270 g). The acetone extract was preadsorbed on silica gel and chromatographed (VLC) over silica gel 60, eluting with hexane and EtOAc in mixtures of increasing polarity. Fractions of 500 mL were collected and combined according to the TLC characteristics to afford twelve major fractions (A-L). Combined fraction “H” [53-57], eluted with hexane-EtOAc (2:3), was purified by repeated flash chromatography, Sephadex filtration chromatography,

preparative TLC and crystallization to furnish an isomorphous crystalline equilibrium mixture of **8** and **9**.

Equilibrium mixture of **8** and **9**

Liquid recovered by filtration after washing with methanol the scraped silica gel from the preparative TLC of combined fraction “H”, was concentrated to dryness yielding a pale yellow solid, whose chromatographic behavior was typical of a flavonoid. Crystallization in methanol gave yellow flakes; m.p. = 262-264 °C. $[\alpha]_D = -2.8^\circ$ (c, 0,5 MeOH). UV, λ_{\max} : (CH₃OH, nm) 225, 287, 330 (sh); IR, ν_{\max} . (cm⁻¹): 3363 (OH), 1636 (C=O), 1604 (C=C), 1521, 1447, 1160 and 1085 (C-O). EI-MS, *m/z* (%): 288 [M⁺] (69), 270 [M⁺-H₂O] (14), 179 (24), 166 (40), 163 (18), 153 (100), 152 (12), 136 (70), 123 (16).

^1H NMR (400 MHz, $\text{DMSO-}d_6$) Data corresponding to 2',3,4,4',6'-pentahydroxychalcone (**8**) $^{\bullet}$ δ_{H} : 7.94 (1H, d, $J = 15.5$ Hz, H- α), 7.61 (1H, d, $J = 15.5$ Hz, H- β), 5.88 (2H, s, H-3' and H-5'), 7.11 (1H, d, $J = 1.7$ Hz, H-2), 6.83 (1H, d, $J = 8.1$ Hz, H-5), 7.02 (1H, dd, $J = 8.1$ and 1.7 Hz, H-6), 12.60 (1H, s, OH), 10.45 (1H, s, OH), 9.66 (1H, s, OH), 9.31 (1H, s, OH); Data corresponding to eriodictyol (**9**) $^{\blacksquare}$ δ_{H} : 5.41 (1H, dd, $J = 12.4$ and 3.0 Hz, H-2), 3.22 (1H, dd, $J = 17.1$ and 12.4 Hz, H-3 α ax), 2.71 (1H, dd, $J = 17.1$ and 3.0 Hz, H-3 β ec), 5.91 (2H, s, H-6 and H-8), 6.91 (1H, bs, H-2'), 6.78 (2H, bs, H-5' and H-6'), 9.08 (1H, s, OH), 9.08 (1H, s, OH), 12.18 (1H, s, OH). ^{13}C NMR (100 MHz, $\text{DMSO-}d_6$) δ_{C} : 192.5 ($>\text{C}=\text{O}$, C- β'), 124.5 ($=\text{CH-}$, C- α), 143.7 ($=\text{CH-}$, C- β), 105.1 ($>\text{C=}$, C-1'), 165.3 ($=\text{C-O-}$, C-2'), 95.8 ($=\text{CH-}$, C-3'), 165.6 ($=\text{C-O-}$, C-4'), 95.8 ($=\text{CH-}$, C-5'), 165.3 ($=\text{C-O-}$, C-6'), 127.6 ($>\text{C=}$, C-1), 115.2 ($=\text{CH-}$, C-2), 146.6 ($=\text{C-O-}$, C-3), 149.4 ($=\text{C-O-}$, C-4), 116.8 ($=\text{CH-}$, C-5), 122.9 ($=\text{CH-}$, C-6); $\delta_{\text{C}}^{\blacksquare}$: 79.3 ($>\text{CH-}$, C-2), 43.0 ($>\text{CH}_2$, C-3), 197.2 ($>\text{C}=\text{O}$, C-4), 164.4 ($=\text{C-O-}$, C-5), 96.7 ($=\text{CH-}$, C-6), 167.5 ($=\text{C-O-}$, C-7), 96.7 ($=\text{CH-}$, C-8), 163.8 ($=\text{C-O-}$, C-8a), 102.7 ($>\text{C=}$, C-4a), 130.4 ($>\text{C=}$, C-1'), 115.3 ($=\text{CH-}$, C-2'), 146.1 ($=\text{C-O-}$, C-3'), 146.6 ($=\text{C-O-}$, C-4'), 116.3 ($=\text{CH-}$, C-5'), 118.8 ($=\text{CH-}$, C-6').

Results and Discussion

The equilibrium isomeric flavonoid mixture crystallized as yellow flakes; m.p. = 262-264 °C. The EI-MS shows a peak corresponding to a molecular ion at m/z : 288 [M^+] which, according to the NMR data, can be assigned to the molecular formula $\text{C}_{15}\text{H}_{12}\text{O}_6$. The presence in its UV spectrum of three bands at λ_{max} : 225, 287, 330 (sh) indicates that it is a chalcone-flavanone mixture⁴². An overview of the ^1H NMR and ^{13}C NMR spectra (Fig. 2), clearly shows that it is a mixture of two compounds whose ratio is 5:3 (**8/9**). Analyzing both spectra in detail, together with the ^1H - ^1H COSY, HMQC, and HMBC (Fig. 3) 2D-NMR spectra, the chalcone-related signals (identified with the sign " \bullet ") and those of the flavanone (identified with the sign " \blacksquare ") could be unequivocally assigned (see experimental; Fig. 2). Next, we will comment and discuss individually the data of each of the compounds that make up the mixture.

Compound (**8**): The molecular formula $\text{C}_{15}\text{H}_{12}\text{O}_6$ revealed ten unsaturation degrees. ^{13}C NMR data analysis (Fig. 2B) allow the characterization in the molecule of functional groups completing eight degrees of unsaturation, a ketone [δ_{C} 192.5 ($>\text{C}=\text{O}$, C-1'')] and fourteen sp^2 carbons that form seven double bonds and therefore, the remaining unsaturation should correspond to two rings, forming a system composed of two benzene nuclei. The remaining spectral data allowed to conclude that this compound is the 2',3,4,4',6'-pentahydroxychalcone (eriodictyol chalcone).

This chalcone is not very abundant in Nature, having been sporadically isolated only from four species: *Galium verum* (Rubiaceae)⁴³, *Limonium cv. 'Gold Coast'* (Plumbaginaceae)⁴⁴, *Tulipa aff. 'Apeldoorn'* (Liliaceae)⁴⁵ and *Sorghum bicolor* (Poaceae)⁴⁶, however, it has been detected by GC/MS in the peel of tomato fruits *Lycopersicon esculentum* (Solanaceae)⁴⁷, and in some medicinal plants: *Artemisia argyi* (Asteraceae)⁴⁸, *Populus sieboldii* (Salicaceae)⁴⁹ and *Pouteria lucuma* (Sapotaceae)⁵⁰. An interesting aspect to highlight is the frequency with which its biosynthesis from caffeoyl-CoA (**6**) has been induced in some plants by genetic engineering, using the enzyme chalcone synthase (CHS); in general, plants of nutritional interest have been used: Tomato (*Lycopersicon esculentum*)⁵¹, barley (*Hordeum vulgare*)⁵², rice (*Oryza sativa*)⁵³, carrot (*Daucus carota*)⁵⁴, apple (*Malus x domestica*)⁵⁵ and pear (*Pyrus communis*)⁵⁵, which support the possible interest of this chalcone as a phytochemical additive in the diet to be ingested in human consumption⁵⁶.

The spectrum of biological activity of this chalcone has been studied, highlighting its antioxidant activity (Rancimat Test // induction periods in lard at concentrations of 0.025, 0.05 and 0.1% = 20, 40, and 73 h, respectively)⁵⁷ and its relative DPPH radical scavenging equivalence (DPPH assay in acetone = 3.8 at 30 min. when tocopherol in ethanol was used as standard)⁵⁸. In an "in vitro" study on antimalarial activity, antiplasmodial effects were evaluated observing that (**8**), when it is given at 1 mM concentration, caused a relevant growth inhibition (88%) of 3D7 *Plasmodium falciparum* strain, after 48 hours incubation; in another experiment, the growth of the parasite was investigated when artemisinin, in sublethal doses ranging from 0.625 to 5.0 nM, was added to a 1.0 mM sample of (**8**), detecting considerable additive effects that induce a substantial potentiation of the antimalarial action of artemisinin⁵⁹.




Fig. 3: HMBC Spectrum Correlations of 2',3,4,4',6'-Pentahydroxychalcone (Eriodictyol-Chalcone) (**8**) and Eriodictyol (**9**).

Another activity of notable interest is the capacity of (**8**) as an antidepressant agent, which was evaluated "in vivo" in mice; the activity of the compound, at a dose of 10 mg/kg, was measured using the forced swimming test, in which it was detected that the mean and standard error of the mean (\pm SEM) for the duration of immobility of ten mice was 98.5 ± 11.0 sec; this value qualifies (**8**) as a potent antidepressant since it has a $P < 0.01$ against Fluoxetine, a reference antidepressant drug⁶⁰. Finally, it is particularly notable that this chalcone exhibits an inhibitory effect on several enzymes such as phosphoenolpyruvate carboxylase (PEPC) ($IC_{50} = 2.5$ and $8.2 \mu\text{M}$)⁶¹, protein tyrosine phosphatase 1B (PTP1B) ($IC_{50} = 1.26 \mu\text{M}$)⁶², aromatase (CYP19A1) ($IC_{50} = 2.8 \mu\text{M}$)⁶³, 5-lipoxygenase (5-LOX) ($IC_{50} = 0.043 \mu\text{M}$)⁶⁴ and cyclooxygenase (COX) ($IC_{50} = 34.0 \mu\text{M}$)⁶⁴. Consequently, it could be stated that chalcone (**8**) seems to be a potential herbicide for its ability to inhibit PEPC, a promising candidate for the treatment of a variety of inflammatory and allergic diseases due to its action on 5-LOX and COX or a therapeutic approach to prevent the development and progression of breast cancer based on its activity on CYP19A1.

Compound (**9**): The signals (typified as signals ■) in the ¹H NMR and ¹³C NMR spectra corresponding to the second component of the mixture are congruent with those of a flavanone, which was identified with eriodictyol (3',4',5,7-tetrahydroxyflavanone). This flavanone was isolated for the first time in 1906 from *Eriodictyon californicum* (Hooker and Arnott) Greene, a California medicinal plant commonly known as "Yerba Santa"⁶⁵, but its structure was not definitively established until 1929, which was also unequivocally supported by its synthesis^{66,67}. Since then, this flavanone has been reported in many higher plants⁶⁸ and particularly in species of the genus *Citrus* (Rutaceae)⁶⁹.

From the biological and pharmacological point of view, eriodictyol has a wide spectrum of activities, as can be contrasted when analyzing the recent review of Islam *et al.*⁶⁸; however, it is appropriate to highlight the most recent papers on its anti-oxidants⁷⁰, anti-inflammatory⁷¹, cardiovascular⁷², and anti-carcinogenic^{73,74} activities, as well as the critical role it plays in diabetes mellitus⁷⁵. At this time, it is particularly interesting to note that eriodictyol is described as an interesting molecule for treating COVID-19, as it is revealed by rigorous studies using a molecular docking approach, via the world's most powerful supercomputer SUMMIT⁷⁶. In these studies, it was confirmed that eriodictyol might affect the life cycle of the SARS-CoV-2 virus by interacting with various amino acids and proteins, binding to multiple targets with efficient binding energy (between -6.7 and -8.8 kcal/mol) and within the active site⁷⁷; consequently, eriodictyol can become a multi-target molecule which can be a strong candidate for the treatment of coronavirus infection when repurposed against SARS-CoV-2.

In accordance with all the above and in harmony with the title of this article, it is appropriate to make some comments about the chalcone-flavanone equilibrium. The scientific literature reveals that it is rare to find this equilibrium type in the same plant species; however, the presence of the equilibrium 2',3,4,4',6'-pentahydroxychalcone (eriodictyol chalcone)-eriodictyol [(**8**) \rightleftharpoons (**9**)], has been reported in certain common plants such as tomato (*Lycopersicon esculentum*, Solanaceae)⁴⁷, sorghum (*Sorghum bicolor*, Poaceae)⁴⁶ and poplar *Populus sieboldii*, Salicaceae)⁴⁹ and also in incubations with plant enzyme preparations obtained from fresh vegetable materials such as flowers of defined genotypes of carnation (*Dianthus caryophyllus*, Caryophyllaceae)⁷⁸ or anthers and pollen of tulips (*Tulipa* *cf.* "Apeldoorn", Liliaceae)⁷⁹.

The synthetic cyclization of chalcones into flavanones and vice versa has been widely studied⁸⁰⁻⁸². Both processes have been carried out chemically using acids [acetic acid, trifluoroacetic acid, methane sulphonic acid, amino acids, Lewis acids and mineral acids such as HCl, H₂SO₄, H₃PO₄, polyphosphoric acid, etc.], bases [alkali metal hydroxides, sodium acetate, potassium carbonate in acetone under reflux condition, amine bases, cobalt(II) Schiff-base], aqueous buffers at variable pH, acidic ion-exchange resin, phase-transfer catalysts, catalysis by Co(III)-salts complexes, bovine serum albumin (BSA), oxidizing agent as potassium ferricyanide, silica gel, zeolites, celite supported potassium fluoride, Amberlyst A-21, photochemical processes, thermal reactions at 60 °C in the solid-state or in the melt at 230 °C, microwave irradiation or under electrochemical conditions. It should be noted that the chalcone-flavanone equilibrium conforms a scaffold that can function as a dynamic covalent molecular switch, that is, a molecule that is capable of being reversibly biased towards two states in response to an external stimulus⁸³, the development of these systems (the so-called "molecular switches") have been a mature area of research during the last decade, as they have application in a variety of fields including sensors, chemical biology, and functional materials⁸⁴. Finally, the cyclization of chalcones can lead to flavonoids other than flavanones; an example of this is the transformation of the eriodictyol chalcone (**8**) into the aurone aureusidin, through a crude peroxidase, polyphenol oxidase (PPO)⁸⁵.

As conclusion, it is undeniable to say that in our case since we never use acids or bases in the extraction and isolation processes, the coexistence of the chalcone-flavanone equilibrium is natural, and must be regulated by the action of the enzyme chalcone isomerase (CHI). This equilibrium, being photo-synthetically originated, could be considered a dynamic covalent photoswitch.

Acknowledgments

The authors are grateful to CDCHT-ULA for financial support (Grant C-1935-15-08-ED). Thank are also due to Eng. Juan Carmo- na Arzola, Department of Pharmacognosy and Organic Medicaments, Faculty of Pharmacy and Bioanalysis, University of Los

Andes (ULA) for the identification of plant material. We also would like to thank Dr. Ángel E. Amesty, Instituto Universitario de Bio-Organica, Universidad de La Laguna. Tenerife, Spain, for the EIMS.

Conflicts of Interest: The authors have no conflicts of interest to declare.

References

- BH Havsteen. The biochemistry and medical significance of the flavonoids. **Pharmacology & Therapeutics**, **96**, 67-202 (2002)
- CR Ferraz, TT Carvalho, MF Manchope, NA Artero, FS Rasquel-Oliveira, V Fattori, R Casagrande, WA Verri Jr. Therapeutic potential of flavonoids in pain and inflammation: Mechanisms of action, pre-clinical and clinical data, and pharmaceutical development. **Molecules**, **25**, 762-797 (2020).
- Â Bisol, P Santos de Campos, M Lazzaron Lamers. Flavonoids as anticancer therapies: A systematic review of clinical trials. **Phytother. Res.**, **34**, 568-582 (2020).
- SS Costa, JNSS Couceiro, ICV Silva, DC Malvar, MAS Coutinho, LMM Camargo, MF Muzitano, FA Vanderlinde. Flavonoids in the therapy and prophylaxis of flu: A patent review. **Expert Opin. Ther. Patents**, **22**, 1111-1121 (2012).
- MM Jucá, FMS Cysne Filho, J Cunha de Almeida, D da Silva Mesquita, J Rodrigues de Moraes Barriga, KC Ferreira Dias, TM Barbosa, L Costa Vasconcelos, LK Almeida Moreira Leal, JE Ribeiro, SM Mendes Vasconcelos. Flavonoids: Biological activities and therapeutic potential. **Nat. Prod. Res.**, **34**, 692-705 (2020).
- AM González-Paramás, B Ayuda-Durán, S Martínez, S González-Manzano, C Santos-Buelga. The mechanisms behind the biological activity of flavonoids. **Curr. Med. Chem.**, **26**, 6976-6985 (2019).
- S Kumar, AK Pandey. Chemistry and biological activities of flavonoids: An overview. **The Scientific World Journal**, **13**, ID 162750 (2013).
- AK Verma, R Pratap. The biological potential of flavones. **Nat. Prod. Rep.**, **27**, 1571-1593 (2010).
- RJ Nijveldt, E van Nood, DEC van Hoorn, PG Boelens, K van Norren, PAM van Leeuwen. Flavonoids: A review of probable mechanisms of action and potential applications. **Am. J. Clin. Nutr.**, **74**, 418-425 (2001).
- A Ahmad, M Kaleem, Z Ahmed, H Shafiq. Therapeutic potential of flavonoids and their mechanism of action against microbial and viral infections-A review. **Food Res. Internat.**, **77**, 221-235 (2015).
- D Ravishankar, AK Rajora, F Greco, HMI Osborn. Flavonoids as prospective compounds for anti-cancer therapy. **Intern. J. Biochem. Cell Biol.**, **45**, 2821-2831 (2013).
- KE Heim, AR Tagliaferro, DJ Bobilya. Flavonoid antioxidants: Chemistry, metabolism and structure-activity relationships. **J. Nutrit. Biochem.**, **13**, 572-584 (2002).
- NK Sahu, SS Balbhadra, J Choudhary, DV Kohli. Exploring pharmacological significance of chalcone scaffold: A review. **Curr. Med. Chem.**, **19**, 209-225 (2012).
- AM Katsori, D Hadjipavlou-Litina. Recent progress in therapeutic applications of chalcones. **Expert Opin. Ther. Patents**, **21**, 1575-1596 (2011).
- DI Batovska, IT Todorova. Trends in utilization of the pharmacological potential of chalcones. **Curr. Clin. Pharmacol.**, **5**, 1-29 (2010).
- G Brahmachari. Naturally occurring flavanones: An overview. **Nat. Prod. Commun.**, **3**, 1337-1354 (2008).
- K Patel, GK Singh, DK Patel. A review on pharmacological and analytical aspects of naringenin. **Chin. J. Integr. Med.**, **24**, 551-560 (2018).
- A Roohbakhsh, H Parhiz, F Soltani, R Rezaee, M Iranshahi. Neuropharmacological properties and pharmacokinetics of the citrus flavonoids hesperidin and hesperetin-A mini-review. **Life Sciences**, **113**, 1-6 (2014).
- G Forkmann, W Heller. Biosynthesis of flavonoids. In: *Comprehensive Natural Products Chemistry, Vol 1 Polyketides and Other Secondary Metabolites Including Fatty Acids and Their Derivatives*. D Barton, K Nakanishi, O Meth-Cohn, U Sankawa (Eds), pp 713-748. Pergamon Press, London (1999).
- K Yonekura-Sakakibara, Y Higashi, R Nakabayashi. The origin and evolution of plant flavonoid metabolism. **Front. Plant Sci.**, **10**, 943 (2019).
- B Winkel-Shirley. Flavonoid biosynthesis, a colorful model for genetics, biochemistry, cell biology and biotechnology. **Plant Physiol.**, **126**, 485-493 (2001).
- JL Ferrer, MB Austin, C Stewart Jr., JP Noel. Structure and function of enzymes involved in the biosynthesis of phenylpropanoids. **Plant Physiol. Biochem.**, **46**, 356-370 (2008).
- H Grisebach, K Hahlbrock. Enzymology and regulation of flavonoid and lignin biosynthesis in plants and plant cell suspension cultures. In: *Recent Advances in Phytochemistry, Vol. 8 Metabolism and Regulation of Secondary Plant Products*. VC Runeckles, EE Conn (Eds), pp 21-52. Academic Press. New York (1974).
- Y Wang, S Chen, O Yu. Metabolic engineering of flavonoids in plants and microorganisms. **Appl. Microbiol. Biotechnol.**, **91**, 949-956 (2011).
- TTH Dao, HJM Linthorst, R Verpoorte. Chalcone synthase and its functions in plant resistance. **Phytochem. Rev.**, **10**, 397-412 (2011).
- YC Yin, XD Zhang, ZQ Gao, T Hu, Y Liu. The research progress of chalcone isomerase (CHI) in plants. **Mol. Biotechnol.**, **61**, 32-52 (2019).
- O Hokche, PE Berry, O Huber. *Nuevo Catálogo de la Flora Vasculare de Venezuela*. Fundación Instituto Botánico de Venezuela Dr. Tobías Lasser. Caracas, Venezuela (2008).
- JM Amaro, M Hung. Estudio fitoquímico de la flora andina venezolana III. Flavonoides y esteroides de la *Stevia lucida* Lagasca. **Rev. Fac. Farm. ULA**, **25**, 125-134 (1985).

29. P Chacón-Morales, JM Amaro-Luis, A Bahsas. Isolation and characterization of (+)-mellein, the first isocoumarin reported in *Stevia* genus. **Av. Quím.**, **8**, 145-151 (2013).
30. JM Amaro, M Hung. Estudio fitoquímico de la flora andina venezolana IV. Estructura del ácido labd-13(Z)-en-8 α -ol-15-oico, un componente de la *Stevia lucida* Lagasca. **Acta Cient. Venez.**, **39**, 21-24 (1988).
31. JM Amaro-Luis. An *ent*-kaurenolide from *Stevia lucida*. **Phytochemistry**, **32**, 1611-1613 (1993).
32. JM Amaro, M Adrián, CM. Cerda, P Joseph-Nathan. Longipinene derivatives from *Stevia lucida* and *S. triflora*. **Phytochemistry**, **27**, 1409-1412 (1988).
33. PA Chacón-Morales, C Santiago Dugarte, JM Amaro-Luis. Helenin from *Stevia lucida*. The first report of this natural eudesmanolide mixture in Eupatorieae tribe. **Nat. Prod. Res.**, 1-4 (2020) DOI: 10.1080/14786419.2020.1739677.
34. P Chacón-Morales, JM Amaro-Luis. Síntesis de un derivado diosfenólico a partir de la 7 β ,9 α -dihidroxi-longipin-2-en-1-ona. **Av. Quím.**, **10**, 41-47 (2015).
35. PA Chacón-Morales, C Santiago-Dugarte, JM Amaro-Luis. Unexpected reduction-allylic oxidation of hemisynthetic diosphenols from longipinene derivatives. **Phytochem. Lett.**, **39**, 105-110 (2020).
36. PA Chacón-Morales, JM Amaro-Luis. Meridane and uladane, two unprecedented sesquiterpene skeletons obtained by Wagner-Meerwein rearrangements of longipinane derivatives. **Tetrah. Lett.**, **57**, 2713-2716 (2016).
37. DS Reddy, AG Kutateladze. Computational structure revision of a longipinane derivative meridane. **Tetrah. Lett.**, **57**, 4727-4729 (2016).
38. PA Chacón-Morales, JM Amaro-Luis, AG Kutateladze. Structure determination and mechanism of formation of a *secomoreliane* derivative supported by computational analysis. **J. Nat. Prod.**, **80**, 1210-1214 (2017).
39. M Salmón, A Ortega, G García de la Mora, E Angeles. A Diterpenic Acid from *Stevia lucida*. **Phytochemistry**, **22**, 1512-1513 (1983).
40. D Guerra-Ramírez, CM Cerda-García-Rojas, AM Puentes, P Joseph-Nathan. Longipinene diesters from *Stevia lucida*. **Phytochemistry**, **48**, 151-154 (1998).
41. PA Chacón Morales, C Santiago Dugarte, JM Amaro Luis. 2',3,4-trihydroxychalcone, phloretin and calomelanone from *Stevia lucida*. The first chalcones reported in *Stevia* genus. **Biochem. System. Ecol.**, **77**, 57-60 (2018).
42. TJ Mabry, KR Markham, MB Thomas. *The Systematic Identification of Flavonoids*, pp 165-174; 227-230. Springer-Verlag. Berlin (1970).
43. A Shafaghat, F Salimi, N Alaniyan, Z Shoaiei. Flavonoids and an ester derivative isolated from *Galium verum* L. **World Appl. Sci. J.**, **11**, 473-477 (2010).
44. S Asen, JR Plimmer. 4,6,4'-Trihydroxyaurone and other flavonoids from *Limonium*. **Phytochemistry**, **11**, 2601-2603 (1972).
45. L Quast, R Wiermann. Über das vorkommen verschieden substituierter chalkone während der mikrosporogenese bei *Tulipa*. **Experientia**, **29**, 1165-1166 (1973).
46. AE Kambal, EC Bate-Smith. A genetic and biochemical study on pericarp pigments in a cross between two cultivars of grain sorghum, *Sorghum bicolor*. **Heredity**, **37**, 413-416 (1976).
47. Y Iijima, K Suda, T Suzuki, K Aoki, D Shibata. Metabolite profiling of chalcones and flavanones in tomato fruit. **J. Japan. Soc. Hort. Sci.**, **77**, 94-102 (2008).
48. B Han, Z Xin, S Ma, W Liu, B Zhang, L Ran, L Yi, D Ren. Comprehensive characterization and identification of antioxidants in *Folium Artemisiae argyi* using high-resolution tandem mass spectrometry. **J. Chromatogr. B**, **1063**, 84-92 (2017).
49. W Greenaway, S English, J May, FR Whatley. Analysis of phenolics of bud exudate of *Populus sieboldii* by GC-MS. **Phytochemistry**, **30**, 3005-3008 (1991).
50. P Guerrero-Castillo, S Reyes, J Robles, MJ Simirgiotis, B Sepulveda, R Fernandez-Burgos, C Areche. Biological activity and chemical characterization of *Pouteria lucuma* seeds: A possible use of an agricultural waste. **Wast. Manag.**, **88**, 319-327 (2019).
51. M Alberstein, M Eisenstein, H Abeliovich. Removing allosteric feedback inhibition of tomato 4-coumarate:CoA ligase by directed evolution. **The Plant Journal**, **69**, 57-69 (2012).
52. AB Christensen, PL Gregersen, J Schröder, DB Collinge. A chalcone synthase with an unusual substrate preference is expressed in barley leaves in response to UV light and pathogen attack. **Plant Mol. Biol.**, **37**, 849-857 (1998).
53. A Peters, HAW Schneider-Poetsch, H Schwarz, G Weissenböck. Biochemical and immunological characterization of chalcone synthase from rye leaves. **J. Plant Physiol.**, **133**, 178-182 (1988).
54. W Hinderer, HU Seitz. Chalcone synthase from cell suspension cultures of *Daucus carota* L. **Arch. Biochem. Biophys.**, **240**, 265-272, (1985).
55. C Gosch, H Halbwirth, B Schneider, D Hölscher, K Stich. Cloning and heterologous expression of glycosyltransferases from *Malus x domestica* and *Pyrus communis*, which convert phloretin to phloretin 20-O-glucoside (phloridzin). **Plant Sci.**, **178**, 299-306 (2010).
56. FA Tomás-Barberán, MN Clifford. Flavanones, chalcones and dihydrochalcones-Nature, occurrence and dietary burden. **J. Sci. Food Agric.**, **80**, 1073-1080. (2000).
57. SZ Dzedzic, BJF Hudson. Polyhydroxy chalcones and flavanones as antioxidants for edible oils. **Food Chem.**, **12**, 205-212 (1983).
58. J Nishida, J Kawabata. DPPH radical scavenging reaction of hydroxy- and methoxychalcones. **Biosci. Biotechnol. Biochem.**, **70**, 193-202 (2006).
59. AR Bilia, AR Sannella, FF Vincieri, L Messori, A Casini, C Gabbiani, C Severini, G Majori. Antiplasmodial effects of a few selected natural flavonoids and their modulation of artemisinin activity. **Nat. Prod. Commun.**, **12**, 1999-2002 (2008).

60. X Sui, YC Quan, Y Chang, RP Zhang, YF Xu, LP Guan. Synthesis and studies on antidepressant activity of 2',4',6'-trihydroxychalcone derivatives. **Med. Chem. Res.**, **21**, 1290-1296 (2012).
61. GTT Nguyen, G Erlenkamp, O Jäck, A Küberl, M Bott, F Fiorani, H Gohlke, G Groth. Chalcone-based selective inhibitors of a C4 plant key enzyme as novel potential herbicides. **Sci. Rep.**, **6**, 27333 (2016).
62. LP Sun, LX Gao, WP Ma, FJ Nan, J Li, HR Piao. Synthesis and biological evaluation of 2,4,6-trihydroxychalcone derivatives as novel protein tyrosine phosphatase 1B inhibitors. **Chem. Biol. Drug. Des.**, **80**, 584-590 (2012).
63. JC Le Bail, C Pouget, C Fagnere, JP Basly, AJ Chulia, G Habrioux. Chalcones are potent inhibitors of aromatase and 17 β -hydroxysteroid dehydrogenase activities. **Life Sci.**, **68**, 751-761 (2001).
64. S Sogawa, Y Nihro, H Ueda, A Izumi, T Miki, H Matsumoto, T Satoh. 3,4-Dihydroxychalcones as potent 5-lipoxygenase and cyclooxygenase inhibitors. **J. Med. Chem.**, **36**, 3904-3909 (1993).
65. FB Power, Tutin F. Chemical Examination of Eriodictyon. **Proc. Am. Pharm. Assoc.**, **54**, 352-369 (1906); **Pharmac. Rev.**, **24**, 300-304 (1907).
66. J Shinoda, I Sato. New synthesis of polyoxy-chalcone, -hydrochalcone and -flavanone, Part 4: On the structure of eriodictyol and homoeriodictyol in the synthetic rod of 3', 4'-dimethoxy-5, 7-bioxyflavanone. **Yakugaku Zasshi**, **49**, 64-70 (1929).
67. J Shinoda, I Sato. New synthesis of polyoxy-chalcone, -hydrochalcone and -flavanone, Part 5: Synthesis of eryodictyol and homoeryodictyol. **Yakugaku Zasshi**, **49**, 71-78 (1929).
68. A Islam, S Islam, K Rahman, N Uddin, R Akanda. The pharmacological and biological roles of eriodictyol. **Arch. Pharm. Res.**, **43**, 582-592 (2020).
69. D Barreca, G Gattuso, E Bellocco, A Calderaro, D Trombetta, A Smeriglio, G Laganá, M Daglia, S Meneghini, SM Nabavi. Flavanones: *Citrus* phytochemical with health-promoting properties. **BioFactors**, **43**, 495-506 (2017).
70. S Belli, M Rossi, N Molasky, L Middleton, C Caldwell, C Bartow-McKenney, M Duong, J Chiu, E Gibbs, A Caldwell, C Gahn, F Caruso. Effective and novel application of hydrodynamic voltammetry to the study of superoxide radical scavenging by natural phenolic antioxidants. **Antioxidants**, **8**, 14 (2019).
71. X Wang, R Deng, J Dong, L Huang, J Li, B Zhang. Eriodictyol ameliorates lipopolysaccharide-induced acute lung injury by suppressing the inflammatory COX-2/NLRP3/NF- κ B pathway in mice. **J. Biochem. Mol. Toxicol.**, **34**, e22434 (2020).
72. P. Rajesh, V Manju. Eriodictyol imparting cardioprotection and lipid modulation in isoproterenol-induced myocardial infarcted rats. **Res. J. Chem. Environ.**, **24**, 112-119 (2020).
73. W Li, Q Du, X Li, X Zheng, F Lv, X Xi, G Huang, J Yang, S Liu. Eriodictyol inhibits proliferation, metastasis and induces apoptosis of glioma cells via PI3K/Akt/NF- κ B signaling pathway. **Front. Pharmacol.**, **11**, 114 (2020).
74. L Yu, X Liu. Eriodictyol suppresses survival of cervical cancer cells through mediation of PTEN/Akt signaling pathway. **Curr. Top. Nutra. Res.**, **18**, 196-200 (2020).
75. JL Liu, YC Kong, JY Miao, XY Mei, SY Wub, YC Yan, XY Cao. Spectroscopy and molecular docking analysis reveal structural specificity of flavonoids in the inhibition of α -glucosidase activity. **Int. J. Biol. Macromol.**, **152**, 981-989 (2020).
76. M Smith, JC Smith. Repurposing therapeutics for COVID-19: Supercomputer-based docking to the SARS-CoV-2 viral spike protein and viral spike protein-human ACE2 interface. **ChemRxiv**TM, DOI:10.26434/chemrxiv.11871402.v4 (2020). [this article is a preprint; it may not have been peer reviewed].
77. RR Deshpande, AP Tiwari, N Nyayanit, M Modak. *In silico* molecular docking analysis for repurposing therapeutics against multiple proteins from SARS-CoV-2. **Eur. J. Pharmacol.**, **886**, 173430 (2020).
78. R Spribille, G Forkmann. Chalcone synthesis and hydroxylation of flavonoids in 3'-position with enzyme preparation from flowers of *Dianthus caryophyllus* L. (carnation). **Planta**, **155**, 176-182 (1982).
79. R Sütfeld, R Wiermann. The Formation of flavanones from hydroxycinnamoyl-coenzyme A thiol esters and malonyl-coenzyme A by enzyme extracts from anthers. **Z. Pflanzenphysiol.**, **79**, 467-472 (1976).
80. C Dittmer, G Raabe, L Hintermann. Asymmetric cyclization of 2-hydroxychalcones to flavanones: Catalysis by chiral Brønsted acids and bases. **Eur. J. Org. Chem.**, 5886-5898 (2007) and reference here cited.
81. P Kulkarni, P Wagh, P Zubaidha. An improved and eco-friendly method for the synthesis of flavanone by the cyclization of 2'-hydroxy chalcone using methane sulphonic acid as catalyst. **Chem. J.**, **2**, 106-110 (2012) and reference here cited.
82. DG Bhosale, PS Kulkarni. Ferric sulphate solid acid catalyst for cyclization of 2'-hydroxychalcone to flavanone under microwave condition. **Iran. J. Org. Chem.**, **5**, 1061-1064 (2013) and reference here cited.
83. J Mai, E Hoxha, CE. Morton, BM Muller, MJ Adler. Towards a dynamic covalent molecular switch: Substituent effects in chalcone/flavanone isomerism. **Org. Biomol. Chem.**, **11**, 3421-3423 (2013).
84. BM Muller, TJ Litberg, RA Yocum, CA Pniewski, MJ Adler. Extended aromatic and heteroaromatic ring systems in the chalcone-flavanone molecular switch scaffold. **J. Org. Chem.**, **81**, 5775-5781 (2016).
85. S Moussouni, A Detsi, M Majdalani, DP Makris, P Kefalas. Crude peroxidase from onion solid waste as a tool for organic synthesis. Part I: Cyclization of 2',3,4,4',6'-pentahydroxy-chalcone into aureusidin. **Tetrahedron Lett.**, **51**, 4076-4078 (2010).