

Low Serum Free Triiodothyronine Levels Mark Familial Longevity: The Leiden Longevity Study

Maarten P. Rozing,¹ Rudi G. J. Westendorp,^{1,2} Anton J. M. de Craen,¹ Marijke Frölich,³ Bastiaan T. Heijmans,⁴ Marian Beekman,⁴ Carolien Wijnsman,¹ Simon P. Mooijaart,¹ Gerard-Jan Blauw,¹ P. Eline Slagboom,^{2,4} and Diana van Heemst¹; on behalf of the Leiden Longevity Study (LLS) Group

¹Department of Gerontology and Geriatrics, Leiden University Medical Center, The Netherlands.

²Netherlands Consortium of Healthy Aging (NCHA), The Hague, The Netherlands.

³Department of Clinical Chemistry and ⁴Department of Molecular Epidemiology, Leiden University Medical Center, The Netherlands.

Address correspondence to Diana van Heemst, PhD, Department of Gerontology and Geriatrics, Leiden University Medical Center, PO Box 9600, 2300 RC, Leiden, The Netherlands. Email: d.van_heemst@lumc.nl

Background. The hypothalamo-pituitary-thyroid axis has been widely implicated in modulating the aging process. Life extension effects associated with low thyroid hormone levels have been reported in multiple animal models. In human populations, an association was observed between low thyroid function and longevity at old age, but the beneficial effects of low thyroid hormone metabolism at middle age remain elusive.

Methods. We have compared serum thyroid hormone function parameters in a group of middle-aged offspring of long-living nonagenarian siblings and a control group of their partners, all participants of the Leiden Longevity Study.

Results. When compared with their partners, the group of offspring of nonagenarian siblings showed a trend toward higher serum thyrotropin levels (1.65 vs 1.57 mU/L, $p = .11$) in conjunction with lower free thyroxine levels (15.0 vs 15.2 pmol/L, $p = .045$) and lower free triiodothyronine levels (4.08 vs 4.14 pmol/L, $p = .024$).

Conclusions. Compared with their partners, the group of offspring of nonagenarian siblings show a lower thyroidal sensitivity to thyrotropin. These findings suggest that the favorable role of low thyroid hormone metabolism on health and longevity in model organism is applicable to humans as well.

Key Words: Thyroid—Longevity—Endocrinology.

Received September 11, 2009; Accepted November 20, 2009

Decision Editor: Luigi Ferrucci, MD, PhD

THE hypothalamo-pituitary-thyroid axis has been widely implicated in modulating the aging process. Life extension effects associated with low thyroid hormone levels have been reported in multiple animal models. In neonatal rats, induction of hypothyroidism results in a moderate extension of life span (1). Similarly, low thyroid hormone levels are characteristic of murine pituitary mutants with delayed aging: Long-lived Ames and Snell dwarf mice show traits that are hypothesized to be related to thyroid hormone deficiency, including hypothermia and delayed maturation (2,3). Administration of thyroid hormone during adulthood partly diminishes longevity in Snell dwarf mice (4). Another very long-living mammal, the naked mole rat (*Heterocephalus glaber*), also has very low serum thyroxine levels (5).

In agreement with the findings in animals, various studies have shown an association between low thyroid function and improved longevity in elderly humans. In the general population of the oldest old, high levels of thyrotropin are

associated with a prolonged life span (6,7). In contrast, low serum thyrotropin and higher serum-free thyroxine levels are related to an increased risk of cardiovascular mortality (8). These findings suggest a favorable effect of thyroid hypofunction on healthy aging in humans.

However, comparative cross-sectional studies involving long-lived participants are hampered by the lack of proper controls. These studies remain inconclusive as to whether thyroid hypofunction in extreme old age represents an adaptive mechanism or is the result of selective survival of participants with lifelong thyroid hypofunction. We designed the Leiden Longevity Study in order to identify familial determinants of healthy longevity in nonagenarian siblings and their offspring, who are enriched for heritable influences on morbidity and mortality (9). The aim of this study was to assess whether low thyroid function observed in extreme old age is already present in middle-aged individuals with higher than average life expectancy. To this end, we have compared thyroid hormone function parameters in a

group of middle-aged offspring of long-living nonagenarian siblings and a control group of their partners of the Leiden Longevity Study.

METHODS

Leiden Longevity Study

In the Leiden Longevity Study, 420 families were recruited consisting of long-lived Caucasian siblings together with their offspring and the partners thereof. Families were recruited if at least two long-lived siblings were alive and fulfilled the age criterion of 89 years or older for men and 91 years or older for women. There were no selection criteria on health or demographic characteristics. For 2,465 of the offspring and their partners, nonfasted serum samples taken at baseline were available for the determination of endocrine and metabolic parameters. Between November 2006 and May 2008, for 2,235 of the offspring and their partners, information on medical history was obtained from the participants' treating physicians (response: 90.7%). For 2,255 of the offspring and their partners, information on the use of medication was obtained from the participants' pharmacist (response: 91.5%). For 2,184 of the offspring and their partners, a general questionnaire containing information on lifestyle and self-reported height and weight was obtained (response: 89.0%). For the present study, for a total of 1,738 of the offspring and their partners, serum as well as information on medical history and information on medication use and the general questionnaire were available (inclusion: 69.5%). The Medical Ethical Committee of the Leiden University Medical Center approved the study, and informed consent was obtained from all participants.

For the current study, participants using thyroid medication were excluded from the analyses: 32 (2.7%) offspring using thyroid medication were excluded and 11 (2.0%) partners were excluded from the analyses. Thyroid hormone medication was defined as thyroid or antithyroid preparations (ATC code H03). Outliers were defined as serum thyroid parameters (thyrotropin, thyroxine, and triiodothyronine) beyond 3 *SDs* below or above the standard error of the mean. Outliers were excluded from the analyses. Thirty-four offspring with serum thyroid parameters beyond 3 *SDs* from the mean were excluded from the analyses, of which one individual was clinically hyperthyroid (thyrotropin < 0.3 mU/L and free thyroxine > 24 pmol/L) and three individuals clinically hypothyroid (thyrotropin > 4.8 mU/L and free thyroxine < 10 pmol/L). Nine partners with serum thyroid parameters beyond 3 *SDs* from the mean were excluded from the analyses, of which two individuals were clinically hyperthyroid (thyrotropin < 0.3 mU/L and free thyroxine > 24 pmol/L). In our laboratory, the reference values for thyrotropin were 0.3–4.8 mIU/L, free thyroxine were 10–24 pmol/L, and free triiodothyronine were 2.5–5.5 pmol/L.

Biochemical Analysis

All serum measurements were performed with fully automated equipment. For thyrotropin, free thyroxine, and free triiodothyronine, the Modular E170 was used from Roche (Almere, The Netherlands). The coefficients of variation of these measurements were all less than 5%.

Statistical Analysis

Distributions of continuous variables were examined for normality and logarithmically transformed, when appropriate, and used in all calculations. Geometric means (with 95% confidence intervals) are reported for thyrotropin. All differences between offspring and partner categories were assessed with the use of linear mixed modeling, adjusted for age and correlation of sibling data. Differences in age and body mass index between the two groups of offspring and partners were tested using a Mann–Whitney rank sum test. Differences in smoking behavior and gender distribution between the group of offspring and the group of partners were calculated using a chi-square test. The Statistical Package for the Social Sciences program for Windows, version 16.0, was used for data analysis.

RESULTS

Table 1 shows the baseline characteristics for the two study populations. In total, we used data on 1,119 middle-aged offspring of nonagenarian siblings and 533 of their middle-aged partners. The group of female offspring was slightly older than that of female partners, whereas the group of male offspring was younger than that of male partners. No significant differences between the two groups were observed with regard to body mass index and current smoking status.

Table 2 displays the mean serum levels of various thyroid function parameters in the group of middle-aged offspring of nonagenarian siblings compared with the group of their middle-aged partners adjusted for age and body mass index. A trend was observed toward higher serum thyrotropin levels in the group of offspring when compared with that of partners ($p = .11$). The free serum thyroxine levels were lower in the group of offspring than that of partners ($p = .045$). Likewise, mean free serum triiodothyronine levels were lower in the group of offspring in comparison with that of partners ($p = .024$). Results were not materially different when analyses were adjusted for smoking behavior.

DISCUSSION

The secretion of thyroid hormone from the thyroid gland is regulated by thyrotropin, which in turn is controlled by the hypothalamic-derived thyroid-releasing hormone. The main thyroid hormone produced in the thyroid gland is thyroxine (3,5,3',5' regular-tetraiodothyronine), which has a low affinity for thyroid hormone receptors in target tissues.

Table 1. Baseline Characteristics of Study Populations

	Offspring	Partners	<i>p</i> Value
Females (<i>n</i>)	596	293	
Age (y)	58.9 (54.9–64.0)	57.4 (52.4–61.9)	<.001
Height (cm)	166.8 (166.2–167.3)	167.0 (166.2–167.7)	.62
Weight (kg)	69.6 (68.6–70.7)	70.8 (69.4–72.2)	.17
Body mass index (kg/m ²)	25.0 (24.7–25.4)	25.4 (24.9–25.9)	.20
Currently smoking, <i>n</i> (%)	75 (12.8)	45 (15.4)	.30
Males (<i>n</i>)	523	240	
Age (y)	59.4 (55.0–64.2)	61.4 (56.2–66.3)	.001
Height (cm)	178.6 (178.0–179.3)	179.1 (178.3–180.0)	.32
Weight (kg)	82.4 (81.3–83.4)	82.7 (81.1–84.2)	.73
Body mass index (kg/m ²)	25.8 (25.5–26.1)	25.7 (25.3–26.1)	.76
Currently smoking, <i>n</i> (%)	78 (15.1)	37 (15.5)	.91

Note: Age is presented as median age with interquartile range. Height, weight, and body mass index are presented as estimated means with 95% confidence intervals. Results for weight, height, and body mass index were adjusted for age.

Thyroxine can be converted peripherally to the more biologically active free 3,5,3'-triiodothyronine or the inactive reverse triiodothyronine (3,3',5'-triiodothyronine). When compared with their partners, the group of offspring of nonagenarian siblings showed a trend toward higher serum thyrotropin levels in conjunction with lower free thyroxine levels and lower free triiodothyronine levels. These findings indicate a lower thyroidal sensitivity to thyrotropin in the group of offspring of nonagenarian siblings.

In middle-aged human populations, the effect of low thyroid hormone metabolism on health is unclear. At middle age, overt hypothyroidism is considered a risk factor for the development of atherosclerosis and myocardial infarction (10,11). Paradoxically, in euthyroid middle-aged participants, lower triiodothyronine serum levels are associated with a beneficial cardiometabolic profile (12,13). In the current study, we demonstrate lower thyroid hormone levels in a middle-aged population, which was previously shown to have a lower prevalence of cardiovascular disease (14).

These data suggest that selective survival of participants with a lifelong thyroid hypofunction may contribute to the association between decreased thyroidal sensitivity to thyrotropin and a longer life span (6,7).

Our results agree with observations in several animal studies showing that lower activity of the thyroid hormone axis is beneficial during the aging process (1,3,15). Active triiodothyronine primarily regulates the basal metabolic rate of cells, thereby increasing thermogenesis and the production of free radicals (16). Data from model organisms show that low triiodothyronine is associated with lower production of reactive oxygen species (ROS), and ROS inflicted genomic damage (17). The more efficient transport of electrons through the respiratory chain under conditions of low triiodothyronine might reduce the production of ROS and slow aging. Additionally, previous studies in euthyroid participants have shown an association between higher levels of thyroid hormones and higher serum glucose levels, higher serum insulin levels, as well as increased serum triglyceride

Table 2. Serum Levels of Thyroid Hormone Axis Parameters for Offspring and Partners

	Offspring	Partners	<i>p</i> Value
All			
Thyrotropin (0.3–4.8 mU/L)	1.65 (1.59–1.71)	1.57 (1.49–1.66)	.11
Free thyroxine (10–24 pmol/L)	15.0 (14.9–15.2)	15.2 (15.0–15.4)	.045
Free triiodothyronine (2.5–5.5 pmol/L)	4.08 (4.04–4.12)	4.14 (4.09–4.20)	.024
Ratio triiodothyronine thyroxine	0.28 (0.27–0.28)	0.28 (0.27–0.28)	.84
Females			
Thyrotropin (0.3–4.8 mU/L)	1.72 (1.63–1.80)	1.64 (1.52–1.76)	.28
Free thyroxine (10–24 pmol/L)	14.8 (14.6–14.9)	15.1 (14.8–15.3)	.034
Free triiodothyronine (2.5–5.5 pmol/L)	3.89 (3.84–3.94)	4.00 (3.93–4.07)	.007
Ratio triiodothyronine thyroxine	0.27 (0.26–0.27)	0.27 (0.26–0.27)	.48
Males			
Thyrotropin (0.3–4.8 mU/L)	1.60 (1.52–1.69)	1.53 (1.42–1.65)	.26
Free thyroxine (10–24 pmol/L)	15.2 (15.0–15.4)	15.5 (15.2–15.7)	.12
Free triiodothyronine (2.5–5.5 pmol/L)	4.26 (4.20–4.31)	4.34 (4.26–4.42)	.048
Ratio triiodothyronine thyroxine	0.28 (0.28–0.29)	0.28 (0.28–0.29)	.95

Note: Data are presented as estimated means with 95% confidence intervals. Results for all were adjusted for age, sex, and body mass index. Results for men and women separately were adjusted for age and body mass index. Reference values are given between parentheses.

levels in men (18,19). Furthermore, higher triiodothyronine levels have been associated with higher blood pressure. Increases in heart rate, cardiac output, myocardial contractility, and blood volume possibly underlie this association between triiodothyronine levels and blood pressure (20).

In conclusion, our data demonstrate that the middle-aged offspring of nonagenarian siblings have lower serum free triiodothyronine levels compared with their middle-aged partners. These findings hint at a role of the thyrotroph axis in the regulation of human health and longevity.

FUNDING

This work was supported by the Innovation Oriented research Program on Genomics (SenterNovem; IGE01014 and IGE5007), the Centre for Medical Systems Biology, the Netherlands Genomics Initiative/Netherlands Organization for scientific research (05040202 and 050-060-810. NCHA), and the European Union funded Network of Excellence Lifespan (FP6 036894). R.G.J.W. is supported by an unrestricted grant from The Netherlands Genomics Initiative (NCHA 050-060-810).

REFERENCES

- Ooka H, Fujita S, Yoshimoto E. Pituitary-thyroid activity and longevity in neonatally thyroxine-treated rats. *Mech Ageing Dev.* 1983;22(2):113–120.
- Tatar M, Bartke A, Antebi A. The endocrine regulation of aging by insulin-like signals. *Science.* 2003;299(5611):1346–1351.
- Brown-Borg HM, Borg KE, Meliska CJ, Bartke A. Dwarf mice and the ageing process. *Nature.* 1996;384(6604):33.
- Vergara M, Smith-Wheelock M, Harper JM, Sigler R, Miller RA. Hormone-treated snell dwarf mice regain fertility but remain long lived and disease resistant. *J Gerontol A Biol Sci Med Sci.* 2004;59(12):1244–1250.
- Buffenstein R. The naked mole-rat: a new long-living model for human aging research. *J Gerontol A Biol Sci Med Sci.* 2005;60(11):1369–1377.
- Atzmon G, Barzilai N, Hollowell JG, Surks MI, Gabrieli I. Extreme longevity is associated with increased serum thyrotropin. *J Clin Endocrinol Metab.* 2009;94(4):1251–1254.
- Gussekkoo J, van Exel E, de Craen AJ, Meinders AE, Frolich M, Westendorp RG. Thyroid status, disability and cognitive function, and survival in old age. *JAMA.* 2004;292(21):2591–2599.
- Parle JV, Maisonneuve P, Sheppard MC, Boyle P, Franklyn JA. Prediction of all-cause and cardiovascular mortality in elderly people from one low serum thyrotropin result: a 10-year cohort study. *Lancet.* 2001;358(9285):861–865.
- Schoenmaker M, de Craen AJ, de Meijer PH et al. Evidence of genetic enrichment for exceptional survival using a family approach: the Leiden Longevity Study. *Eur J Hum Genet.* 2006;14(1):79–84.
- Kvetny J, Heldgaard PE, Bladbjerg EM, Gram J. Subclinical hypothyroidism is associated with a low-grade inflammation, increased triglyceride levels and predicts cardiovascular disease in males below 50 years. *Clin Endocrinol (Oxf).* 2004;61(2):232–238.
- Walsh JP, Bremner AP, Bulsara MK et al. Subclinical thyroid dysfunction as a risk factor for cardiovascular disease. *Arch Intern Med.* 2005;165(21):2467–2472.
- De Pergola G, Ciampolillo A, Paolotti S, Trerotoli P, Giorgino R. Free triiodothyronine and thyroid stimulating hormone are directly associated with waist circumference, independently of insulin resistance, metabolic parameters and blood pressure in overweight and obese women. *Clin Endocrinol (Oxf).* 2007;67(2):265–269.
- Ortega E, Koska J, Pannaciuoli N, Bunt JC, Krakoff J. Free triiodothyronine plasma concentrations are positively associated with insulin secretion in euthyroid individuals. *Eur J Endocrinol.* 2008;158(2):217–221.
- Westendorp RG, van Heemst D, Rozing MP et al. Nonagenarian siblings and their offspring display lower risk of mortality and morbidity than sporadic nonagenarians: the Leiden Longevity Study. *J Am Geriatr Soc.* 2009;57(9):1634–1637.
- Hauck SJ, Hunter WS, Danilovich N, Kopchick JJ, Bartke A. Reduced levels of thyroid hormones, insulin, and glucose, and lower body core temperature in the growth hormone receptor/binding protein knockout mouse. *Exp Biol Med (Maywood).* 2001;226(6):552–558.
- Harper ME, Seifert EL. Thyroid hormone effects on mitochondrial energetics. *Thyroid.* 2008;18(2):145–156.
- Lopez-Torres M, Romero M, Barja G. Effect of thyroid hormones on mitochondrial oxygen free radical production and DNA oxidative damage in the rat heart. *Mol Cell Endocrinol.* 2000;168(1–2):127–134.
- Bakker SJ, ter Maaten JC, Popp-Snijders C, Heine RJ, Gans RO. Triiodothyronine: a link between the insulin resistance syndrome and blood pressure? *J Hypertens.* 1999;17(12 Pt 1):1725–1730.
- Kim BJ, Kim TY, Koh JM et al. Relationship between serum free T4 (FT4) levels and metabolic syndrome (MS) and its components in healthy euthyroid subjects. *Clin Endocrinol (Oxf).* 2009;70(1):152–160.
- Klein I. Thyroid hormone and the cardiovascular system. *Am J Med.* 1990;88(6):631–637.