

# Distribution of Sea Urchins in Punta Dumalag, Matina Aplaya, Davao City

Marnie Grace I. Sonico<sup>1</sup>

<sup>1</sup>Natural Science Department, College of Arts and Sciences, University of Southeastern Philippines  
marniegracesonico@yahoo.com

Received: July 18,2017; Accepted: December 12, 2017

## Abstract

This study investigated the distribution and diversity of sea urchins along the intertidal zone of Punta Dumalag, Matina Aplaya, Davao City. A total of 100 quadrats (1m x 1 m) spaced 5 m apart, in 3 transects laid perpendicular to the shoreline, recorded a total of 347 individuals, resolved into 5 species, namely, *Echinometra mathaei* (rock- boring or common sea urchin), *Echinothrix calamaris* (Hatpin urchin), *Diadema setosum* (long-spine or “tuyom”), *Prionocidaris verticillata* (rough spine urchin), and *Diadema savignyi*. Among the 347 individuals recorded, *Echinometra mathaei* was the most abundant in the site with a total of 286 individuals. Shannon-Weiner Diversity Analysis (0.602) suggests low species diversity. Physico-chemical measurements such as water temperature, pH, dissolved oxygen, and salinity revealed that the study site is suitable for the culture of sea urchins.

*Keywords: sea urchin, Punta Dumalag, distribution, diversity*

Sea urchins also known as sea hedgehogs are small, spiny, and globular animals comprises the class Echinoidea of Phylum Echinoderm with their close relatives, such as sand dollars (Animal Diversity Web, 2012). Their colors are black and dull shades of red, green, brown, purple, and olive. It plays an important role in marine food chains, consuming various algae, and invertebrates, and being consumed by fishes, sea stars, crabs, birds as well as mammals. They can be found in ocean floors worldwide and also inhabiting coral reefs (Hibberd, 2009).

The biodiversity of sea urchins are ecologically important. Echinoids, specifically the sea urchins have been found to retain levels of metals in their bodies. Recent studies demonstrated that Echinoids are valuable biological indicator of heavy metal contamination (Freeman, 2004). Even public aquariums used sea urchins as an indicator for water quality. Sea urchins are one of the first organisms to show signs of stress and leads to death. Thus, it makes them excellent organisms to use for studies on pollution. Sea urchins act as important keystone species within an ecosystem. By feeding on algae, sea urchins control the algal growth that if allowed to proliferate, would cover coral surfaces, smothering, and killing the coral. Through their burrowing and feeding behaviors, they moderate the balance between coral erosion,

algal growth, and considered sensitive bio-indicators of various contaminants (Zizka, 2012).

Sea urchins also have economic importance. They serve as source of livelihood for the residents in the area. Their gonad is a high valued fishery commodity sold as fresh roe in central wholesale market of Tokyo, Japan. *Tripteneustes gratilla* is an economically important fishery resource generating multi-million peso earning per annum. It is widely collected for food in many Pacific countries. Their shells serve as decorations like seashells but it will not last because their color will fade and their spines will fall off from the test (Prado et al., 2012). Tomsic et al. (2101) also stated that the gonads of sea urchin as well as their roe can be eaten boiled, baked, or raw. They also used as tools in laboratory in embryology. Their gametes are the same size with human gametes (Freeman, 2004).

Researchers nowadays are using sea urchin study with regard to diseases such as Alzheimer’s diseases, cancer, Parkinson’s diseases, and muscular dystrophy. Sea urchins, though invertebrates, share common ancestor with humans and have more than 7,000 of the same genes. With a complete map of DNA of sea urchins, scientist can learn how to treat and prevent diseases in humans better (Freeman, 2004).

Punta Dumalag is known for its high diver-

sity of marine species and beautiful spot that makes it one of the visited places in Davao City. It is located in Barangay Matina Aplaya 75-A, eastern part of Davao City with seashores and natural resources, such as mangroves, sea grasses, sea urchins, corals and nesting area for turtles. It is bounded on the north by Barangay 76-A, on the east by Davao Gulf, on the south by Barangay Talomo and on the west by Barangay Matina (LGU-Dumalag, 2008). Punta Dumalag is a Marine Protected Area (MPA). Marine Protected Area (MPA) sustains fisheries utilization in the fishing areas, and also protects and conserves ecosystems. They provide habitat and protection to large and reproductive mature individuals of various marine species. MPA also protects and conserves the nesting sites for the endangered marine turtles and birds thus; the study was conducted to gather ecological information as well as identification and distribution of sea urchins in the Southside of Marine Protected Area (MPA) in Punta Dumalag, Matina Aplaya, Davao City.

This study was conducted to gather important ecological information of sea urchins in Punta Dumalag, Matina Aplaya, Davao City. Specifically, this study sought to (i) identify the different species of sea urchins found in the study area based on morphological characteristics, (ii) determine species frequency, species density, species richness, species evenness, abundance, diversity, and (iii) determine the physico-chemical characteristics of the study site such as water temperature, pH, dissolved oxygen, and salinity.

## Materials and Method

### Study Area

This study was conducted in Punta Dumalag, Matina Aplaya, Davao City. The place is recognized as one of the Marine Protected Areas (MPA) in the Gulf of Davao. The site was declared as a protected area due to the presence of mangroves, seagrasses and coral reef resources within the 37-hectare MPA, which boundaries start from the shoreline and extend to a distance of about 300 meters seaward (Barangay Punta Dumalag 5-Year Marine Protected Area Management Plan, 2008-2012). The study covered an overall total sampling area of 1,360 m<sup>2</sup>.

Geographic coordinates were determined using Garmin GPS receiver to geotag the location of the area, namely, Transect 1 (7° 1'28.30"N, 125° 34'35.10"E); Transect 2 (7° 1'28. 16"N, 125° 34'34.81"E), and Transect 3 (7° 1'28.00"N, 125°

34'34. 43"E).

### Sampling Design

Three (3) transect lines were established perpendicular to the shoreline in the southern part of Punta Dumalag Marine Protected Area (MPA). Each transect consisted of 100 quadrats measuring 1m x 1m, spaced 5 m apart. The total area covered 1,360 m<sup>2</sup> while the total area of the three (3) transects measured 300 m<sup>2</sup>. Specimens were collected in shallow water by snorkeling and by and picking along the sampling area. All sea urchins found in the quadrat were photographed and counted. Species length and weight measurements were taken using a ruler and digital balance.

### Collection of Data

Field data were gathered during the lowest low tide, in July, September and October, 2014, where species were visible. Collected sea urchins were soaked in 70 % isopropyl alcohol. All morphometric measurements were taken in millimeter (mm).

Physico-chemical parameters such as pH, water temperature, dissolved oxygen, and salinity were measured *in situ* using a thermometer and portable water quality meters, respectively.

### Species Identification

Sea urchins were identified using morphological markers (spines and colors) using field guides, books, journals and previous theses, and taxonomic verification was carried out by local and international experts (Rach Mooi, Ph.D., Center of Echinoderm, Dept. of Invertebrate Zoology and Geology, California Academy of Science, San Francisco, USA).

### Spicule Determination

Spicules of sea urchins found in the study area were obtained by scrapping the inner test or shell using a scalpel or blade. The scrapped part was mounted on the slide, added with a small amount of *zonrox* (chlorine) solution. The slide was viewed in the microscope under high power magnification. Identification of spicules was determined using a field guide. Verification of the initial species ID was done by sending the data to Dr. Rach Mooi.

## Results and Discussion

Five (5) species of sea urchins were found in Punta Dumalag Marine Protected Area - *Echinometra mathae* (rock-boring or common sea urchin), *Diadema setosum* (long-spine or "tuyom"), *Echinothrix calamaris* (Hatpin urchin), *Diadema savignyi* (long-spine urchin), and *Prionocidaris*

*verticillata* (rough spine urchin), resolved in three (3) Families. Three (3) species were found under Diadematidae, one (1) species under Echinometridae, and one (1) species under Cidaridae. *Echinometra mathaei* from Echinometridae recorded the highest number of individuals among the five (5) species, while, *Prionocidaris verticillata* showed the least number. As observed, representative *Diadema* are bigger and heavier compared to the other two families.

Table 1 shows that there were 347 individuals found in the sampling area, dominated by *Echinometra mathaei* with a total count of 296 individuals.

As shown in Table 2, *Echinometra mathaei* has the highest frequency and density observed in seventy six (76) quadrats. Followed by *Diadema setosum*, then *Echinothrix calamaris*, then *Diadema savignyi*, while *Prionocidaris verticillata* has the lowest frequency and density.

Table 3 shows species diversity analysis of sea urchins in the study area. Results show that Shannon-Weiner Diversity Index in Punta Dumalag is 0.602, suggesting low diversity because an area is considered diverse if the value of the Shannon-

Weiner Diversity Index is greater than 1.5.

Table 4 shows the results of the physico-chemical parameter in the sampling area. Mean value of pH in Punta Dumalag is 8.07. This falls within the normal pH range of 6.5 to 8.5 pH (Tehranifard, 2011). This pH helps sperm motility and respiration rate. *Diadema savignyi* can tolerate the pH 7.5 to 8.2. Their optimal pH is 7.8. *Diadema setosum* and *Echinometra mathei* preferred 8.1 to 8.4 pH. Below pH 6, sea urchin larval motility weakens and growth stops. pH must be maintained between 6.0 to 9.0 pH. Result shows that the mean water temperature in Punta Dumalag is 25.1°C. According to Tehranifard (2011), sea urchin can maintain high growth rate between 24°C to 25°C. Dissolved oxygen has a mean value of 17.98 mg/L which is suitable for the survival of sea urchin. Rivero et al. (1996) stated that Oxygen concentrations of 1- 2 mg/L reduce fertilization success to below 50% and can cause mass mortality (Riveros et al., 1996). The mean salinity in Punta Dumalag is 26 ppt. Hendler (2013) attested that Sea urchin cannot tolerate less than 21 ppt salinity because, if reduced it can cause mortality. Result revealed that the physico-chemical parameters in

Table 1. Sea urchins identified in Punta Dumalag, Matina Aplaya, Davao City.

Family	Scientific Name	Common Name	Counts	Length (mm)	Weights (g)
Echinometridae	<i>Echinometra mathaei</i>	Common Urchin	296	70	50.2
Diadematidae	<i>Diadema setosum</i>	Long spine urchin	21	89	90.1
Diadematidae	<i>Echinothrix calamaris</i>	Hatpin urchin	15	124	170.4
Diadematidae	<i>Diadema savignyi</i>	Long spine urchin	11	158	150.5
Cidaridae	<i>Prionocidaris verticillata</i>	Rough spine urchin	4	48	30.9

Table 2. Ecological distribution of sea urchins in Punta Dumalag, Matina Aplaya, Davao City.

Species Name	Total number of individuals	Species Frequency	Relative Frequency (%)	Population Density	Relative Density (%)
<i>Echinometra mathaei</i>	296	0.8530	85.3	0.99	85.30
<i>Diadema setosum</i>	21	0.0605	6.05	0.07	6.05
<i>Echinothrix calamaris</i>	15	0.0432	4.32	0.05	4.32
<i>Diadema savignyi</i>	11	0.0317	3.17	0.04	3.17
<i>Prionocidaris verticillata</i>	4	0.0115	1.52	0.01	1.15

Table 3. Species diversity of sea urchin in the study area.

Family	Species	Total number of individuals
Echinometridae	<i>Echinometra mathaei</i>	296
Diadematidae	<i>Diadema setosum</i>	21
Diadematidae	<i>Echinothrix calamaris</i>	15
Diadematidae	<i>Diadema savignyi</i>	11
Cidaridae	<i>Prionocidaris verticillata</i>	4
Total		347
Shannon-Weiner Diversity Index		<b>0.602</b>
H max		1.609
Species Evenness		0.374
Species Richness		5

Table 4. Physico-chemical parameters in Punta Dumalag.

Transect	pH	Water Temperature	Dissolved Oxygen	Salinity
1	8.05	25.1	17.95	25
2	8.09	25.3	18.35	27
3	8.07	24.9	17.64	26
Mean	8.07	25.1	17.98	26

Punta Dumalag is suitable for the survival of sea urchin.

**Family Echinometridae - *Echinometra mathaei***

Collected *Echinometra mathaei* (Figure 1A) measures 70 mm and weighs 50.2 g. The color is quite variable but the test is usually a dark color. During the lowest low tide, they are found along the shoreline in varieties of color.

The test is small, strong, and oval shape. Primary spines are long as half as the test diameter, usually stout, tapering, and colors are getting lighter at tips. Secondary spines are short, flat tips like rolling pins. It has a pale ring at the base of the spine (Jeng, 1998). The spicules found in the inner test are tylote, sigma, chela, and isochela (Figure 1C-F). It thrives in shallow waters.




Figure 1. A. *Echinometra methai*, common sea urchin collected in Punta Dumalag, Matina Aplaya, Davao City. B. Measurement of specimen. C-F. Different spicules of the species: C. tylote, D. sigma, E. chela, F. isochela micrographs taken at 100X using (microscope brand).

#### Family Diadematidae - *Diadema setosum*

The collected specimen measures 89 mm and weighs 90.1g (Figure 2). They are herbivores and can therefore play a vital role as an interesting member of the clean-up crew (Coppard & Cambell, 2006).

This urchin is well known for having long, sharp, brittle spines capable of painful sting when stepped upon, but slightly venomous. It can be dis-

tinguished from other *Diadema* species by the presence of five white spots on the test located between the ambulacral grooves. It also includes the presence of a bright orange ring around the urchin's periproctal cone, a structure called "anus" (Jeng, 1998). The only spicule found is called strongyle (Figure 2C-F) that was found in the inner test of these species (Lee & Shin, 2013).




Figure 2.A. *Diadema setosum*, long spine urchin collected in Punta Dumalag, Matina Aplaya, Davao City. B. Measurement of specimen. C-F. Spicules of the species: strongyles micrographs taken at 110X using (microscope brand).

### Family Diadematidae - *Echinothrix calamris*

The collected specimen measures 124 mm and weighs 170.4 g. *E. calamris* (Figure 3) occur from surface to 70 cm depth and can be found in lagoons and channels (Kroh & Mooi, 2014). They are active during night and hide in crevices or under rocks during the day (LeBris & Maran, 2010). The spicules (Figure 3C-E) found in the inner test of this species are chela, style, and strongyle (Lee

& Shin, 2013).

*Echinothrix calamris* are found in an intertidal zone along the horizontal coral rocky area during night time. It has a number of color patterns in the spines which move smoothly, white for the primary and yellow for the shorter spines that can deliver a nasty sting, while the longer open ended spines are banded with light and dark color.




Figure 3. A. *Echinothrix calamris*, hatpin urchin collected in Punta Dumalag, Matina, Aplaya, Davao City. B. Measurement of specimen. C-E. Different spicules of the species: C. chela, D. style, E. strongyle micrographs taken at 100X using (microscope brand).

### Family Diadematidae - *Diadema savignyi*

It was observed that the long spines of *Diadema savignyi* (Figure 4) and other species of Echinoidea oftentimes can cause injury when people accidentally steps onto an urchin. Alender & Russel (1966) stated that the once the spine detaches and penetrate the body tissue, an infection can occur. Collected specimen measures 158 mm and weighs 150.5 g. It has solid iridescent blue or sometimes green lines that run along its black test and periproct, area surrounding anus (Coppard & Cambell, 2006).

This species lives in shallow waters (Muthiga, 2003). It is an urchin that displays pentamerism, which is a quality of the class Echinodermata. It has many long spines, tube feet and its body is round. Its anal sac is dark. According to Muthiga et al. (2007), given suitable

condition, the test can grow as long as 90 mm in diameter. They live in sand flats and coral reef areas, which are warm shallow areas near coasts (Hoey, 2008). Species of *Diadema* are prey by 15 species of finfish, the spiny lobster, and 2 species of gastropods (Shafir & McClanahan, 1990). Spicules (Figure 4C-D) found in the inner test of *Diadema savignyi* are tylote and toxaspire.

### Family Cidaridae - *Prionocidaris verticillata*

*Prionocidaris verticillata* (Figure 5) is found in shallow waters in rocks and patch reefs. It varies in color from place to place. In Punta Dumalag, Matina Aplaya, *Prionocidaris verticillata* mimics the color of the rock. The specimen found measures 48 mm and weighs 30.9 g. Spicules found in the inner test in *P. verticillata* is only rod (Figure 5C-F). They feed on invertebrates, sponges and even dead fish. To move, they use




Figure 4. A. *Diadema savignyi*, long spine urchin collected in Punta Dumalag, Matina Aplaya, Davao City. B. Measurement of specimen. C-D. Different spicules of the species: C. tylote, D. toxaspire micrographs taken at 100X using (microscope brand).




Figure 5. A. *Prionocidaris verticillata*, rough spine urchin collected in Punta Dumalag, Matina Aplaya, Davao City. B. Measurement of specimen. C-F. Different spicules of the species: C. sanidaster, D. buttons, E. tylote, F. toxaspire micrographs taken at 100X using (microscope brand).

their tube feet but there were times that they hitch a lift on the back of crabs. Kroh & Mooi (2014) observed that the anus of this species is located on top, their jaws are so strong which they used for grinding their prey and they are also sensitive to light. At a short distance, they have the capacity to shoot venom loaded spines.

This echinoid has a small globular test with an armature of disproportionately long thorn-like spines. Rough long spines are used as walking stilts and the points of the thorns are often worn (Taylor,

1996). Spicules (Figure 5C-F) found in the inner test are sanidaster, buttons, tylote, and toxaspire (Lee & Shin, 2013).

### Conclusion

It was concluded that there were five (5) species of sea urchin observed in the study area.

The species diversity is low; therefore Punta Dumalag, being a marine protected area (MPA)

is not an assurance for the diversity of species in the area. Meanwhile, the physico-chemical parameters such as water temperature, pH, dissolved oxygen and salinity are suitable for the survival of sea urchin species.

It is recommended that annual assessment must be conducted in the area to further evaluate the status of sea urchin. It is also recommended to study the variations in environmental factors, such as, temperature, salinity, pH, and dissolve oxygen that has an important effect on the physiological and biochemical rates of an organism to survive and reproduce.

### Acknowledgments

The research study was made possible with the financial support of University of Southeastern Philippines.

### References

- Arakaki, Y., Uehara, T., & Fagoonee, I. (1998). Comparative studies of the genus *Echinometra* for Okinawa and Mauritius. *Zoological Science*, 15, 159-168.
- Coppard, S., & Cambell, A. (2006). Taxonomic significance of test morphology in the echinoid genera *Diadema* Gray, 1825 and *Echinothrix* Peters, 1854 (Echinoder- mata). *Zoosystema* 28, 93-112.
- Corrales, J. A. & Tuban, E. C. (2008). Preliminary study on the polyculture of sea urchin (*Tripteneustes gratilla*) and sea cucumber (*Actinopyga* spp.) in concrete tank. *UNP Research Journal*, 17, 25-37.
- Freeman, S.M. (2004). Echinoidea. In B. Grzimek, S. F. Craig, D. A. Thoney, N. Schlager, and M. Hutchins. *Grzimek's Animal Life Encyclopedia*, 2<sup>nd</sup> Edition. Detroit, MI: Thompson/Gale. ISBN 0787657786.
- Gotshall, D.W. (1994). *Guide to Marine Invertebrate - Alaska to Baja California*. Sea Challengers Publication, Monterey, California.
- Gray, J.E. (1855). An arrangement of the families of *Echinida*, with description of some new genera and species. *Proceedings of the Zoological Society London* 1855, 35-39. Retrieved from [http://www.nhm.ac.uk/research-curation/research/](http://www.nhm.ac.uk/research-curation/research/projects/echinoid-directory/taxa/taxon.jsp?id=1187)
- Hendler, G. (2013). Recent mass mortality of *Strongylocentrotus purpuratus* (Echinoder- mata: Echinopidea) at Malibu and a review of purple sea urchin kills elsewhere in California. *Bulletin of the Southern California Academy of Sciences* 112, 19-37.
- Hibberd, T. (2009). Class Echinoidea. *Field Identification Guide to Heard Island and McDonald Island Benthic Invertebrates: a guide for scientific observers aboard fishing vessels* 117-118. Australia, Fisheries research and Development Corporation.
- Hickman, Jr., C. P., Roberts, L. S., & Larson, A. (2001). Echinoderms. *Integrated principles of zoology* (11<sup>th</sup> ed., 458-480pp.). NY: New York, The McGraw-Hill Companies Incorporated.
- Hoey, J. (2008). The effect of herbivory by the long-spined sea urchin, *Diadema saving- yi* on algae growth in the coral reefs of Moorea, French Polynesia.
- James, P. (2007). The effects of environmental factors and husbandry techniques on roe enhancement of the New Zealand sea urchin, *Evechinus chlo- roticus*. Doctor of Philosophy in Marine Biology Thesis, Victoria University of Wellington.
- Jeng, M-S. (1998). Shallow-water echinoderms of Taiping Island in the South China Sea. *Zoological Studies*, 37(2), 137-153.
- Kroh, A., & Mooi, R. (2014). *World Echinoidea Database*. Retrieved from <http://www.marinespecies.org/aphia.php?p=taxdetails&id=213377>.
- Kroh, A., & Smith, A.B. (2010). The phylogeny and classification of post-Paleozoic echinoids. *Journal of Systematics Paleontology* 8, 147-212.
- LeBris, S., & Maran, V (2010). *E. calamaris* (Pallas, 1774). Retrieved from <http://www.marinespecies.org/aphia.php?p=taxdetails&id=213377>.
- Lee, T. & Shin, S. (2013). Echinoderm fauna of Kosrae, The Federation States of Micronesia. *Animal Systematics, Evolution, and Diversity*, 29(1), 1-17.



- LGU-Dumalag (2008). *Marine Protected Area Management Plan of Punta Dumalag 2008-2012*.
- Mann, K., Poustka, A. J., & Mann, M. (2008). The sea urchin (*Strongylocentrotus purpuratus*) test and spine proteomes. *Proteome Science*, 6(22), 1-10. doi: 10.1186/1477-5956-6-22.
- Mortillero, H. J. (2009). The distribution of sea urchin (*Echinoide*) in Baranagay Tigpan A, Babak District, Island Garden City of Samal. Bachelor of Science in Biology Undergraduate Thesis, University of Southeastern Philippines.
- Muthiga, N. (2003). Coexistence and reproductive isolation of the sympatric echinoids *Diadema savi-gnyi* Michelin and *Diadema setosum* (Leske) on Kenyan coral reefs. *Marine Biology* 143, 669-677.
- Muthiga, N., McClanahan, T., & Lawrence, J. (2007). *Edible Sea Urchins: Biology and Ecology*. New York: Elsevier Science.
- Prado, V. V., Galvez, G. N., & Rivera, R. (2012). Size structure and density of sea urchin, *Tripteneustes gratilla* along Balaoan waters, La Union, Philippines. *E-International Scientific Research Journal*, 4(3), 197-203.
- Pomroy, C. M. (2003). A guide to the shallow-water *Echinodermata* of the Texas Coast. *Contributions in Marine Science*, 36, 1-188.
- Riveros, A., Zuniga, M., Larrain, A., & Becerra, J. (1996). Relationships between fertilization of the Southeastern Pacific sea urchin *Arbacia spatuligera* and environmental variables in polluted coastal waters. *Marine Ecology Progress Series*, 134, 159-169.
- Salas, A., Diaz, F., Re, A. D., Gonzalez, M., & Galindo, C. (2012). Thermoregulatory behavior of red sea urchin *Strongylocentrotus Franciscanus* (Agassiz, 1863) and purple sea urchin *Strongylocentrotus Purpuratus* (Stimpson, 1857) (Echinodermata: Echinoidea). *The Open Zoology Journal*, 5, 42-46.
- Shafir, S., & McClanahan, T. (1990). Causes and consequences of sea urchin abundance and diversity of in Kenyan coral reef lagoons. *Oecologia* 8, 362-370.
- Shimda, A. & Wilson, M. (1985). Biological determinants of species diversity. *Journal of Biogeography*. 12, 1-20.
- Taylor, B. J. (1996). Taxonomy and morphology of *Echinodermata* from the Aptian of Alexander Island. *British Antarctic Survey Bulletin*, 8, 1-18.
- Tehranifard, A., Oryan, S., & Vosoghi, G. (2011). Introducing a holothuria sea cucumber species *Stichnopus hermani* from Kish Island in the Persia Gulf in Iran. *International Conference on Environment and Industrial Innovating* 12, 1-6.
- Tomsic, S., Conides, A., Dupric I., & Glamuniza, B. (2010). Growth, size class frequency and reproduction of puplar sea urchin, *Paracentrotus lividus* (Lamarck, 1816) in Brastina Bay (Adriatic Sea, Croatia). *Acta Adriata* 51, 65-74. Retrieved from <http://www.google.com.ph/url?sa=t&rct=j&q=&esrc=s&source=web&cd=1&ved=>
- Zizka, M. D. (2012). Sea urchins on the move: Distribution change of *Echinometra* in Mo'orea French Polynesia. *Berkeley Scientific Journal*, 15(1), 1-10.