

Reviewing the world's edible mushroom species: A new evidence-based classification system

Huili Li^{1,2,3}  | Yang Tian⁴ | Nelson Menolli Jr^{5,6} | Lei Ye^{1,2,3} | Samantha C. Karunarathna^{1,2,3} | Jesus Perez-Moreno⁷ | Mohammad Mahmudur Rahman⁸ | Md Harunur Rashid⁸ | Pheng Phengsintham⁹ | Leela Rizal¹⁰ | Taiga Kasuya¹¹ | Young Woon Lim¹² | Arun Kumar Dutta¹³ | Abdul Nasir Khalid¹⁴ | Le Thanh Huyen¹⁵ | Marilen Parungao Balolong¹⁶ | Gautam Baruah¹⁷ | Sumedha Madawala¹⁸ | Naritsada Thongklang^{19,20} | Kevin D. Hyde^{19,20,21} | Paul M. Kirk²² | Jianchu Xu^{1,2,3} | Jun Sheng²³ | Eric Boa²⁴ | Peter E. Mortimer^{1,3}

¹ CAS Key Laboratory for Plant Diversity and Biogeography of East Asia, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming, Yunnan, China

² East and Central Asia Regional Office, World Agroforestry Centre (ICRAF), Kunming, Yunnan, China

³ Centre for Mountain Futures, Kunming Institute of Botany, Kunming, Yunnan, China

⁴ College of Food Science and Technology, Yunnan Agricultural University, Kunming, Yunnan, China

⁵ Núcleo de Pesquisa em Micologia, Instituto de Botânica, São Paulo, Brazil

⁶ Departamento de Ciências da Natureza e Matemática (DCM), Subárea de Biologia (SAB), Instituto Federal de Educação, Ciência e Tecnologia de São Paulo (IFSP), São Paulo, Brazil

⁷ Colegio de Postgraduados, Campus Montecillo, Texcoco, México

⁸ Global Centre for Environmental Remediation (GCER), Faculty of Science, The University of Newcastle, Callaghan, NSW 2308, Australia

⁹ Biology Department, National University of Laos, Lao PDR

¹⁰ The University of Queensland, School of Biological Sciences, Brisbane, Queensland, Australia

¹¹ Department of Biology, Keio University, Yokohama, Kanagawa, Japan

¹² School of Biological Sciences and Institute of Microbiology, Seoul National University, Seoul, Korea

¹³ Department of Botany, West Bengal State University, Barasat, West Bengal, India

¹⁴ Department of Botany, University of the Punjab, Lahore, Pakistan

¹⁵ Department of Toxicology and Environmental Monitoring, Faculty of Environment, Hanoi University of Natural Resources and Environment, Tu Liem North District, Hanoi, Vietnam

¹⁶ Department of Biology, College of Arts and Sciences, University of the Philippines, Manila, the Philippines

¹⁷ Balipara Tract and Frontier Foundation, Guwahati, Assam, India

¹⁸ Department of Botany, Faculty of Science, University of Peradeniya, Peradeniya, Sri Lanka

¹⁹ Center of Excellence in Fungal Research, Mae Fah Luang University, Chiang Rai, Thailand

²⁰ School of Science, Mae Fah Luang University, Chiang Rai, Thailand

²¹ Mushroom Research Foundation, Chiang Mai, Thailand

²² Biodiversity Informatics and Spatial Analysis, Jodrell Laboratory, Royal Botanic Gardens Kew, Surrey, UK

²³ Key Laboratory for Agro-biodiversity and Pest Control of Ministry of Education, Yunnan Agricultural University, Kunming, China

²⁴ Institute of Biology, University of Aberdeen, Aberdeen, UK

Correspondence

Peter E. Mortimer, Centre for Mountain Futures, Kunming Institute of Botany, Kunming 650201, Yunnan, China.
Email: peter@mail.kib.ac.cn;
Eric Boa, Institute of Biology, University of Aberdeen, United Kingdom.
Email: e.boa@abdn.ac.uk;
Jun Sheng, Key Laboratory for Agrobiodiversity and Pest Control of Ministry of Education, Yunnan Agricultural University, Kunming, 650201, China.
Email: shengjunpuer@qq.com;
Jianchu Xu, Center for Mountain Futures, Kunming Institute of Botany, Kunming 650201, Yunnan, China.
Email: jxu@mail.kib.ac.cn;
Nelson Menolli Jr, Núcleo de Pesquisa em Micologia, Instituto de Botânica, Av. Miguel Stefano 3687, Êgua Funda, São Paulo, SP 04301-012, Brazil.
Email: menollijr@yahoo.com.br

Funding information

Yunnan Provincial Key Programs of Yunnan Eco-friendly Food International Cooperation Research Center Project, Grant/Award Number: 2019ZG00908; Key Research Program of Frontier Sciences “Response of Asian mountain ecosystems to global change,” CAS, Grant/Award Number: QYZDY-SSW-SMC014; São Paulo Research Foundation, Grant/Award Number: #2018/15677-0; National Science Foundation of China, Grant/Award Numbers: 41761144055, 41771063

Abstract

Wild mushrooms are a vital source of income and nutrition for many poor communities and of value to recreational foragers. Literature relating to the edibility of mushroom species continues to expand, driven by an increasing demand for wild mushrooms, a wider interest in foraging, and the study of traditional foods. Although numerous case reports have been published on edible mushrooms, doubt and confusion persist regarding which species are safe and suitable to consume. Case reports often differ, and the evidence supporting the stated properties of mushrooms can be incomplete or ambiguous. The need for greater clarity on edible species is further underlined by increases in mushroom-related poisonings. We propose a system for categorizing mushroom species and assigning a final edibility status. Using this system, we reviewed 2,786 mushroom species from 99 countries, accessing 9,783 case reports, from over 1,100 sources. We identified 2,189 edible species, of which 2,006 can be consumed safely, and a further 183 species which required some form of pretreatment prior to safe consumption or were associated with allergic reactions by some. We identified 471 species of uncertain edibility because of missing or incomplete evidence of consumption, and 76 unconfirmed species because of unresolved, differing opinions on edibility and toxicity. This is the most comprehensive list of edible mushrooms available to date, demonstrating the huge number of mushrooms species consumed. Our review highlights the need for further information on uncertain and clash species, and the need to present evidence in a clear, unambiguous, and consistent manner.

KEYWORDS

edibility, foraging, mycology, poisonous mushrooms, wild foods

1 | INTRODUCTION

The earliest reports of mushroom consumption are from Spain (18,700 years ago), China (5,000 to 6,000 years ago), and Egypt (4,600 years ago) (Chang, 2006; Power, Salazar-García, Straus, Morales, & Henry, 2015; Straus, Morales, Carretero, & Marín-Arroyo, 2015; Zhang, Chen, Huang, Gao, & Qu, 2015). The exact process by which early humans identified edible mushroom species—those that were safe and suitable to eat—is unclear, but there is little doubt it was by trial and error, a common approach used for wild plants and other living things that were hunted or gathered for food. Small amounts were tentatively tasted, before smell, texture, and lack of any adverse reaction decided which mushrooms could be eaten. Other species were either ignored because they lacked taste or were difficult to digest, or actively avoided because they were toxic. The “Universal Edibility Test” published in the US Army Survival Manual FM 21–76 is a modern version of this process (Survival Use of Plants, 2020). The introduction of

cooking (Yang, Belwal, Devkota, Li, & Luo, 2019) expanded what plants and mushrooms were “safe and suitable” to consume, through softening of tissues and detoxification. As Arora (1986) points out: “The only way to determine the edibility of a mushroom is to eat it.”

Modern diets depend heavily on cultivated foods, yet foraging for wild plants and mushrooms is still widespread, both of necessity and for pleasure. The traditional knowledge of rural and indigenous communities remains a historically significant source of knowledge on edibility of wild mushrooms. Ethnomycological studies from around the world continue to document which species are considered edible and poisonous. Notable examples by region include Africa (Buyck, 2008; Buyck & Nzigidahera, 1995; Degreef, Malaisse, Rammeloo, & Baudart, 1997; Morris, 1984; Pegler & Pearce, 1980), Asia (Brown, 2019; Chamberlain, 1996; Fui, Saikim, Kulip, & Seelan, 2018; Kang et al., 2016; Kumar, Harsh, Prasad, & Pandey, 2017), Europe (Cai, Pectenella, & Vidale, 2011; Kasper-Pakosz, Pietras, & Łuczaj, 2016; Łuczaj,

Stawarczyk, Kosiek, Pietras, & Kujawa, 2015; Pieroni, Nebel, Santoro, & Heinrich, 2005), Australia (Kalotas, 1997), North America (Álvarez-Farías, Diaz-Godínez, Téllez-Téllez, Villegas, & Acosta-Urdapilleta, 2016; de Avila, Welden, & Guzmán, 1980; Garibay-Orijel et al., 2020; Ruan-Soto, 2018), Central America (Guzmán, 2001b; Morales, Bran, & Cáceres, 2010), and South America (Fidalgo & Prance, 1976; Gamboa-Trujillo et al., 2019; Sanuma et al., 2016; Vasco-Palacios, Suaza, Castañón-Betancur, & Franco-Molano, 2008). An increasing knowledge of how mushrooms grow has led to the cultivation of over 90 species (Boa, 2004), of which around 30 are grown commercially for food and consumed widely (Chang & Miles, 2004). The remaining cultivated species are grown for medicinal purposes (Lu, Lou, Hu, Liu, & Chen, 2020; Ma et al., 2018; Roncero-Ramos & Delgado-Andrade, 2017; Yang, Belwal, Devkota, Li, & Luo, 2019), as functional foods (Jabłońska-Ryś, Skrzypczak, Sławińska, Radzki, & Gustaw, 2019; Reis, Martins, Vasconcelos, Morales, & Ferreira, 2017) and extracts used as food additives (Hadar & Dosoretz, 1991; Sun et al., 2020).

The overall result is that mushroom consumption has risen 21-fold over the last 56 years (Food and Agriculture Organization Statistical, 2017). Harvesting of wild mushrooms, the vast majority of which cannot be cultivated, remains hugely important, both as a source of food and income (Boa, 2004; Hall, Buchanan, Cole, Yun, & Stephenson, 2003). The first significant attempt to list edible species from around the world was by Chandra (1989). She listed 628 edible mushroom species, with notes on taste and how to prepare before eating. An extensive global review of the literature, including gray literature and personal contacts, by Boa (2004) identified 2166 edible mushroom species collected across 85 countries. Peintner et al. (2013) listed 268 wild mushroom species of commercial importance from 27 European countries. Wu et al. (2019) list 1,020 edible mushroom species and 480 poisonous mushroom species from China. Some authors note edible species that contain toxins when raw and require pretreatment before safe consumption is possible (Niksic, Klaus, & Argyropoulos, 2016). The published information on edible properties varies considerably, however, with field guides intended for a wider, non-academic readership, providing much more detail.

Interest in edible mushrooms continues to grow as the demand for new types of food increases and alternative sources of income for rural communities are sought (Pilz & Molina, 2002). In Asia, Europe and North America more people than ever are “foraging,” reviving mushroom traditions diluted by migrations to cities and creating new challenges for sustainable production (Egli, Peter, Buser, Stahel, & Ayer, 2006). Harvesting wild species of commercial importance, such as matsutake (*Tricholoma*

spp.), boletes (*Boletus* spp.), truffles (*Tuber* spp.), morels (*Morchella* spp.), and various *Lactarius* species (e.g., *L. deliciosus*), is big business in many countries and essential income for collectors and their families (Boa, 2004; de-Román & Boa, 2006; Yeh, 2000).

Regular reports of mushroom poisoning continue to occur. There is no evidence to suggest that the frequency of such reports has significantly increased on a global scale, but there is little doubt that the greater ease of sharing information through social media and the internet has heightened concerns about the danger of eating wild mushrooms. Govorushko, Rezaee, Duanov, and Tsatsakis (2019) report that between 200 and 250 people are poisoned by mushrooms each year. Furthermore, Chen, Ping, and Zhang (2014) reported that between the years 2000 and 2014, there were 1,954 poisoning cases and 409 deaths related to mushroom use in China.

Reports on edible and poisonous mushrooms appear regularly in many different publications, including peer-reviewed papers, field guides, newsletters of mycological and nature societies, and “gray literature,” such as project reports and informal publications. A plethora of websites offer opinions on what mushrooms to eat and what to avoid yet reports are often inconsistent and contradictory. Opinions vary on the culinary merits of a species, but, more importantly, on what species can be eaten and which ones are poisonous. These disagreements occur partly because of different cultural attitudes and biases (Härkönen, 2002), but also because of the degree of risk people are willing to accept, which is often tied to the depth of knowledge regarding the use of wild mushrooms. Those with a rich tradition of eating mushrooms from the wild are often more adventurous in the species they will eat. Where this tradition is weak or absent, fewer species from the wild are consumed, and there is an increased tendency to label species as “inedible,” “avoid,” or “suspect.” Such terms imply potential toxicity, yet Shernoff (2015) suggested that many species labeled as “inedible” in US field guides were harmless.

In an attempt to distinguish reports that showed clear evidence of a species having been consumed (either by the author(s) or reliable cited source), Boa (2011) separated “food” species from “edible” species, with an additional category for poisonous species. These categories were an attempt to standardize reports on properties and to enable comparisons from different sources. It was also an attempt to emphasize the quality of evidence presented for a report and therefore the reliability of the report.

The most reliable reports on edibility are those that provide evidence of consumption, which could include personal experiences of authors or those of colleagues (e.g., Arora, 1986) or cited sources (e.g., Chandra, 1989). A surprising number of peer-reviewed publications do not give

the source of evidence, as discussed by Rubel and Arora (2008). No source is given for the 1,323 edible species reported from China (Wu et al., 2019), for example. Field guides written for naturalists and foragers are a rich source of first-hand information on a wide range of edible and poisonous species, often over a large physical area (e.g., Rodríguez et al., 1999; Tudor, 2010; Wang, Liu, & Yu, 2004). Ethnomycology studies provide in-depth local knowledge on edible mushrooms for a defined region (Brown, 2019; Comandini & Rinaldi, 2020).

The increasing interest in foraging and the commercial importance of edible mushrooms collected from the wild emphasize the need for reliable information on the properties of species. Medical staff need accurate information on properties to diagnose cases of suspected mushroom poisonings. There is also a wider imperative to ensure that consumers understand any risks associated with edible species, such as the need to prepare in a particular way or avoid consuming with alcohol. Current confusions and disagreements about the properties of edible mushrooms gathered from the wild limits their potential both as food and a source of income. Of the 2,166 edible mushrooms listed in Boa (2011), 589 species recorded as edible lacked clear and credible evidence of consumption.

The uncertainty surrounding the edibility of a significant number of mushroom species and the need for consistent reporting has prompted the current review of published case reports from around the world. This paper describes an evidence-based approach for weighing up published information and establishing greater clarity of current knowledge on the edibility of wild mushrooms.

2 | DATA RESOURCES UTILIZED IN THE REVIEW PROCESS

Extensive search was carried out in peer-reviewed publications, field guides, books, conference proceedings, project reports, and other information available on the web or from personal knowledge of mycologists for information concerning the edibility of mushroom species. Each record of a species property is referred to here as a case report. We included two records of slime molds in our list of edible mushrooms (*Fuligo septica* var. *septica* and *Reticularia lycoperdon*) as these have been considered as edible fungi in the past, and many collectors of wild fungi still treat them accordingly. The original records varied in length from one-word labels to more detailed discussions of properties.

This study builds on an original summary of case reports that appeared in Boa (2004), updated to include information from new sources published up to April 2020. The earliest source of case reports is from a 1849 publi-

cation by Lindley (Lindley, 1849). The complete data set contains 9,783 case reports from 99 countries and over 1,100 sources. We incorporated the following languages in our search: Bengali, Bhutanese, Burmese, English, Filipino, Japanese, Korean, Laotian, Malayalam, Mandarin, Nepali, Portuguese, Russian, Spanish, Tamil, Thai, Vietnamese, and Urdu. Scientific names follow indexfungorum.org (Index Fungorum, 2020) and speciesfungorum.org (Species Fungorum, 2020). Current names are used and spelling errors that occurred in the case reports have been corrected. The published scientific names used in the case reports appear in Supporting Information Table S1 together with their current names.

3 | SYSTEM FOR CATEGORIZING MUSHROOM EDIBILITY

Boa (2004) assigned case reports to a series of categories defining the key characteristics of mushrooms and, crucially, distinguishing between reports which provided clear evidence of consumption. The original scheme has been simplified (Table 1) and four categories have been used to classify all case reports analyzed based on available information: edible (E1); edible with conditions (E2); edible, unconfirmed (E3); and poisonous (P).

The advantage of this classification is that it enables a cross-comparison of case reports from different sources and establishes a consistent approach for determining the final edibility status (FES) of a species. The disadvantage of using the four categories listed in Table 1 is that the finer details of extended case reports are lost, including comments on taste and texture and possible toxicity to people. Taste and texture are subjective features and are not part of the current analysis. Where information is unclear, ambiguous or missing, case reports are assigned to category E3. This category may also include species which are safe to eat but are deemed unsuitable because of taste, smell, texture or some other feature that people find undesirable. Such species were often described as “inedible.” Rubel and Arora (2008) explain how lack of awareness of different cultural attitudes and practices has influenced and distorted the perception of *Amanita muscaria* as a highly poisonous mushroom, despite the fact that it can, with suitable treatment, be safely consumed.

Almost half of the 2,786 species we report on had only one case report (Table 3). This becomes the *de facto* FES. For the remaining species, we separated those with consistent case reports from those with mixed case reports (Figure 1), which were then assessed and given an FES based on a “summary judgement” (Table 2). Note that all species with a combination of at least one poisonous and at least one E1, E2, or E3 case report were given an

TABLE 1 Edibility categories (E1, E2, E3, and P) assigned to classify reported properties of mushrooms

Edibility category	Code	Description
Edible, confirmed	E1	Clear evidence that a species has been consumed without adverse or harmful effects and is safe and suitable to eat
Edible, confirmed but with conditions	E2	Clear evidence that a species has been consumed after it has been cooked or prepared in such a way as to make it safe and suitable to eat. Includes species that may result in allergic or adverse reactions when consumed with alcohol, for example
Edible, unconfirmed	E3	Evidence of safe consumption is uncertain or incomplete
Poisonous	P	Causes adverse and harmful reactions when consumed




FIGURE 1 Flow chart for determination of final edibility status

“unconfirmed” (U) FES. These U species were then given an additional level of scrutiny by the authors, being reviewed and classified as E1, E2, or P depending on the balance and strength of the available evidence. For exam-

ple, a U species sold at a market provided compelling evidence that it was edible. When no FES could be made using reliable evidence, the status remained U, an indication that further research is required to clarify the edibility of these species. Species with mixed case reports of E3 and P only were described as poisonous (Table 2).

TABLE 2 Assessing the final edibility status of mushrooms with differing (mixed) case reports

Combination of case reports	Final edible status	Notes
E1 and E2	E2	
E2 and E3	E2	
E1 and E3	E1	
E1, E2 and E3	E2	
P and E3 only	E3	
P and E1, E2 or E3 or combination	U (unconfirmed)	Species were reclassified (E1, E2, or P) if supported by the balance of evidence contained in case reports

The number of consistent case reports obtained indicates the general strength of the supporting evidence. This strength is also, however, related to the number of people who have eaten a mushroom of a particular species without ill-effect, information that varied considerably in the original source or was simply missing. No distinction was made between case reports that described varying degrees or intensity of adverse reactions to consumption of poisonous species. Accordingly, the poisonous category includes species described as mildly poisonous, potentially lethal or deadly, or other variations on a toxic theme. All reports of poisonous species were taken at face value, regardless of the strength of the supporting evidence.

Consistent efforts were made to obtain at least two case reports for each species and to ensure that these were from different countries or major regions within larger

TABLE 3 Final edibility status of 2,786 mushrooms based on an analysis of 9,783 case reports from 99 countries

Final edibility status (FES)	Taxa	One case report	More than one case report, consistent	More than one case report, mixed
E1: edibility confirmed	2,006	964	692	350
E2: edible but with conditions	183	16	0	167
E3: edibility uncertain	471	453	16	2
P: poisonous	50	0	0	50
U: unconfirmed	76	0	0	76
Total	2,786	1,433	708	645




FIGURE 2 Final edibility status of E1 and E2 taxa from different regions

countries. This was often not possible, as noted in the results and discussed later in this paper. We did not systematically attempt to document all ostensibly poisonous species. Sources of information on edible mushrooms usually contained case reports of poisonous species, but it was clear that lists were often incomplete. Laessøe and Petersen (2019) list 112 poisonous species for temperate Europe that are not yet included in our data set, for example.

Further information and wider evaluation are required for those species with mixed case reports, especially those with conflicting reports of edibility and toxicity. In addition to giving extra weight to those species sold in markets when reviewing U species, we also looked carefully at first-hand information provided in a number of key sources from different regions: North America (e.g., Arora, 1986; McIlvaine & Macadam, 1900; Pérez-Moreno, Lorenzana-Fernández, Carrasco-Hernández, & Yescas-Pérez, 2010; Villarreal & Perez-Moreno, 1989a), Europe (e.g., Afonkin, 2013; Papetti, Consiglio, & Simonini, 2017), and Asia (e.g., Adhikari, 2000; Wang, Liu, & Yu, 2004).

4 | OVERVIEW OF WORLD'S EDIBLE MUSHROOMS

We report on the properties of 2,786 taxa, including four forms, two subspecies, and one variety (Table 3), obtained from 9,783 case reports (the source of individual case reports is shown in Supporting Information SI). We obtained case reports of wild mushroom species from 33 countries in Africa, 26 from Asia, 20 from Europe, 18 from the Americas, and two from Oceania. The highest number of edible mushroom species (E1 and E2) were reported in Asia (1,493), followed by Europe (629), North America (487), Africa (351), South America (204), Central America (100), and Oceania (19) (Figure 2). There are 614 edible mushroom species (E1 and E2) found on two or more continents.

The vast majority of species (72%) had an FES of E1, that is those mushrooms with clear evidence of being safe and suitable to eat. This group included 350 (17%) taxa with an FES of E1 which had mixed case reports, mainly “confirmed” and “uncertain or incomplete” edibility. Concrete




FIGURE 3 The number of taxa from genera with 10 or more case reports and their final edibility status. E1: confirmed edible; E2: edible with conditions; E3: uncertain; U: unconfirmed edible; P: poisonous

evidence of consumption here carries most weight. A lack of evidence does not mean that a species cannot be eaten or is poisonous. It may simply indicate that a species is not eaten in a particular place.

Of the 230 taxa originally classified as “unconfirmed,” a small but notable number of taxa (76) could not be reclassified (Table 3) because of unresolved clashes on reported properties. A total of 154 taxa with contradictory evidence on safe consumption were reclassified, of which 28 taxa were given an FES of E1 after consulting reliable sources of evidence (Table 4), and 76 taxa given an FES of E2. Precedence was given in mixed case reports to at least one stating preconditions for consumption or allergic reactions.

The sum of taxa with an FES of E3 or U accounts for 20% of all the taxa recorded in our review of global sources, and further research is needed to decide whether these are edible or poisonous. The majority of taxa with an FES of E3 had only one case report. The depth of reporting on the properties of taxa with an FES of E1 and E2 is also weak, with 908 species, 35% of the combined taxa, represented by one case report.

Data on the FES of species from major genera with 10 or more taxa are shown in Figure 3 and Supporting Informa-

tion Table S2. *Russula* had the greatest number of recorded species (161), with 143 confirmed as edible (E1 or E2), 11 as E3, one as P, and six of unconfirmed FES. *Amanita* had the second highest number of taxa (100 species and 1 forma specialis), including 78 edible taxa (E1 or E2), 5 as E3, 8 as P, and 10 species of unconfirmed FES. *Lactarius* was third with 88 species, comprising 76 edible species (E1 or E2), 2 as E3, and 10 species as U. Nine genera contained only confirmed edible species (E1 or E2): *Auricularia* (21 spp.), *Butyriboletus* (10 spp.), *Chroogomphus* (11 spp.), *Clavariadelphus* (10 spp.), *Hygrophorus* (39 spp.), *Laccaria* (32 spp.), *Melanoleuca* (16 spp.), *Termitomyces* (29 spp.), and *Tuber* (38 spp.).

We calculated the number of edible species (FES of E1 or E2) for major genera as a proportion of all the species shown in speciesfungorum.org (Species Fungorum, 2020). Although *Russula* had the highest number of edible species, these represented only 11% of all species (1322 spp.). The same proportion of edible species was found for *Amanita* (Supporting information Table S2 & S3). Edible species of *Auricularia*, *Retiboletus*, *Sparassis*, and *Termitomyces* represent more than 50% of all the species recorded in speciesfungorum.org. *Entoloma* and *Inocybe*

TABLE 4 Revised final edibility status of 154 taxa with mixed case reports, including at least one poisonous, and originally classified as U (unconfirmed) *. E1: Edibility confirmed; E2: Edibility with conditions; E3: Edibility uncertain; P: Poisonous

Taxa	Number of case reports					Final edibility status (FES)	Source(s) used to revise FES
	E1	E2	E3	P	All		
<i>Agaricus abruptibulbus</i>	2	0	0	1	3	E2	(Arora, 1986; Zhou & Zhang, 2005)
<i>Agaricus arvensis</i>	27	0	2	1	30	E2	(Afonkin, 2013; Arora, 1986)
<i>Agaricus phaeolepidotus</i>	0	0	1	1	2	P	
<i>Agaricus semotus</i>	3	0	2	1	6	E1	(Afonkin, 2013)
<i>Agaricus sylvicola</i>	24	0	2	1	27	E2	(Afonkin, 2013; Arora, 1986; McIlvaine & Macadam, 1900)
<i>Amanita castanopsidis</i>	0	0	1	2	3	P	
<i>Amanita citrina</i>	1	0	2	8	11	P	(Bresinsky & Bresl, 1990)
<i>Amanita excelsa</i>	4	2	2	3	11	E2	(McIlvaine & Macadam, 1900)
<i>Amanita flavipes</i>	0	0	1	1	2	P	
<i>Amanita fuliginea</i>	0	0	1	4	5	P	
<i>Amanita ovoidea</i>	3	0	0	2	5	E1	(Afonkin, 2013; Papetti et al., 2017)
<i>Amanita porphyria</i>	0	0	2	5	7	P	
<i>Amanita pseudogemmata</i>	0	0	1	2	3	P	
<i>Amanita pseudoporphyria</i>	2	0	0	5	7	E2	(Afonkin, 2013; Wang et al., 2004)
<i>Amanita rubrovolvata</i>	0	0	1	4	5	P	
<i>Amanita rufoferruginea</i>	0	0	1	4	5	P	
<i>Amanita vaginata</i>	19	4	10	2	35	E2	(Afonkin, 2013; Arora, 1986)
<i>Ampulloclitocybe clavipes</i>	7	4	2	1	14	E2	(Afonkin, 2013; Arora, 1986; McIlvaine & Macadam, 1900)
<i>Astraeus hygrometricus</i>	7	1	1	1	10	E2	(Afonkin, 2013; Arora, 1991)
<i>Baorangia pseudocalopus</i>	1	0	0	2	3	P	(Li et al., 2019)
<i>Bolbitius titubans</i>	0	0	1	4	5	P	
<i>Boletus sensibilis</i>	2	0	1	4	7	E1	(Wang et al., 2004)
<i>Boletus subvelutipes</i>	3	0	0	3	6	E1	(Rubel & Arora, 2008)
<i>Caloboletus calopus</i>	3	0	1	6	10	E2	(Wang et al., 2004)
<i>Chlorophyllum molybdites</i>	5	0	1	21	27	E2	(Arora, 1986)
<i>Chlorophyllum rhacodes</i>	19	2	0	2	23	E2	(Afonkin, 2013)
<i>Clathrus archeri</i>	2	0	0	2	4	E2	(Arora, 1986)

(Continues)

TABLE 4 (Continued)

Taxa	Number of case reports					Final edibility status (FES)	Source(s) used to revise FES
	E1	E2	E3	P	All		
<i>Clavariadelphus pistillaris</i>	8	1	1	2	12	E1	(Arora, 1986)
<i>Clitocybe dealbata</i>	1	0	0	11	12	P	(Afonkin, 2013; Papetti et al., 2017)
<i>Clitocybe fragrans</i>	3	0	1	4	8	E2	(McIlvaine & Macadam, 1900)
<i>Clitocybe nebularis</i>	14	4	2	3	23	E2	(Afonkin, 2013; Arora, 1986; Zhou & Zhang, 2005)
<i>Clitocybe phyllophila</i>	2	0	1	11	14	P	(Afonkin, 2013)
<i>Clitocybe rivulosa</i>	0	0	1	6	7	P	
<i>Conocybe apala</i>	0	0	1	3	4	P	
<i>Conocybe tenera</i>	0	0	1	2	3	P	
<i>Coprinellus micaceus</i>	5	6	3	1	15	E2	(Arora, 1986)
<i>Coprinopsis atramentaria</i>	11	6	2	6	25	E2	(Afonkin, 2013; Arora, 1986; Zhou & Zhang, 2005)
<i>Coprinopsis picacea</i>	1	0	0	1	2	E2	(Arora, 1986; McIlvaine & Macadam, 1900)
<i>Coprinus comatus</i>	31	3	4	1	39	E1	(Afonkin, 2013; Arora, 1986)
<i>Cortinarius bolaris</i>	0	0	1	1	2	P	
<i>Cortinarius cinnamomeus</i>	4	0	0	4	8	E2	(McIlvaine & Macadam, 1900)
<i>Cortinarius gentilis</i>	0	0	1	1	2	P	
<i>Cortinarius sanguineus</i>	0	0	1	4	5	P	
<i>Cortinarius semisanguineus</i>	0	0	1	2	3	P	
<i>Cortinarius traganus</i>	0	0	1	2	3	P	
<i>Cyanoboletus pulverulentus</i>	8	0	0	2	10	E1	(Afonkin, 2013; Papetti et al., 2013)
<i>Deconica merdaria</i>	0	0	1	2	3	P	
<i>Dictyophora multicolor</i>	1	0	0	1	2	E2	(Arora, 1986)
<i>Entoloma omiense</i>	0	0	1	2	3	P	
<i>Entoloma rhodopolium</i>	1	0	1	14	16	P	(Afonkin, 2013)
<i>Entoloma vernum</i>	1	0	0	4	5	P	(Afonkin, 2013)
<i>Gliophorus psittacinus</i>	2	0	1	1	4	E2	(McIlvaine & Macadam, 1900)
<i>Gymnopilus aeruginosus</i>	0	0	1	3	4	P	
<i>Gymnopilus junonius</i>	2	2	0	8	12	E2	(Afonkin, 2013; Deschamps, 2002)
<i>Gyromitra esculenta</i>	3	3	5	13	24	E2	(Afonkin, 2013; Arora, 1986)
<i>Gyromitra gigas</i>	3	1	0	5	9	E2	(Afonkin, 2013; Arora, 1986)

(Continues)

TABLE 4 (Continued)

Taxa	Number of case reports					Final edibility status (FES)	Source(s) used to revise FES
	E1	E2	E3	P	All		
<i>Gyroporus castaneus</i>	11	0	4	1	16	E1	(Arora, 1986; Zhou & Zhang, 2005)
<i>Hebeloma fastibile</i>	2	0	0	1	3	E2	(Pérez-Moreno, Martínez-Reyes, Yescas-Pérez, Delgado-Alvarado, & Xoconostle-Cázares, 2008)
<i>Hebeloma mesophaeum</i>	1	0	0	1	2	E2	(Pérez-Moreno et al., 2008)
<i>Hebeloma sacchariolens</i>	0	0	1	1	2	P	
<i>Hebeloma sinapizans</i>	0	0	1	3	4	P	
<i>Helvella acetabulum</i>	7	0	0	1	8	E1	(Montoya-Esquivel, Estrada-Torres, Kong, & Juárez-Sánchez, 2001)
<i>Helvella crispa</i>	19	0	0	2	21	E2	(McIlvaine & Macadam, 1900; Pérez-Moreno et al., 2008)
<i>Helvella elastica</i>	8	0	2	2	12	E2	(McIlvaine & Macadam, 1900; Pérez-Moreno et al., 2008)
<i>Helvella lacunosa</i>	18	1	2	2	23	E2	(Arora, 1986; Pérez-Moreno et al., 2008)
<i>Hygrocybe nigrescens</i>	1	0	1	2	4	E2	(Pérez-Moreno et al., 2008)
<i>Hygrocybe ovina</i>	0	0	1	1	2	P	
<i>Hygrocybe punicea</i>	8	0	2	1	11	E2	(Arora, 1986)
<i>Hygrophoropsis aurantiaca</i>	9	1	3	1	14	E2	(Arora, 1986)
<i>Hypholoma capnoides</i>	8	0	1	1	10	E2	(Arora, 1986; McIlvaine & Macadam, 1900)
<i>Imleria badia</i>	20	0	3	1	24	E2	(Arora, 1986; McIlvaine & Macadam, 1900)
<i>Imperator rhodopurpureus</i>	0	1	0	2	3	E2	(McIlvaine & Macadam, 1900)
<i>Infundibulicybe gibba</i>	15	0	1	1	17	E2	(Arora, 1986; Pérez-Moreno et al., 2008)
<i>Inocybe flavobrunnea</i>	0	0	1	2	3	P	
<i>Inocybe geophylla</i>	0	0	1	7	8	P	

(Continues)

TABLE 4 (Continued)

Taxa	Number of case reports					Final edibility status (FES)	Source(s) used to revise FES
	E1	E2	E3	P	All		
<i>Inonotus obliquus</i>	2	0	1	1	4	P	
<i>Lactarius piperatus</i>	17	2	3	1	23	E2	(Arora, 1986; Zhou & Zhang, 2005)
<i>Lactarius pubescens</i>	6	0	1	2	9	E1	(Afonkin, 2013)
<i>Lactarius repraesentaneus</i>	2	0	2	1	5	E1	(Afonkin, 2013)
<i>Lactarius rufus</i>	3	1	1	2	7	E1	(Afonkin, 2013)
<i>Lactarius scrobiculatus</i>	5	0	2	2	9	E1	(Afonkin, 2013; Papetti et al., 2017)
<i>Lactarius torminosus</i>	4	1	2	6	13	P	(Afonkin, 2013; Li et al., 2020; Papetti et al., 2017)
<i>Lactarius turpis</i>	5	0	1	1	7	E2	(McIlvaine & Macadam, 1900)
<i>Lactarius uvidus</i>	2	0	2	1	5	E1	(Afonkin, 2013)
<i>Lactarius vellereus</i>	9	0	3	1	13	E2	(McIlvaine & Macadam, 1900)
<i>Lactarius zonarius</i>	3	0	2	4	9	E1	(Afonkin, 2013)
<i>Lepiota clypeolaria</i>	2	0	3	2	7	P	(Afonkin, 2013; Papetti et al., 2017)
<i>Lepiota cristata</i>	0	0	1	8	9	P	
<i>Lepista irina</i>	8	0	2	1	11	E1	(Afonkin, 2013)
<i>Leucoagaricus americanus</i>	4	0	3	2	9	E2	(Afonkin, 2013; Mendoza, 1938)
<i>Leucoagaricus badhamii</i>	0	0	1	4	5	P	
<i>Leucoagaricus leucothites</i>	17	0	2	1	20	E1	(Afonkin, 2013; Papetti et al., 2017)
<i>Leucocoprinus birnbaumii</i>	0	0	1	3	4	P	
<i>Leucocoprinus cepistipes</i>	2	0	2	1	5	E2	(Afonkin, 2013; Zhou & Zhang, 2005)
<i>Lysurus arachnoideus</i>	0	0	1	2	3	P	
<i>Meiorganum curtisii</i>	0	0	1	2	3	P	
<i>Meripilus giganteus</i>	12	2	0	1	15	E2	(Afonkin, 2013)
<i>Mutinus bambusinus</i>	0	0	2	1	3	P	
<i>Mutinus caninus</i>	0	0	1	1	2	P	
<i>Mycena haematopus</i>	1	0	1	1	3	E2	(Arora, 1986)
<i>Mycena pura</i>	4	0	2	10	16	E2	(Arora, 1986; Zhou & Zhang, 2005)
<i>Neoboletus erythropus</i>	5	0	0	1	6	E2	(Arora, 1986)
<i>Neoboletus luridiformis</i>	11	2	3	1	17	E1	(Afonkin, 2013)
<i>Neolentinus lepideus</i>	9	2	0	1	12	E1	(Afonkin, 2013; McIlvaine & Macadam, 1900)

(Continues)

TABLE 4 (Continued)

Taxa	Number of case reports					Final edibility status (FES)	Source(s) used to revise FES
	E1	E2	E3	P	All		
<i>Omphalotus olearius</i>	1	0	1	14	16	P	(Bresinsky & Bresl, 1990; Papetti et al., 2017)
<i>Otidea cochleata</i>	1	0	0	1	2	E1	(Afonkin, 2013)
<i>Paxillus involutus</i>	2	3	2	14	21	E2	(McIlvaine & Macadam, 1900)
<i>Phaeolepiota aurea</i>	8	3	1	1	13	E2	(Arora, 1986)
<i>Phallus impudicus</i>	11	2	1	2	16	E2	(McIlvaine & Macadam, 1900)
<i>Phallus tenuis</i>	0	0	1	2	3	P	
<i>Pholiota flammans</i>	2	0	2	1	5	E2	(Arora, 1986; McIlvaine & Macadam, 1900)
<i>Pholiota squarrosa</i>	6	2	2	3	13	E2	(Afonkin, 2013)
<i>Pholiota terrestris</i>	2	0	0	3	5	E2	(Arora, 1986)
<i>Pluteus petasatus</i>	5	0	0	1	6	E1	(Afonkin, 2013)
<i>Pluteus salicinus</i>	2	0	1	3	6	E1	(Afonkin, 2013; Papetti et al., 2017)
<i>Protostropharia semiglobata</i>	2	0	3	1	6	E2	(McIlvaine & Macadam, 1900)
<i>Psathyrella corrugis</i>	0	0	1	1	2	P	
<i>Psilocybe coronilla</i>	4	0	1	4	9	E2	(Pérez-Moreno et al., 2008)
<i>Pulveroboletus ravenelii</i>	1	0	3	2	6	E1	(Wang et al., 2004)
<i>Ramaria flava</i>	13	1	2	3	19	E2	(Adhikari, 2000)
<i>Ramaria flavobrunnescens</i>	4	0	1	1	6	E2	(Pérez-Moreno et al., 2008)
<i>Ramaria formosa</i>	3	0	1	14	18	E2	(Afonkin, 2013; Wang et al., 2004)
<i>Rubroboletus satanas</i>	1	0	0	15	16	P	(Afonkin, 2013; Arora, 1991; Papetti et al., 2017)
<i>Russula adusta</i>	16	1	3	3	23	E2	(McIlvaine & Macadam, 1900; Wang et al., 2004)
<i>Russula densifolia</i>	8	1	1	4	14	E2	(Arora, 1986; Wang et al., 2004)
<i>Russula fellea</i>	1	0	1	1	3	E1	(Afonkin, 2013)
<i>Russula foetens</i>	3	0	2	7	12	E2	(McIlvaine & Macadam, 1900)
<i>Russula sardonia</i>	2	0	1	1	4	P	(Afonkin, 2013)
<i>Sarcosphaera coronaria</i>	3	1	1	2	7	E2	(Arora, 1986)
<i>Scleroderma cepa</i>	1	0	0	1	2	P	(Afonkin, 2013; Li et al., 2020)
<i>Scleroderma citrinum</i>	4	0	3	9	16	E2	(Wang et al., 2004)
<i>Scleroderma flavidum</i>	2	0	0	2	4	E1	(Wang et al., 2004)

(Continues)

TABLE 4 (Continued)

Taxa	Number of case reports					Final edibility status (FES)	Source(s) used to revise FES
	E1	E2	E3	P	All		
<i>Strobilomyces strobilaceus</i>	10	0	1	1	12	E1	(Arora, 1986; McIlvaine & Macadam, 1900)
<i>Stropharia aeruginosa</i>	9	0	0	3	12	E2	(Dai, Yang, Cui, Yu, & Zhou, 2009; Zhou & Zhang, 2005)
<i>Suillellus luridus</i>	10	2	3	2	17	E2	(McIlvaine & Macadam, 1900)
<i>Suillus americanus</i>	9	0	0	2	11	E2	(Pérez-Moreno et al., 2008)
<i>Suillus granulatus</i>	34	3	4	1	42	E2	(Arora, 1986; Pérez-Moreno et al., 2008)
<i>Suillus luteus</i>	30	2	0	1	33	E2	(Arora, 1986; Pérez-Moreno et al., 2008)
<i>Suillus placidus</i>	8	0	1	3	12	E2	(Wang et al., 2004)
<i>Suillus spraguei</i>	6	1	1	1	9	E2	(Arora, 1986)
<i>Tapinella panuoides</i>	2	0	1	2	5	P	(Afonkin, 2013)
<i>Trametes versicolor</i>	5	0	1	1	7	E2	(Arora, 1986)
<i>Tricholoma album</i>	6	0	1	3	10	E2	(McIlvaine & Macadam, 1900)
<i>Tricholoma equestre</i>	24	1	1	8	34	E2	(Arora, 1986; McIlvaine & Macadam, 1900)
<i>Tricholoma focale</i>	2	0	0	1	3	E1	(Afonkin, 2013)
<i>Tricholoma pessundatum</i>	3	1	1	3	8	E2	(Arora, 1986)
<i>Tricholoma saponaceum</i>	1	2	2	5	10	E2	(McIlvaine & Macadam, 1900; Zhou & Zhang, 2005)
<i>Tricholoma sejunctum</i>	8	0	1	2	11	E2	(McIlvaine & Macadam, 1900)
<i>Tricholoma sulphureum</i>	2	0	0	2	4	P	(Afonkin, 2013; Papetti et al., 2017)
<i>Tricholoma ustale</i>	2	1	1	4	8	E2	(McIlvaine & Macadam, 1900)
<i>Tricholomopsis rutilans</i>	4	4	3	3	14	E2	(Arora, 1986)
<i>Tylopilus neofelleus</i>	0	0	1	1	2	P	
<i>Verpa bohemica</i>	11	2	2	1	16	E2	(Arora, 1986)
<i>Volvopluteus gloiocephalus</i>	23	0	2	2	27	E1	(Afonkin, 2013; Jones, Whalley, & Hywel-Jones, 1994)

*Mixed case reports comprising P and E3 only were given an FES of P.

had the lowest proportion of edible species (< 0.6%) of all species.

The species listed in our study for *Auricularia*, *Butyriboletus*, *Chroogomphus*, *Clavariadelphus*, *Hygrophorus*, *Laccaria*, *Melanoleuca*, *Termitomyces*, and *Tuber* were all considered edible in their FES, without any preconditions or uncertainties, an indication that these species are undoubt-

edly safe to eat. Alternatively, there are 100 *Amanita* species included in our study, representing 14% of the species in this group, with 78 (11%) species listed as confirmed edible (Species Fungorum, 2020; Zhang, Tang, Cai, & Xu, 2015). The work of Zhang et al. (2015) listed approximately 50 *Amanita* species as being edible; thus, our research updates this number. *Amanita* species should

be treated with caution, as there are about 100 considered poisonous, and of these, about 50 species are considered to be deadly (Cai, Cui, & Yang, 2016), some of which could be easily confused with the edible *Amanita* species listed in our study. Similarly, although with fewer deadly species, *Russula* is another group of mushrooms to be treated with a degree of caution, as there are numerous poisonous species, often occurring alongside, and sharing similar morphologies as edible *Russula* species (143 listed in our study). Of the species listed within *Entoloma* and *Inocybe*, < 0.6% are confirmed edible species, and thus when eating mushrooms from these genera, extreme caution is recommended. Although not considered deadly to humans, there are a number of poisoning cases resulting from the consumption of *Entoloma* species (Chen, Yang, Bau, & Li, 2016; de Oliveira, 2009; Hossain & Park, 2016; Puschner, 2018; Schenk-Jaeger et al., 2012). Furthermore, *Inocybe erubescens* is considered deadly to humans, and a number of additional *Inocybe* species are poisonous, due to the presence of muscarine (Brown, Malone, Stuntz, & Tyler Jr, 1962; Lurie et al., 2009).

5 | THE IMPORTANCE OF DEVELOPING AN EVIDENCE-BASED CLASSIFICATION SYSTEM

5.1 | Assessing the viability of the evidence-based classification system

This review is the first systematic attempt to document and categorize edible mushroom species using an evidence-based approach. Much of the available information regarding species edibility is from unverified sources, word-of-mouth, or publications that have not been subject to any form of review, and it is unsurprising that reports on the edibility of a species often vary. The use of standard categories to describe case reports is an essential part of the method we have used to determine the FES of species, combined with a proposed set of rules for interpreting mixed case reports (Table 2; Figure 1).

The Universal Edibility Test is the only published method for determining which species can be safely eaten (Survival Use of Plants, 2020). There are no formal laboratory standards for establishing if a species is edible. Known toxins can be detected in mushroom species by chemical analyses (Bever et al., 2020; Bresinsky & Besl, 1990), although their presence does not preclude consumption. Cooking and other pretreatment help to destroy and eliminate poisons present in certain species of raw mushrooms, as demonstrated for *Amanita muscaria* and discussed more widely for other species by Rubel and Arora (2008).

Our review of mushroom properties has highlighted the reliance on subjective, often anecdotal, sources of information on the edibility of a mushroom, and is one possible reason why case reports differ. Our proposed classification system allows researchers and practitioners to weigh up the available evidence for a species, and then categorize the species according to a set of rules (Table 3, Figure 1). This classification system enables users to easily understand the certainty with which a mushroom can be considered edible and provides a verifiable chain of evidence for decisions made on the FES.

This review of edible mushrooms is a work in progress, and our list of edible species and related judgments on FES are likely to change and be updated as new evidence is presented. One of the challenges we faced when determining the FES of a species was the limited number of case reports showing direct evidence of consumption. We hope that in the future, this situation will be improved upon and that subsequent judgements will be more robust as a result. The classification system will need further testing and discussions to ensure that it is fit for purpose and to encourage wider adoption of a standard method for confirming and categorizing the edibility of species and the strength of the supporting evidence.

We have identified 2,189 edible mushroom species (FES = E1 or E2), a slight increase in the 2,166 edible species reported by Boa (2004), despite the greater number of species analyzed and case reports used in the current study. However, we have also identified 471 species of unknown edibility (E3), of which a substantial proportion could be reclassified once additional information is made available.

With the exception of the E3 group, we speculate that it is unlikely that there are many more edible species of value to be documented. We also acknowledge that limitations of our review based on the literature that was available for this analysis. The majority of species within *Auricularia*, *Retiboletus*, *Sparassis*, and *Termitomyces* were found to be edible, yet there are other species less well known and regions where published information on mushroom properties is limited. It is likely that all species of *Lentinula* and *Pleurotus* are in fact edible but there is no current documentation confirming this. We speculate that all mushroom species from a genus such as *Tuber* are edible, though of varying value by taste.

We estimate that there are about 27,000 mushroom species (Hawksworth, 2001; Cannon et al., 2018), of which our review shows about 10% to be edible. According to the estimates of Hawksworth and Lücking (2017), many unknown mushroom species have yet to be documented, and though we can expect a high number of new edible species to be discovered, their value as food remains to be confirmed. Documenting these new species and

determining the edibility of them will take generations of mycologists, incorporating studies on taxonomy, ecology, ethnomycology, and pharmacology.

5.2 | Distribution of wild edible mushrooms

The number of case reports by species and country varied hugely with a strong bias toward countries with a well-established research community, not only in mycology but also forestry and allied areas where mushrooms are of economic importance. For example, over 60% of the data were retrieved from just eight countries in Asia, Europe, and North America: China (26.8%), Japan (7.4%), Germany (4.6%), Mexico (6.7%), Russia (4.2%), the UK (4.1%), India (3.5%), South Korea (3.2%), the USA (3.1%), and Italy (2.4%). This disproportionate representation does not necessarily suggest that these countries have a high diversity of edible mushrooms, but mainly that they have a strong culture surrounding their use, documentation, and research.

There were many more reports from North America and Europe, for example, compared with Africa, even though the consumption of wild mushrooms is common and widespread in large areas of the African continent (Härkönen, Niemelä, & Mwasumbi, 2003; Härkönen, Niemelä, Mbindo, Kotiranta, & Pearce, 2015; Rammeloo & Walley, 1993). China has seen an explosion of publications on mushrooms, with a strong emphasis on their useful properties to people (Wang et al., 2004; Wu et al., 2019). Mexico is another notable source of information on edible mushrooms (Garibay-Orijel et al., 2020). As a result of this bias, there are gaps in our data, for example, we present very little information on edible mushrooms consumed in Eastern Europe. However, we remain confident that the data captured on the edible properties of mushroom species are representative of all the major genera, though it is also clear that stronger evidence is needed to support decisions concerning the FES of many species.

5.3 | Food safety and regional variation in opinions on the edibility of wild mushrooms

There is ongoing debate and a degree of uncertainty regarding the edibility of a number of mushroom species, most of which are classified as E2 in our results. These are species that require a degree of preparation before being considered safe to eat and often the opinions regarding edibility differ regionally. For example, *Gyromitra esculenta*, known as the false morel, is commonly eaten in many countries, including Poland, Lithuania, Estonia, Fin-

land, and Sweden, where it is sold in cans under the brand name Fammarp (https://www.natmat.se/produkt/murklor-190-g-fammarp). Elsewhere, for example, in Italy and Spain, field guides consistently advise that *G. esculenta* is poisonous and that it should be avoided at all costs. *Gyromitra* species contain toxic compounds such as Gyromitrin and poisoning incidents do occur (Leathem & Dorran, 2007; Michelot & Toth, 1991; White et al., 2019). Another example of a mushroom that is widely considered toxic in some regions and edible in others is *Chlorophyllum molybdites*. This species was responsible for more poisoning incidents in China in 2019 than any other species of mushroom (Li et al., 2020). Numerous other sources confirm that this mushroom is poisonous and should not be eaten (Bijeesh, Vrinda, & Pradeep, 2017; Lehmann & Khazan, 1992; Meijer, Amazonas, Rubio, & Curial, 2007), yet a wider literature search reveals that *C. molybdites* is consumed in nine states of Mexico (Villarreal & Perez-Moreno, 1989a). Rammeloo and Walley (1993), in their review of edible mushrooms in Africa, found case reports of *C. molybdites* being eaten in Madagascar, DR Congo, Republic of Congo, and Nigeria. Although there is no doubt that mushrooms such as *G. esculenta* and *C. molybdites* are toxic, these mushrooms can be, and are, eaten following thorough and careful cooking (Jarvinen, Kosonen, & Joutjarvi, 2004). Humans have evolved to eat cooked food, and few mushrooms are eaten raw; thus, when considering the edibility of mushrooms, greater emphasis should be given to the properties of cooked specimens.

Opinions on edibility vary with region; species that are prized in some cultures may not be sought after in others. The regional preferences for *Gyromitra esculenta* have already been discussed above; other notable examples include *Schizophyllum commune*, *Phallus indusiatus* and *Tricholoma equestre*. *Schizophyllum commune* is a popular edible mushroom in China and India, yet few consider it as an edible in the West. Similarly, *P. indusiatus*, also known as the Long net stinkhorn, is sought after for soups in Asia, whereas yet many in the West consider it inedible. *Tricholoma equestre* remains a controversial choice as an edible mushroom, a study conducted by Bedry et al. (2001) reported that this species contains numerous toxins and is considered as potentially deadly to humans (Bedry et al., 2001); however, recent research has found no evidence that this mushroom is in fact toxic (Rzymiski & Klimaszuk, 2018). As a result of this conflicting evidence, *T. equestre* is not eaten in many places; for example, it is listed as a poisonous mushroom in Spain, Italy, and France (Rzymiski, Klimaszuk, & Benjamin, 2019). However, it remains sought after in parts of Asia and Europe. Although the debate on this species' edibility continues, our data support it as

an edible species, though to be treated with a degree of caution.

6 | CONCLUSION AND FUTURE TRENDS

A rising awareness of the health benefits associated with eating mushrooms, and the recognition of mushrooms as a functional food, have resulted in an unprecedented increase in the consumption of mushrooms. Wild mushroom consumption, as well as other wild foods, is not without risks, and as such, public safety and awareness form an important part in managing the collection of wild mushrooms. Unfortunately, inaccurate taxonomy, lack of knowledge, or simply personal opinions have resulted in numerous conflicting reports on which mushrooms are edible and which are not safe to eat, creating a need for consensus of how to determine edibility and which species are in fact safe to eat. Our work proposes a standard method for addressing these issues. We recommend that all future reports should clearly indicate the source of evidence to support their assessments. An edible mushroom is one that has been safely consumed by a known person or population; however, some form of documentation is required to carry this information to a broader audience.

An edible mushroom is one that is safe and suitable to eat. Opinions differ on taste, texture, and smell, and it is important to separate subjective judgments on what is worthwhile to consume from an objective assessment of whether a mushroom can be consumed without adverse effects.

ACKNOWLEDGMENT

This work was supported by Yunnan Provincial Key Programs of Yunnan Eco-friendly Food International Cooperation Research Center Project under grant 2019ZG00908 and Key Research Program of Frontier Sciences "Response of Asian mountain ecosystems to global change," CAS, Grant No. QYZDY-SSW-SMC014." Peter E. Mortimer thanks the National Science Foundation of China for financial support under the following grants: 41761144055 and 41771063. Samantha C. Karunarathna would like to thank the CAS President's International Fellowship Initiative (PIFI) young staff under grant number 2020FYC0002 and the National Science Foundation of China (NSFC) project code 31750110478 for funding this work. Austin Smith at World Agroforestry (ICRAF), Kunming Institute of Botany, China, is thanked for English editing. Nelson Menolli Jr. thanks the São Paulo Research Foundation (FAPESP) grant #2018/15677-0; Massimo Tasselli for helping with some references on edibility of *Russula* species; Giuliana Furci for helping with some additional data from

Chile; as well as Altielys C. Magnago, Bárbara L.B. Schüemann, Larissa Trierweiler-Pereira, Marcelo A. Sulzbacher, Mariana P. Drewinski, Melissa Palacio, Rosana Maziero, Tatiana B. Gibertoni, and Vitor X. de Lima for helping with additional data from Brazil. We would also like to thank Miriam de Román for her thoughts and contributions towards developing the classification system.

AUTHOR CONTRIBUTIONS

Huili Li, Peter E. Mortimer, and Eric Boa designed and drafted the review; Nelson Menolli Jr, Lei Ye, Samantha C. Karunarathna, Jesus Perez-Moreno, Mohammad Mahmudur Rahman, Md Harunur Rashid, Pheng Phengsintham, Leela Rizal, Taiga Kasuya, Young Woon Lim, Arun Kumar Dutta, Abdul Nasir Khalid, Le Thanh Huyen, Marilen Parungao Balolong, Gautam Baruah, Sumedha Madawala, and Naritsada Thongklang contributed to the global data of wild edible mushroom. Paul M. Kirk contributed to the classification of wild edible mushroom. All authors contributed to the writing of the manuscript. The main revisions were carried out by Huili Li, Eric Boa, and Peter E. Mortimer; Nelson Menolli Jr, Kevin D. Hyde, Jianchu Xu, and Jun Sheng were involved in final review and completion of the paper.

ORCID

Huili Li  <https://orcid.org/0000-0002-6229-3780>

REFERENCES

- Acar, İ., Blando, F., Gul, B., Greco, A., Mukemre, M., Uzun, Y., & Dalar, A. (2020). The phenolic profile and biological activities of the wild-edible mushrooms *Helvella leucopus* and *Morchella pulchella*. *Journal of Food Measurement and Characterization*, 1–12. <https://doi.org/10.1007/s11694-020-00654-3>
- Acharya, K., Das, K., Paloi, S., Dutta, A. K., Hembrom, M. E., Khatua, S., & Parihar, A. (2017). Exploring a novel edible mushroom *Ramaria subalpina*: Chemical characterization and antioxidant activity. *Pharmacognosy Journal*, 9, 30–34.
- Acharya, K., Ghosh, S., & Kundu, I. (2016). Pharmacognostic standardization of a well-known edible mushroom, *Volvariella volvacea*. *Journal of Applied Pharmaceutical Science*, 6, 185–190.
- Adhikari, K. S., & Adhikari, M. K. (1996). Collection and consumption of wild edible mushrooms sold in Kathmandu valley, Nepal. *The Geographer's Point*, 1, 1–9.
- Adhikari, M. K. (1999). Wild relatives of some arable mushrooms found in Nepal. *National conference on wild relatives of cultivated plants in Nepal* (pp. 149–155). Kathmandu: Green Energy Mission.
- Adhikari, M. K. (2000). *Mushrooms of Nepal*. Kathmandu: Privately published.
- Adhikari, M. K., & Durrieu, G. (1996). Ethnomycologie Nepalaïse. *Bulletin Société Mycologique de France*, 112, 31–41.
- Afiukwa, C. A., Ebem, E. C., & Igwe, D. O. (2015). Characterization of the proximate and amino acid composition of edible wild mushroom species in Abakaliki, Nigeria. *AASCIT J. Biosci*, 1, 20–25.
- Afonkin, S. U. (2013). *Grand illustrated encyclopedia. Mushrooms of Russia*. Vilnius: Bestiary.

- Afyon, A. (1997). Macrofungi of Seydisehir district (Konya). *Turkish Journal of Botany*, 21, 173–176.
- Aguilar-Cruz, Y., & Villegas, M. (2010). Especies de Gomphales comestibles en el municipio de Villa del Carbón Estado de México. *Revista Mexicana de Micología*, 31, 1–8.
- Ahmad, S. (1956). Fungi of West Pakistan. *Monograph*, 1, 40–56.
- Ahmad, S., Iqbal, S., & Khalid, A. (1997). *Fungi of Pakistan*. Lahore, Pakistan: Sultan Ahmad Mycological Society of Pakistan. Department of Botany, University of the Punjab.
- Ajith, T. A., & Janardhanan, K. K. (2007). Indian medicinal mushrooms as a source of antioxidant and antitumor agents. *Journal of Clinical Biochemistry and Nutrition*, 40, 157–162.
- Akavia, E., Beharav, A., Wasser, S., & Nevo, E. (2009). Disposal of agro-industrial by-products by organic cultivation of the culinary and medicinal mushroom *Hypsizygus marmoreus*. *Waste Management*, 29, 1622–1627.
- Akul, H., Sevindik, M., Coban, C., Alli, H., & Selamoglu, Z. (2017). New approaches in traditional and complementary alternative medicine practices: *Auricularia auricula* and *Trametes versicolor*. *J Tradit Med Clin Natur*, 6, 1–4.
- Akyuz, M., & Kirbag, S. (2009). Antimicrobial activity of *Pleurotus eryngii* var. *ferulae* grown on various agro-wastes. *EurAsian Journal of BioSciences*, 3, 58–63.
- Alanne, A.-L., Issakainen, J., Pihlaja, K., Jokioja, J., & Sinkkonen, J. (2019). Metabolomic discrimination of the edible mushrooms *Kuehneromyces mutabilis* and *Hypholoma capnoides* (Strophariaceae, Agaricales) by NMR spectroscopy. *Zeitschrift für Naturforschung C*, 74, 201–210.
- Al-Naama, N. M., Ewaze, J. O., & Nema, J. H. (1988). Chemical constituents of Iraqi truffles. *Iraqi Journal of Agricultural Sciences*, 6, 51–56.
- Alofe, F. V., Odeyemi, O., & Oke, O. L. (1996). Three edible wild mushrooms from Nigeria: Their proximate and mineral composition. *Plant Foods for Human Nutrition*, 49, 63–73.
- Alsheikh, A. M., & Trappe, J. M. (1983). Desert truffles: The genus *Tirmania*. *Transactions of the British Mycological Society*, 81, 83–90.
- Álvarez-Farias, Z. J., Diaz-Godinez, G., Téllez-Téllez, M., Villegas, E., & Acosta-Urdapilleta, M. L. (2016). Ethnomycological knowledge of wild edible mushrooms in Tlayacapan, Morelos. *Mycosphere*, 7(10), 1491–1499.
- Ammirati, J. F. (1985). *Poisonous mushrooms of the northern United States and Canada*. Minneapolis, MN: University of Minnesota Press.
- Anand, J. S., & Chwaluk, P. (2010). Acute intoxication with *Tricholoma Equestre*—clinical course. *Przegląd Lekarski*, 67, 617–618.
- Anand, J. S., Chwaluk, P., & Sut, M. (2009). Acute poisoning with *Tricholoma equestre*. *Przegląd Lekarski*, 66, 339–340.
- Angajchariya, A., Naranong, S., Phichairat, D., Kaewsongsang, L., Phupong, W., & Mahae, N. (2017). Mycelial growth, antioxidant and antibacterial properties of *Boletus griseipurpureus* from south Thailand. *International Journal of Agricultural Technology*, 13, 521–529.
- Antonin, V., & Fraiture, A. (1998). *Marasmius heinemannianus*, a new edible species from Benin, West Africa. *Belgian Journal of Botany*, 131, 127–132.
- Ao, T., Seb, J., Ajungla, T., & Deb, C. R. (2016). Diversity of wild mushrooms in Nagaland, India. *Open Journal of Forestry*, 6, 404–419.
- Aphshahana, K., & Sharma, A. (2018). Wild edible mushrooms traded by ethnic communities of Senapati and Kangpokpi district of Manipur, India. *Journal of Pharmacognosy and Phytochemistry*, 7, 2303–2306.
- Arlukowicz-Grabowska, M., Wójcicki, M., Raszeja-Wyszomirska, J., Szydłowska-Jakimiuk, M., Piotuch, B., & Milkiewicz, P. (2019). Acute liver injury, acute liver failure and acute on chronic liver failure: A clinical spectrum of poisoning due to *Gyromitra esculenta*. *Annals of hepatology*, 18, 514–516.
- Aroche, R. M., Cifuentes, J., Lorea, F., Fuentes, P., Bonavides, J., Galicia, H., ... Valenzuela, V. (1984). Poisonous and edible macromycetes in a communal region of the Valle de Mexico, I. *Boletín de la Sociedad Mexicana de Micología*, 19, 291–318.
- Arora, D. (1986). *Mushrooms demystified*. Berkeley, CA: Ten Speed Press.
- Arora, D. (1991). All that the rain promises and more. *A hip pocket guide to Western mushrooms*. Berkeley, CA: Ten Speed Press.
- Aryal, H. P., & Budhathoki, U. (2013). Mycodiversity at Sankarnagar Community Forest, Rupandehi District. *Nepal Journal of Science and Technology*, 14, 75–80.
- Aslim, B., & Ozturk, S. (2011). Phenolic composition and antimicrobial and antioxidant activities of *Leucoagaricus leucothites* (Vittad.) Wasser. *Journal of Medicinal Food*, 14, 1419–1424.
- Avci, E., Çağatay, G., Alp Avci, G., Suiçmez, M., & Coşkun Cevher, Ş. (2016). An edible mushroom with medicinal significance, *Auricularia polytricha*. *Hittite Journal of Science and Engineering*, 3, 111–116.
- Azeem, U., Hakeem, K. R., & Ali, M. (2020). Toxigenic fungi. In *Fungi for human health* (pp. 41–48). Cham, Switzerland: Springer International Publishing.
- Badalyan, S. M. (2003). Edible and medicinal higher basidiomycetes mushrooms as a source of natural antioxidants. *International Journal of Medicinal Mushrooms*, 5, 153–162.
- Bal, A., Anil, M., Yilmaz, I., Akata, I., & Atilla, O. D. (2016). An outbreak of non-fatal mushroom poisoning with *Omphalotus olearius* among Syrian refugees in Izmir, Turkey. *Toxin Reviews*, 35, 1–3.
- Bandala, V. M., Montoya, L., Villegas, R., Cabrera, T. G., de Jesús Gutiérrez, M., & Acero, T. (2014). “Nangañaña” (*Tremelloscypha gelatinosa*, Sebacinaceae), hongo silvestre comestible del bosque tropical deciduo en la depresión central de Chiapas, México. *Acta Botanica Mexicana*, 106, 149–159.
- Bandara, A. R., Karunarathna, S. C., Mortimer, . E., Hyde, K. D., Khan, S., Kakumyan, P., & Xu, J. (2017). First successful domestication and determination of nutritional and antioxidant properties of the red ear mushroom *Auricularia thailandica* (Auriculariales, Basidiomycota). *Mycological Progress*, 16, 1029–1039.
- Barroetaveña, C., & Toledo, C. (2019). Diversity and ecology of edible wild mushrooms from native forests of Patagonia, Argentina. In J. Pérez-Moreno, A. Guerin-Laguette, R. Flores-Arzù, & F. Q. Yu (Eds.), *Mushrooms, humans and nature in a changing world: Perspectives from ecological, agricultural and social sciences* (pp. 297–318). Cham, Switzerland: Springer International Publishing.
- Barros, L., Baptista, P., Correia, D. M., Casal, S., Oliveira, B., & Ferreira, I. C. (2007). Fatty acid and sugar compositions, and nutritional value of five wild edible mushrooms from Northeast Portugal. *Food Chemistry*, 105, 140–145.

- Basumatary, M., & Gogoi, M. (2016). Uses of wild edible macro fungi by Bodo community of Kokrajhar district, Assam, India. *Tropical Plant Research*, 3, 176–181.
- Batra, L. R. (1983). Edible discomycetes and gasteromycetes of Afghanistan, Pakistan and north-western India. *Biologia (Lahore)*, 29, 293–304.
- Bautista-Nava, E., & Moreno-Fuentes, Á. (2009). Primer registro de *Calostoma cinnabarina* (Sclerodermatales) como especie comestible. *Revista mexicana de biodiversidad*, 80, 561–564.
- Beaumier, M., Rioult, J.-P., Georges, M., Brocheriou, I., Lobbedez, T., & Lanot, A. (2019). Mushroom poisoning presenting with acute kidney injury and elevated transaminases. *Kidney International Reports*, 4, 877–881.
- Becker, U., Erkel, G., Anke, T., & Sterner, O. (1997). Puraquinonic acid, a novel inducer of differentiation of human HL-60 promyelocytic leukemia cells from *Mycena pura* (Pers. ex Fr.). *Natural Product Letters*, 9, 229–236.
- Bedry, R., Baudrimont, I., Deffieux, G., Creppy, E. E., Pomies, J. P., Ragnaud, J. M., ... Godeau, P. (2001). Wild mushroom intoxication as a cause of rhabdomyolysis. *New England Journal of Medicine*, 345, 798–802.
- Berger, K. J., & Guss, D. A. (2005). Mycotoxins revisited: Part II. *The Journal of Emergency Medicine*, 28, 175–183.
- Beug, M. W. (2017). Amatoxin Mushroom Poisoning in North America 2015–2016. *The Mycophile*, 57, 1–24.
- Bever, C. S., Swanson, K. D., Hamelin, E. I., Filigenzi, M., Poppenga, R. H., Kaae, J., ... Stanker, L. H. (2020). Rapid, sensitive, and accurate point-of-care detection of lethal amatoxins in urine. *Toxins*, 12, 1–10.
- Bhatt, R., Singh, U., & Stephenson, S. (2016). Wild edible mushrooms from high elevations in the Garhwal Himalaya-I. *Current Research in Environmental & Applied Mycology*, 6, 118–131.
- Bi, H., Gao, T., Li, Z., Ji, L., Yang, W., Iteku, B. J., ... Zhou, Y. (2013). Structural elucidation and antioxidant activity of a water-soluble polysaccharide from the fruit bodies of *Bulgaria inquinans* (Fries). *Food Chemistry*, 138, 1470–1475.
- Bijeesh, C., Vrinda, K., & Pradeep, C. (2017). Mushroom poisoning by *Chlorophyllum molybdites* in Kerala. *Journal of Mycopathological Research*, 54, 477–483.
- Birks, A. A. (1991). Fungi in folk medicine. *McIlvainea*, 10, 89–94.
- Boa, E. R. (2004). *Wild edible fungi: A global overview of their use and importance to people*. Rome: Food and Agriculture Organization of The United Nations.
- Boa, E. R. (2011). From Chipho to Msika: An introduction to mushrooms, trees and forests. In A. B. Cunningham & X. Yang (Eds.), *Mushrooms in forests and woodlands* (pp. 19–38). Routledge.
- Boa, E. R., Ngulube, M., Meke, G., & Munthali, C. (2000). *First regional workshop on sustainable use of forest products: Miombo wild edible fungi* (pp. 61). Zomba, Malawi: Forest Research Institute of Malawi and CABI Bioscience.
- Bokhary, H. A., & Parvez, S. (1993). Chemical composition of desert truffles *Terfezia clavaryi*. *Journal of Food Composition and Analysis*, 6, 285–293.
- Bononi, V. L. (1984). Basidiomicetos do Parque Estadual da Ilha do Cardoso. IV. Adições às famílias Hymenochaetaceae, Stereaceae e Thelephoraceae. *Rickia*, 11, 43–52.
- Bononi, V. L., Capelari, M., Maziero, R., & Trufem, S. (1995). *Cultivo de cogumelos comestíveis*. Ícone.
- Boruah, P., Adhikary, R. K., Kalita, P., & Bordoloi, D. (1996). Some edible fungi growing in the forest of East Khasi Hills (Meghalaya). *Advances in Forestry Research in India*, 14, 214–219.
- Bofaris, M. S. M., Alzand, K., Ünal, S., Karadeniz, M., & Bartouh, M. (2019). Trace elements concentrations in Turkey species of wild growing edible mushrooms: a review. *World Journal of Pharmaceutical Research*, 8, 47–62.
- Bouriquet, G. (1943). Notes de mycologie malgache. *Bulletin Academie Malgache*, 25, 12–24.
- Bouriquet, G. (1970). Les principaux champignons de Madagascar. *Terre Malgache*, 7, 10–37.
- Braeuer, S., Goessler, W., Kameník, J., Konvalinková, T., Žigová, A., & Borovička, J. (2018). Arsenic hyperaccumulation and speciation in the edible ink stain bolete (*Cyanoboletus pulverulentus*). *Food chemistry*, 242, 225–231.
- Bresinsky, A., & Besl, H. (1990). *A colour atlas of poisonous fungi*. London: Wolfe.
- Brown, M. (2019). Yi ethnomycology: Wild mushroom knowledge and use in Yunnan, China. *Journal of Ethnobiology*, 39, 131–157.
- Brown, J., Malone, M., Stuntz, D., & Tyler Jr, V. (1962). Paper chromatographic determination of muscarine in *Inocybe* species. *Journal of Pharmaceutical Sciences*, 51, 853–856.
- Brvar, M., Možina, M., & Bunc, M. (2006). Prolonged psychosis after *Amanita muscaria* ingestion. *Wiener klinische wochenschrift*, 118, 294–297.
- Büchel, E., Martini, U., Mayer, A., Anke, H., & Sterner, O. (1998). Omphalotins B, C and D, nematicidal cyclopeptides from *Omphalotus olearius*. Absolute configuration of omphalotin A. *Tetrahedron*, 54, 5345–5352.
- Burkhill, I. H. (1935). *A dictionary of the economic products of the Malay Peninsula* (Vol. 2). London: Crown Agents for the Colonies.
- Burrola-Aguilar, C., Garibay-Orijel, R., & Hernández Téllez, M. (2012). Los hongos comestibles silvestres del estado de México: Propuesta para su aprovechamiento. In J. Sánchez & G. Mata (Eds.), *Hongos comestibles y medicinales en Iberoamerica, investigación y desarrollo en un entorno multicultural* (pp. 39–49). El Colegio de la Frontera Sur, Chiapas.
- Butkhup, L., Samappito, W., & Jorjong, S. (2018). Evaluation of bioactivities and phenolic contents of wild edible mushrooms from northeastern Thailand. *Food Science and Biotechnology*, 27, 193–202.
- Buyck, B. (1994a). Ectotrophy in tropical African ecosystems. In J. H. Seyani & A. C. Chikuni (Eds.), *Proceedings of the XIIIth Plenary meeting of AETFAT, Zomba, Malawi* (pp. 705–718). 2–11 April 1991. Zomba: National Herbarium and Botanic Gardens of Malawi.
- Buyck, B. (1994b). *Ubwoba: Les Champignons Comestibles de l'Ouest du Burundi*. Bruxelles: Administration Generale de la Cooperation au Developpement.
- Buyck, B. (2008). The edible mushrooms of Madagascar: An evolving enigma. *Economic Botany*, 62(3), 509–520.
- Buyck, B., & Nzigidahera, B. (1995). Ethnomycological notes from Burundi. *Belgian Journal of Botany*, 128, 131–138.
- Çağlarlırmak, N., Unal, K., & Otles, S. (2002). Nutritional value of edible wild mushrooms collected from the Black Sea region of Turkey. *Micologia Aplicada Internacional*, 14, 1–5.
- Cai, Q., Cui, Y.-Y., & Yang, Z. L. (2016). Lethal amanita species in China. *Mycologia*, 108, 993–1009.

- Cai, M., Pettenella, D., & Vidale, E. (2011). Income generation from wild mushrooms in marginal rural areas. *Forest Policy and Economics*, 13(3), 221–226.
- Campos, J. (1998). *Principales productos forestales no madereros en Chile*. Santiago, Chile: FAO.
- Cannon, P. F., Aguirre-Hudson, B., Aime, M. C., Bidartondo, M. I., Gaya, E., Hawksworth, D., Kirk, P., Leitch, L. J., & Lücking, R. (2018). *State of the World's Fungi* (pp. 1–92). Royal Botanic Gardens, Kew.
- Cao, H., Qin, D., Guo, H., Cui, X., Wang, S., Wu, Y., ... Yu, J. (2020). The shaggy ink cap medicinal mushroom, *Coprinus comatus* (Agaricomycetes), a versatile functional species: A review. *International Journal of Medicinal Mushrooms*, 22, 245–255.
- Cao, J. (1991). A new wild edible fungus – *Wynnella silvicola*. *Zhongguo Shiyongjun (Edible Fungi of China: A bimonthly journal)*, 10, 27–28.
- Cardoso, D. B. O. S., De Queiroz, L. P., Bandeira, F. P., & Góes-Neto, A. (2010). Correlations between indigenous Brazilian folk classifications of fungi and their systematics. *Journal of Ethnobiology*, 30, 252–264.
- Carvalho, L. M., Carvalho, F., de Lourdes, Bastos, M., Baptista, P., Moreira, N., ... de Pinho, P. G. (2014). Non-targeted and targeted analysis of wild toxic and edible mushrooms using gas chromatography–ion trap mass spectrometry. *Talanta*, 118, 292–303.
- Cervera, M., & Colinas, C. (1997). Comercialización de seta silvestre en la ciudad de Lleida. In F. Puertas & M. Rivas (Eds.), *Actas del I Congreso Forestal Hispano LUSo, II Congreso Forestal Espanol-IRATI 97* (425–429). Pamplona.
- Chamberlain, M. (1996). Ethnomycological experiences in South West China. *Mycologist*, 10, 13–16.
- Chandra, A. (1989). *Elsevier's dictionary of edible mushrooms. Botanical and common names in various languages of the world*. Amsterdam, Netherlands: Elsevier.
- Chang, S.-T., & Hayes, W. A. (1978). *The biology and cultivation of edible mushrooms*. New York: Academic Press.
- Chang, S.-T. (2006). The world mushroom industry: Trends and technological development. *International Journal of Medicinal Mushrooms*, 8(4), 297–314.
- Chang, S.-T., & Mao, X. L. (1995). *Hong Kong mushrooms*. Hong Kong: The Chinese University of Hong Kong Press.
- Chang, S.-T., & Miles, P. G. (2004). *Mushrooms: Cultivation, nutritional value, medicinal effect and environmental impact* (2nd ed.). Boca Raton, FL: CRC Press.
- Chen, X.-N., Zhang, M., Li, T., & Zeng, N.-K. (2019). A new species of *Heimioporus* (Boletaceae) from southern China. *Phytotaxa*, 415, 179–188.
- Chen, Z. (1987). Domesticating cultivation of wild *Pleurotus sapidus* in Xinjiang. *Zhongguo Shiyongjun (Edible Fungi of China)*, 6(25), 19.
- Chen, Z., Ping, Z., & Zhang, Z. (2014). Investigation and analysis of 102 mushroom poisoning cases in Southern China from 1994 to 2012. *Fungal Diversity*, 64, 123–131.
- Chen, Z., Yang, Z., Bau, T., & Li, T. (2016). *Poisonous mushrooms: Recognition and poisoning treatment*. Beijing: Science Press.
- Cheung, P. C. (1996). The hypocholesterolemic effect of two edible mushrooms: *Auricularia auricula* (tree-ear) and *Tremella fuciformis* (white jelly-leaf) in hypercholesterolemic rats. *Nutrition Research*, 16, 1721–1725.
- Chevalier, G., & Frochot, H. (1990). Ecology and possibility of culture in Europe of the Burgundy truffle (*Tuber uncinatum* Chatin). *Agriculture, Ecosystems & Environment*, 28, 71–73.
- Chodorowski, Z., Anand, J. S., & Grass, M. (2003). Acute poisoning with *Tricholoma equestre* of five-year old child. *Przegląd lekarski*, 60, 309–310.
- Chodorowski, Z., Waldman, W., & Sein Anand, J. (2002). Acute poisoning with *Tricholoma equestre*. *Przegląd lekarski*, 59, 386–387.
- Chong, E., Sia, C., Chang, S., Yim, H., & Khoo, H. (2014). Antioxidative properties of an extract of *Hygrocybe conica*, a wild edible mushroom. *Malaysian Journal of Nutrition*, 20, 101–111.
- Christensen, M., Bhattarai, S., Devkota, S., & Larsen, H. O. (2008). Collection and use of wild edible fungi in Nepal. *Economic Botany*, 62, 12–23.
- Chu, T. N. (2018). *Study on biodiversity of Pluteaceae in Cuc Phuong National Park, Vietnam*. University of Natural Resources and Environment.
- Chwaluk, P. (2010). Acute intoxication with *Tricholoma equestre*—clinical course. *Przegląd Lekarski*, 67, 617–618.
- Clemente, R. F., Liwanag, J. M., Santos, E. E., Flores, F. R., & Dulay, R. M. (2017). Species listing of macrofungi in Angat Watershed Reservation, Bulacan Province, Luzon Island, Philippines. *International Journal of biology, Pharmacy, and Allied Sciences*, 6, 1060–1068.
- Colavolpe, M. B., & Albertó, E. (2014). Cultivation requirements and substrate degradation of the edible mushroom *Gymnopilus pampeanus*—A novel species for mushroom cultivation. *Scientia Horticulturae*, 180, 161–166.
- Comandini, O., & Rinaldi, A. C. (2020). Ethnomycology in Europe: The past, the present, and the future. In J. Perez-Moreno, A. Guerin-Laguette, R. Flores Arzu, & F. Q. Yu (Eds.), *Mushrooms, humans and nature in a changing world* (pp. 341–364). Cham, Switzerland: Springer International Publishing.
- Consiglio, G., & Papetti, C. (2009). *Atlante fotografico dei Funghi d'Italia*. Trento: Associazione Micologica Bresadola.
- Cooke, M. C. (1891). *British edible fungi*. Tench, Trubner: Kegan Paul.
- Crowe, A. (2016). *A field guide to the native edible plants of New Zealand*. Auckland, New Zealand: Penguin Books.
- Dabrowski, W. M., & Sikorski, Z. E. (2004). *Toxins in food*. CRC Press.
- Dai, Y. C., & Xing-Liang, W. U. (2004). A new edible Polypore—*Bondarzewia berkeleyi*. *Edible Fungi of China*, 23, 3–4.
- Dai, Y.-C., Yang, Z.-L., Cui, B.-K., Yu, C.-J., & Zhou, L.-W. (2009). Species diversity and utilization of medicinal mushrooms and fungi in China (review). *International Journal of Medicinal Mushrooms*, 11, 287–302.
- Das, K. (2009). *Mushrooms of Sikkim I: Barsey Rhododendron Sanctuary, Sikkim State Biodiversity Board, Gangtok and Botanical Survey of India, Kolkata, India*.
- Das, K., & Chakraborty, D. (2014). *Lactarius vesterholtii*, a new species from India. *Mycotaxon*, 129, 477–484.
- Das, N., Chowdhury, P., & Pisman, B. (2010). Cultivation practice of *Pleurotus fossulatus* on rice straw. *Journal of Life Sciences*, 4, 20–25.
- de Avila, B. A., Welden, A. L., & Guzmán, G. (1980). Notes on the ethnomycology of Hueyapan, Morelos, Mexico. *Journal of Ethnopharmacology*, 2(4), 311–321.
- De-Kesel, A., Kasongo, B., & Degreef, J. (2017). *Champignons comestibles du Haut-Katanga (RD Congo)*. Royal Belgian Institute of Natural Sciences.

- De Leon, A., Kalaw, S., Dulay, R., Undan, J., Alfonso, D., Undan, J., & Reyes, R. (2016). Ethnomycological survey of the Kalanguya indigenous community in Caranglan, Nueva Ecija, Philippines. *Current Research in Environmental & Applied Mycology*, 6, 61–66.
- De Leon, A., Luangsa-Ard, J., Karunarathna, S., Hyde, K., Reyes, R., & Dela-Cruz, T. (2013). Species listing, distribution, and molecular identification of macrofungi in six Aeta tribal communities in Central Luzon, Philippines. *Mycosphere*, 4, 478–494.
- De Leon, A., Reyes, R., & dela Cruz, T. (2012). An ethnomycological survey of macrofungi utilized by Aeta communities in Central Luzon, Philippines. *Mycosphere*, 3, 251–259.
- de Oliveira, P. (2009). Mushroom poisoning. *Medicina Interna*, 16, 232–238.
- Degreef, J., Malaisse, F., Rammeloo, J., & Baudart, E. (1997). Edible mushrooms of the Zambezi woodland area: A nutritional and ecological approach. *BASE (Biotechnologi, Agronomie, Societe et Environnement)*, 1, 221–231.
- De-Kesel, A. (2002). *Wild edible fungi from Benin*. Personal Communication.
- Demirbas, A. (2000). Accumulation of heavy metals in some edible mushrooms from Turkey. *Food Chemistry*, 68, 415–419.
- Demirel, K., Kaya, A., & Uzun, Y. (2003). Macrofungi of Erzurum Province. *Turkish Journal of Botany*, 27, 29–36.
- Denchev, C. (2002). *Wild edible fungi in Bulgaria*. Personal Communication.
- de-Román, M., & Boa, E. (2006). The marketing of *Lactarius deliciosus* in Spain. *Economic Botany*, 60, 284–290.
- Deschamps, J. R. (2002). *Hongos silvestres comestibles del Mercosur con valor gastronómico: Documentos de trabajo*. No. 86. Argentina: Universidad de Belgrano.
- Diamandis, S. (2002). *Greek edible fungi*. Personal Communication.
- Diaz, J. H. (2016). Mistaken mushroom poisonings. *Wilderness & Environmental Medicine*, 27, 330–335.
- Díaz-Talamantes, C., Burrola-Aguilar, C., Aguilar-Miguel, X., & Mata, G. (2017). In vitro mycelial growth of wild edible mushrooms from the central Mexican highlands. *Revista Chapingo Serie Ciencias Forestales y del Ambiente*, 23, 3.
- Diez, C. (2003). *Collecting Boletus edulis for commercial purposes in Peru*. Personal Communication.
- Dimitrijevic, M. V., Mitic, V. D., Nikolic, J. S., Djordjevic, A. S., Mutic, J. J., Stankov Jovanovic, V. P., & Stojanovic, G. S. (2019). First report about mineral content, fatty acids composition and biological activities of four wild edible mushrooms. *Chemistry & Biodiversity*, 16, e1800492.
- Dong, M., & Shen, A. (1993). Studies on *Lactarius camphoratus*. 1 Biological characteristics of *L. camphoratus*. *Zhongguo Shiyongjun (Edible Fungi of China)*, 12, 3–5.
- Dörfelt, H., Kiet, T., & Berg, A. (2004). Neue Makromyceten-Kollektionen von Vietnam und deren systematische und ökogeographische Bedeutung. *Feddes Repertorium: Zeitschrift für botanische Taxonomie und Geobotanik*, 115, 164–177.
- Doskocil, I., Havlik, J., Verlotta, R., Tauchen, J., Vesela, L., Macakova, K., ... Rada, V. (2016). In vitro immunomodulatory activity, cytotoxicity and chemistry of some central European polypores. *Pharmaceutical Biology*, 54, 2369–2376.
- Doyungan, S. (1990). *Philippine edible mushrooms*. Technician (Philippines).
- Ducouso, M., Ba, A. M., & Thoen, D. (2002). Ectomycorrhizal fungi associated with native and planted tree species in West Africa: A potential source of edible mushrooms. In I. R. Hall, Y. Wang, A. Zambonelli, & E. Danell (Eds.), *Proceedings of the second international conference on edible mycorrhizal mushrooms*. Christchurch: CD-ROM. New Zealand Institute for Crop & Food Research Limited.
- Dutta, A. K., & Acharya, K. (2014). Traditional and ethno-medicinal knowledge of mushrooms in West Bengal, India. *Asian Journal of Pharmaceutical and Clinical Research*, 7, 36–41.
- Egli, S., Peter, M., Buser, C., Stahel, W., & Ayer, F. (2006). Mushroom picking does not impair future harvests – Results of a long-term study in Switzerland. *Biological Conservation*, 129, 271–276.
- El'chibaev, A. A. (1964). *S'edobnye griby Kirgizii [Edible mushrooms of the Kirghiz SSR]*. Kirgizskoi SSR: Izdatel'stvo Akademii Nauk.
- Ellen, R. (2008). Ethnomycology among the Nualu of the Moluccas: Putting Berlin's 'general principles' of ethnobiological classification to the test. *Economic Botany*, 62, 483–496.
- Ellingsen, H. J. (1982). Some gasteromycetes from northern Thailand. *Nordic Journal of Botany*, 2, 283–285.
- Emsen, B., & Guven, B. (2019). Activities of two edible macrofungi, *Coprinus comatus* and *Leucoagaricus leucothites* in human lymphocytes: Cytogenetic and biochemical study. *Plant Biosystems*, 154(3), 361–368.
- Endo, N., Fangfuk, W., Kodaira, M., Sakuma, D., Hadano, E., Hadano, A., ... Fukuda, M. (2017). Reevaluation of Japanese *Amanita* section *Caesareae* species with yellow and brown pileus with descriptions of *Amanita kitamagotake* and *A. chatamagotake* spp. nov. *Mycoscience*, 58, 457–471.
- Endo, N., Fangfuk, W., Sakuma, D., Phosri, C., Matsushita, N., Fukuda, M., & Yamada, A. (2016). Taxonomic consideration of the Japanese red-cap Caesar's mushroom based on morphological and phylogenetic analyses. *Mycoscience*, 57, 200–207.
- Endo, N., Kawamura, F., Kitahara, R., Sakuma, D., Fukuda, M., & Yamada, A. (2014). Synthesis of Japanese *Boletus edulis* ectomycorrhizae with Japanese red pine. *Mycoscience*, 55, 405–416.
- Endo, N., Tokoo, R., Fukuda, M., & Yamada, A. (2018). *Hygrophorus yukishiro* sp. nov., a new vernal edible mushroom from Nagano Prefecture, Japan. *Mycoscience*, 59, 449–454.
- Ereifej, K. I., & Al-Raddad, A. M. (2000). Identification and quality evaluation of two wild mushrooms in relation to *Agaricus bisporus* from Jordan. In L. Van-Griensven (Ed.), *Science and cultivation of edible fungi*. Proceedings of the 15th International Congress on the Science and Cultivation of Edible Fungi, Maastricht, the Netherlands, 15–19 May 2000 (pp. 721–724).
- Ertuğ, F. (2000). An ethnobotanical study in central Anatolia (Turkey). *Economic Botany*, 54, 155.
- Escobar-Aguirre, G., & Toledo-Ascencio, J. (1977). *El "Tenquique", hongo comestible de El Salvador* (pp. 1). Comunicaciones de la Universidad de El Salvador.
- Estrada-Martinez, E., Guzman, G., Tovar, D. C., & Paczka, R. O. (2009). Contribution to ethnomycological knowledge of wild edible mushrooms from regional markets and communities of Sierra Nevada (Mexico). *Interciencia*, 34, 25–33.
- Ezuka, A. (1990). Notes on some species of *Exobasidium* in Japan (1). *Transactions of the Mycological Society of Japan*, 31, 375–388.
- Fangkrathok, N., Junlatat, J., & Sripanidkulchai, B. (2013). In vivo and in vitro anti-inflammatory activity of *Lentinus polychrous* extract. *Journal of Ethnopharmacology*, 147, 631–637.

- Fidalgo, O. (1965). Conhecimento micológico dos índios brasileiros. *Rickia*, 2, 1–10.
- Fidalgo, O. (1968). Conhecimento micológico dos índios brasileiros. *Revista de Antropologia*, 15–16, 27–34.
- Fidalgo, O., & Fidalgo, M. (1970). A poisonous *Ramaria* from Southern Brazil. *Rickia*, 5, 71–91.
- Fidalgo, O., & Hirata, J. M. (1979). Etnomicología Caiabi, Txicao e Txucarramae. *Rickia*, 8, 1–5.
- Fidalgo, O., & Prance, G. T. (1976). The ethnomycology of the Sanama Indians. *Mycologia*, 68, 201–210.
- Fischer, D. W., & Bessette, A. E. (1992). *Edible wild mushrooms of North America*. Austin, TX: University of Texas Press.
- Flores, R. (2002). *Guatemala edible fungi*. Personal Communication.
- Flores Arzú, R., Comandini, O., & Rinaldi, A. (2012). A preliminary checklist of macrofungi of Guatemala, with notes on edibility and traditional knowledge. *Mycosphere*, 3, 1–21.
- Food and Agriculture Organization Statistical. (2017). FAOSTAT—Data-crops-visualized-mushrooms and truffles.
- Franquemont, C., Plowman, T., Franquemont, E., King, S. R., Niezgodá, C., Davis, W., & Sperling, C. R. (1990). The ethnobotany of Chinchero, an Andean community in southern Peru. *Fieldiana*, 24, 1–126.
- Fu, W. J., Wu, G. B., Yang, S. R., Liang, Y. J., & Li, Y. Z. (1998). Research on the soil status of pine mushroom occurring in the Changbai Mountains. *Acta Pedologica Sinica*, 35, 140–143.
- Fu, Z., Wang, J., & Aisa, H. (2016). Determination of chemical composition and antioxidant properties of a novel wild edible mushroom. *Toxicological & Environmental Chemistry*, 98, 439–449.
- Fui, F. S., Saikim, F. H., Kulip, J., & Seelan, J. S. S. (2018). Distribution and ethnomycological knowledge of wild edible mushrooms in Sabah (Northern Borneo), Malaysia. *Journal of Tropical Biology & Conservation (JTBC)*, 15, 203–222.
- Furci, G. (2013). *Guía de campo. Hongos de Chile*. v. 1. Santiago: Fundación Fungi.
- Furci, G. (2018). *Guía de campo. Hongos de Chile*, v. 2. Santiago: Fundación Fungi.
- Furusawa, E., Chou, S., Furusawa, S., Hirazumi, A., & Dang, Y. (1992). Antitumour activity of *Ganoderma lucidum*, an edible mushroom, on intraperitoneally implanted Lewis lung carcinoma in synergetic mice. *Phytotherapy Research*, 6, 300–304.
- Galli, R. (1996). *Le Russule*. Edinatura, Milano.
- Gamboa-Trujillo, J. P., Wartchow, F., Cerón, C., Aules, E., Aigaje, C., Calvalcanti, L., & Gibertoni, T. (2014). Traditional use of *Gymnopus nubicola* as food resource in a Kichwa community, Pichincha, Ecuador. *Mycosphere*, 5, 180–186.
- Gamboa-Trujillo, P., Wartchow, F., Cerón-Martínez, C., Andi, D., Uwinjin, P., Grefa, G., ... García-Aranda, M. (2019). Some edible, toxic and medicinal mushrooms from temperate forests in the north of Mexico. Instituto de Ciencias Biomédicas. In K. R. Sridhar & S. K. Deshmukh (Eds.), *Advances in macrofungi: Diversity, ecology and biotechnology*. Boca Raton, FL: CRC Press.
- Garnet, S. (2003). *Economically important macrofungi in British Columbia: Species descriptions, distribution and commercial status*. In Forest Investment Account (FIA), Forest Science Program. British Columbia: Mycology Resources.
- Gamundí, I. (2002). *Edible fungi collected in Argentina*. Personal Communication.
- Gamundí, I., & Horak, E. (2002). *Fungi of the Andean-Patagonian forests*. Buenos Aires: Vazquez Mazzini Editores.
- Gardezi, H. (2003). Contemporary sociology in Pakistan. In S. A. Zaidi (Ed.), *Social Science in Pakistan in the 1990s* (pp. 103–114). Islamabad, Pakistan: Council of Social Sciences, Pakistan.
- Gardezi, R. A. (1993). Agaric fungi from Rawalakot, Azad Kashmir. *Sarhad Journal of Agriculture*, 8, 225–226.
- Garibay-Orijel, R., & Ruan-Soto, F. (2014). Listado de los hongos silvestres consumidos como alimento tradicional en México. *La etnomicología en México, estado del arte*. México: CONACYT-UAEH-UNAM.
- Garibay-Orijel, R., Argüelles-Moyao, A., Álvarez-Manjarrez, J., Ángeles-Argáiz, R. E., García-Guzmán, O. M., & Hernández-Yáñez, H. (2020). Diversity and importance of edible mushrooms in ectomycorrhizal communities in Mexican neotropics. In J. Pérez-Moreno, A. Guerin-Laguette, R. Flores Arzú, & F. Q. Yu. (Eds.), *Mushrooms, humans and nature in a changing world* (pp. 407–424). Cham, Switzerland: Springer International Publishing.
- Garibay-Orijel, R., Ruan-Soto, F., & Estrada-Martínez, E. (2010). El conocimiento micológico tradicional, motor para el desarrollo del aprovechamiento de los hongos comestibles y medicinales. In D. Martínez-Carrera, N. Curvetto, M. Sobal, P. Morales, & M. Mora (Eds.), *Hacia un desarrollo sostenible del sistema de producción-consumo de los hongos comestibles y medicinales en Latinoamérica: Avances y perspectivas en el siglo XXI*. Red Latinoamérica de hongos comestibles y medicinales. Colegio de Postgraduados and CONACYT. Puebla, Mexico, pp. 243–270.
- Garza-Ocañas, F. (2019). *Some edible, toxic and medicinal mushrooms from temperate forests in the north of Mexico*. Instituto de Ciencias Biomédicas.
- Gennari, A. (2000). *401 Funghi*. Arezzo, Italy: Artigiana Ezechielli.
- Gerhardt, E. (1994). *Przewodnik Grzyby*. Warsaw, Poland: Multico.
- Gerhardt, E. (2001). *Der grosse BLV Pilzfürher für unterwegs*. Munich: BLV.
- Giraldo, L. R. D., Arteaga, J. J. M., & Murillo, W. (2018). Cytotoxic activity of ethanolic extracts of a selection of macromycetes. *Caryologia*, 71, 191–209.
- Góes-Neto, A. (2001–2003). A review of ethnomycology of indigenous people in Brazil and its relevance to the comparative ethnomycological studies in Latin America. *Revista Mexicana de Micología*, 17, 11–16.
- Govorushko, S., Rezaee, R., Duanov, J., & Tsatsakis, A. (2019). Poisoning associated with the use of mushrooms: A review of the global pattern and main characteristics. *Food and Chemical Toxicology*, 128, 267–279.
- Gupta, H. K. (1999). *The morel mushrooms of the northwest Himalayas, India: Collection, trade and experimentation*. 2 page typescript for project proposal with general information.
- Gursoy, N., Sarikurkcu, C., Tepe, B., & Solak, M. H. (2010). Evaluation of antioxidant activities of 3 edible mushrooms: *Ramaria flava* (Schaeef.: Fr.) Quél., *Rhizopogon roseolus* (Corda) TM Fries., and *Russula delica* Fr. *Food Science and Biotechnology*, 19, 691–696.
- Guzmán, G. (1987). Un caso especial de envenenamiento mortal producido por hongos en el Estado de Veracruz. *Revista Mexicana de Micología*, 3, 203–209.
- Guzmán, G. (2001a). Hallucinogenic, medicinal, and edible mushrooms in Mexico and Guatemala: Traditions, myths, and knowledge. *International Journal of Medicinal Mushrooms*, 3, 399–408.

- Guzmán, G. (2001b). Medicinal fungi in Mexico: Traditions, myths and knowledge. *International Journal of Medicinal Mushrooms*, 3, 95.
- Guzmán, G., Medel, R., & Ramírez, F. (2009). Hongos. In G. Ceballos, R. List, G. Garduño, R. López, M. Muñozcano, E. Collado, & J. Eivin (Eds.), *La diversidad biológica del estado de México* (pp. 251–260). Toluca: Estudio de Estado.
- Habtemariam, S. (1996). Cytotoxicity of extracts from the mushroom *Paxillus involutus*. *Toxicon*, 34, 711–713.
- Hadar, Y., & Dosoretz, C. G. (1991). Mushroom mycelium as a potential source of food flavour. *Trends in Food Science & Technology*, 2, 214–218.
- Hall, I. R., Buchanan, P. K., Wang, Y., & Cole, A. L. J. (1998). *Edible and poisonous mushrooms: An introduction*. Christchurch, New Zealand: New Zealand Institute for Crop and Food Research Ltd.
- Hall, I. R., Lyon, A. J. E., Wang, Y., & Sinclair, L. (1998). Ectomycorrhizal fungi with edible fruiting bodies 2. *Boletus edulis*. *Economic Botany*, 52, 44–56.
- Hall, I. R., Stephenson, S., Buchanan, P., Wang, Y., & Cole, A. L. J. (2003). *Edible and poisonous mushrooms of the world*. Portland: Timber Press.
- Hall, I., Lyon, T., Wang, Y., & Buchanan, P. (2007). Truffles and Mushrooms. A list of putative edible or medicinal ectomycorrhizal mushrooms. Truffles & Mushrooms (Consulting) Ltd.
- Härkönen, M. (2002). Mushroom collecting in Tanzania and Hunan (southern China): Inherited wisdom and folklore of two different cultures. In R. Watling, J. C. Frankland, A. M. Ainsworth, S. Isaac, & C. H. Robinson (Eds.), *Tropical mycology. Vol. 1 Macromycetes* (pp. 149–165). Wallingford: CAB International.
- Härkönen, M., Niemelä, T., & Mwasumbi, L. (2003). *Tanzanian mushrooms. Edible, harmful and other fungi*. Helsinki: Botanical Museum, Finnish Museum of Natural History.
- Härkönen, M., Niemelä, T., Mbindo, K., Kotiranta, H., & Pearce, G. D. (2015). *Zambian mushrooms and mycology: Finnish museum of natural history*. Helsinki: University of Helsinki.
- Härkönen, M., Saarimäki, T., & Mwasumbi, L. (1994a). Edible and poisonous mushrooms of Tanzania. *The African Journal of Mycology and Biotechnology*, 2, 99–123.
- Härkönen, M., Saarimäki, T., & Mwasumbi, L. (1994b). Setting up a research project on Tanzanian mushrooms and their uses. In J. H. Seyani & A. C. Chikuni (Eds.), *Proceedings of the XIIIth Plenary meeting of AETFAT, Zomba, Malawi, 2–11 April 1991* (Vol. 1, 729–734). Zomba, Malawi: National Herbarium and Botanic Gardens of Malawi.
- Härkönen, M., Saarimäki, T., & Mwasumbi, L. (1994c). Tanzanian mushrooms and their uses 4. Some reddish edible and poisonous *Amanita* species. *Karstenia*, 34, 47–60.
- Harmaja, H. (1976). Another poisonous species discovered in the genus *Gyromitra*: *G. ambigua*. *Karstenia*, 15, 36–37.
- Haro-Luna, M. X., Ruan-Soto, F., & Guzmán-Dávalos, L. (2019). Traditional knowledge, uses, and perceptions of mushrooms among the Wixaritari and mestizos of Villa Guerrero, Jalisco, Mexico. *IMA Fungus*, 10, 1–14.
- Harsh, N. S. K., Tiwari, C. K., & Rai, B. K. (1996). Forest fungi in the aid of tribal women of Madhya Pradesh [India]. *Sustainable Forestry*, 1, 10–15.
- Hawksworth, D. L. (2001). Mushrooms: The extent of the unexplored potential. *International Journal of Medicinal Mushrooms*, 3, 333–337.
- Hawksworth, D. L., & Lücking, R. (2017). Fungal diversity revisited: 2.2 to 3.8 million species. *Microbiol Spectrum*, 5, 1–17.
- He, M.-Q., Zhao, R.-L., Hyde, K. D., Begerow, D., Kemler, M., Yurkov, A., ... Sánchez-Ramírez, S. (2019). Notes, outline and divergence times of Basidiomycota. *Fungal Diversity*, 99, 105–367.
- He, X. (1991). *Verpa bohemica* – A seldom known and delicious edible fungus. *Zhongguo Shiyongjun (Edible Fungi of China)*, 10, 19.
- He, Z., Su, Y., Li, S., Long, P., Zhang, P., & Chen, Z. (2019). Development and evaluation of isothermal amplification methods for rapid detection of lethal *Amanita* species. *Frontiers in Microbiology*, 10, 1523.
- Heleno, S. A., Barros, L., Martins, A., Morales, P., Fernández-Ruiz, V., Glamoclija, J., ... Ferreira, I. C. (2015). Nutritional value, bioactive compounds, antimicrobial activity and bioaccessibility studies with wild edible mushrooms. *LWT-Food Science and Technology*, 63, 799–806.
- Heleno, S. A., Barros, L., Sousa, M. J., Martins, A., & Ferreira, I. C. (2009). Study and characterization of selected nutrients in wild mushrooms from Portugal by gas chromatography and high performance liquid chromatography. *Microchemical Journal*, 93, 195–199.
- Heleno, S. A., Barros, L., Sousa, M. J., Martins, A., & Ferreira, I. C. (2010). Tocopherols composition of Portuguese wild mushrooms with antioxidant capacity. *Food Chemistry*, 119, 1443–1450.
- Heleno, S. A., Ferreira, I. C., Ćirić, A., Glamoclija, J., Martins, A., Queiroz, M. J. R., & Soković, M. (2014). *Coprinopsis atramentaria* extract, its organic acids, and synthesized glucuronated and methylated derivatives as antibacterial and antifungal agents. *Food & Function*, 5, 2521–2528.
- Henkel, T. W., Aime, M. C., Chin, M., & Andrew, C. (2004). Edible mushrooms from Guyana. *Mycologist*, 18, 104–111.
- Hernández del Valle, J. F., García-Jiménez, J., Ayala-Vásquez, O. & de la Fuente-López, J. (2017). Edible ectomycorrhizal mushrooms from the montane cloud forest from El Cielo Biosphere Reserve, Tamaulipas, Mexico, pp. 101. In J. Pérez-Moreno & A. Guerin-Laguette (Eds.), *Mushrooms, humans and nature in a changing world*. Proceedings of the 9th International Workshop on Edible Mycorrhizal Mushrooms (IWEMM9). Colegio de Postgraduados, Texcoco, Mexico: 174 p.
- Hobbs, C. (1995). *Medicinal mushrooms: An exploration of tradition, healing, & culture* (2nd ed.). Santa Cruz, CA: Botanica Press.
- Hongo, T., & Izawa, M. (1994). *Yama-Kei field books. Vol. 10, Mushrooms*. Tokyo: Yamakei Publishers.
- Horowitz, K. M., Long, E. L. & Horowitz, B. Z. (2018). *Gyromitra* mushroom toxicity. In *StatPearls [Internet]*. StatPearls Publishing.
- Hosaka, H. (2002). *Laos edible fungi*. Personal Communication.
- Hossain, M. A., & Park, S.-C. (2016). A review on mushroom toxins. In D. Bagchi & A. Swaroop (Eds.), *Food Toxicology* (pp. 275–302). Boca Raton: CRC Press.
- Huang, N. (1989). New method of increasing production on *Rhizoglyphus piceus* in the south of Fujian Province. *Zhongguo Shiyongjun (Edible Fungi of China)*, 8 (39), 8.
- Huang, N. L. (1994). *Agaricus blazei* deserving of attention and popularization. *Zhongguo Shiyongjun (Edible Fungi of China)*, 13, 8–9.
- Hyde, K. D., Xu, J., Rapior, S., Jeewon, R., Lumyong, S., Niego, A. G. T., ... Stadler, M. (2019). The amazing potential of fungi: 50 ways we can exploit fungi industrially. *Fungal Diversity*, 97, 1–136.

- Imai, S. (1954). Elvellaceae Japoniae. *Science Reports of the Yokohama National University*, 3, 1–35.
- Imazeki, R., Hongo, T., & Otani, Y. (2011). *Fungi of Japan*: Yamakei Publishers, Tokyo.
- Index Fungorum. (2020). Retrieved from <http://www.indexfungorum.org>.
- Iordanov, D., Vanev, S. G., & Fakirova, V. I. (1978). *Gubite v Bulgariya: Opredelitel na nai-razprostraneniye yadlivi i otrovni gubi [Fungi of Bulgaria: Keys to the identification of the most widely distributed edible and poisonous fungi]*. Sofiya: Izd-vo na Bulg. Akad. na Naukite.
- Iqbal, M. (1993). International trade in non-wood forest products: An overview. *Misc/93/II Working Paper*. Rome: FAO.
- Iqbal, S., & Khalid, A. (1996). *Materials for the fungus flora of Pakistan*. 1. Check list of agarics, their distribution and association with surrounding vegetation. *Science International (Lahore)*, 8, 51–64.
- Irwin, A., & Leech, A. (2014). Fatal poisoning of a dog by the fungus *Clitocybe rivulosa*. *Veterinary Record*, 175, 122–123.
- Ishizuka, K. (2013). Osanpo Kinoko. *Shinano-mainichi Shimbun*, Nagano.
- Iwafuchi, Y., Morita, T., Kobayashi, H., Kasuga, K., Ito, K., Nakagawa, O., ... Kamimura, A. (2003). Delayed onset acute renal failure associated with *Amanita pseudoporphyria* Hongo ingestion. *Internal medicine*, 42, 78–81.
- Jabeen, S., Niazi, A. R., & Khalid, A. N. (2016). First record of *Russula anthracina* and its ectomycorrhiza associated with Himalayan cedar from South Asia. *Mycotaxon*, 131, 31–44.
- Jabłońska-Ryś, E., Skrzypczak, K., Sławińska, A., Radzki, W., & Gustaw, W. (2019). Lactic acid fermentation of edible mushrooms: Tradition, technology, current state of research: A review. *Comprehensive Reviews in Food Science and Food Safety*, 18(2), 655–669.
- Jang, M.-J., Lee, Y.-H., Ju, Y.-C., & Koo, H.-M. (2010). Cultural characteristics by sawdust and liquid spawn for the cultivation of *Neolentinus lepideus*. *The Korean Journal of Mycology*, 38, 125–129.
- Jarvinen, I., Kosonen, L., & Joutjarvi, M. (2004). *Parimad soogiseened ja maitsvaimad seenehorgutised*. Tallinn, Estonia: Sinisukk.
- Jayasuriya, W., Suresh, T., Abeytunga, D. T. U., Fernando, G., & Wanigatunga, C. (2012). Oral hypoglycemic activity of culinary-medicinal mushrooms *Pleurotus ostreatus* and *P. cystidiosus* (higher basidiomycetes) in normal and alloxan-induced diabetic Wistar rats. *International Journal of Medicinal Mushrooms*, 14, 347–355.
- Jo, W.-S., Hossain, M. A., & Park, S.-C. (2014). Toxicological profiles of poisonous, edible, and medicinal mushrooms. *Mycobiology*, 42, 215–220.
- Jones, E. B. G., Whalley, A. J. S., & Hywel-Jones, N. L. (1994). A fungus foray to Chiang Mai market in Northern Thailand. *Mycologist*, 8, 87–90.
- Jung, H., Park, Y., Yoo, Y.-B., Jeon, D.-H., & Park, K.-M. (2013). Analysis of nutritional contents and physiological activities of *Neolentinus lepideus*. *Journal of Mushroom*, 11, 261–268.
- Kabere, F. (2003). *Ate sorghum smut as child in Uganda*. Personal communication.
- Kaewnarin, K., Suwannarach, N., Kumla, J., & Lumyong, S. (2016). Phenolic profile of various wild edible mushroom extracts from Thailand and their antioxidant properties, anti-tyrosinase and hyperglycaemic inhibitory activities. *Journal of Functional Foods*, 27, 352–364.
- Kalotas, A. (1997). Aboriginal knowledge and use of fungi. In K. Mallet & C. Grgurinovic (Eds.), *Fungi of Australia. Volume 1B. Introduction – Fungi in the environment* (pp. 269–295). Canberra, Australia: Australian Biological Resources Study, CSIRO Publishing.
- Kalyoncu, F., Oskay, M., Sağlam, H., Erdoğan, T. F., & Tamer, A. Ü. (2010). Antimicrobial and antioxidant activities of mycelia of 10 wild mushroom species. *Journal of Medicinal Food*, 13, 415–419.
- Kamal, A., Fauzia, B., & Abul, K. (2009). Mushroom production in Bangladesh: Present scenario and potentialities. *SAARC Journal of Agriculture*, 7, 91–105.
- Kamalebo, H. M., & De-Kesel, A. (2020). Wild edible ectomycorrhizal fungi: An underutilized food resource from the rainforests of Tshopo province (Democratic Republic of the Congo). *Journal of Ethnobiology and Ethnomedicine*, 16, 1–82.
- Kamalebo, H. M., Malale, H. N. S. W., Ndabaga, C. M., Degreef, J., & De-Kesel, A. (2018). Uses and importance of wild fungi: Traditional knowledge from the Tshopo province in the Democratic Republic of the Congo. *Journal of Ethnobiology and Ethnomedicine*, 14, 13.
- kan Bakır, T., Boufars, M., Karadeniz, M., & Ünal, S. (2018). Amino acid composition and antioxidant properties of five edible mushroom species from Kastamonu, Turkey. *African Journal of Traditional, Complementary and Alternative Medicines*, 15, 80–87.
- Kanagasabapathy, G., Malek, S., Kuppasamy, U., & Vikineswary, S. (2011). Chemical composition and antioxidant properties of extracts of fresh fruiting bodies of *Pleurotus sajor-caju* (Fr.) Singer. *Journal of Agricultural and Food Chemistry*, 59, 2618–2626.
- Kang, J., Kang, Y., Ji, X., Guo, Q., Jacques, G., Pietras, M., ... Łuczaj, Ł. (2016). Wild food plants and fungi used in the mycophilous Tibetan community of Zhagana (Tewo County, Gansu, China). *Journal of Ethnobiology and Ethnomedicine*, 12(1), 21.
- Kaplaner, E., Singeç, M. H., & Öztürk, M. (2017). Fatty acid composition and antioxidant activity of *Tricholoma imbricatum* and *T. Focale*. *Turkish Journal of Agriculture-Food Science and Technology*, 5, 1080–1085.
- Karadelev, M., Miteva, S., & Stojkoska, K. (2003). Checklist of humano-toxic macromycetes in the Republic of Macedonia. In *Proceedings of the 2nd Congress of Ecologists of the Republic of Macedonia with International Participation, 25–29.10 2003*. (pp. 472–478). Special issues of Macedonian Ecological Society, Vol. 6, Skopje.
- Karhula, P., Harkonen, M., Saarimaki, T., Verbeken, A., & Mwasumbi, I. (1998). Tanzanian mushrooms and their uses, *Lactarius Karstenia*, 38, 49–69.
- Karun, N. C., & Sridhar, K. R. (2017). Edible wild mushrooms of the Western Ghats: Data on the ethnic knowledge. *Data in Brief*, 14, 320–328.
- Karun, N. C., Sridhar, K. R., & Ambarish, C. N. (2018). Nutritional potential of *Auricularia auricula-judae* and *Termitomyces umkowaan*—The wild edible mushrooms of southwestern India. In V. K. Gupta, H. Treichel, V. Shapaval, L. Antonio de Oliveira, M. G. Tuohy (Eds.), *Microbial Functional Foods and Nutraceuticals* (pp. 281–301). Hoboken, NJ: Wiley.
- Kasper-Pakosz, R., Pietras, M., & Łuczaj, Ł. (2016). Wild and native plants and mushrooms sold in the open-air markets of south-eastern Poland. *Journal of Ethnobiology and Ethnomedicine*, 12, 45.
- Kasuya, T. (2008). *Phallus luteus* comb. nov., a new taxonomic treatment of a tropical phalloid fungus. *Mycotaxon*, 106, 7–13.

- Kasuya, T., & Katumoto, K. (2008). Notes on Japanese Lycoperdaceae. 4. Validation of Japanese giant puffball, *Calvatia nipponica*. *Mycoscience*, *49*, 271–275.
- Kasuya, T., Maruyama, T., Fuse, K., Hosaka, K., & Minowa, K. (2017). *Desarmillaria ectypa*, a paludal fungus newly found in Niigata Prefecture, Central Honshu, Japan. *Annual Report of the Kashiwazaki City Museum*, *31*, 99–108.
- Katende, A. B., Segawa, P., & Birnie, A. (1999). *Wild food plants and mushrooms of Uganda*. Nairobi, Kenya: Regional Land Management Unit, Swedish International Development Cooperation Agency.
- Kawagoe, S. (1924). The market fungi of Japan. *Transactions of the British Mycological Society*, *10*, 201–206.
- Kawamura, S. (1954). *Genshoku Nihon kinrui zukan: Icons of Japanese fungi* (Vol. 6). Tokyo: Kazama Shobō.
- Kaya, A., Kılıçel, F., Karapınar, H. S., & Uzun, Y. (2017). Mineral contents of some wild edible mushrooms. *The Journal of Fungus*, *8*, 178–183.
- Kayano, T., Kitamura, N., Miyazaki, S., Ichianagi, T., Shimomura, N., Shibuya, I., & Aimi, T. (2014). Gymnopilins, a product of a hallucinogenic mushroom, inhibit the nicotinic acetylcholine receptor. *Toxicon*, *81*, 23–31.
- Kerrigan, R. W. (2005). *Agaricus subrufescens*, a cultivated edible and medicinal mushroom, and its synonyms. *Mycologia*, *97*, 12–24.
- Khobar, L., & Najim, L. (2001). Les truffes au Maroc: Biodiversité et circuit de commercialisation. In Science et Culture de la Truffe. Actes du Ve Congrès International, 4 au 6 Mars 1999, Aix-en-Provence, France (pp. 511–514). Federation Française des Trufficulteurs.
- Khatua, S., & Acharya, K. (2017). Alkaline extractive crude polysaccharide from *Russula senecis* possesses antioxidant potential and stimulates innate immunity response. *Journal of Pharmacy and Pharmacology*, *69*, 1817–1828.
- Khatua, S., & Acharya, K. (2014). Antioxidant and antimicrobial potentiality of quantitatively analysed ethanol extract from *Macrocybe crassa*. *International Journal of Pharmaceutical Sciences Review and Research*, *29*, 53–60.
- Khatua, S., Dutta, A. K., Chandra, S., Paloi, S., Das, K., & Acharya, K. (2017). Introducing a novel mushroom from mycophagy community with emphasis on biomedical potency. *PLoS One*, *12*, e0178050.
- Kho, Y., Vikineswary, S., Abdullah, N., Kuppusamy, U. R., & Oh, H. (2009). Antioxidant capacity of fresh and processed fruit bodies and mycelium of *Auricularia auricula-judae* (Fr.) Qué. *Journal of Medicinal Food*, *12*, 167–174.
- Kim, T., Lee, D., Lee, J. H., Lee, Y.-S., Oh, B. J., Lim, K. S., & Kim, W. Y. (2017). Predictors of poor outcomes in patients with wild mushroom-induced acute liver injury. *World Journal of Gastroenterology*, *23*, 1262.
- Kirchmair, M., Carrilho, P., Pfab, R., Haberl, B., Felgueiras, J., Carvalho, F., ... Neuhauser, S. (2012). *Amanita* poisonings resulting in acute, reversible renal failure: New cases, new toxic *Amanita* mushrooms. *Nephrology Dialysis Transplantation*, *27*, 1380–1386.
- Kirk, P. M., Cannon, P. F., David, J. C., & Stalpers, J. A. (2001). *Dictionary of the fungi* (9th ed.). Wallingford: CAB International.
- Klimaszyn, P., & Rzymyski, P. (2018). The yellow knight fights back: Toxicological, epidemiological, and survey studies defend edibility of *Tricholoma equestre*. *Toxins*, *10*, 468.
- Kodama, N., Komuta, K., Sakai, N., & Nanba, H. (2002). Effects of D-Fraction, a polysaccharide from *Grifola frondosa* on tumor growth involve activation of NK cells. *Biological and Pharmaceutical Bulletin*, *25*, 1647–1650.
- Kovacs, G. M., Trappe, J. M., Alsheikh, A. M., Bóka, K., & Elliott, T. F. (2008). Imaia, a new truffle genus to accommodate *Terfezia gigantea*. *Mycologia*, *100*, 930–939.
- Kumar, M., Harsh, N. S. K., Prasad, R., & Pandey, V. V. (2017). An ethnomycological survey of Jaunsar, Chakrata, Dehradun, India. *Journal of Threatened Taxa*, *9*(9), 10717–10725.
- Kumar, R., Tapwal, A., Pandey, S., Borah, R. K., Borah, D., & Borghain, J. (2013). Macro-fungal diversity and nutrient content of some edible mushrooms of Nagaland, India. *Nusantara Bioscience*, *5*, 1–7.
- Kumla, J., Suwannarach, N., Jaiyasen, A., Bussaban, B., & Lumyong, S. (2013). Development of an edible wild strain of Thai oyster mushroom for economic mushroom production. *Chiang Mai Journal of Science*, *40*, 161–172.
- Kytovuori, I. (1989). The *Tricholoma caligatum* group in Europe and North Africa. *Karstenia*, *28*, 65–77.
- Laessoe, T., & del-Conte, A. (1996). *The mushroom book*. London: Dorling Kindersley.
- Laessøe, T., & Petersen, J. H. (2019). *Fungi of temperate Europe (Volumes 1 and 2)*. Princeton, NJ: Princeton University Press.
- Lalrinawmi, Z., Vabeikhochei, J., & Zothanzama, J. (2017). Edible mushrooms of Mizoram. *Science Vision*, *17*, 172–181.
- Lamus, V., Franco, S., Montoya, L., Endara, A. R., Caballero, L. A., & Bandala, V. M. (2015). Mycorrhizal synthesis of the edible mushroom *Turbinellus floccosus* with *Abies religiosa* from central Mexico. *Mycoscience*, *56*, 622–626.
- Le, T. H., Nguyen, T. P. M., & Doan, T. K. N. (2018). Biodiversity assessment of mushroom's family Polyporeacea in Cuc Phuong national park of Vietnam. In The first National conference on mushroom 2018, 03 Dec 2018. Vietnam.
- Leathem, A. M., & Dorran, T. J. (2007). Poisoning due to raw *Gyromitra esculenta* (false morels) west of the Rockies. *Canadian Journal of Emergency Medicine*, *9*, 127–130.
- Lee, S.-S., Kim, D.-H., & Chung, H.-C. (2002). Observations on the ectomycorrhizas of edible fungi collected from the central mountains of SoBaek, Korea. In I. R. Hall, Y. Wang, A. Zambonelli, & E. Danell (Eds.), *Edible ectomycorrhizal mushrooms and their cultivation. Proceedings of the second international conference on edible ectomycorrhizal mushrooms*. Christchurch: New Zealand Institute for Crop and Food Research Limited.
- Legg, A. (1991). Your top twenty fungi – The final list. *Mycologist*, *4*, 23–24.
- Lehmann, P. F., & Khazan, U. (1992). Mushroom poisoning by *Chlorophyllum molybdites* in the Midwest United States. *Mycopathologia*, *118*, 3–13.
- Li, G., Bono, L., Onofrio, S., Franca, S., Camillo, C., Carmela, Z., ... Flavia, C. (2019). Acute renal failure after *Amanita ovoidea* eating. *Indian Journal of Nephrology*, *29*, 73–74.
- Li, H., Lee, H. S., Kim, S. H., Moon, B., & Lee, C. (2014). Antioxidant and anti-inflammatory activities of methanol extracts of *Tremella fuciformis* and its major phenolic acids. *Journal of Food Science*, *79*, C460–C468.
- Li, H., Zhang, H., Zhang, Y., Zhang, K., Zhou, J., Yin, Y., ... Zhang, Y. (2020). Mushroom poisoning outbreaks—China, 2019. *China CDC Weekly*, *2*, 19–24.

- Li, T., Yang, Z., Chen, Z., Song, B., & Deng, W. (2005). Poisonous mushrooms known from China – Species resources and distribution. *ACTA Edulis Fungi* 2005 vol.12 Supplement: Proceedings of the Fifth International Conference on Mushroom Biology and Mushroom Products.
- Lima, A. D., Fortes, R. C., Novaes, M. G., & Percário, S. (2012). Poisonous mushrooms: A review of the most common intoxications. *Nutricion Hospitalaria*, 27, 402–408.
- Lincoff, G., & Mitchel, D. H. (1977). *Toxic and hallucinogenic mushroom poisoning: A handbook for physicians and mushroom hunters*. New York: Van Nostrand Reinhold Company.
- Lindley, J. (1849). *Medical and oecconomical botany*. London: Bradbury and Evans.
- Liu, G. (2002). *Eating Sporisorium cruenta in China*. Personal Communication.
- Liu, K., Wang, J., Zhao, L., & Wang, Q. (2013). Anticancer, antioxidant and antibiotic activities of mushroom *Ramaria flava*. *Food and Chemical Toxicology*, 58, 375–380.
- Liu, P. G. (1990). Investigation of the edible mushroom resources of Mt. Daqing of Inner Mongolia. *Zhongguo Shiyongjun (Edible Fungi of China)*, 9, 26–27.
- Liu, W. P., & Yang, H. R. (1982). An investigation of mushroom poisoning in Ninghua county during the last 20 years. *Chinese Journal of Preventative Medicine*, 16, 226–228.
- Locquin, M. (1954). Une chanterelle comestible de la Cote d'Ivoire: *Hygrophoropsis mangelotii* sp. nov. *Journal d'Agriculture Tropicale et de Botanique Appliquée*, 1, 359–261.
- Locsmándi-Vasas, A. (1995). *Ismerjuk fel a gombákat (Let's recognize mushrooms)*. Budapest (probably): Gabo.
- Logemann, H., Argueta, J., Guzman, G., Montoya-Bello, L., Bandala-Munoz, V. M., & de-Leon-Chocooj, R. (1987). Lethal poisoning by mushrooms in Guatemala. *Revista Mexicana de Micología*, 3, 211–216.
- Longvah, T., & Deosthale, Y. G. (1998). Compositional and nutritional studies on edible wild mushroom from northeast India. *Food Chemistry*, 63, 331–334.
- López García, A., Jiménez-Ruiz, M., & Pérez-Moreno, J. (2017). Vocablos relacionados con el recurso micológico en el idioma de la cultura chinanteca de la Sierra Norte del estado de Oaxaca, México. *Revista Mexicana de Micología*, 46, 9–18.
- Lopez, G. A., Manzola-Cruz, J. M., & Zamora-Martinez, M. C. (1992). Evaluacion de la produccion de hongos comestibles silvestres en San Juan Tetla, Puebla. Ciclo 1992. In Reunion Cientifica Forestal y Agropecuaria (pp. 182–191). Coyocan, Mexico.
- Lu, T., Bau, T., & Ohga, S. (2017). Physiological study of the wild edible mushroom *Leucocalocybe mongolica*. *Journal of the Faculty of Agriculture*, 62, 1–8.
- Lu, H., Lou, H., Hu, J., Liu, Z., & Chen, Q. (2020). Macrofungi: A review of cultivation strategies, bioactivity, and application of mushrooms. *Comprehensive Reviews in Food Science and Food Safety*, 2020(11), 1–24.
- Luangharn, T., Hyde, K. D., & Chukeatirote, E. (2014). Proximate analysis and mineral content of *Laetiporus sulphureus* strain MFLUCC 12–0546 from northern Thailand. *Chiang Mai Journal of Science*, 41, 765–770.
- Łuczaj, Ł., Stawarczyk, K., Kosiek, T., Pietras, M., & Kujawa, A. (2015). Wild food plants and fungi used by Ukrainians in the western part of the Maramureş region in Romania. *Acta Societatis Botanicorum Poloniae*, 84, 339–346.
- Luginina, E., & Egoshina, T. 2020. Productivity of edible fungi in taiga communities. IOP Conference Series: Earth and Environmental Science.
- Lurie, Y., Wasser, S. P., Taha, M., Shehade, H., Nijim, J., Hoffmann, Y., ... Suaed, S. (2009). Mushroom poisoning from species of genus *Inocybe* (fiber head mushroom): A case series with exact species identification. *Clinical toxicology*, 47, 562–565.
- Ma, G., Yang, W., Zhao, L., Pei, F., Fang, D., & Hu, Q. (2018). A critical review on the health promoting effects of mushrooms nutraceuticals. *Food Science and Human Wellness*, 7, 125–133.
- Maiti, S., Bhutia, S. K., Mallick, S. K., Kumar, A., Khadgi, N., & Maiti, T. K. (2008). Antiproliferative and immunostimulatory protein fraction from edible mushrooms. *Environmental Toxicology and Pharmacology*, 26, 187–191.
- Malaisse, F., De-Kesel, A., N'gasse, G., & Lognay, G. (2008). Diversité des champignons consommés par les pygmées Bofi de la Lobaye (République centrafricaine). *Geo-Eco-Trop*, 32, 1–8.
- Mallikarjuna, S., Ranjini, A., Haware, D. J., Vijayalakshmi, M., Shashirekha, M., & Rajarathnam, S. (2013). Mineral composition of four edible mushrooms. *Journal of Chemistry*, 2013, 1–5.
- Malyi, L. P. (1987). Resources of edible fungi in Belorussia [Belarus] and the possibility of their utilization. *Rastitelo'nye Resursy*, 23, 532–536.
- Mao, X. L. (1991). Poisonous fungi of Amanitaceae and their toxins in China. *Microbiology China*, 18, 160–165.
- Mao, X. L. (1998). *Economic fungi of China (in Chinese)*. Beijing: Science Press.
- Mao, X. L. (2000). *The Macrofungi of China*. Zhengzhou, Henan: Henan Science and Technology Press.
- Maravalhas, N. (1965). O pao de indio. *Ciencia e Cultura*, 17, 18–20.
- Marles, R. J., Clavelle, C., Monteleone, L., Tays, N., & Burns, D. (2000). *Aboriginal plant use in Canada's northwest boreal forest*. Vancouver: University of British Columbia.
- Martínez, A., Oria-de-Rueda, J. A., & Martínez, P. (1997). *Estudio sobre la potencialidad de los diferentes usos del bosque para la creación de empleo y actividad económica en el medio rural de Castilla y León. La producción de mayor potencialidad: Hongos silvestres comestibles*. Palencia: Junta de Castilla y León y Fondo Social Europeo.
- Martinez-de-Aragón, J., Florit, E., & Colinas, C. (1998). Producción de setas micorrízicas y comestibles en la comarca del Solsones en 1997. In R.B. Eduardo (Ed.), *III Forum de Política Forestal* (pp. 322–328). Solsona (Lleida): Centre Tecnologic Forestal de Catalunya.
- Martins, X. (2004). *Cogumelos*. Viseu: Mirandela.
- Mata, J. L., & Mishra, N. T. (2015). Comparative study of worldwide species of genus *Lentinus* (= *Lentinula*, higher Basidiomycetes) based on linear mycelium growth. *International Journal of Medicinal Mushrooms*, 17(5), 481–489.
- Mattick, F. (1969). Not known but information on *Umbilicaria esculenta* (rock lichen). *Nova Hedwigia*, 16, 511.
- McIlvaine, C., & Macadam, R. K. (1900). *One thousand American fungi*. AbeBooks.
- Meijer, A. A. R. d., Amazonas, M. A. L. d. A., Rubio, G. B. G., & Curial, R. M. (2007). Incidences of poisonings due to *Chlorophyllum molybdites* in the state of Paraná, Brazil. *Brazilian Archives of Biology and Technology*, 50, 479–488.
- Meijer, A. d. (2008). *Macrofungos notáveis das Florestas de Pinheiro-do-Paraná*. Colombo: Embrapa.

- Mendoza, J. M. (1938). Philippines mushrooms. *Philippine Journal of Science*, 65, 1–128 (and 179 plates).
- Meuninck, J. (2017). *Foraging mushrooms oregon: Finding, identifying, and preparing edible wild mushrooms*. Helena, Montana: Falcon Guides.
- Michelot, D., & Toth, B. (1991). Poisoning by *Gyromitra esculenta*—a review. *Journal of Applied Toxicology*, 11, 235–243.
- Minato, K.-i., Laan, L. C., Ohara, A., & van Die, I. (2016). Pleurotus citrinopileatus polysaccharide induces activation of human dendritic cells through multiple pathways. *International Immunopharmacology*, 40, 156–163.
- Ministry of Agriculture and Forestry of Finland. (2020). Retrieved from <http://www.mmm.fi>.
- Ministry of Forestry. (2020). Retrieved from <http://www.ogm.gov.tr>.
- Minter, D. (2002). *Edible fungi in Chile, Cuba and Argentina*. Personal Communication.
- Minter, D., Cannon, P. F., & Peredo, H. L. (1987). South America species of *Cyrtaria* (a remarkable and beautiful group of edible ascomycetes). *Mycologist*, 1, 7–11.
- Mleczek, M., Magdziak, Z., Goliński, P., Siwulski, M., & Stuper-Szablewska, K. (2013). Concentrations of minerals in selected edible mushroom species growing in Poland and their effect on human health. *Acta Scientiarum Polonorum Technologia Alimentaria*, 12, 203–214.
- Molnár, P., Ósz, E., Turcsi, E., & Deli, J. (2019). Carotenoid composition of the mushroom Scarlet elf cup (*Sarcoscypha coccinea*). *Heliyon*, 5, e01883.
- Mizuno, T. (1995). Mushrooms: The versatile fungus-food and medicinal properties. *Food Reviews International*, 11 (1), 151–166.
- Montoya, A., Briones-Dumas, E., Núñez-López, R. A., Kong, A., Ortiz-Hernández, V., & Moreno-Fuentes, Á. (2019). Los hongos conocidos por la comunidad Yuhmu de Ixtenco, Tlaxcala, México. *Scientia fungorum*, 49, e1230.
- Montoya-Esquivel, A. (1998). Ethnomycology of Tlaxcala, Mexico. *McIlvainea*, 13, 6–12.
- Montoya-Esquivel, A., Estrada-Torres, A., Kong, A., & Juarez-Sanchez, L. (2001). Commercialization of wild mushrooms during market days of Tlaxcala, Mexico. *Micologia Aplicada Internacional*, 13, 31–40.
- Morales, O., Bran, M., & Cáceres, R. (2010). Los hongos comestibles de uso tradicional en Guatemala. Hacia un Desarrollo Sostenible del Sistema de Producción-Consumo de los Hongos Comestibles y Medicinales en Latinoamérica: *Avances y Perspectivas en el Siglo*, 21, 437–464.
- Morales, O., Bran, M., Cáceres, R., & Flores, R. (2003). *Contribución al conocimiento de los hongos comestibles de Guatemala. Documento interno, Facultad de Ciencias Químicas y de Farmacia, Guatemala, Universidad de San Carlos*.
- Morales, O., Bran, M., Cáceres, R., & Flores, R. (2015). Las especies del género *Laccaria* (Agaricales) en Guatemala. *Revista Científica de la Facultad de Ciencias Químicas y Farmacia*, 16, 89–94.
- Moreno-Arroyo, B., Recio, J. M., Gomez, J., & Pulido, E. (2001). *Tuber oligospermum* from Morocco. *Mycologist*, 15, 41–42.
- Moreno-Fuentes, A., Cifuentes, J., Bye, R., & Valenzuela, R. (1996). Kute-mo'ko-a: An edible fungus of the Raramuri Indians of Mexico. *Revista Mexicana de Micología*, 12, 31–39.
- Moreno-Solis, S. (2018). *Importancia biocultural de los hongos comestibles con potencial biotecnológico para la producción de especies forestales de importancia en México*. Guerrero, Mexico: Instituto Tecnológico de Ciudad Altamirano.
- Morris, B. (1984). Macrofungi of Malawi: Some ethnobotanical notes. *Bulletin of the British Mycological Society*, 18, 48–57.
- Mortimer, P., Xu, J., Karunaratna, S., & Hyde, K. D. (2014). *Mushrooms for trees and people: A field guide to useful mushrooms of the Mekong region*. Kunming: The World Agroforestry Centre.
- Mukerji, K. G., & Manoharachary, C. (2010). *Taxonomy and ecology of Indian fungi*. New Delhi: IK International Pvt Ltd.
- Murakami, Y. (1993). *Larger fungi from northern Pakistan*. In T. Nakaike & S. Malik (Eds.), *Cryptogamic flora of Pakistan* (Vol. 2, pp. 105–147). VolTokyo: National Science Museum.
- Mushroom Atlas. (2020). Retrieved from <http://www.grzyby.pl/>.
- Mustonen, A.-M., Määttä, M., Kärjä, V., Puukka, K., Aho, J., Saarela, S., & Nieminen, P. (2018). Myo- and cardiotoxic effects of the wild winter mushroom (*Flammulina velutipes*) on mice. *Experimental Biology and Medicine*, 243, 639–644.
- Mycoweb. (2020). Retrieved from <http://www.mykoweb.com/>.
- Nagasawa, E., & Redhead, S. (1988). *A new edible agaric from Japan: Clitocybula esculenta*. Reports of the Tottori Mycological Institute (Japan).
- Namgyel, P. (2000). The story of Buddha mushroom. *Tricholoma matsutake. Unpublished manuscript* (pp. 1–14). Thimpu, Bhutan.
- Nanagulyan, S., Perevedenceva, L., Margaryan, L., Hovhannisyan, E. K., & Hovhannisyan, H. (2017). Poisoning fungi and poisonings with macromycetes. *Chemistry and Biology*, 51, 129–131.
- Nanagulyan, S. G. (2002). *Edible fungi in Armenia*. Personal Communication.
- Niazi, A. R. K. (2008). *Biodiversity of ectomycorrhizas in conifers from Himalayan moist temperate forests of Pakistan*. Lahore, Pakistan: University of the Punjab.
- Niksic, M., Klaus, A., & Argyropoulos, D. (2016). Safety of foods based on mushrooms. In V. Prakash O. Martín-Belloso L. Keener S. Astley H. McMahon & H. Lelieveld (Eds.), *Regulating Safety of Traditional and Ethnic Foods* (pp. 421–439). Amsterdam: Elsevier.
- Nomura, C., Masayama, A., Yamaguchi, M., Sakuma, D., & Kajimura, K. (2017). PCR-based method for the detection of toxic mushrooms causing food-poisoning incidents. *Shokuhin Eiseigaku Zasshi Journal of the Food Hygienic Society of Japan*, 58, 132–142.
- Obodai, M., & Apetorgbor, M. (2001). An ethnobotanical study of mushroom germplasm and its domestication in the Bia Biosphere Reserve of Ghana. Report presented to UNESCO through Environmental Protection Agency of Ghana, Accra.
- Ogawa, W., Endo, N., Fukuda, M., & Yamada, A. (2018). Phylogenetic analyses of Japanese golden chanterelles and a new species description, *Cantharellus anzutake* sp. nov. *Mycoscience*, 59, 153–165.
- Omar, N. A. M., Abdullah, N., Kuppusamy, U. R., Abdulla, M. A., & Sabaratnam, V. (2011). Nutritional composition, antioxidant activities, and antiulcer potential of *Lentinus squarrosulus* (Mont.) mycelia extract. Evidence-based Complementary and Alternative Medicine (ECAM).
- Ori, F., Trappe, J., Leonardi, M., Iotti, M., & Pacioni, G. (2018). Crested porcupines (*Hystrix cristata*): Mycophagist spore dispersers of the ectomycorrhizal truffle *Tuber aestivum*. *Mycorrhiza*, 28, 561–565.
- Osemwegie, O., Okhuoya, J., & Dania, T. (2014). Ethnomycological conspectus of West African mushrooms: An awareness document. *Advances in Microbiology*, 4, 39–54.

- Oso, B. (1975). Mushrooms and the Yoruba people of Nigeria. *Mycologia*, 67, 311–319.
- Ota, Y., Matsushita, N., Nagasawa, E., Terashita, T., Fukuda, K., & Suzuki, K. (1998). Biological species of *Armillaria* in Japan. *Plant Disease*, 82, 537–543.
- Otani, Y. (1981). *Termitomyces clypeatus* Heim collected from Iriomote Island, Okinawa, Japan. *Bulletin of the National Science Museum. Series B: Botany*, 7, 131–134.
- Paloi, S., & Acharya, K. (2013). Antioxidant activities and bioactive compounds of polyphenol rich extract from *Amanita vaginata* (Bull.) Lam. *International Journal of PharmTech Research*, 5, 1645–1654.
- Panda, M. K., & Tayung, K. (2015). Documentation and ethnomedicinal knowledge on wild edible mushrooms among ethnic tribes of northern Odisha, India. *Asian J Pharm Clin Res*, 8, 139–143.
- Papetti, C., Consiglio, G., & Simonini, G. (2017). *Atalante fotografica dei funghi d' Italia* (Vol. 1). Trento: Associazione Micologica Bresadola.
- Park, W. H., & Lee, H. D. (1999). *Korean mushrooms*. Seoul: Kyo-Hak-Sa.
- Park, W., & Lee, J. (2011). *New wild fungi of Korea*. Seoul: Kyo-Hak Publishing Co.
- Patocka, J. (2018). Bolesatine, a toxic protein from the mushroom *rubroboletus satanas*. *Military Medical Science Letters*, 87, 14–20.
- Pegler, D. N., & Pearce, G. D. (1980). The edible mushrooms of Zambia. *Kew Bulletin*, 35, 475–491.
- Pegler, D. N., & Vanhaecke, M. (1994). *Termitomyces* of southeast Asia. *Kew Bulletin*, 49, 717–736.
- Peintner, U., Schwarz, S., Mešić, A., Moreau, P.-A., Moreno, G., & Saviuc, P. (2013). Mycophilic or mycophobic? Legislation and guidelines on wild mushroom commerce reveal different consumption behaviour in European countries. *PloS one*, 8, e63926.
- Pelkonen, R., Alfthan, G., & Järvinen, O. (2006). *Cadmium, lead, arsenic and nickel in wild edible mushrooms*. Helsinki: Finnish Environment Institute.
- Pereira, E., Barros, L., Martins, A., & Ferreira, I. C. (2012). Towards chemical and nutritional inventory of Portuguese wild edible mushrooms in different habitats. *Food Chemistry*, 130, 394–403.
- Pereira, L. T. (2019). FANCs de Angatuba: Fungos Alimentícios Não Convencionais de Angatuba e região: Simplíssimo.
- Pérez-Moreno, J., Lorenzana-Fernández, A., Carrasco-Hernández, V., & Yescas-Pérez, A. (2010). *Los hongos comestibles silvestres del parque nacional Izta-Popo, Zoquiapan y anexos*. Montecillo, Texcoco, Estado de México: Colegio de Postgraduados, SEMARNAT, Conacyt.
- Pérez-Moreno, J., Martínez-Reyes, M., Yescas-Pérez, A., Delgado-Alvarado, A., & Xoconostle-Cázares, B. (2008). Wild mushroom markets in Central Mexico and a case study at Ozumba. *Economic botany*, 62, 425–436.
- Pha Khao, Lao (2020). Retrieved from <https://www.phakhaolao.la/en/kb/0000086>.
- Phengsintham, P., Souvvanasane, T., & Keokaen, T. (2018). Fungi from Phouchomwoy Provincial Protected Area (Report).
- Phillips, R. (1981). *Mushrooms and other fungi of Great Britain* (pp. 288). Europe: Pan Books.
- Phongseun, S., Somsanith, B., & Thaviphone, S. (2017). *Some mushrooms found in Houay Yang Reserves*. Vientiane, Laos. Ministry of Science and Technology, Lao PDR.
- Pearce, G. D. (1981). Zambian mushrooms – Customs and folklore. *Bulletin of the British Mycological Society*, 15, 139–142.
- Pieroni, A., Nebel, S., Santoro, R. F., & Heinrich, M. (2005). Food for two seasons: Culinary uses of non-cultivated local vegetables and mushrooms in a south Italian village. *International Journal of Food Sciences and Nutrition*, 56(4), 245–272.
- Pilz, D., & Molina, R. (2002). Commercial harvests of edible mushrooms from the forests of the Pacific Northwest United States: Issues, management, and monitoring for sustainability. *Forest Ecology & Management*, 155, 3–16.
- Pires, A. d. R. A., Ruthes, A. C., Cadena, S. M. S. C., Acco, A., Gorin, P. A. J., & Iacomini, M. (2013). Cytotoxic effect of *Agaricus bisporus* and *Lactarius rufus* β -D-glucans on HepG2 cells. *International Journal of Biological Macromolecules*, 58, 95–103.
- Poudel, D., Kaphle, K., & Shrestha, S. M. (2018). Status of mushroom cultivation and wild mushroom poisoning in Chitwan, Nepal. Wild harvests, governance, and livelihoods in Asia.
- Power, R. C., Salazar-García, D. C., Straus, L. G., Morales, M. R. G., & Henry, A. G. (2015). Microremains from El Mirón Cave human dental calculus suggest a mixed plant–animal subsistence economy during the Magdalenian in Northern Iberia. *Journal of Archaeological Science*, 60, 39–46.
- Prance, G. (1984). The use of edible fungi by Amazonian Indians. *Advances in Economic Botany*, 1, 127–139.
- Prance, G. T. (1972). An ethnomycological comparison of four tribes of Amazonian Indians. *Acta Amazonica*, 2, 7–27.
- Prance, G. T. (1987). Etnobotánica de algumas tribos amazônicas. *Suma etnológica brasileira*, 1, 119–133.
- Prasher, I., Sharma, M., & Chander, H. (2007). Edible and medicinal macrofungi of Chandigarh. In I. Prasher & M. Sharma (Eds.), *Advances in Mycology and Plant Pathology* (pp. 317–330). Dehra Dun: Bishen Singh Mahendra Pal Singh.
- Prospects of Mushroom in Bangladesh. (2020). Internship report on prospects of mushroom in bangladesh: A case study of mushroom products (www.assignmentpoint.com).
- Purkayastha, R. P., & Chandra, A. (1985). *Manual of edible mushrooms*. New Delhi: Today and Tomorrow's Printers and Publishers.
- Puschner, B. (2018). Mushroom toxins. In R.C. Gupta (Ed.), *Veterinary Toxicology: Basic and Clinical Principles: Third Edition* (pp. 955–966). United State: Academic Press.
- Pushpa, H., & Purushothama, K. (2012). Biodiversity of mushrooms in and around Bangalore (Karnataka), India. *American-Eurasian Journal of Agricultural & Environmental Sciences*, 12, 750–759.
- Putzke, J. (2014). *Cogumelos no Sul do Brasil*, v. 1. Casa das Letras, Estrela.
- Rai, M. K. (1997). Wild edible mushrooms of tribal areas of Seoni district, Madhya Pradesh. *Mushroom Research*, 6, 107.
- Ramírez-Carbajal, E. (2017). *Etnomicrología en la zona Tlahuica-Pjiekakjoo del estado de México*. San Felipe del Progreso, Mexico: Universidad Intercultural del Estado de México.
- Ramkumar, L., Ramanathan, T., Thirunavukkarasu, P., & Arivuselvan, N. (2010). Antioxidant and radical scavenging activity of nine edible mushrooms extract. *International Journal of Pharmacology*, 6, 950–953.
- Rammeloo, J., & Walley, R. (1993). The edible fungi of Africa south of the Sahara: A literature survey. *Scripta Botanica Belgica*, 5, 1–62.

- Razaq, A., Ilyas, S., & Khalid, A. N. (2014). Taxonomic and phylogenetic affinities of *Hygrophorus chrysodon* from western Himalayan forests. *Austrian Journal of Mycology*, 23, 21–29.
- Razaq, A., & Shahzad, S. (2016). New records of order boletales from Pakistan. *Pak. J. Bot.*, 48, 1313–1317.
- Razzaq, A., & Shahzad, S. (2004). *Pisolithus tinctorius*, a new record from Pakistan. *Pakistan Journal of Botany*, 36, 449–452.
- Redeuilh, G. (2004). *Larousse des Champignons*. Place of publication not indicated: Larousse.
- Reis, F. S., Martins, A., Vasconcelos, M. H., Morales, P., & Ferreira, I. C. (2017). Functional foods based on extracts or compounds derived from mushrooms. *Trends in Food Science & Technology*, 66, 48–62.
- Remotti, C. D., & Colan, J. A. (1990). Identification of wild edible fungi in Dantas Forest, Huanuco. *Revista Forestal del Peru*, 17, 21–37.
- Reverberi, M., Fabbri, A. A., Zjalic, S., Ricelli, A., Punelli, F., & Fanelli, C. (2005). Antioxidant enzymes stimulation in *Aspergillus parasiticus* by *Lentinula edodes* inhibits aflatoxin production. *Applied Microbiology and Biotechnology*, 69, 207–215.
- Richardson, D. H. S. (1991). Lichens and man. In D. L. Hawksworth (Ed.), *Frontiers in mycology* (pp. 187–210). Wallingford: CAB International.
- Ridwan, A. Y., Matoba, R., Wu, J., Choi, J.-H., Hirai, H., & Kawagishi, H. (2018). A novel plant growth regulator from *Pholiota lubrica*. *Tetrahedron Letters*, 59, 2559–2561.
- Ríos-García, U. (2018). *Saberes etnomicológicos en la comunidad mazateca de Eloxochitlán de Flores Magón, Oaxaca*. BSc Dissertation. Oaxaca, Mexico: Instituto Tecnológico del Valle de Oaxaca.
- Rizal, L. M., Hyde, K. D., Chukeatirote, E., Karunarathna, S. C., Kakumyan, P., & Chamyuang, S. (2016). First successful cultivation of the edible mushroom *Macrolepiota dolichaula* in Thailand. *The Chiang Mai Journal of Science*, 43, 959–971.
- Robinson, R. (2003). *Fungi of the South-West Forests*. Kensington, West Australia: Department of Conservation and Land Management.
- Robles-García, D., Suzán-Azpiri, H., Montoya-Esquivel, A., García-Jiménez, J., Esquivel-Naranjo, E. U., Yahia, E., & Landeros-Jaime, F. (2018). Ethnomycological knowledge in three communities in Amealco, Querétaro, México. *Journal of Ethnobiology and Ethnomedicine*, 14, 1–13.
- Rodríguez Hernández, A. (2018). *Aspectos bioculturales de Saproamanita thiersii (Fungi: Agaricales)*. en el municipio de Apaxco de Ocampo, Estado de México.
- Rodríguez, J. A., Llamas-Frade, B., Terrón-Alfonso, A., Sánchez-Rodríguez, J. A., García-Prieto, O., Arrojo-Martín, E., & Jarauta, T. P. (1999). *Guía de Hongos de la Península Ibérica* (3rd ed.). León: Celarayn.
- Rodríguez-Gutiérrez, I. C.-H. L., Delgado-Zúñiga, P., & López-Gómez, F. D. (2017). *Traditional ethnomycological knowledge among the Mexican ethnic group Hiú Hu or Otomí in the communities of Huixquilucan, Mexico*. Texcoco, Mexico: Colegio de Postgraduados.
- Roncero-Ramos, I., & Delgado-Andrade, C. (2017). The beneficial role of edible mushrooms in human health. *Current Opinion in Food Science*, 14, 122–128.
- Roux, J. (2002). Eating *Termitomyces umkowaanii* in South Africa. Personal Communication.
- Ruan-Soto, F. (2018). Sociodemographic differences in the cultural significance of edible and toxic mushrooms among Tsotsil towns in the Highlands of Chiapas, Mexico. *Journal of Ethnobiology and Ethnomedicine*, 14(1), 32.
- Ruan-Soto, F., & Ordaz-Velázquez, M. (2015). Aproximaciones a la etnomicología maya. *Revista pueblos y fronteras digital*, 10, 44–69.
- Ruan-Soto, F., Caballero, J., Martorell, C., Cifuentes, J., González-Esquinca, A. R., & Garibay-Orjiz, R. (2013). Evaluation of the degree of mycophilia-mycophobia among highland and lowland inhabitants from Chiapas, Mexico. *Journal of Ethnobiology and Ethnomedicine*, 9, 36.
- Ruan-Soto, F., Cifuentes, J., Mariaca, R., Limón, F., Pérez-Ramírez, L., & Sierra, S. (2009). Uso y manejo de hongos silvestres en dos comunidades de la Selva Lacandona, Chiapas, México. *Revista Mexicana de Micología*, 29, 61–72.
- Ruan-Soto, F., Pérez, R., Cifuentes, B., Ordaz, V., Cruz, S., García, V., ... del Mariaca, M. (2017). *Hongos de los lacandones de Nahá y Metzabok: Guía ilustrada de macromicetos*. Chiapas, Mexico: San Cristóbal de Las Casas.
- Rubel, W., & Arora, D. (2008). A study of cultural bias in field guide determinations of mushroom edibility using the iconic mushroom, *Amanita muscaria*, as an example. *Economic botany*, 62, 223–243.
- Rzyski, P., & Klimaszyk, P. (2018). Is the yellow knight mushroom edible or not? A systematic review and critical viewpoints on the toxicity of *Tricholoma equestre*. *Comprehensive Reviews in Food Science and Food Safety*, 17, 1309–1324.
- Rzyski, P., Klimaszyk, P., & Benjamin, D. (2019). Comment on “Study of biological activity of *Tricholoma equestre* fruiting bodies and their safety for human”. *European Food Research and Technology*, 245, 963–965.
- Sabra, A., & Walter, S. (2001). Non-wood forest products in the Near East: A regional and national overview. Working paper FOPW/01/2. Rome: FAO.
- Sáenz, J. A., Lizano, A. V. M., & Nassar, M. C. (1983). Edible, poisonous and hallucinatory fungi in Costa Rica. *Revista de Biología Tropical*, 31, 201–207.
- Saha, A. K., Acharya, S., & Roy, A. (2012). Antioxidant level of wild edible mushroom: *Pleurotus djamor* (Fr.) Boedijn. *Journal of Agricultural Technology*, 8, 1343–1351.
- Saito, H. (2006). Wild mushroom use in Japan and its ecological background. *Biostory*, 6, 106–121.
- Saito, T., Tonouchi, A., & Harada, Y. (2014). Biological characteristics and molecular phylogeny of *Sarcomyxa edulis* comb. nov. and *S. serotina*. *Japanese Journal of Mycology*, 55, 19–28.
- Salam, S., & Jamir, N. (2018). Wild Edible Fungi Sold In Local Markets of Ukhrul District of Manipur, India. *International Journal of Scientific and Research Publications*, 8, 208–219.
- Sánchez, C. (2004). Modern aspects of mushroom culture technology. *Applied Microbiology and Biotechnology*, 64, 756–762.
- Sanmee, R., Dell, B., Lumyong, P., Izumori, K., & Lumyong, S. (2003). Nutritive value of popular wild edible mushrooms from northern Thailand. *Food Chemistry*, 82, 527–532.
- Sanon, K. B., Ba, A. M., & Dexheimer, J. (1997). Mycorrhizal status of some fungi fruiting beneath indigenous trees in Burkina Faso. *Forest ecology and Management*, 98, 61–69.
- Santos, F. R. M. (2017). *Manual de Cogumelos comestíveis no Distrito Federal*. Brasília: IESB.

- Sanuma, O. I.; Tokimoto, K., Sanuma, C., Autuori, J., Sanuma, L. R., Sanuma, M., ... Apiamo, R. M. (2016). *Enciclopédia dos alimentos Yanomami (Sanõma): Cogumelos*. São Paulo: Instituto Socioambiental.
- Sarkar, B. B., Chakraborty, D. K., & Bhattacharjee, A. (1988). Wild edible mushroom flora of Tripura. *Indian Agriculturist*, 32, 139–143.
- Sarma, T., Sarma, I., & Patiri, B. (2010). Wild edible mushrooms used by some ethnic tribes of Western Assam. *The Bioscan*, 3, 613–625.
- Sarwar, S., Jabeen, S., Khalid, A. N., & Dentinger, B. M. (2016). Molecular phylogenetic analysis of fleshy pored mushrooms: *Neobolus luridiformis* and *Hortiboletus rubellus* from western Himalayan range of Pakistan. *Pakistan Journal of Botany*, 48, 2077–2083.
- Sarwar, S., Saba, M., Khalid, A., & Dentinger, B. (2018). *Suillus himalayensis* (Boletales: Basidiomycota: Fungi) and its symbiotic association with roots of *Pinus wallichiana*, first report from coniferous forests of Pakistan. *Journal of Animal and Plant Sciences*, 28, 576–583.
- Sato, H., & Hattori, T. (2015). New species of *Boletellus* section *Boletellus* (Boletaceae, Boletales) from Japan, *B. aurocontextus* sp. nov. and *B. areolatus* sp. nov. *PLoS One*, 10, e0128184.
- Saupe, S. G. (1981). Occurrence of psilocybin/psilocin in *Pluteus salicinus* (Pluteaceae). *Mycologia*, 73, 781–784.
- Saviuc, P., Moreau, P., Fouilhe, S.-L. N., Gamelin, L., Pulce, C., Butler, J., & Danel, V. (2008). Four cases of poisoning by *Entoloma vernum*. *Clinical Toxicology*, 46, 408–408.
- Schenk-Jaeger, K. M., Rauber-Lüthy, C., Bodmer, M., Kupferschmidt, H., Kullak-Ublick, G. A., & Ceschi, A. (2012). Mushroom poisoning: A study on circumstances of exposure and patterns of toxicity. *European Journal of Internal Medicine*, 23, e85–e91.
- Schmeda-Hirschmann, G., Razmilic, I., Gutierrez, M. I., & Loyola, J. I. (1999b). Proximate composition and biological activity of food plants gathered by Chilean Amerindians. *Economic Botany*, 53, 177–187.
- Schmeda-Hirschmann, G., Razmilic, I., Reyes, S., Gutierrez, M. I., & Loyola, J. I. (1999a). Biological activity and food analysis of *Cyttaria* spp. (Discomycetes). *Economic Botany*, 53, 30–40.
- Secretariat of Environment and Natural Resources. (2020). Retrieved from <https://www.gob.mx/semarnat>.
- Şelem, E., Keleş, A., İsmail, A., & Demirel, K. (2019). Edible macrofungi determined in Gürpınar (Van) district. *Anatolian Journal of Botany*, 3, 7–12.
- Semwal, K., Stephenson, S., Bhatt, V., & Bhatt, R. (2014). Edible mushrooms of the Northwestern Himalaya, India: A study of indigenous knowledge, distribution and diversity. *Mycosphere*, 5, 440–461.
- Seok, S.-J., Jin, Y.-J., Kwon, S.-W., Kim, Y.-S., & Kim, W.-G. (2013). A taxonomic study of genus *Melanophyllum* in Korea. *The Korean Journal of Mycology*, 41, 205–211.
- Seok, S.-J., Kim, Y.-S., Weon, H.-Y., Lee, K.-H., Park, K.-M., Min, K.-H., & Yoo, K.-H. (2002). Taxonomic study on *Volvariella* in Korea. *Mycobiology*, 30, 183–192.
- Sergeeva, M. (2000). *Fungi. 250 species of edible, poisonous and medicinal fungi*. Moscow: Culture and Traditions.
- Sevindik, M. (2020). Poisonous mushroom (nonedible) as an antioxidant source. In H. M. Ekiert & K. G. Ramawat (Eds.), *Plant Antioxidants and Health* (pp. 1–25). Cham, Switzerland: Springer International Publishing.
- Sevindik, M., & Akata, I. (2020). Antioxidant, oxidant potentials and element content of edible wild mushroom *Helvella leucopus*. *Indian Journal of Natural Products and Resources (IJNPR)[Formerly Natural Product Radiance (NPR)]*, 10, 266–271.
- Sevindik, M., Rasul, A., Hussain, G., Anwar, H., Zahoor, M. K., Sarfraz, I., ... Selamoglu, Z. (2018). Determination of anti-oxidative, anti-microbial activity and heavy metal contents of *Leucoagaricus leucothites*. *Pakistan Journal of Pharmaceutical Sciences*, 31, 2163–2168.
- Sharda, R. M., Kaushal, S. C., & Negi, G. S. (1997). Edible fungi of Garhwal Himalayas. *Mushroom Journal*, 1997, 11–13.
- Sharma, J. (2018). *Medicinal plants and mushrooms of India*. New Delhi: Ane Books Pvt. Ltd.
- Sharma, S., & Singh, R. (2016). Mushroom poisoning: From toxicity to forensic analysis. *Journal of the Indian Society of Toxicology*, 12, 43–48.
- Shaw, D. (1984). Microorganisms in Papua New Guinea. Research Bulletin No.33, Department of Primary Industry, Port Moresby.
- Shepard, G. H., Arora, D., & Lampman, A. (2008). The grace of the flood: Classification and use of wild mushrooms among the highland Maya of Chiapas. *Economic botany*, 62, 437–470.
- Shernoff, L. (2015). Mushroom edibility, by the numbers. *Mushroom the Journal*, 2015, 59–65.
- Shibata, H. (1992a). Higher basidiomycetes from Pakistan. In T. Nakaike & S. Malik (Eds.), *Cryptogamic flora of Pakistan* (Vol. 1, pp. 145–163). Tokyo: National Science Museum.
- Shibata, H. (1992b). Higher basidiomycetes from Pakistan. *Cryptogamic flora of Pakistan*, 1, 145–164.
- Shirouzu, T., Inaba, S., Ushijima, S., Okuda, Y., & Nagasawa, E. (2018). Taxonomic study of Japanese “*Auricularia auricula-judae*” and “*A. polytricha*” based on molecular phylogeny and morphological observation. *Japanese Journal of Mycology*, 59, 7–20.
- Shmidt, O. (2006). *Wood and Tree Fungi. Biology, damage, protection and use*. Germany: Springer-Verlag Berlin Heidelberg.
- Shrestha, S., & Rai, S. K. (2012). Survey of marketable vegetables and edible fruits in Dharan, eastern Nepal. *Nepalese Journal of Biosciences*, 2, 134–147.
- Sibounnavong, P., Cynthia, C., Kalaw, S., Reyes, R., & Soyong, K. (2008). Some species of macrofungi at Puncan, Carranglan, Nueva Ecija in the Philippines. *Journal of Agricultural Technology*, 4, 105–115.
- Siddiqi, N. A. (1998). *Ethnobotany of non-timber forest products of Chittagong hill tracts*. In R. L. Banik, M. K. Alam, S. J. Pei, & A. Rastogi (Eds.), *Applied ethnobotany* (pp. 52–55). Chittagong, Bangladesh: Bangladesh Forest Research Institute.
- Sillitoe, P. (1995). Ethnoscience observations on entomology and mycology in the southern highlands of Papua New Guinea. *Science in New Guinea*, 21, 3–26.
- Šíma, J., Vondruška, J., Svoboda, L., Šeda, M., & Rokos, L. (2019). The accumulation of risk and essential elements in edible mushrooms *Chlorophyllum rhacodes*, *Suillus grevillei*, *Imleria badia*, and *Xerocomellus chrysenteron* growing in the Czech Republic. *Chemistry & Biodiversity*, 16, e1800478.
- Simmons, C., Henkel, T., & Bas, C. (2002). The genus *Amanita* in the Pakaraima mountains of Guyana. *Persoonia*, 17, 563–582.
- Singer, R. (1953). Four years of mycological work in southern South America. *Mycologia*, 45, 865–891.
- Singer, R. (1986). *The Agaricales in modern taxonomy* (3rd ed.). Koenigstein: Koeltz Scientific Books.

- Skariyachan, S., Prasanna, A., Manjunath, S. P., Karanth, S. S., & Nazre, A. (2016). Exploring the medicinal potential of the fruit bodies of oyster mushroom, *Pleurotus ostreatus* (Agaricomycetes), against multidrug-resistant bacterial isolates. *International Journal of Medicinal Mushrooms*, 18(3), 245–252.
- Smith, D. (2020). Mycology lessons, articles, photos and descriptions of northeastern fungi. Retrieved from <http://www.fungikingdom.net/fungi-photos/basidiomycota/boletales-order/boletaceae-family/Lanmaoa-pseudosensibilis.html>.
- Species Fungorum. (2020). Retrieved from <http://www.speciesfungorum.org>.
- Spegazzini, C. (1891). Fungi guaranitici nonnulli novi v. critici: Coni é hijos. *Revista Argent. Hist. Nat.*, 1, 101–111.
- Spooner, B., & Laessoe, T. (1994). The folklore of ‘Gasteromycetes’. *Mycologist*, 8, 119–123.
- Stamets, P. (1993). *Growing gourmet and medicinal mushrooms*. Berkeley, CA: Ten Speed Press.
- Stojković, D., Reis, F. S., Barros, L., Glamočlija, J., Ćirić, A., van Griensven, L. J., ... Ferreira, I. C. (2013). Nutrients and non-nutrients composition and bioactivity of wild and cultivated *Coprinus comatus* (OF Müll.) Pers. *Food and Chemical Toxicology*, 59, 289–296.
- Stojkovic, D., Smiljkovic, M., Ćiric, A., Glamočlija, J., Van Griensven, L., Ferreira, I. C., & Sokovic, M. (2019). An insight into antidiabetic properties of six medicinal and edible mushrooms: Inhibition of α -amylase and α -glucosidase linked to type-2 diabetes. *South African Journal of Botany*, 120, 100–103.
- Straus, L. G., Morales, M. R. G., Carretero, J. M., & Marín-Arroyo, A. B. (2015). “The Red Lady of El Mirón”. Lower Magdalenian life and death in Oldest Dryas Cantabrian Spain: An overview. *Journal of Archaeological Science*, 60, 134–137.
- Suabjakyong, P., Saiki, R., Van Griensven, L. J., Higashi, K., Nishimura, K., Igarashi, K., & Toida, T. (2015). Polyphenol extract from *Phellinus igniarius* protects against acrolein toxicity in vitro and provides protection in a mouse stroke model. *PLoS One*, 10, e0122733.
- Sultana, K., Rauf, C. A., Riaz, A., Naz, F., Irshad, G., & Haque, M. I. U. (2011). Checklist of Agarics of Kaghan valley-1. *Pakistan Journal of Botany*, 43, 1777–1787.
- Sun, L.-b., Zhang, Z.-y., Xin, G., Sun, B.-x., Bao, X.-j., Wei, Y.-y., ... Xu, H.-r (2020). Advances in umami taste and aroma of edible mushrooms. *Trends in Food Science & Technology*, 96, 176–187.
- Survival Use of Plants, (2020). Retrieved from <http://www.survivalebooks.com/09.htm#par1>.
- Swenberg, M.-L. L., Kelleher, W., & Schwarting, A. (1967). Muscarine: Isolation from cultures of *Clitocybe rivulosa*. *Science*, 155, 1259–1259.
- Syed-Riaz, A. G., & Mahmood-Khan, S. (1999). Edible mushrooms from Azad Jammu and Kashmir. *Pakistan Journal of Phytopathology*, 11, 163–165.
- Takabatake, K. (2015). Current trends and future prospects for mushroom production in Japan. *Mokuzai Gakkaishi*, 61, 243–249.
- Takeuchi, T. (1997). Studies on cultivation of *Pholiota lubrica* -on the trial cultivation on the raw wood and saw dust medium. *Applied Forest Science*, 6, 205–206.
- Tantengco, O., & Rragio, E. (2018). Ethnomycological survey of macrofungi utilized by Ayta communities in Bataan, Philippines. *Journal of Fungal Biology*, 8, 104–108.
- Taylor, F. W., Thamage, D. M., Baker, N., Roth-Bejerano, N., & Kagan-Zur, V. (1995). Notes on the Kalahari desert truffle, *Terfezia pfeillii*. *Mycological Research*, 99, 874–878.
- Tedder, S., Mitchell, D., & Farran, R. (2000). *Seeing the forest beneath the trees: The social and economic potential of non-timber forest products and services in the Queen Charlotte Islands/Haida Gwaii*. British Columbia: Mitchell Consulting and the B.C. Ministry of Forests.
- Tel, G., Deveci, E., Küçükaydın, S., Özler, M. A., Duru, M. E., & Harmandar, M. (2014). Evaluation of antioxidant activity of *Armillaria tabescens*, *Leucopaxillus gentianeus* and *Suillus granulatus*: The mushroom species from Anatolia. *Eurasian Journal of Analytical Chemistry*, 8, 136–147.
- Tham, L. X., Chuong, N. N., Duong, P. N., & Thiem, B. H. (2011). The biogeographical speciations of shiitake *Lentinula edodes* and a new species *Lentinula platinedodes* sp. nov. found in Cat Tien, South Vietnam. *Tap Chi Sinh Hoc*, 33, 29–39.
- Thawthong, A., Hapuarachchi, K., Wen, T., Raspé, O., Thongklang, N., Kang, J., & Hyde, K. (2017). *Ganoderma sichuanense* (Ganodermataceae, Polyporales) new to Thailand. *MycKeys*, 22, 27–43.
- Thawthong, A., Karunarathna, S. C., Thongklang, N., Chukeatirote, E., Kakumyan, P., Chamyuang, S., ... Callac, P. (2014). Discovering and domesticating wild tropical cultivatable mushrooms. *Chiang Mai Journal of Science*, 41, 731–764.
- Thoen, D., & Ba, A. M. (1989). Ectomycorrhizas and putative ectomycorrhizal fungi of *Azelia africana* and *Uapaca senegalensis* in southern Senegal. *New Phytologist*, 113, 549–559.
- Thongbai, B., Rapior, S., Hyde, K. D., Wittstein, K., & Stadler, M. (2015). *Hericium erinaceus*, an amazing medicinal mushroom. *Mycological Progress*, 14, 91.
- Thongklang, N., Thongbai, B., Chamyuang, S., Callac, P., Chukeatirote, E., Hyde, K. D., ... Stadler, M. (2017). Blazeispirol A, a chemotaxonomic marker from mycelia of the medicinal mushroom *Agaricus subrufescens*. *Chiang Mai Journal of Science*, 44, 298–308.
- Thu, Z. M., Myo, K. K., Aung, H. T., Clericuzio, M., Armijos, C., & Vidari, G. (2020). Bioactive phytochemical constituents of wild edible mushrooms from southeast Asia. *Molecules*, 25, 1972.
- Timm, J. F. (2018). *O livro Primavera Fungi – Guia de Fungos do Sul do Brasil*. Porto Alegre: Via Sapiens.
- Toledo, C., Barroetaveña, C., & Rajchenberg, M. (2016). Hongos comestibles silvestres de los bosques nativos de la región Andino Patagónica de Argentina. Manual-CIEFAP (Chubut-Argentina). (2016). (no 16) 71 p.
- Tom Volk's Fungi (2020). TomVolkFungi.net.
- Tominaga, Y., Xian, M. Y., Liu, F. X., & Tan, W. (1989). Matsu-take mushrooms in the People's Republic of China. III Matsu-take mushrooms in the Yanyuan and Dechjang districts of Sichuan Province. *Bulletin of the Hiroshima Agriculture College*, 8, 733–742.
- Toshinungla, A., Deb, C. R., & Neilazonuo, K. (2016). Wild edible mushrooms of Nagaland, India: A potential food resource. *Journal of Experimental Biology and Agricultural Sciences*, 4, 59–65.
- Trappe, J. M. (1990). Use of truffles and false truffles around the world. In M. Bencivenga & B. Granetti (Eds.), *Proceedings, Atti*

- del Segundo Congreso Internacional sul Tartufo. Spoleto 1988 (pp. 19–30). Spoleto, Italy: Comunita Montana dei Monti Martini e del Serano.
- Trierveiler-Pereira, L., Sulzbacher, M., & Baltazar, J. (2018). Diversidade de fungos brasileiros e alimentação: O que podemos consumir? In Conference III Fórum Ambiental de Angatuba At: Angatuba-SP. Retrieved from <https://www.researchgate.net/publication/325736818>.
- Trudell, S. A., Xu, J., Saar, I., Justo, A., & Cifuentes, J. (2017). North American matsutake: Names clarified and a new species described. *Mycologia*, *109*, 379–390.
- Tu, G. L. (1987). Using bagasse and waste cotton as substrate for bag cultivation of *Pleurotus sajor-caju*. *Zhongguo Shiyongjun (Edible Fungi of China)*, *6*(23), 30–34.
- Tudor, I. (2010). *Ciupercile din flora spontană a României (Mushrooms from the spontaneous flora of Romania)*. Bucharest: Gramen Editura.
- Tulloss, R., Iqbal, S., Khalid, A., Bhatt, R., & Bhat, V. (2001). Studies in *Amanita* (Amanitaceae) from southern Asia. I. Some species of Pakistan's Northwest Frontier Province. *Mycotaxon*, *77*, 455–490.
- Tuno, N. (2001). Mushroom utilization by the Majangir, an Ethiopian tribe. *The Mycologist*, *15*, 78–79.
- Uaciquete, A., Dai, M. d. L., & Motta, H. (1996). *Distribuição, valor econômico e uso sustentável do cogumelo comestível em Moçambique [Distribution, economic value and sustainable use of edible mushrooms in Mozambique]*. Maputo, Moçambique: Grupo de Trabalho Ambiental [Environmental Working Group].
- Ullah, T. S., Firdous, S. S., Mehmood, A., Shaheen, H., & Dar, M. E. U. I. (2017). Ethnomycological and nutritional analyses of some wild edible mushrooms from Western Himalayas, Azad Jammu and Kashmir (Pakistan). *International Journal of Medicinal Mushrooms*, *19*(10), 949–955.
- Umagat, M. R., Dulay, R. M. R., Kalaw, S. P., Abon, M. D., & Reyes, R. G. (2016). Mineral composition of different species of *Pleurotus* commercially grown on rice straw-based formulation in Central Luzon, Philippines. *Advances in Environmental Biology*, *10*, 51–56.
- Ushijima, S., Shimomura, N., Nagasawa, E., & Maekawa, N. (2012). Taxonomic reevaluation of a fungus described as "*Mucidula mucida*" in Japan. *Mushroom Science and Biotechnology*, *20*, 22–30.
- Van-Dijk, H., Onguene, N. A., & Kuyper, T. W. (2003). Knowledge and utilization of edible mushrooms by local populations of the rain forest of South Cameroon. *AMBIO: A Journal of the Human Environment*, *32*, 19–23.
- Ványolós, A., Muszyńska, B., Chuluunbaatar, B., Gdula-Argasińska, J., Kała, K., & Hohmann, J. (2020). Extracts and steroids from the edible mushroom *Hypholoma lateritium* exhibit anti-inflammatory properties by inhibition of COX-2 and activation of Nrf2. *Chemistry & Biodiversity*, *17*, e2000391.
- Varev, S., Fakirova, V., & Iordanov, D. (1998). Edible and poisonous mushrooms (fungi) in Bulgaria. *An identification key* (revised and supplemented edition). Sofiya: Bulgarian Academy of Sciences, Botanic Institute.
- Vargas, N., & Restrepo, S. (2020). A checklist of ectomycorrhizal mushrooms associated with *Quercus humboldtii* in Colombia. In J. Perez-Moreno A. Guerin-Laguette R. Flores Arzu & F. Q. Yu (Eds.), *Mushrooms, humans and nature in a changing world* (pp. 425–450). Cham, Switzerland: Springer International Publishing.
- Vargas, N., Bernal, A., Sarria, V., Franco-Molano, A., & Restrepo, S. (2011). Amatoxin and phallotoxin composition in species of the genus *Amanita* in Colombia: A taxonomic perspective. *Toxicon*, *58*, 583–590.
- Vargas-Isla, R., Ishikawa, N. K., & Py-Daniel, V. (2013). Contribuições etnomicológicas dos povos indígenas da Amazônia. *Biota Amazônia (Biota Amazonie, Biota Amazonia, Amazonian Biota)*, *3*, 58–65.
- Varghese, S. P., Pradeep, C., & Vrinda, K. (2010). Mushrooms of tribal importance in Wayanad area of Kerala. *Journal of Mycopathological Research*, *48*, 311–320.
- Vasco-Palacios, A. M., Suaza, S. C., Castañõ-Betancur, M., & Franco-Molano, A. E. (2008). Conocimiento etnoecológico de los hongos entre los indígenas Uitoto, Muinane y Andoke de la Amazonía Colombiana. *Acta Amazonica*, *38*, 17–30.
- Vasil'eva, L. N. (1978). *Edible mushrooms of the Far East*. Vladivostock: Far Eastern Publishing Houes.
- Verma, R. K., Pandro, V., & Pyasi, A. (2017). Diversity and distribution of *Russula* in India with reference to Central Indian species. *International Journal of Current Microbiology and Applied Sciences*, *7*, 3078–3103.
- Vidal, J. (1959). Noms vernaculaires de plantes (Lao, Meo, Kha) en usage au Laos. *Bulletin de l'École française d'Extrême-Orient*, *49*, 435–608.
- Villarreal, L., & Perez-Moreno, J. (1989a). Los hongos comestibles silvestres de Mexico, un enfoque integral. *Micologia Neotropica Aplicada*, *2*, 77–114.
- Villarreal, L., & Pérez-Moreno, J. (1989b). Aprovechamiento y conservación del 'Matsutake Americano' (*Tricholoma magnelivare*) en los bosques de Mexico. *Micologia Neotropica Aplicada*, *2*, 131–144.
- Vishwakarma, M. P., Bhatt, R. P., & Joshi, S. (2012). Macrofungal diversity in moist temperate forests of Garhwal Himalaya. *Indian Journal of Science & Technology*, *5*, 1928–1932.
- Walley, R., & Rammeloo, J. (1994). The poisonous and useful fungi of Africa south of the Sahara: A literature survey. *Scripta Botanica Belgica*, *6*, 1–56.
- Walter, S. (2001). *Non-wood forest products in Africa: A regional and national overview*. Rome: FAO.
- Walton, J. D., Hallen-Adams, H. E., & Luo, H. (2010). Ribosomal biosynthesis of the cyclic peptide toxins of *Amanita* mushrooms. *Peptide Science*, *94*, 659–664.
- Wang, Q., Sun, M., Lv, H., Lu, P., Ma, C., Liu, Y., ... Gao, Y. (2020). *Amanita fuliginea* poisoning with thrombocytopenia: A case series. *Toxicon*, *174*, 43–47.
- Wang, T.-t., Zhao, C.-y., Chen, X., Guo, X., & Wu, S.-r. (2017). Analysis and recommendations on current status of Yunnan Province edible fungi industry during the "12th five year" period. *Edible fungi of China*, *36*, 70–75.
- Wang, X., & Liu, P. (2002). Resources investigation and studies on the wild commercial fungi in Yunnan. *Biodiversity Science*, *10*, 318–325.
- Wang, X., Liu, P., & Yu, F. (2004). *Color atlas of wild commercial mushrooms in Yunnan*. Yunnan: Yunnan Science and Technology Press.
- Wang, Y., Hall, I. R., & Evans, L. A. (1997). Ectomycorrhizal fungi with edible fruiting bodies. 1. *Tricholoma matsutake* and related fungi. *Economic-Botany*, *51*, 311–327.
- Wasser, S. (2002). Medicinal mushrooms as a source of antitumor and immunomodulating polysaccharides. *Applied Microbiology and Biotechnology*, *60*, 258–274.

- Wasser, S. P. (2002). *Edible fungi Israel, Ukraine and others*. Personal Communication.
- White, J., Weinstein, S. A., De Haro, L., Bédry, R., Schaper, A., Rumack, B. H., & Zilker, T. (2019). Mushroom poisoning: A proposed new clinical classification. *Toxicon*, *157*, 53–65.
- Wieland, T. (2012). *Peptides of poisonous Amanita mushrooms*. New York, Berlin: Springer Science & Business Media.
- Wijayawardene, N. N., Hyde, K. D., Rajeshkumar, K. C., Hawksworth, D. L., Madrid, H., Kirk, P. M., ... Kukwa, M. (2017). Notes for genera: Ascomycota. *Fungal Diversity*, *86*, 1–594.
- Wilson, K., Cammack, D., & Shumba, F. (1989). *Food provisioning amongst Mozambican refugees in Malawi*. A study of aid, livelihood and development. A report prepared for the World Food Programme. Oxford: Oxford University.
- Winkler, D. (2002). Forest use and implications of the 1998 logging ban in the Tibetan prefectures of Sichuan: Case study on forestry, reforestation and NTFP in Litang County, Ganzi TAP, China. In Z. Ziang, M. Centritto, S. Liu, & S. Zhang (Eds.), *The ecological basis and sustainable management of forest resources* (pp. [in press]): *Informatore Botanico Italiano 134* (Supplemento 2).
- Winkler, D. (2002). Matsutake Initiative – Proposal for action. Open Letter.
- Wu, F., Zhou, L.-W., Yang, Z.-L., Bau, T., Li, T.-H., & Dai, Y.-C. (2019). Resource diversity of Chinese macrofungi: Edible, medicinal and poisonous species. *Fungal Diversity*, *2019*, 1–76.
- Wu, X., Dai, Y., Li, T., Yang, Z., & Song, B. (2011). *Fungi of tropical China*. Beijing: Science Press.
- Xiang, Y. T., & Han, Z. (1987). Using sun-cured bed to increase temperature in the early spring for culturing straw mushroom (*Volvariella esculenta*). *Zhongguo Shiyongjun (Edible Fungi of China)*, *6*(23), 16–17.
- Yamada, A. (2002). Utility of mycorrhizal mushrooms as food resources in Japan. *Journal of the Faculty of Agriculture*, *38*, 1–17.
- Yamada, M., Tokumitsu, N., Saikawa, Y., Nakata, M., Asano, J., Miyairi, K., ... Hashimoto, K. (2012). Molybdophyllysin, a toxic metalloendopeptidase from the tropical toadstool, *Chlorophyllum molybdites*. *Bioorganic & Medicinal Chemistry*, *20*, 6583–6588.
- Yanaga, K., Sotome, K., Ushijima, S., & Maekawa, N. (2015). *Hydnum* species producing whitish basidiomata in Japan. *Mycoscience*, *56*, 434–442.
- Yang, M., Belwal, T., Devkota, H. P., Li, L., & Luo, Z. (2019). Trends of utilizing mushroom polysaccharides (MPs) as potent nutraceutical components in food and medicine: A comprehensive review. *Trends in Food Science & Technology*, *92*, 94–110.
- Yang, S. (1990). Effect of boron and zinc on growth of mycelia of *Pleurotus florida*. *Zhongguo Shiyongjun (Edible Fungi of China)*, *9*, 19–20.
- Yang, Z. (1990). A delicious tropical mushroom – *Termitomyces heimii* occurring in Yunnan, China. *Zhongguo Shiyongjun (Edible Fungi of China)*, *9*, 28–30.
- Yang, Z. (1992). *Polyozellus multiplex* – A rare edible fungus. *Zhongguo Shiyongjun (Edible Fungi of China)*, *11*, 1–4.
- Yang, Z. L., & Yang, C. (1992). Recognition of *Hypsizygus marmoratus* and its cultivation. *Zhongguo Shiyongjun (Edible Fungi of China)*, *11*, 19–20.
- Yaseen, T., Khan, M., Shakeel, M., & Ara, H. (2016). Morphological study of edible and non-edible mushrooms, district Swabi, Khyber Pakhtunkhwa, Pakistan. *Pure and Applied Biology*, *5*, 889–894.
- Ye, L., Zheng, X., Zhang, J., Tang, Q., Yang, Y., Wang, X., ... Pan, Y. (2011). Biochemical characterization of a proteoglycan complex from an edible mushroom *Ganoderma lucidum* fruiting bodies and its immunoregulatory activity. *Food Research International*, *44*, 367–372.
- Yeh, E. (2000). Forest claims, conflicts and commodification: The political ecology of Tibetan mushroom-harvesting villages in Yunnan Province, China. *China Quarterly*, *161*, 225–278.
- Yetter, G. (2002). *Edible fungi from Haiti; for sale in Brooklyn; link to eating Psathyrella in Africa*. Personal Communication.
- Yilmaz, F., Oder, N., & Isiloglu, M. (1997). The macrofungi of the Soma (Manisa) and Savastepe (Balikesir) districts. *Turkish Journal of Botany*, *21*, 221–230.
- Yin, X., Yang, A.-A., & Gao, J.-M. (2019). Mushroom Toxins: Chemistry and Toxicology. *Journal of agricultural and food chemistry*, *67*, 5053–5071.
- Yongabi, K., Agho, M., & Martínez-Carrera, D. (2004). Ethnomycological studies on wild mushrooms in Cameroon, Central Africa. *Micologia Aplicada International*, *16*, 34–36.
- Yorou, S. (2002). *Selling wild edible fungi in Benin*. Personal Communication.
- Yorou, S. N., & De-Kesel, A. (2002). Connaissances ethnomycologiques des peuples Nagot du centre du Bénin (Afrique de l'Ouest). In E. Robbrecht, J. Degreef, & I. Friis (Eds.), *Plant systematics and phytogeography for the understanding of African Biodiversity*. Proceedings of the XVth AETFAT Congress (2000, Meise, National Botanic Garden of Belgium) (Vol. 71, pp. 627–637): *Systematics and Geography of Plants*.
- Yorou, S. N., De-Kesel, A., Sinsin, B., & Codjia, J. T. C. (2002). Diversité et productivité des champignons comestibles de la forêt classée de Wari Maro (Benin, Afrique de l'Ouest). In E. Robbrecht, J. Degreef, & I. Friis (Eds.), *Plant systematics and phytogeography for the understanding of African biodiversity*. Proceedings of the XVIth AETFAT Congress (2000, Meise, National Botanic Garden of Belgium) (Vol. 71, pp. 613–625): *Systematics and Geography of Plants*.
- Zahid, M. K., Barua, S., & Haque, S. I. (2009). Proximate composition and mineral content of selected edible mushroom varieties of Bangladesh. *Bangladesh Journal of Nutrition*, *22–23*, 61–68.
- Zakhary, J. W., Abo-Bakr, T. M., El-Mahdy, A. R., & El-Tabery, S. A. M. (1983). Chemical composition of wild mushrooms collected from Alexandria, Egypt. *Food Chemistry*, *11*, 31–41.
- Zaklina, M. (1998). Edible mycorrhizal mushrooms in Serbia - problems with protection. 2nd International Conference on Mycorrhiza, Upsala, Sweden, 5–10 July 1998. Retrieved from www.mycorrhiza.ag.utk.edu.
- Zamora-Martinez, M. C., Alvarado, G., & Dominguez, J. M. (2000). *Hongos Silvestres Comestibles region de Zacualtipan, Hidalgo*. Pachuca, Hidalgo, Mexico: INIFAP CIR-CENTRO#.
- Zamora-Martinez, M. C., Reygadas, G. F., & Cifuentes, J. (1994). *Hongos comestibles silvestres de la subcuena Arroyo El Zorrillo, Distrito Federal*. Coyoacan, DF Mexico: INIFAP.
- Zamora-Martinez, M., Nieto de Pascual-Pola, M. C. d. C., Sánchez-Colín, R., García Campusano, F., & Zamora-Morales, B. (2018). *Hongos comestibles de Zacatlán de Las Manzanas, Puebla (II)*. INIFAP. Folleto técnico 29. Mexico City, 81.
- Zang, M. (1984). Mushroom distribution and the diversity of habitats in Tibet, China. *McIlvainea*, *6*, 15–20.

- Zang, M. (1988). An interesting edible mushroom: *Agaricus gennadii*. *Zhongguo Shiyongjun (Edible Fungi of China)*, 7(32), 3–4.
- Zang, M., & Doi, Y. (1995). *Secotium jimalaicum* sp. nov. from Nepal – A folklore concerning the food of abominable snowman. *Acta Botanica Yunnanica*, 17, 30–32.
- Zang, M., & Petersen, R. (1990). An endemic and edible fungus – *Endophyllus yunnanensis* from China. *Zhongguo Shiyongjun (Edible Fungi of China)*, 9, 3–5.
- Zang, M., & Pu, C. (1992). Confirmatory *Tuber indica* distributed in China. *Zhongguo Shiyongjun (Edible Fungi of China)*, 11, 19.
- Zang, M., & Yang, Z. L. (1991). *Agrocybe salicicola*, a delicious edible mushroom newly discovered from Yunnan. *Zhongguo Shiyongjun (Edible Fungi of China)*, 10, 18.
- Zent, E. L., Zent, S., & Iturriaga, T. (2004). Knowledge and use of fungi by a mycophilic society of the Venezuelan Amazon. *Economic Botany*, 58, 214–226.
- Zero emissions research and initiatives. (2020). Retrieved from <http://www.zeri.org/>.
- Zerova, M. Y., & Rozhenko, G. L. (1988). *Atlas s'edobnykh i yadovitykh gribov [Atlas of edible and poisonous fungi – Ukraine]*. Kiev, Ukraine: Radyans'ka shkola.
- Zervakis, G. (2003). *Wild edible fungi collected and eaten in Kalamata, southwestern Peloponnese*. Personal Communication.
- Zervakis, G. I., Bekiaris, G., Tarantilis, P. A., & Pappas, C. S. (2012). Rapid strain classification and taxa delimitation within the edible mushroom genus *Pleurotus* through the use of diffuse reflectance infrared Fourier transform (DRIFT) spectroscopy. *Fungal Biology*, 116, 715–728.
- Zervakis, G. I., Polemis, E., & Dimou, D. M. (2002). Mycodiversity studies in selected ecosystems of Greece: III Macrofungi recorded in *Quercus* forests from Southern Peloponnese. *Mycotaxon*, 84, 141–162.
- Zhang, J., Chen, Q., Huang, C., Gao, W., & Qu, J. (2015). History, current situation and trend of edible mushroom industry development. *Mycosystema*, 34, 524–540.
- Zhang, N., Chen, H., Zhang, Y., Xing, L., Li, S., Wang, X., & Sun, Z. (2015). Chemical composition and antioxidant properties of five edible Hymenomycetes mushrooms. *International Journal of Food Science & Technology*, 50, 465–471.
- Zhang, P., Tang, L.-P., Cai, Q., & Xu, J.-P. (2015). A review on the diversity, phylogeography and population genetics of *Amanita* mushrooms. *Mycology*, 6, 86–93.
- Zhou, L.-W., Ghobad-Nejhad, M., Tian, X.-M., Wang, Y.-F., & Wu, F. (2020). Current status of 'Sanghuang' as a group of medicinal mushrooms and their perspective in industry development. *Food Reviews International*, 1–19. <https://doi.org/10.1080/87559129.2020.1740245>
- Zhou, X., Lin, J., Li, Q., Sun, X., & Tang, K. (2007). Study progress on bioactive proteins from *Ganoderma* spp. *Natural Product Research & Development*, 19, 917–924.
- Zhou, W., & Zhang, D. (2005). *Mushrooms of Taiwan*. Taipei: Yuan-Liou Publishing.
- Zhuang, Y. (1993). Characterization and textual criticism of Huai'er (*Trametes robiniophila*). *Zhongguo Shiyongjun (Edible Fungi of China)*, 13, 22–23.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

How to cite this article: Li H, Tian Y, Menolli N, et al. Reviewing the world's edible mushroom species: A new evidence-based classification system. *Compr Rev Food Sci Food Saf*. 2021;1–33. <https://doi.org/10.1111/1541-4337.12708>