


REVIEW ARTICLE

Nutritional and Therapeutic Potential of Garlic and Onion (*Allium* sp.)

Prakash S. Bisen^{1*} and Mila Emerald²

¹*School of Studies in Biotechnology, Jiwaji University, Gwalior 474005, India;* ²*PHYTOCEUTICALS International & NOVOTEK Global Solutions, London, Ontario, Canada*

Abstract: Genus *Allium* belongs to the family Liliaceae, which contains more than 600 species. Garlic (*Allium sativum*) and onion (*Allium cepa*) are two the most popular food ingredients widely used all over the world. During the last few decades, garlic and onion have received tremendous attention for their wide range of therapeutic properties and great health benefits. The variety of garlic and onion species' considerable differences in manufacturing process cause discrepancies in a spectrum of the ingredients derived. Current garlic preparations available on the market, including garlic powder, garlic oil, raw or cooked garlic and aged garlic extract. Garlic and onion extracts possess many therapeutic properties including antimicrobial, antiviral, antifungal, anti-protozoal, hepatoprotective, cardioprotective, anti-inflammatory, neuroprotective, anti-amnesic, anticarcinogenic, antimutagenic, antiasthmatic, immunomodulatory, hypolipidemic, anti-hypertensive, anti-diabetic and antioxidant. These therapeutic properties are caused by the combination and biological activity of organo-sulphur compounds such as S-allyl-l-cysteine, diallyl disulfide, diallyl trisulfide, ajoene, and allicin. Allicin, which is one of the most researched therapeutic compounds of garlic and onion, is extremely unstable and rapidly degrades with time, even at low temperatures, which causes its prompt degradation during contact with stomach acid during oral consumption. Present review discusses biochemical, pharmacological, therapeutic properties and nutritional value of garlic and onion, their use for prevention disease and maintenance of good health, as well as novel potential nanoparticles drug delivery systems for more effective oral and topical administration of natural organo-sulphur compounds.



ARTICLE HISTORY

Received: September 21, 2015
Revised: April 29, 2016
Accepted: May 26, 2016

DOI:
10.2174/1573401312666160608121
954

Keywords: *Allium cepa*, *Allium sativum*, garlic, onion, allicin, organosulfur, flavonoids, sterols.

INTRODUCTION

Genus *Allium* belongs to the family Liliaceae having more than 700 species, differing in taste, form and colour, but have quite similar nutraceutical, phytochemical and biochemical properties [1, 2]. Garlic (*Allium sativum*) and onion (*Allium cepa*) are the two most popular food ingredients used widely all over the world. Crops are usually used fresh after harvest and/or dried and are ready for storage. During the last few decades, garlic and onion have received tremendous attention to their wide range of therapeutic properties and great health benefits as functional food [3-11]. The variety of garlic and onion species' considerable differences in manufacturing process cause discrepancies in a spectrum of the ingredients derived. Numerous meta studies have shown a protective relationship between high consumption of *Allium* sp. vegetable and the relative risk of both prevention of cancer and cardiovascular disorders [12-16]. Onion with estimated annual production over 47 million tons a year

is usually consumed as fresh, and its commercial products are less abundant than those of garlic. Current garlic preparations available on the market include garlic powder, garlic oil, raw or cooked garlic and aged garlic extract. Garlic and onion extracts demonstrate strong antioxidant, hypocholesterolemic, hypolipidemic, anti-hypertensive, anti-diabetic, antithrombotic, anti-inflammatory, anti-spasmodic, anti-allergic, anti-viral, anti-protozoa, anti-hyperhomocysteinemic, anti-parasitic and neuroprotective properties [1, 15, 17]. Both plants are used in the prevention of vascular disease, and cancer of the bladder, brain, breast, colon, lungs, ovaries, and stomach [18-22]. Other potential benefits include, but are not limited to, kidney function [23], atherosclerosis [24] anti-bacterial/fungal activity [25-28], cataractogenesis [29], immune function [30, 31] and prebiotic effect [32, 33]. The therapeutic properties of onion and garlic are caused by the combination and biological activity of organo-sulfur compounds such as S-allyl-l-cysteine, diallyl disulfide, diallyl trisulfide, ajoene, and allicin. Studies suggest that the biological and therapeutic properties of garlic and onions are mainly due to their high organo-sulfur compounds [1].

*Address correspondence to this author at the School of Studies in Biotechnology, Jiwaji University, Gwalior 474005, India; Tel: +91 751 2462500; Fax: +91 751 2341768; E-mail: psbisen@gmail.com




Fig. (1). Sulphur-containing constituents of onion and garlic.

Phytochemicals in *Allium* Genera

The primary sulphur containing constituents in both plants are the S-alkyl-L-cysteine sulphoxides (ACSOs), such as (a) Methiin and alliin (b) allicin (c) diallyl sulphide (DAS), and (d) diallyl disulphide (DADS) (Fig. 1). These compounds provide to garlic and onion their characteristic odor and flavor, as well as most of their biological properties.

Of the countless phenolic compounds found in the onion, flavonoid formed by oxidation of C3 units, have persistently been assigned an important nutritional and therapeutic role [34]. The degree of oxidation identifies them in different subgroups like flavonols, flavanones, steroidal saponins, and anthocyanins (Fig. 2 and 3). One of the major sources of two forms of dietary flavonoids found in onions are (a) antho-

cyanins in red and purple skinned bulbs and (b) quercetin and its derivatives in yellow skinned onion bulbs. Flavonoids are abundant in onion (Fig. 2 and 3) while is completely lacking in garlic. Garlic contains a small amount of sallyl cysteine (SAC), non-volatile water-soluble sulphur compound (Fig. 1). Fruto-kestose (GF2), nystose (GF3) fructofuranosyl-nystose (GF4), thiosulfonates are other sulphur containing bioactive compounds (Fig. 3). Glycerophospholipids, lectins, saponins (β -sitosterol, gitogenin, oleanolic acid, amyirin, diosgenin, cepagenin), glucosides, fructan, pectin, vitamins A, B1, B2, B3, B6, C, E, allixin and organo-selenium compounds such as γ -glutamyl-Se-methylselenocysteine, Se-methylselenocysteine, "Se-alliins", Se-methionine, Se-cystine/Se-cysteine are the additional most abundant functional constituents in garlic and onion. Sulphur compounds do not exist as such in the intact cells. They are




Fig. (2). Some common flavonoids of onion.

formed as a result of enzymic reaction between allinase and volatile precursors substrate S-alk(en)yl cysteine sulphoxide and sulphonic acid. Thiosulphinates and sulphonic acid compounds are derived when the cells are ruptured [35, 36]. Sulfur compounds are found in onion bulbs alongside with seleno derivatives thiosulfinates, thiosulfonates cepaenes, S-oxides, S,S'-dioxides, monosulfides, disulfides, trisulfides, zwiebelanes, and soluble sulphur (γ -glutamylcysteine peptides) (Fig. 1 and 4).

A wide range of different flavonols, differentiated on the basis of degree of hydroxylation, are found in onions consisting of the aglycones and glycosylated derivatives. They are quercetin glucosides viz. the 4'-glucoside, the 7,4'-diglucoside, the 3,4'-diglucoside, the 3-glucoside, the 7-glucoside, the 3,7-diglucoside, the 3-rutinoside (rutin), the 3-rhanmoside (quercetrin), the 7,4'-, and 3-glucosides of kaempferol, isorhamnetin 4'-glucoside and rest are anthocyanins (Fig. 2 and 3).

An abundance of flavonols is typical for flesh scale tissue compared to fresh white skinned onions, garlic and leek containing only trace amounts of flavonols. Onion is also a potential source of prostaglandins [37]. Bulbs and leaves of alliums possess flavonoids, sterols and steroidal glycosides. In garlic, 0.021% and in onion, 0.095%, saponin level has been reported [38, 39]. Fructans and fructosyl are the great source of soluble dietary fiber and the main stored carbohydrates in onion plant. The accumulation of fructans is associated with high concentration of thiosulfinate, soluble carbohydrates and pungency according to onion phenotype. Fructans, with maximum degree of polymerisation are typical for high dry matter onion bulbs. They are found in high concentration in onions bulbs (35-40% dry weight) constituting a major portion of the water-soluble carbohydrates. The chain




Fig. (3). Some common phytochemicals of onion (*Allium cepa*).

of fructans / fructosyl polymers of D-fructose molecules bonded by β (2 \rightarrow 1) linkages is terminated by a D-glucose molecule linked to fructose by an α (1 \rightarrow 2) bond during polymerisation [1, 39, 40]. Maximum concentration of lower molecular weight fructans with degree of polymerization 5 are found in the onion inner, younger leaf bases. Flavonols, anthocyanins and dihydroflavonols are found to occur in onion bulbs [1, 40]. Flavonols are the predominant pigments with 270-1187 mg per kilogram (FW) in yellow onions and 415-1917 mg of flavonols per kilogram in red onions (FW). Twenty five different flavonols are characterized with quercetin derivatives (Quercetin 4'-glucoside and quercetin 3,4'-diglucoside) as the most abundant dietary flavonoids with glycosyl moieties having glucose as exclusive moiety in all onion cultivars (Fig. 3). Glycosyl moieties are mainly attached to the 4', 3, and/or 7- positions of the aglycones. Analogous derivatives of kaempferol (Fig. 2) and isorhamnetin are minor pigments. Oligomeric structures of quercetin and the condensation products of quercetin and protocatechuic acid are found in the outer dry layers of onion bulbs. The anthocyanins and cyanidin glucosides acylated with malonic acid or nonacylated are found in red onions [40-42]. Essential and vegetable oils derived from the onion and garlic seeds have potential therapeutic value. The cold pressed onion seed oil contains fatty acids; palmitic acid (C16:0, 6.4-7.1%), oleic acid (C18:1, 24.8-26 %) and linoleic acid (C18:2, 65.2-64 %) [1, 43]. Oleic and linoleic acids are recognized for their anti-inflammatory, wound healing, pro-coagulant, cardioprotective, antiarrhythmic, hypotensive,




Fig. (4). Enzymic reaction of allinase.

neuroprotective properties and reduce the severity of both acute and the relapsing phases of chronic multiple sclerosis [1, 44]. Essential oil from different species of *Allium* viz. *Allium cepa*, *Allium sativum* L., *Allium jesdianum* Boiss, *Allium roseum* L., *Allium sphaerocephalon* L., *Allium nigrum*, *Allium rotundum* and others demonstrate strong antibacterial activity [45-49]. The essential oil distilled from onion seeds contains about 48 active compounds, including 2-phenylethanol (0.42 % to 41.17 %), 1-pentanol, 2-heptanol, 2-octanol, 1-octen-3-ol, 1-heptanol, phenylmethanol, 1-dodecanol, 1-pentanol, 2-heptanol, 1-octen-3-ol and phenylmethanol (up to 0.66 %), aromatic hydrocarbons (p-xylene and mesitylene), cyclic monoterpene α -phellandrene, bicyclic monoterpene, p-cymene, esters (hexylacetate, nonylacetate, ethyloctanoate, methylhexanoate and nonylacetate) [49].

Garlic (*Allium sativum*)

Garlic (*Allium sativum*) is the herb known for its flavour and pungency and most widely quoted for medicinal properties [1, 9, 35, 36, 50]. Garlic ranks second best selling vegetable in the United States and used extensively as medicinal plant all over the world for over 5000 years for different ailments [51, 52]. Therapeutic potential of alliums was first recorded in collection of Zoroastrian holy writings followed by Sumerian and the ancient Egyptians. The Olympics in Greece used raw garlic to feed the athletes and increasing stamina [51]. In ancient medicine alliums and especially garlic was used to aid digestion, respiration and to treat leprosy [53]. Garlic gain especial attention as a plant with exceptional disease preventative and therapeutic properties used by Avicenna in treatment of variety of disease such as toothache, constipation, parasitic infestation, arthritis, insect bites, and infectious diseases as natural antibiotic [53]. Garlic demonstrates a number of pharmacological effects viz. cancer chemoprevention, antibiotic, antihypertensive, anti-

diabetic, antithrombotic and cholesterol-lowering properties [5]. The thiosulfinate extracts from garlic inhibit the tumor cells growth [20-22]. Abundance of natural oil and water-soluble sulfur-containing organosulfur compounds are responsible for the characteristic flavor and pungency of garlic having several pharmacological significance whereas intact, undisturbed bulbs of garlic contain only a few bioactive components. An odorless amino acid, alliin (Fig. 4) is found in intact garlic which gets converted by an enzyme allinase into alliin responsible for the characteristic odor when cloves are crushed [37]. The spontaneous decomposition results to form numerous sulfur-containing compounds, some of which are having chemopreventive activity (Fig. 4).

Onion (*Allium cepa*)

The onion (*Allium cepa*) (Latin 'cepa' = onion), most widely cultivated species of the genus *Allium*, commonly known as bulb onion, is used as a vegetable for its nutritive value (Table 1). They are cultivated by many cultures and used globally for their medicinal qualities. The top global producers of onion during last 5 years are China and India with higher yields followed by Uzbekistan, Egypt, China, Korea and USA. They are usually eaten raw or used to make pickles or chutneys, served cooked as food stuff, as a vegetable or part of a prepared savoury dish.

Table 1. Nutritional value of raw onion in each 100 g. (Source USDA www.onions-usa.org)

Energy	166kJ (40 kcal)
Carbohydrate	9.34 g
Sugars	4.24 g
Dietary fiber	1.7 g
Fat	0.1 g
Protein	1.1 g
Vitamins	
Thiamine (B ₁)	(4%) 0.046 mg
Riboflavin (B ₂)	(2%) 0.027 mg
Niacin (B ₃)	(1%) 0.116 mg
Pantothenic acid (B ₅)	(2%) 0.123 mg
Vitamin B ₆	(9%) 0.12 mg
Folate (B ₉)	(5%) 19 µg
Vitamin C	(9%) 7.4 mg
Trace Metals	
Calcium	(2%) 23 mg
Iron	(2%) 21 mg
Magnesium	(3%) 10 mg
Manganese	(6%) 0.129 mg
Phosphorus	(4%) 29 mg
Potassium	(3%) 146 mg
Zinc	(2%) 0.17 mg
Other constituents	
Water	89.11 g
Fluoride	1.1 µg

The plant has a fan like hollow bluish-green leaves with the bulb at the base. On maturation, bulb swells, the foliage dies down in the autumn and the outer layers of the bulb become dry and brittle. The crop is harvested, dried and are ready for use or storage. There are several other species; in this genera cultivated for food, such as shallots and potato onions with multiple bulbs, variously referred to as onions having a unique rich combination of fructans, dietary flavonoids and organosulfur compounds (Fig. 3). These three families of compounds express salutary effects on human health with functional benefits against life style diseases including reduced risk of coronary heart disease, different types of cancer, anti-oxidative activity, metal ion chelation, lipid peroxidation inhibition, asthma, allergies, arthritis, diabetes, neurodegenerative disorders, osteoporosis and others. [54]. Onion pigments like 4'-glycosylation promote unique and unusual substitution patterns of sugar moieties. There are variation in structural identification of dihydroflavonols from onions compared to flavonols and anthocyanins and may be at high concentrations in some cultivars.

Shallot (*Allium ascalonicum*)

Shallot is used traditionally from the ancient times demonstrating a variety of health benefits including antioxidant, hypocholesterolemic, hepatoprotective, antidermatophytic, anti-angiogenic, anti-inflammatory, antibacterial (potential inhibitory activity against *Helicobacter pylori*), anti-fungal and peroxynitrite-scavenging properties [55]. The most effective pharmacological compounds of *Allium ascalonicum* are mannose-specific lectin, furostanol saponins, selenium, sulfur, antifungal peptide and flavonol glucosides [55]. The shallot extract inhibits proliferation and growth of tumor cell lines as HeLa and MCF-7 cells *in vivo* and *in vitro*, in combination with garlic is effective against human leukemia cells (HL-60) and human cervical carcinoma HeLa cells [56]. *A. ascalonicum* significantly decreases the inflammation *in vivo* via reduction of the vascular permeability or through inhibition of inflammatory mediators [55, 56].

Therapeutic Potential of Allium

Alliums are used as medicine for many major and minor disorders. Variety of data presently confirms the therapeutic properties and low toxicity of active ingredients in different cultivars of Allium by enhancing the blood's fibrinolytic activity and suppressing blood fibrinogen concentration, and increase the blood coagulation time [57, 58]. Garlic and onion also play a role in lowering serum cholesterol levels both in animals and humans [59]. Garlic and onion bulbs can also reduce the level of blood sugar [60, 61], thus they are beneficial to diabetes, obesity, respiratory and skin infections, insect bites, memory impairment, demonstrate strong anthelmintic properties [62] reduction of hypertension [63] and stimulation of the body's immune system [64]. Flavonoids present in the bulbs and leaves of alliums are very effective antioxidants with variable biochemical functions [35, 65, 66] like immune functionality, gene expression, liver function, enzyme activity, platelet aggregation, histamine metabolism, *etc* [34, 65]. Oligofructans are helpful in propagation of bifidobacteria and in reduction of deleterious bacteria in the colon, prevent wound infection and food spoilage, protection

of liver function and detoxification, constipation prevention, reduction of serum cholesterol and blood pressure with various anticancer effects [67].

Anticarcinogenic Activity

Alliums have anticarcinogenic activity, but garlic is known to be more effective due to the interaction of the sulphur components of garlic with the tumour cell metabolism [68-70]. DNA adduct formation, mutagenesis, free radicals scavenging, cell proliferation, differentiation and angiogenesis can be modulated by different garlic derivatives and onion suppress/inhibit the platelet aggregation *i.e.*, induced by collagen, arachidonic acid, ADP, adrenaline and the thrombin and the effect is dose dependent. Studies have demonstrated that organosulfur compounds such as diallyl disulfide (DDS), S-allylcysteine (SAC), and S-methylcysteine (SMC) inhibit colon and renal carcinogenesis *in vitro* and *in vivo* [7, 71]. The steroidal saponins isolated from a variety of different species of onions demonstrate significant cytotoxic activity against murine fibrosarcoma, lung carcinoma, human melanoma and leukemia [72, 73]. The risk of *Helicobacter pylori* infection linked to stomach cancer through ulcer formation is reported to be decreased by providing protection from colon cancer by inhibiting both initiation and proliferation through the effect of both organosulfur and flavonoids compounds [74, 75] with high consumption of allium vegetables [74]. Organosulfur compounds modulate the cytokine pattern to inhibit NFB, the central regulator of pro-inflammatory gene expression and link between inflammation and tumor progression. They exert immunomodulatory effects through alterations in the production of cytokines and immune modulators. Diallyl trisulfide (DATS) has been shown to reduce tumor mitosis, decrease histone deacetylase activity, increase acetylation of H3 and H4, inhibited cell cycle progression, and decrease pro-tumor markers [76]. DATS inhibited cell growth of human melanoma and ajoene has been shown to induce apoptosis in leukemic cells via stimulation of peroxide production, through activation of caspase-3-like and caspase-8 activity [77].

Quercetin is known to heal the gastric ulcers and colonocytes through free radical scavenging by preventing the induction of gastric mucosal injury and relieves muscular spasm due to calcium-antagonistic properties [53, 64-66], highly effective in inducing apoptotic cell death in colorectal tumour cells while sparing normal cells [41, 80], gene transcription inhibition to protection against UV-induced immune-suppression [81,82] and inhibition of kinase activity in the signal transduction pathway as a mode of protective action [83, 84]. Antioxidant property of quercetin is reported to be inversely associated with cancer formation [78, 79]. Quercetin functions in multiple ways *viz.* (a) decrease cancer tumour initiation; (b) promote healing of stomach ulcers (c) inhibit the proliferation of cultured ovarian, breast, and colon cancer cells. There is a significant anti-proliferative effect of quercetin on ovarian carcinoma, breast cancer cells by increasing the activity of reductase enzymes known to inactivate cytotoxic carcinogenic compounds and synergistically increase the activation of quinoid anticancer drugs [41, 52]. The organosulfur compounds are largely responsible for the taste and smell of onion and also (a) reduce symptoms asso-

ciated with diabetes mellitus (b) inhibit platelet aggregation (involved in thrombosis) (c) prevent inflammatory processes associated with asthma. Many of these studies used non-human subjects. Some of the most common organosulfur compounds dipropyl sulfide (DPS) and dipropyl disulfide (DPDS) in onions protect SPF Wistar male rats against cancer at early and late stages of carcinogenesis in by triggering microsomal epoxide hydrolase, glutathione S-transferase, and UDP-glucuronosyltransferase activities in the liver [71]. Other two common organosulfur compounds cysteine and S-methylcysteine of onion showed chemopreventive activity against hepatocarcinogenesis and colon carcinogenesis in a rat model [7, 85]. Both organosulfur compounds and flavonoids found in onions have a protective effect against N-nitroso compounds (NOCs) known for increased risk of brain cancer in China [86]. Dietary intake of *Allium* vegetables intake diminishes the risk of sarcoma and carcinoma in various tissues and organs such as oral, skin, brain, pharyngeal, esophageal, stomach, colorectal, laryngeal, breast, ovarian, prostate, and renal cell cancers [77, 86-90]. Eruboside-B, a steroid saponin, kaempferol, and organo Se compounds [91] are largely responsible for the anticarcinogenic activity [92]. These effects appear to be mediated by various mechanisms such as (a) increasing the detoxification enzymatic systems activity (b) facilitating its excretion from the body (c) inhibiting the procarcinogens activation and (d) inhibition of oxidative damage due to their antioxidant action. The apoptotic effect of water soluble garlic extract on the human metastatic MDA-MB-435 cancer cells in mitosis is due to the modifications of the proapoptotic protein BimEL to mitochondria leading to execution of its proapoptotic function. Several epidemiologic studies show that garlic components may be effective to inhibit tumorigenesis in several experimental models [12] and in human cancer risks [4, 8]. There is a strong inverse relationship between stomach and colon cancer risk and increasing garlic intake [93, 94]. However, not all epidemiological studies have shown garlic to be protective against carcinogenesis. Additional reports have shown garlic to be ineffective [94]. The inconclusive results are likely due to differences in the type of garlic compounds, quantity of organosulfur compounds available in fresh and/or commercially available preparations used by various investigators [95].

Cardiovascular Diseases (CVD)

CVD is affecting 1 billion people and expected to affect over 1.6 billion people by 2025. Garlic has been advocated for the prevention of antihypertensive properties responsible for CVD. However, due to insufficient available evidences, it has not been recommended as a routine clinical therapy for the treatment of hypertensive subjects [96]. The antihypertensive effect of garlic is shown to be related with the hydrogen sulphide production, allicin content, and alliinase, the enzyme which possesses angiotensin II inhibiting and vasodilating properties. Basically, one of the mechanisms of antihypertensive activity of garlic was suggested to be explained through its prostaglandin-like effects leading to eventual decrease of peripheral vascular resistance. Majority of studies suggest that the cardioprotective properties are more likely due to lowering of low-density lipoproteins (LDL) and blood pressure. The metaanalysis results showed that an

average of 800-900 mg garlic/day could decrease total serum cholesterol levels by approximately 9-12% [96, 97]. Long term application of garlic was effective to reduce almost 50% cholesterol and triglycerides in serum and atheromatous lesions in the aorta [98]. Garlic significantly inhibited intracellular Ca²⁺ mobilization, thromboxane-A₂ synthesis and protected against thrombosis. The aged garlic extract can inhibit blood platelet aggregation through suppression of the GPIIb/IIIa receptor and an increase in cAMP [99], can cause decrease of plasma viscosity, and increase the elastic property of blood vessels in mice [100]. A dietary intake of *Allium* is beneficial for normalizing platelet hyperactivation and prevention of cardiovascular diseases. The validity of these reports, however, is reduced by methodological shortcomings, including the fact that dietary intake, weight, and/or exogenous garlic ingestion was not always well-controlled. In a multicenter, randomized, placebo-controlled trial in which dietary assessment and supervision were strictly controlled, 12 weeks of garlic treatment was ineffective in lowering cholesterol levels in subjects with hypercholesterolemia [101]. However, it is yet not clear the specific component of garlic responsible for cholesterol lowering.

Immunomodulatory Properties

Allium sativum and its compounds significantly reduce the leukocyte- cytokine production in inflammatory conditions (IBS) [102]. Some garlic proteins modulate NK cell line activity in the mesenteric lymph node of mouse, and aged garlic extract increases NK activities against different cancerous cell lines [103].

Lipid Metabolism

Total serum cholesterol, LDL and very low density lipoproteins (VLDL) are drastically decreased in rats and rabbits after raw garlic consumption with significant increase in the level of high density lipoproteins (HDL). Allicin and its derivative compounds are the main active substances responsible for the hypolipidemic and hypocholesterolemic effects of onion and garlic [1]. Studies have shown that saponins and flavonoid quercetin are able to reduce level of serum cholesterol and arteriosclerosis severity, diabetes [39], and HIV suppression/AIDS blocking [104]. Most likely, the organosulfur compounds [7, 105], working through sulfur-sulfur or sulfur-oxygen linkages [6], are believed to possess anti-inflammatory, anti-allergic, anti-microbial, and anti-thrombotic activity through inhibition of cyclooxygenase and lipoxygenase [37].

Antimicrobial and Antifungal

Garlic and onion have been used for centuries in folk medicine in several societies against microbial infections. Sulphur compounds characterised from onion and garlic are considered as the main active antimicrobial agents [106-109]. Garlic and onion extracts are known to inhibit the growth of Gram-positive, Gram-negative and acid-fast bacteria [110] including *E. coli*, *Shigella*, *Salmonella*, *Proteus*, *Klebsiella*, *Enterobacter*, *Vibrio cholera*, *V. parahaemolyticus*, *Staphylococcus aureus* [106], *S. typhi*, *Pseudomonas pyocaneus*, *P. vulgaris*, *Bacillus subtilis*, *Cryptococcal meningitis*, *Shigella dysenterae* and *Aspergillus* [111]. Garlic

suppresses beneficial intestinal microflora as well. However, probably due to a greater sensitivity of enterobacteria to allicin, it is more effective against potentially harmful enterobacteria [106-115].

Garlic extract with high content of ajoene is significantly effective against *Candida albicans*, *Paracoccidioides*, *Aspergillus* sp., *Botrytis cinerea* and *Trichoderma harzianum*, *Scedosporium prolificans*, *Tinea pedis*, *Opalina ranarum*, *Entamoeba histolytica*, *Leishmania*, *Trypanosomes*, *Leptomonas*, *Crithidia*, etc. [38, 116, 117]. The identified main active antifungal agents from onion and garlic extracts are the breakdown products of allicin which is able of inhibit sulfhydryl enzymes, DAS, DADs, diallyl trisulphide (DATS) and ajoene having greater antifungal effect than allicin [38, 115]. The active antifungal compound of garlic and onion destroy fungal cells by (a) reducing the oxygen uptake potentiality (b) reduce cellular multiplication and (c) by inhibiting synthesis of lipids, proteins and nucleic acids [38]. In addition to effectiveness against variety of pathogens, allicin stimulates the endogenous immune system via immune-correlated signalling pathways in cells. Allicin is an electrophile molecule with positive charge attract towards cysteine, activate channel protein via modification of cysteine residues and activation of TRPA1 leading to cause alteration in the conductance of the ion channels in the cell membranes.

Antiparasitic and Antiprotozoal

About 1.5 million people, especially children die of the malaria initiated by *Plasmodium* sporozoites every year [http://www.who.int/malaria/media/world_malaria_report_2014/en/]. Cysteine protease and ajoene present in freshly crushed garlic cloves inhibits target preerythrocytic and erythrocytic stages of the *Plasmodium* development. Pure allicin inhibits cytopathological effects and the growth of the intestinal parasites *Entamoeba histolytica* and *Plasmodium berghei*. Garlic extract is effective against *Leishmania* protozoan which infects nearly 12 million people globally and have no effective vaccine yet. Aged garlic extract modulates cytokine patterns towards a Th1-type response, increasing nitric oxide production and macrophage activation, which are important for eradication of parasite [118].

Antiviral Activity

The antiviral activities of various commercial garlic products have been developed against influenza A and B viruses [119] human cytomegalovirus [120, 121], human immunodeficiency virus (HIV), viral pneumonia and rotavirus [122, 123]. Quercetin, lectins and sulphur compounds are reported to possess antiviral activity and enhance the bioavailability of some anti-viral drugs with pronounced anti-HIV activity [120, 123].

Antioxidant Activity

Quercetin, which is the major dietary flavonoid, found in onion, show the highest antioxidative activity comparable to α -tocopherol [124-126]. The dry outer layers of onion contains large amount of quercetin. S-allyl cysteine (SAC) and S allylmercaptocysteine (SAMC), bioavailable water-soluble organo-sulphur compounds of garlic, along with stable lipid-

soluble allyl sulphides, as DAS, DATS, DADS and diallyl polysulphides, flavonoids (as allixin); saponins, essential micronutrients (selenium) and macronutrients, as lectins, are known to express potent antioxidant activity [125, 126].

Antihyperglycemic Activity

Diabetes mellitus (NIDDM, type 2 diabetes) is characterized by hyperglycemia, which is expressed with a rapid rise in blood glucose level due to hydrolysis of starch by pancreatic α -amylase and absorption of glucose in the small intestine by α -glucosidases [127]. The aqueous juice of various forms of raw onions and garlic and isolated compounds possess significant antihyperglycemic effect in laboratory settings. The root of the Welsh onion (*Allium fistulosum* L.) alleviates hyperglycemia via inhibiting α -glucosidase *in vitro*, as well as in animal model *in vivo* [127]. Quercetin, isoquercitrin and rutin demonstrate significant inhibitory activity on α -glucosidase from the rat intestine [127]. The Korean onions (*Allium cepa* L.) have strong inhibitory potential against rat α -glucosidases (sucrase, maltase, porcine pancreatic α -amylase) *in vitro* and *in vivo*. The consistent dietary intake of the onion (*Allium cepa* L.) extract in rat improves diabetes by decreasing triglyceride and free fatty acid levels. Aged garlic extract increase insulin secretion and sensitivity, increase nitric oxide production and significantly decrease inflammatory cytokines, glycosylated hemoglobin (HbA1c), postprandial blood glucose (PPBG) and fasting blood glucose (FBG) level in human studies [128]. Garlic oil demonstrates antioxidant activity in diabetic skeletal muscle via reversing insulin resistance, lowering the nitric oxide production and improving GLUT4 expression.

Effect on Respiratory System

A wide range of different flavonols, differentiated on the basis of degree of hydroxylation, are found in onions consisting of the aglycones and certain onion-derived compounds (not garlic), particularly thiosulfinates and cepaenes, show an anti-inflammatory and antiasthmatic properties *in vivo*. The main compounds found in *Allium cepa* L. extract (AcE) are the natural phenolics, flavonoids such as quercetin, exhibiting anti-inflammatory and antioxidant effects. There is a reduction in the production of inflammatory cytokines, a relaxation of tracheal rings, and a reduction in the total number of cells in BAL and EPO in the lungs by treatment with onion extract or quercetin [129]. Along with flavanols, the major bioactive constituents in *Allium cepa* L. are sulfurous compounds [130-133].

Effect on Skin

The nuclear transcription factor kappaB which is known to be linked to psoriasis can be interrupted by garlic compounds, such as diallyl sulfide, sallylmercaptocysteine and ajoene. These compounds play important role in development and progression of alopecia areata, wound healing, keloid scars, leishmaniasis. Garlic stimulates the proliferation of macrophages and lymphocytes, protecting against the suppression of immunity by ultraviolet radiation. *Allium cepa* improves the appearance of scars, their softness, diminish redness and improve the skin texture. Ingested fresh

Allium sativum delays formation of skin papillomas and protect against skin cancer [134].

Allium Products

Due to therapeutic properties of alliums, onion and garlic, diet supplements with capsules, tablets and lozenges are widely available on local and international markets. Onion products include dehydrated onion pieces, onion powder, oleoresins, onion salts, pickled onions, canned, frozen and packaged onions, etc. Dehydrated garlic powder, garlic juice, garlic salt, garlic flavouring are some of the advantageous garlic products being marketed [134]. Essential oil of garlic is extracted by distillation in boiling water in a still or by introducing live steam generated in a separate steam boiler [134, 135].

CONFLICT OF INTEREST

The authors confirm that this article content has no conflict of interest.

ACKNOWLEDGEMENTS

Declared none.

REFERENCES

- Osman SA.-M, Ata A.-TM, El-Hak SE.-NH.G. Morphological, germination, bolting and cytogenetical characteristics of fourteen promising garlic genotypes. *Afri Crop Sci Confer Proceed* 2007; 8: 2005-12.
- Jo MH, Ham IK, Moe KT, et al. Classification of genetic variation in garlic (*Allium sativum* L.) using SSR markers. *Aus Jour Crop Sci* 2012; 6: 625-31.
- Block G, Patterson B, Subar A. Fruit, vegetables, and cancer prevention: A review of the epidemiological evidence. *Nutr Cancer* 1992; 18: 1-29.
- Dorant E, van den Brandt PA, Goldbohm RA, et al. Garlic and its significance for the prevention of cancer in humans: A critical review. *Br J Cancer* 1993; 67: 424-9.
- Srivastava KC, Bordia A, Verma SK. Garlic (*Allium sativum*) for disease prevention. *S Afr J Sci* 1995; 91: 68-77.
- Augusti K. Therapeutic values of onion and garlic. *Ind J Expt Bio* 1996; 34: 634-40.
- Fukushima S, Takada N, Hori T, et al. Cancer prevention by organosulfur compounds from garlic and onion. *J Cell Biochem Suppl* 1997; 27: 100-5.
- Ernst E. Can allium vegetables prevent cancer? *Phytomedicine* 1997; 4: 79-83.
- Nagourney RA. Garlic: Medicinal food or nutritious medicine? *J Medicinal Food* 1998; 1: 13-28.
- Hasler CM. Functional foods: Their role in disease prevention and health promotion. *Food Tech* 1998; 52: 63-70.
- The institute of food technologists' expert panel on food safety and nutrition. Functional Foods: Their role in disease prevention and health promotion. *Food Technol Mag* 1998; 52(11): 63-70. Available From: <http://www.ift.org/knowledge-center/read-ift-publications/science-reports/scientific-status-summaries/functional-foods.aspx>
- Reuter HD, Koch HP, Lawson LD. Therapeutic effects and applications of garlic and its preparations. In: Koch HP, Lawson LD, Eds. *Garlic. The science and therapeutic application of Allium sativum L. and related species*, 2nd ed. Williams & Wilkins: Baltimore 1996.
- Mulbauer RC, Li F. Effect of vegetables on bone metabolism. *Nature* 1999; 401: 343-4.
- Ames B, Gold L. The prevention of cancer. In: Shibamoto T, Terao J, Osawa T, Eds. *Functional foods for disease prevention I*. American Chemical Society: San Francisco 1998.
- Cao G, Sofic E, Prior R. Antioxidant capacity of tea and common vegetables. *J Agric Food Chem* 1996; 44: 3426-31.
- Song K, Milner JA. The influence of heating on the anticancer properties of garlic. *J Nutrition* 2001; 131:1054s-7s.
- Yin M, Cheng W. Antioxidant activity of several Allium members. *J Agric Food Chem* 1998; 46: 4097-101.
- Dorant E, Van Din Brandt P, Goldbohm R. A prospective cohort study on Allium vegetable consumption, garlic supplement use, and the risk of lung carcinoma in the Netherlands. *Cancer Res* 1994; 54: 6148-53.
- Dorant E, Van Den Breandt P, Goldbohm A. Allium vegetable consumption, garlic supplement intake, and female breast carcinoma incidence. *Breast Cancer Res Treat* 1995; 33: 163-70.
- Dorant E, Van Den Breandt P, Goldbohm R, et al. Consumption of onions and a reduced risk of stomach carcinoma. *Gastroenterology* 1996; 110: 12-20.
- Challier B, Pernau J, Viel J. Garlic, onion and cereal fibre as protective for breast cancer: A French case-study. *Eur J Epidem* 1998; 14: 739-47.
- de Groot H, Rauhen U. Tissue injury by reactive oxygen species and the protective effects of flavonoids. *Fundam Clin Pharmacol* 1998; 12: 249-55.
- Shoskes D. Effect of bioflavonoids quercetin and curcumin on ischemic renal injury. *Transplantation* 1998; 66: 147-52.
- Sainani G, Desai D, Sainine P. Onion in prevention of atherosclerosis. *J Ind Med Assoc* 1978; 71: 109.
- Dankert J, Tromp T, De Vries H et al. Antimicrobial activity of crude juices of *Allium ascalonicum*, *Allium cepa*, and *Allium sativum*. *Zentrabl Bakteriol Parasitenkd Infektkrankh Hyg Ab* 1979; 245: 229-39.
- Jain RC. Antitubercular activity of garlic oil. *Indian Drugs* 1993; 30: 73-5.
- Kim J. Anti-bacterial action of onion extracts against oral pathogenic bacteria. *J Nihon Univ Sch Dent* 1997; 39: 136-41.
- Fan J, Chen J. Inhibition of aflatoxin producing fungi by Welsh onion extracts. *J Food Protec* 1999; 62: 414-7.
- Sanderson J, Mclauchlin W, Williamson G. Quercetin inhibits hydrogen peroxide-induced oxidization of the rat lens. *Free Radical Biol Med* 1999; 26: 639-45.
- Chisty M, Quddus R, Islam B, et al. Effect of onion extract on immune response in rabbits. *Bangladesh Med Res Counc Bull* 1996; 22: 81-5.
- Steenberg PA, Garssen J, Dortant P, et al. Protection of UV-induced suppression of skin contact hypersensitivity: A common feature of flavonoids after oral administration? *Photochem Photobiol* 1998; 67: 456-61.
- Gibson G. Dietary modulation of the human gut microflora using prebiotics. *British J Nutr* 1998; 80: S209-12.
- Jaime L, Martin-Cabrejas MA, Molla E, et al. Effect of storage on fructan and fructo-oligosaccharide of onion (*Allium cepa* L.). *J Ag Food Chem* 2001; 49: 982-8.
- Slimestad R, Fossen T, Vågen IM. Onions: A source of unique dietary flavonoids. *J Agric Food Chem* 2007; 55: 10067-80.
- Hertog M, Katan M. Quercetin in foods, cardiovascular disease, and cancer. In: Rice- Evans CA, Packer L, Eds. *Flavonoids in health and disease*. Marcel Decker Inc. New York 1998; pp. 447-67.
- Lancaster JE, Shaw ML, Randle WM. Differential hydrolysis of alk(en)yl cysteine sulphoxides by alliinase in onion macerates: Flavour implications. *J Sci Food Agric* 1998; 78: 367-72.
- Block E, Gulati H, Putman D, et al. Allium chemistry: Synthesis of 1-[alk(en)ylsulfanyl]propyl alk(en)yl disulfides (cepaenes), anti-thrombotic flavorants from homogenates of onion (*Allium cepa*). *J Agric Food Chem* 1997; 45: 4414-22.
- Lanzotti V, Barile E, Antignani V, et al. Antifungal saponins from bulbs of garlic, *Allium sativum* L var. Voghiera. *Phytochemistry* 2012; 78: 126-34.
- Lanzotti V. The analysis of onion and garlic. *J Chromatogr A* 2006; 1112: 3-22.
- Sheela CG, Kumud K, Augusti KT. Anti-diabetic effects of onion and garlic sulfoxide amino acids in rats. *Planta Med* 1995; 61: 356-7.
- Deschner E, Ruperto J, Wong G, et al. Quercetin and rutin as inhibitors of azoxymethanol-induced colonic neoplasia. *Carcinogenesis* 1991; 12: 1193-6.
- Formica JV, Regelson W. Review of the biology of quercetin and related bioflavonoids. *Fd Chem Toxic* 1995; 33: 1061-80.

- [43] Parry J, Hao Z, Luther M, *et al.* Characterization of cold-pressed onion, parsley, cardamom, mullein, roasted pumpkin, and milk thistle seed oils. *J Am Oil Chem Soc* 2006; 83: 847-54.
- [44] Harbige LS, Layward L, Morris-Downes MM, *et al.* The protective effects of omega-6 fatty acids in experimental autoimmune encephalomyelitis (EAE) in relation to transforming growth factor-beta 1 (TGF- β 1) up-regulation and increased prostaglandin E2 (PGE2) production. *Clin Exp Immunol* 2000; 122: 445-52.
- [45] Benkeblia N. Antimicrobial activity of essential oil extracts of various onions (*Allium cepa*) and garlic (*Allium sativum*). *Lebensm Wiss Technol* 2004; 37: 263-8.
- [46] Amiri H. Chemical composition and antibacterial activity of the essential oil of *Allium Jesdianum* Boiss. and Buhse from Iran. *J Med Plants* 2007; 6: 39-44.
- [47] Dehpour AA, Yousefian M, Kelarijani SAJ, *et al.* Antibacterial activity and composition of essential oils of flowers *Allium rotundum*. *Adv Environ Biol* 2012; 6: 1020-5.
- [48] Lazarevic JS, Đorđević AS, Zlatković BK, *et al.* Chemical composition and antioxidant and antimicrobial activities of essential oil of *Allium sphaerocephalon* L. subsp. *Sphaerocephalon* (Liliaceae) inflorescences. *J Sci Food Agr* 2011; 91: 322-9.
- [49] Najjaa H, Neffati M, Zouari S, *et al.* Essential oil composition and antibacterial activity of different extracts of *Allium roseum* L. a North African endemic species. *C R Chim* 2007; 10: 820-6.
- [50] Venugopal KJ. New herbal health ingredients and their functional applications: A review. *Ingredient Directory*, 1-7. Available From: www. b5srl.com
- [51] Lawson LD, Bauer R. Garlic: A review of its medicinal effects and indicated active compounds. In: *Phytomedicines of Europe. Chemistry and Biological Activity*. Washington DC: American Chemical Society 1998; Series 691: pp. 176-209.
- [52] Anonymous. Quercetin. *Altern Med Rev* 1998; 3: 140-3.
- [53] Rivlm RS. Patient with hyperlipidemia who received garlic supplements Lipid management. Report from the Lipid Education Council 1998; 3: 6-7.
- [54] Gazzani G, Papetti A, Daglia M, *et al.* Protective activity of water soluble components of some common diet vegetable on rat liver microsome and the effects of the thermal treatment. *J Agric Food Chem* 1998; 46: 4123-7.
- [55] Fattorusso E, Iorizzi M, Lanzotti V, *et al.* Chemical composition of shallot (*Allium ascalonicum* Hort.). *J Agric Food Chem* 2002; 50: 5686-90.
- [56] Azadi GH, Ghaffari SH, Riazi GH, *et al.* Antiproliferative activity of chloroformic extract of Persian Shallot, *Allium hirtifolium*, on tumor cell lines. *Cytotechnology* 2008; 56: 179-85.
- [57] Bordia A, Verma SK, Srivastava KC. Effect of garlic (*Allium sativum*) on blood lipids, blood sugar, fibrinogen and fibrinolytic activity in patients with coronary artery disease. *Prostaglandins Leukot Essent Fatty Acids* 1998; 58: 257-63.
- [58] Sobenin I, Andrianova I, Ionova V, *et al.* Anti-aggregatory and fibrinolytic effects of time-released garlic powder tablets. *Appl Technol Innovations* 2012; 10: 40-7.
- [59] Banerjee SK, Maulik SK. Effect of garlic on cardiovascular disorders: a review *Nutr J* 2002; 1: 4.
- [60] Thomson M, Al-Qattan KK, Bordia, T, *et al.* Including garlic in the diet may help lower blood glucose, cholesterol, and triglycerides. *J Nutr* 2006; 136: 800S-802S
- [61] Jelodar GA, Maleki M, Motadayen MH, *et al.* Effect of fenugreek, onion and garlic on blood glucose and histopathology of pancreas of alloxan-induced diabetic rats. *Indian J Med Sci* 2005; 59: 64-69.
- [62] Bidkar A, Ghadiali M, Patel C, *et al.* Anthelmintic Activities of the Crude Extracts of *Allium cepa* Bulbs and *Elletatria cardomomum* Seeds. *Res J Phar Bio Che Sci* 2012; 3: 50-58.
- [63] Ried K, Fakler P. Potential of garlic (*Allium sativum*) in lowering high blood pressure: Mechanisms of action and clinical relevance. *Integr Blood Press Control* 2014; 7: 71-82.
- [64] Mirabeau TY, Samson ES. Effect of *Allium Cepa* and *Allium Sativum* on some immunological cells in rats. *Afr J Tradit Complement Altern Med* 2012; 9: 374-79.
- [65] Wardlaw GM. *Perspective in nutrition*. 4th ed. McGraw-Hill Companies, USA, 1999; pp. 520-2.
- [66] Rice-Evans CA, Miller NJ, Bolwell PG, *et al.* The relative antioxidant activities of plant derived polyphenolic flavonoids. *Free Rad Res* 1995; 22: 375-83.
- [67] Buddington RK, Weiher E. The Application of ecological principles and fermentable fibers to manage the gastrointestinal tract ecosystem. *J Nutr* 1999; 129: 1446S-50S,
- [68] Rodgers HE, Grant HM. The effect of the flavonoids, quercetin, myricetin and epicatechin on the growth and enzyme activities of MCF7 human breast cancer cells. *Chemico-Biological Interactions* 1998; 116: 213-28.
- [69] Shen F, Herenyiova M, Weber G. Synergistic down-regulation of signal transduction and cytotoxicity by tiazofurin and quercetin in human ovarian carcinoma cells. *Life Sci* 1999; 64: 1869-76.
- [70] Shenoy N, Choughuley A. Inhibitory effect of diet related sulphdryl compounds in the formation of carcinogenic nitrosamines. *Cancer Letters* 1992; 65: 227-232.
- [71] Bianchini F, Vainio H. Allium vegetables and organosulfur compounds: Do they help prevent cancer? *Environ Health Perspec* 2001; 109: 893-902.
- [72] Fattorusso E, Lanzotti V, Tagliatalata-Scafati O, *et al.* Cytotoxic saponins from bulbs of *Allium porrum* L. *J Agric Food Chem* 2000; 48: 3455-62.
- [73] Carotenuto A, Fattorusso E, Lanzotti V, *et al.* 12-Ketoporriegenin and the unique 2,3-seco-porriegenin, new antiproliferative saponins from *Allium porrum*. *Tetrahedron* 1997; 53: 3401-6.
- [74] Sivam GP. Protection against *Helicobacter pylori* and other bacterial infections by Garlic. *Am J Nutr* 2001; 131: 11065-85.
- [75] Cellini L, Di Campli E, Masuli M, *et al.* Inhibition of *Helicobacter pylori* by garlic extract (*Allium sativum*). *FEMS Immunol Med Microbiol* 1996; 13: 273-77.
- [76] Wallace IV GC, Haar CP, Vandergrift III WA, *et al.* Multitargeted DATS prevents tumor progression and promotes apoptosis in ectopic glioblastoma xenografts in SCID mice via HDAC inhibition. *J Neurooncol* 2013; 114: 43-50.
- [77] Wang HC, Pao J, Lin SY, *et al.* Molecular mechanisms of garlic-derived allyl sulfides in the inhibition of skin cancer progression. *Ann N Y Acad Sci* 2012; 1271: 44-52.
- [78] Graf BA, Milbury PE, Blumberg JB. Flavonols, flavonones, flavanones and human health: Epidemiological evidence. *J Med Food* 2005; 8: 281-90.
- [79] Arts IC, Hollman PC. Polyphenols and disease risk in epidemiologic studies. *Am J Clin Nutr* 2005; 81: 317S-5S.
- [80] Zhang XA, Zhang S, Yin Q, *et al.* Quercetin induces human colon cancer cells apoptosis by inhibiting the nuclear factor-kappa B Pathway. *Pharmacogn Mag* 2015; 11: 404-9.
- [81] Brozyna A, Zbytek B, Granese J, *et al.* Mechanism of UV-related carcinogenesis and its contribution to nevi/melanoma. *Expert Rev Dermatol* 2007; 2: 451-69.
- [82] Lee CH, Wu SB, Hong CH, *et al.* Molecular Mechanisms of UV-Induced Apoptosis and Its Effects on Skin Residential Cells: The Implication in UV-Based Phototherapy. *Int J Mol Sci* 2013; 14: 6414-35.
- [83] Rajasekaran R, Lu KH, Sheen LY. Recent research progress on garlic as a potential anticarcinogenic agent against major digestive cancers. *J Tradit Complement Med* 2012; 2012: 192-201.
- [84] Jung M, Bu SY, Tak K-H, *et al.* Anticarcinogenic effect of quercetin by inhibition of insulin-like growth factor (IGF)-1 signaling in mouse skin cancer. *Nutr Res Pract* 2013; 7: 439-45.
- [85] Hatono S, Jimenez A, Wargovich M. Chemopreventive effect of sallylcysteine and the relationship to the detoxification enzyme glutathione S-transferase. *Carcinogenesis* 1996; 17: 1041-44.
- [86] Hu J, La Vecchia C, Negri E, *et al.* Diet and brain cancer in adults: a case-control study in northeast China. *Int J Cancer* 1999; 81: 20-23.
- [87] Galeone C, Pelucchi C, Levi F, *et al.* Onion and garlic use and human cancer. *Am J Clin Nutr* 2006; 84: 1027-32.
- [88] Hsing AW, Chokkalingam AP, Gao YT, *et al.* Allium vegetables and risk of prostate cancer: a population-based study. *J Natl Cancer Inst* 2002; 94: 1648-51.
- [89] Tsubura A, Lai YC, Kuwata M, *et al.* Anticancer effects of garlic and garlic-derived compounds for breast cancer control. *Anticancer Agents Med Chem* 2011; 11: 249-53.
- [90] Wang Y, Tian WX, Ma XF. Inhibitory effects of onion (*Allium cepa* L.) extract on proliferation of cancer cells and adipocytes via inhibiting fatty acid synthase. *Asian Pac J Cancer Prev* 2012; 13: 5573-79.
- [91] Arnault I, Auger J. Seleno-compounds in garlic and onion. *J Chromato* 2006; A 1112: 23-30.

- [92] Yu Z, Zhang T, Zhou F, *et al.* Anticancer activity of saponins from *Allium chinense* against the B16 melanoma and 4T1 breast carcinoma cell. *Evid Based Complement Alternat Med* 2015; 2015: 725023.
- [93] You W-C, Blot WJ, Chang Y-S, *et al.* Diet and high risk of stomach cancer in Shandong, China. *Cancer Res* 1988; 48: 3518-23.
- [94] Steinmetz KA, Potter JD. Vegetables, fruit, and cancer prevention: a review. *J Am Diet Assoc* 1996; 96: 1027-39.
- [95] Lawson LD, Hughes, BG. Characterization of the formation of allicin and other thiosulfonates from garlic. *Planta Med* 1992; 58:345-50.
- [96] Warshafsky S, Kamer RS, Sivak SL. Effect of garlic on total serum cholesterol. A meta-analysis. *Ann Int Med* 1993; 119: 599-605.
- [97] Silagy C, Neil A. Garlic as a lipid-lowering agent-a meta-analysis. *J Royal Coll Physicians Lond* 1994; 28: 39-45.
- [98] Jain RC. Effect of garlic on serum lipids, coagulability and fibrinolytic activity of Blood. *Am J Clin Nutr* 1977; 30: 1380-81.
- [99] Allison GL, Lowe GM, Rahman K. Aged garlic extract inhibits platelet activation by increasing intracellular cAMP and reducing the interaction of GPIIb/IIIa receptor with fibrinogen. *Life Sciences* 2012; 91: 25-6.
- [100] Sumiyoshi H, Wargovich M.J. Chemoprevention of 1,2-dimethylhydrazine-induced colon cancer in mice by naturally occurring organosulfur compounds. *Cancer Res* 1990; 50: 5084-87.
- [101] Isaacsohn JL, Moser M, Stein EA, *et al.* Garlic powder and plasma lipids and lipoproteins: A multicenter, randomized, placebo-controlled trial. *Arch Int Med* 1998; 158:1189-1194.
- [102] Hodge G, Hodge S, Han P. *Allium sativum* (garlic) suppresses leukocyte inflammatory cytokine production *in vitro*: potential therapeutic use in the treatment of inflammatory bowel disease. *Cytometry* 2002; 48: 209-15.
- [103] Arreola R, Quintero-Fabián S, López-Roa RI, *et al.* Immunomodulation and anti-inflammatory effects of garlic compounds. *J Immunol Res* 2015; 401630 doi: 10.1155/2015/401630.
- [104] Shimura M, Zhou Y, Asada Y, *et al.* Inhibition of Vpr-induced cell cycle abnormality by quercetin: A novel strategy for searching compounds targeting Vpr. *Biochem Biophys Res Comm* 1999; 261: 308-16.
- [105] Fukushima S, Takada N, Wanibuchi H, *et al.* Suppression of chemical carcinogenesis by water-soluble organosulfur compounds. *J Nutrition* 2001; 131: 1049s-53s.
- [106] Cavallito CJ, Bailey JH. Allicin, the antibacterial principle of *Allium sativum*. I. Isolation, physical properties and antibacterial action. *J Am Chem Soc* 1944; 66: 1950-51.
- [107] De Witt JC, Notermans S, Gorin N, *et al.* Effect of garlic oil or onion oil on toxin production by *Clostridium botulinum* in meat slurry. *J Food Protect* 1979; 42: 222-24.
- [108] Delaha EC, Garagusi VF. Inhibition of mycobacteria by garlic extracts (*Allium sativum*) antimicrob agents. *Chemother* 1985; 27: 485-6.
- [109] Dini C, Fabbri A, Geraci A. The potential role of garlic (*Allium sativum*) against the multi-drug resistant tuberculosis pandemic: A review. *Ann 1st Super Sanita* 2011; 47: 465-73.
- [110] Houshmand B, Mahjour F, Dianat O. Antibacterial effect of different concentrations of garlic (*Allium sativum*) extract on dental plaque bacteria. *Indian J Dent Res* 2013; 24: 71.
- [111] Yoshida S, Kasuga S, Hayashi N, *et al.* Antifungal activity of ajoene derived from garlic. *Appl Environ Microbiol* 1987; 53: 615-17.
- [112] Johnson MG, Vaughn RH. Death of *Escherichia coli* in the presence of freshly reconstituted dehydrated garlic and onion. *Appl Microbiol* 1969; 17: 903-905.
- [113] Fromtling RA, Bulmer GS. *In vitro* effect of aqueous extract of garlic (*Allium sativum*) on the growth and viability of *Cryptococcus neoformans*. *Mycologia* 1978; 70: 397-405.
- [114] Ghannoum MA. Studies on the anticandidal mode of action of *Allium sativum* (garlic). *J Gen Microbiol* 1988; 134: 2917-24.
- [115] Hitokoto H, Morozumi S, Wauke T, *et al.* Inhibitory effects of spices on growth and toxin production of toxigenic fungi. *Appl Environ Microbiol* 1980; 39: 818-22.
- [116] Lemar KM, Turner MP, Lloyd D. Garlic (*Allium sativum*) as an anti-Candida agent: a comparison of the efficacy of fresh garlic and freeze-dried extracts. *J Appl Microbiol* 2002; 93: 398-405.
- [117] Davis SR, Penie R, Apitz-Castro R. The *in vitro* susceptibility of *Scedosporium prolificans* to ajoene, allitridium and a raw extract of garlic (*Allium sativum*). *J Antimicrobiol Chemother* 2003; 5: 1593-97.
- [118] Ahmadi-Renani K, Mahmoodzadeh A, Cheraghali AM, *et al.* Effect of Garlic Extract on *Cutaneous leishmaniasis* and the role of nitric oxide. *IJMS* 2002; 27: 97-100.
- [119] Fenwick GR, Hanley AB. *Allium* species poisoning. *Vet Rec* 1985; 116: 28.
- [120] Weber ND, Andersen DO, North JA, *et al.* *In vitro* virucidal effects of *Allium sativum* (garlic) extract and compounds. *Planta Med* 1992; 58: 417-23.
- [121] Meng Y, Lu D, Guo N, *et al.* Anti-HCMV effect of garlic components. *Virol Sin* 1993; 8: 147-50.
- [122] Tatarintsev AV, Vrzhets PV, Ershov DE, *et al.* The ajoene blockade of integrin-dependent processes in an HIV-infected cell system. *Vestn Ross Akad Med Nauk* 1992; (11-12): 6-10.
- [123] Shoji S, Furuishi K, Yanase R, *et al.* Allyl compounds selectively killed human immunodeficiency virus (type 1)-infected cells. *Biochem Biophys Res Comm* 1993; 30: 610-621.
- [124] Yin MC, Cheng W.-S. Antioxidant activity of several allium members. *J Agric Food Chem* 1998; 46: 4097-01.
- [125] Nencini C, Menchiari A, Franchi GG, *et al.* *In vitro* antioxidant activity of aged extracts of some Italian *Allium* species. *Plant Foods Hum Nutr* 2011; 66: 11-6.
- [126] Capasso A. Antioxidant action and therapeutic efficacy of *Allium sativum* L. *Molecule* 2013; 18: 690-700.
- [127] Kang MJ, Kim JH, Choi HN, *et al.* Hypoglycemic effects of Welsh onion in an animal model of diabetes mellitus. *Nutr Res Pract* 2010; 4: 486-91.
- [128] Kumar R, Chhatwal S, Arora S, *et al.* Antihyperglycemic, anti-hyperlipidemic, anti-inflammatory and adenosine deaminase-lowering effects of garlic in patients with type 2 diabetes mellitus with obesity. *Diabetes Metab Syndr Obes* 2013; 6: 49-56.
- [129] Oliveira T, Figueiredo, CA, Brito, C, *et al.* *Allium cepa* L. and quercetin inhibit RANKL/porphyromonas gingivalis LPS-induced osteoclastogenesis by downregulating NF- κ B Signaling Pathway. *Evid Based Complement Alternat Med* 2015; 2015: ID 704781, 11 pages <http://dx.doi.org/10.1155/2015/704781>
- [130] Kaiser P, Youssouf MS, Tasduq SA, *et al.* Anti-allergic effects of herbal product from *Allium cepa* (bulb). *J Med Food* 2009; 12: 374-82.
- [131] Rogerio AP, Kanashiro A, Fontanari C, *et al.* Anti-inflammatory activity of quercetin and isoquercitrin in experimental murine allergic asthma. *Inflamm Res* 2007; 56: 402-8. doi: 10.1007/s00011-007-7005-6.
- [132] Park HJ, Lee CM, Jung ID, *et al.* Quercetin regulates Th1/Th2 balance in a murine model of asthma. *Int Immunopharmacol* 2009; 9: 261-67.
- [133] Elberry AA, Mufti S, Al-Maghrabi J, *et al.* Immunomodulatory effect of red onion (*Allium cepa* Linn) scale extract on experimentally induced atypical prostatic hyperplasia in Wistar rats. *Mediators Inflamm* 2014; 2014: 640746.
- [134] Guo C, Hou GQ, Li XD, *et al.* Quercetin triggers apoptosis of lipopolysaccharide (LPS)- induced osteoclast and inhibits bone resorption in RAW 264.7 cells. *Cell Physiol Biochem* 2012; 30:123-136.
- [135] Kosseva R Maria. Sources, characterization, and composition of food industry wastes. In: Kosseva RM, Webb C, Eds. *Food industry wastes assessment and recuperation of commodities*. Academic Press, London, 2013; pp. 37-60.