




ELSEVIER

Contents lists available at ScienceDirect

## The Journal of Foot &amp; Ankle Surgery

journal homepage: www.jfas.org



## Original Research

## Bilateral Hallux Valgus: A Utility Outcome Score Assessment

Asim M. Makhdom, MD, MSc<sup>1,2</sup>, Hani Sinno, MD<sup>3</sup>, Sultan Aldebeyan, MBBS, MSc<sup>1,4</sup>,  
Adam Cota, MD<sup>1</sup>, Reggie Charles Hamdy, MD, MSc, FRCS<sup>1</sup>,  
Mohammad Alzahrani, MBBS, MSc<sup>1,5</sup>, Chantal Janelle, MD, FRCS<sup>1</sup>

<sup>1</sup> Orthopedist, Division of Orthopaedic Surgery, Shriners Hospital for Children, Montreal Children Hospital, McGill University, Montreal, Quebec, Canada

<sup>2</sup> Orthopedist, Department of Orthopaedic Surgery, King Abdulaziz University, Jeddah, Saudi Arabia

<sup>3</sup> Plastic Surgeon, Division of Plastic and Reconstructive Surgery, Montreal Children's Hospital, Faculty of Medicine, McGill University, Montreal, Quebec, Canada

<sup>4</sup> Orthopedist, Department of Orthopaedic Surgery, King Fahad Medical City, Riyadh, Saudi Arabia

<sup>5</sup> Orthopedist, Department of Orthopaedic Surgery, University of Dammam, Dammam, Saudi Arabia

## ARTICLE INFO

Level of Clinical Evidence: 4

## Keywords:

bunion  
foot deformity  
hallux valgus  
standard gamble  
time trade-off  
utility score  
visual analog scale

## ABSTRACT

Hallux valgus is the most common forefoot problem in adults. Although it can cause considerable disability and affect the quality of life of those affected, many patients seek medical attention because of cosmetic concerns. Our aim was to objectively measure the perceived health burden of living with bilateral hallux valgus. Previously validated utility outcome measures, including the visual analog scale, time trade-off, and standard gamble tests, were used to quantify the health burden for single-eye blindness, double-eye blindness, and bilateral hallux valgus in 103 healthy subjects using an online survey. The Student *t* test and linear regression analysis were used for statistical analysis. The mean visual analog scale, time trade-off, and standard gamble scores for bilateral hallux valgus were  $0.86 \pm 1.6$ ,  $0.95 \pm 0.5$ , and  $0.95 \pm 0.14$ , respectively. These were significantly greater than the utility scores for single-eye and double-eye blindness ( $p < .05$ ). Age, gender, race, income, and education were not statistically significant independent predictors of the utility scores for hallux valgus. In conclusion, we have objectively demonstrated the effect of living with bilateral hallux valgus deformities. Our sample population reported being willing to undergo a procedure with a 5% mortality rate and sacrifice 1.8 years of life to attain perfect health and avoid the bilateral hallux valgus health state. Our findings will guide us in counseling our patients and understanding how they perceive their foot deformity.

© 2016 by the American College of Foot and Ankle Surgeons. All rights reserved.

Hallux valgus is the most common pathologic entity of the big toe, affecting about 28% of the population with a greater incidence seen in females in their third to fifth decade of life (1,2). Although the prevalence increases steadily with age, some studies have also reported a high incidence of hallux valgus in the juvenile and adolescent groups before skeletal maturity (3). Patients can present with pain over the medial eminence, cosmetic concerns, and problems wearing certain types of shoe wear (4). Furthermore, hallux valgus can be a disabling condition affecting a patient's quality of life and functional mobility (5,6). To the best of our knowledge, no previous studies have objectively investigated the effect of living with the physical appearance of hallux valgus.

Utility outcome scores are standardized measures designed to objectify the disease burden on an individual by assigning a value representing the relative quality of life experienced by the individual compared with a state of perfect health (7–9). Scores range from 0 (clinical death) to 1 (perfect health). Several validated measurement tools exist, and these utilities provide a method for translating qualitative descriptions of an individual's health state to quantitative values that can be used for analysis and comparisons in health economics and resource allocation (7,9). The primary aim of the present study was to measure the perceived burden of hallux valgus among the general population using 3 previously validated utility outcomes scores: standard gamble (SG), time trade-off (TTO), and visual analog scale (VAS) (9–12). The secondary aim was to investigate whether age, gender, race, education, and income were independent predictors of the utility scores for bilateral hallux valgus deformity.

**Financial Disclosure:** None reported.

**Conflict of Interest:** None reported.

Address correspondence to: Sultan Aldebeyan, MBBS, MSc, Division of Orthopaedic Surgery, Shriners Hospital for Children, Montreal Children Hospital, McGill University, 1529 Cedar Avenue, Montreal, QC H3G 1A6, Canada.

E-mail address: [sultan.aldebeyan@gmail.com](mailto:sultan.aldebeyan@gmail.com) (S. Aldebeyan).

## Materials and Methods

After receiving ethics approval for the study from our local institutional review board, an open enrollment internet-based website was created and hosted by McGill University to measure the utility scores for hallux valgus. The online survey was posted

to online classified websites to recruit a sample from the general population. All prospective participants provided electronic consent before they participated in the present study. All the participants completed anonymous demographic questionnaires, health state questionnaires, and utility assessments for scenarios describing a patient with single-eye blindness, double-eye blindness, or hallux valgus. The inclusion of the single- and double-eye blindness scenarios was to familiarize the participants with the survey process and to assess their comprehension of the study objectives. Those who assigned a higher utility score (representing a health state closer to perfect health) to double-eye blindness than to single-eye blindness were excluded from the study. All participants were required to register with a valid electronic mail address, and repeat surveys from the same electronic mail address were excluded. A health-state scenario of bilateral hallux valgus was created based on expert opinion and included a photograph of a patient's feet with bilateral hallux valgus (Fig. 1).

The SG, VAS, and TTO were the 3 utility score measures used to assess for health-related quality of life. Using more than 1 utility score helped to minimize bias and overcome inherent weaknesses associated with the individual utility tests (7,13–15). The 3 utility outcome measures were performed in the same fashion for each of the 3 health states tested (single-eye blindness, double-eye blindness, and bilateral hallux valgus).

Regarding the score assessments for hallux valgus, the VAS component asked participants to visualize themselves as the patient in the provided scenario with the feet presented in Fig. 1 and to rate the perceived effect of this health state by providing a score between 0 (death) and 100 (perfect health). The VAS utility score for bilateral hallux valgus was then calculated using the formula: utility score = participant's score/100.

For the TTO, the participants were asked the number of years they would be willing to trade to have the health state in question corrected. Using a previously described algorithm, the individuals were asked to choose between living 36 more years of their life with bilateral hallux valgus versus trading off some of those years of life to live in perfect health (no hallux valgus) (7). An indifference point representing the maximum amount of life years an individual would sacrifice to avoid bilateral hallux valgus was identified by using  $\leq 6$  iterations of a bisecting algorithm that would present varying amounts of life-years lost to the participant until an acceptable trade-off was reached. The score was then calculated using the formula: TTO utility score = (number of years living with hallux valgus – number of years traded at the indifference point)/number of years living with hallux valgus (7).

For the SG utility score, the participants had to choose between 2 options: to remain in the given health state (e.g., hallux valgus) or to gamble on a treatment with some probability of success (perfect health) and some probability of failure (death). The

initial gamble was presented as a treatment with a 99% chance of perfect health but 1% chance of death. The participants were then offered treatment with an increasing chance of death in percentages until they reached their maximum level of risk. The bisecting algorithm was then used to further define the indifference point at which the participant would accept the gamble to avoid having bilateral hallux valgus. The SG utility score was determined using the formula: SG score = 1.00 – the probability of death at the point of indifference (16). At the end of the questionnaire, the participants were asked to rate their quality of life by generating a utility score for their current health state using the previously described TTO measure.

Statistical analysis was performed using Statistical Package for Social Sciences, version 20 (SPSS Inc., Chicago, IL). A paired *t* test was used to compare the mean utility scores. Furthermore, linear regression analysis was performed using age, race, gender, income, and education as independent factors of all 3 utility measures (SG, TTO, and VAS). A *p* value < .05 was considered statistically significant.

## Results

A total of 131 participants voluntarily enrolled in the study through the website. Of the 131 patients, 28 (21.4%) were excluded, leaving 103 to be included in the analysis. Of the 103 included participants, most were female ( $n = 73$ ; 70.9%) versus male ( $n = 23$ ; 21.7%), with a mean age of  $22.5 \pm 5.6$  years. The participants were predominantly white ( $n = 58$ ; 54.6%). The remaining participant demographics are listed in Table 1.

All participants included in our study rated double-eye blindness lower in all 3 mean utility scores (VAS  $0.33 \pm 1.8$ ; TTO,  $0.62 \pm 0.28$ ; SG,  $0.61 \pm 0.26$ ) compared with single-eye blindness (VAS,  $0.63 \pm 1.7$ ; TTO,  $0.84 \pm 0.16$ ; SG,  $0.85 \pm 0.16$ ;  $p < .0001$ ; Table 2). The mean VAS, TTO, and SG hallux valgus utility scores were  $0.86 \pm 1.6$ ,  $0.95 \pm 0.5$ , and  $0.95 \pm 0.14$ , respectively. All 3 scores for hallux valgus were significantly greater than the utility scores for single-eye ( $p < .0001$ ) and double-eye ( $p < .0001$ ) blindness (Table 2). The participants rated their current quality of life (TTO,  $0.97 \pm 0.02$ ) greater than that of the patient in the hallux valgus scenario (TTO,  $0.95 \pm 0.5$ ;  $p = .02$ ).

The mean bilateral hallux valgus utility scores for the SG ( $0.95 \pm 0.14$ ) and TTO ( $0.95 \pm 0.5$ ) denote that participants were willing to undergo a procedure with a 5% mortality rate and sacrifice 1.8 years of life to attain perfect health. Age, gender, race, income, and education were not statistically significant independent predictors of the utility scores for hallux valgus using logistic regression analysis ( $p > .05$ ).

**Table 1**  
Participant demographics (N = 103)

Variable	Value
Mean age (y)	22.5 ± 5.6
Gender	
Female	73 (70.9)
Male	23 (22.3)
Not reported	7 (6.8)
Race	
African-American	2 (1.9)
Asian	12 (11.7)
White	58 (56.3)
Hispanic	1 (1)
Other	8 (7.8)
Prefer not to answer	22 (21.3)
Education	
Some college	45 (43.7)
College graduate	19 (18.4)
Graduate or professional degree	9 (8.7)
Medical education	5 (4.9)
High school	2 (1.9)
Prefer not to answer	23 (22.3)
Income	
<\$10,000	33 (32)
\$10,000 to 25,000	12 (11.7)
\$25,000 to 50,000	2 (1.9)
\$50,000 to 100,000	3 (2.9)
>\$100,000	0 (0)
Prefer not to answer	53 (51.5)

Data presented as mean ± standard deviation or n (%).



**Imagine your self like Jane:**

I have feet bunions.

I have **no** problems walking about.

I have **no** problems with self-care.

I have **no** problems performing my usual activities.

I have **occasional** pain or discomfort.

I am **minimally** anxious or depressed because of myself image.

**Fig.** Image of bilateral hallux valgus. This image was presented, along with the clinical scenario, to all participants.

**Table 2**  
Utility outcome scores (N = 103)

Test	Utility Score			p Value		Participant Self-Rated Health
	SEB	DEB	HV	SEB Versus HV	DEB Versus HV	
VAS	0.63 ± 1.7	0.33 ± 1.8	0.86 ± 1.6	<.0001	<.0001	
TTO	0.84 ± 0.16	0.62 ± 0.28	0.95 ± 0.5	<.0001	<.0001	0.97 ± 0.02*
SG	0.85 ± 0.16	0.61 ± 0.26	0.95 ± 0.14	<.0001	<.0001	

Abbreviations: DEB, double-eye blindness; HV, hallux valgus; SEB, single-eye blindness; SG, standard gamble; TTO, time trade-off; VAS, visual analog scale.

Data presented as mean ± standard deviation.

\*  $p = .02$ .

## Discussion

Hallux valgus deformity is the most common forefoot problem (17). Although conservative treatment is the first step in managing this deformity, patients with symptomatic hallux valgus who undergo surgery are much more satisfied, have lower pain intensity, and have better functional outcomes compared with those who receive prolonged nonoperative management (18). Radl et al (19) found that cosmetic concerns play an important role in hallux valgus surgery and might lead to postoperative dissatisfaction if these concerns have not been discussed and clarified before surgery. Furthermore, in joint replacement surgery, it is well recognized that patients' expectations correlate highly with postoperative satisfaction (20). The utility scores for bilateral hallux valgus reported in the present study highlight the importance of discussing patient expectations in the preoperative period.

In our study, the perceived burden of living with hallux valgus was measured objectively by using previously established utility outcome scoring models (7–12,15,21,22). Our study demonstrated that participants were willing to undergo a procedure with a 5% mortality rate (mean SG score 0.95 ± 0.14) and sacrifice 1.8 years of life (mean TTO score 0.95 ± 0.5) to attain perfect health and avoid the bilateral hallux valgus health state. Because the utility scores for hallux valgus have now been measured, we can compare them with the established utility scores of other health states. We found that comparable levels of perceived dysfunction and willingness to trade off a number of life-years are seen with other medical conditions such as erectile dysfunction (TTO score 0.96) (23), inflammatory bowel disease in adolescent patients (TTO score 0.92) (24), cystic fibrosis in adolescent patients (TTO score 0.96) (25), and functional and aesthetic nasal deformity after primary rhinoplasty (TTO score 0.90) (21). The similar TTO scores between hallux valgus and these illnesses highlight, not only the negative effect on the study participants' perceived quality of life and self-image due to the presence of bilateral hallux valgus, but also the potential positive outcome operative intervention to correct the deformity can have on an affected individual's self-image, functional status, and satisfaction.

Although age, gender, race, income, and education were not independent predictive factors of the utility outcome score for hallux valgus, our sample population was predominantly female (n = 73; 70.9%), white (58; 56.3%), young (mean age 22.5 ± 5.6 years), and healthy (self-rated TTO score 0.97 ± 0.02). The demographic data indicated a potential selection bias in the participants who volunteered for the survey, ultimately limiting the generalizability of the results to other patient populations (e.g., females in their third to fifth decade of life who are typically affected by hallux valgus).

Another potential limitation of the present study was the calculation of utility scores using participants not affected by the disease state of interest. Previous studies have reported differences in utility scores between individuals asked to visualize themselves

with a particular medical condition and individuals who actually have the condition of interest (23,26,27). This might reflect a habituation process to the health condition and can present with a decreased health burden over time (28). Future studies might focus on establishing utility scores for patients undergoing treatment for bilateral hallux valgus to assess for any differences in score magnitude compared with the healthy participants used in the present study.

In conclusion, we have demonstrated that the impact on the perceived quality of life and functional status for a person with bilateral hallux valgus deformities is comparable to living with erectile dysfunction, inflammatory bowel disease, cystic fibrosis, and a functional and aesthetic nasal deformity after primary rhinoplasty. The objective utility scores for hallux valgus allow for a quantitative measurement of the perceived disability associated with bilateral hallux valgus and can be used for cost-effectiveness analysis studies, secondary comparisons with other disease states, and to provide insight when making healthcare allocation decisions.

## Acknowledgments

Asim M. Makhdom and Hani Sinno contributed equally to this study.

## References

- Pique-Vidal C, Sole MT, Antich J. Hallux valgus inheritance: pedigree research in 350 patients with bunion deformity. *J Foot Ankle Surg* 46:149–154, 2007.
- Roddy E, Zhang W, Doherty M. Prevalence and associations of hallux valgus in a primary care population. *Arthritis Rheum* 59:857–862, 2008.
- Coughlin MJ. Juvenile hallux valgus: etiology and treatment. *Foot Ankle Int* 16:682–697, 1995.
- Mann RA, Rudicel S, Graves S. Repair of hallux valgus with a distal soft-tissue procedure and proximal metatarsal osteotomy: a long-term follow-up. *J Bone Joint Surg Am* 74:124–129, 1992.
- Menz HB, Lord SR. Gait instability in older people with hallux valgus. *Foot Ankle Int* 26:483–489, 2005.
- Menz HB, Roddy E, Thomas E, Croft PR. Impact of hallux valgus severity on general and foot-specific health-related quality of life. *Arthritis Care Res* 63:396–404, 2011.
- Chang WT, Collins ED, Kerrigan CL. An internet-based utility assessment of breast hypertrophy. *Plast Reconstr Surg* 108:370–377, 2001.
- Read JL, Quinn RJ, Berwick DM, Fineberg HV, Weinstein MC. Preferences for health outcomes: comparison of assessment methods. *Med Decis Making* 4:315–329, 1984.
- Torrance GW, Feeny D. Utilities and quality-adjusted life years. *Int J Technol Assess Health Care* 5:559–575, 1989.
- Sinno H, Thibaudeau S, Duggal A, Lessard L. Utility scores for facial disfigurement requiring facial transplantation [outcomes article]. *Plast Reconstr Surg* 126:443–449, 2010.
- Stevens KJ, McCabe CJ, Brazier JE. Mapping between visual analogue scale and standard gamble data: results from the UK Health Utilities Index 2 valuation survey. *Health Econ* 15:527–533, 2006.
- Van Osch SM, Stiggebout AM. The construction of standard gamble utilities. *Health Econ* 17:31–40, 2008.
- Boyd NF, Sutherland HJ, Tritchler DL, Cummings BJ. Whose utilities for decision analysis? *Med Decis Making* 10:58–67, 1990.
- Nord E. EuroQol: health-related quality of life measurement: valuations of health states by the general public in Norway. *Health Policy* 18:25–36, 1991.
- Sinno H, Tahiri Y, Thibaudeau S, Izadpaneh A, Christodoulou G, Lin SJ, Gilardino M. Cleft lip and palate: an objective measure outcome study. *Plast Reconstr Surg* 130:408–414, 2012.
- Liem YS, Bosch JL, Hunink MG. Preference-based quality of life of patients on renal replacement therapy: a systematic review and meta-analysis. *Value Health* 11:733–741, 2008.
- Mann RA, Coughlin MJ. Hallux valgus—etiology, anatomy, treatment and surgical considerations. *Clin Orthop Relat Res* 157:31–41, 1981.
- Torkki M, Malmivaara A, Seitsalo S, Hoikka V, Laippala P, Paavolainen P. Surgery vs orthosis vs watchful waiting for hallux valgus: a randomized controlled trial. *JAMA* 285:2474–2480, 2001.
- Radl R, Leithner A, Zacherl M, Lackner U, Egger J, Windhager R. The influence of personality traits on the subjective outcome of operative hallux valgus correction. *Int Orthop* 28:303–306, 2004.
- Maratt JD, Lee YY, Lyman S, Westrich GH. Predictors of satisfaction following total knee arthroplasty. *J Arthroplasty* 30:1142–1145, 2015.
- Sinno H, Izadpanah A, Thibaudeau S, Christodoulou G, Tahiri Y, Slavina SA, Lin SJ. The impact of living with a functional and aesthetic nasal deformity

- after primary rhinoplasty: a utility outcomes score assessment. *Ann Plast Surg* 69:431–434, 2012.
22. Sinno H, Thibaudeau S, Izadpanah A, Tahiri Y, Christodoulou G, Zukor R, Lin SJ. Utility outcome scores for unilateral facial paralysis. *Ann Plast Surg* 69:435–438, 2012.
  23. Smith KJ, Roberts MS. Quality-of-life utility values for erectile function and sildenafil treatment. *Clin Drug Investig* 25:99–105, 2005.
  24. Yi MS, Britto MT, Sherman SN, Moyer MS, Cotton S, Kotagal UR, Canfield D, Putnam FW, Carlton-Ford S, Tsevat J. Health values in adolescents with or without inflammatory bowel disease. *J Pediatr* 154:527–534, 2009.
  25. Yi MS, Britto MT, Wilmott RW, Kotagal UR, Echman MH, Neilson DW, Kociela VL, Tsevat J. Health values of adolescents with cystic fibrosis. *J Pediatr* 142:133–140, 2003.
  26. Torrance GW. Preferences for health outcomes and cost-utility analysis. *Am J Manag Care* 3(suppl):S8–S20, 1997.
  27. Torrance GW. Preferences for health states: a review of measurement methods. *Mead Johnson Symp Perinat Dev Med*:37–45, 1982.
  28. Barker JH, Furr A, Cunningham M, Grossi F, Vasilic D, Storey B, Wiggins O, Majzoub R, Vossen M, Brouha P. Investigation of risk acceptance in facial transplantation. *Plast Reconstr Surg* 118:663–670, 2006.