

In Silico Profiling of Regulatory Micronatargets in GJB3 Gene

^{1,2}*Khaista Rahman, ¹Aftab Ali Shah, ¹Muhammad Hidayatullah Khan and ³Sana Ullah*

¹Department of Biotechnology, University of Malakand, Pakistan

²Department of Biotechnology, Quaid i Azam University Islamabad, Pakistan

³Department of Animal Sciences, Quaid-i-Azam University Islamabad, Pakistan

Abstract: MicroRNAs are novel class of small non-coding RNAs that target and silence various genes across diverse signalling pathways comprising key physiological networks. Despite an important gene regulator, computational and experimental approaches are still infancy and an overall understanding of the importance of these regulatory transcripts is still far of satisfactory. Currently computer based prediction remains the only source for rapid identification of putative microRNA (miRNA) target. An increasing number of experimentally validated miRNAs targets are now available; utilizing this additional information in search of further targets may help to improve the specificity of bioinformatics based methods for predicting gene target site. An effort has been made in this pilot approach to investigate regulatory miRNA sequence and their location in the candidate gene of hearing impairment (*GJB3* gene) using online bioinformatics tool called miRDB. Using this online database, we identified eleven specific miRNAs (i.e: hsa-miR-466, hsa-miR-3613-3p, hsa-miR-16-1-3p, hsa-miR-144-3p, hsa-miR-561-5p, hsa-miR-326, hsa-miR-2110, hsa-miR-4510, hsa-miR-4419a, hsa-miR-3605-5p and hsa-miR-4738-3p). These targets different regions in the *GJB3* gene. Multiple sequence alignment to investigate whether similarities exists among mature sequences of these selected miRNAs were also performed. This data set will provide concrete bases and will help in experimental validation of these miRNAs.

Key words: Bioinformatics Tools • *GJB3* • Microna • Multiple Alignment • Mirdb

INTRODUCTION

MicroRNAs (miRNAs) are small, endogenous, noncoding RNAs, usually between 18 and 25 nucleotides in length, involved in the regulation of cellular and developmental processes through post-transcriptional gene repression [1-3]. These are expressed either as single transcription unit or as polycistronic transcripts from miRNA clusters, encoded within intronic or intergenic regions of the genome [4, 5]. Primary or pri-miRNAs are formed by polymerase II that drives transcription of miRNAs as inverted repeats embedded in long primary transcripts, which spontaneously fold to form imperfect long hairpins [6-8]. It is then processed into shorter hairpin precursor miRNAs, or pre-miRNAs, in the nucleus by RNase III enzyme complex of DROSHA (RNASEN)/DGCR8 [9]. Pre miRNAs are transferred into cytoplasm by a trans-nuclear membrane protein called exportin 5 (XPO5) [7, 8]. In cytoplasm, the RNase III enzyme DICER1 cut the pre-miRNA to form a

double-stranded miRNA duplex comprising a mature miRNA (guide strand) and a partially complementary passenger, or “star” (*), strand [10,11]. The fully matured miRNA associates with argonaute proteins in the RNA-induced silencing complex (RISC) to direct translational repression by binding to the regions of complementarity in 3'untranslated regions of target mRNAs [12,13].

RISC loading has been shown to be largely asymmetric, with only a single strand of miRNA duplex being incorporated to direct gene silencing [14]. However, some miRNA duplexes encode mature miRNAs on both strands and recent evidence suggests that strand based in miRNA expression may be influenced by tissue-specific processing factors [15]. The altered expression may be due to a variety of mechanisms including transcriptional regulation, amplification, deletion, mutation and epigenetic silencing [16]. Although miRNA signatures were established in tumor cells [17,18], recent studies revealed that the potential capabilities of miRNAs as blood-based biomarkers for cancer and other diseases[19].

Approximately 98% of RNAs in mammalian cells do not code for proteins. The epigenetic factors are associated with hearing loss [20], elevating the possibility that noncoding RNAs, such as miRNAs, might also involve in inner ear development and hearing loss.

In Silico miRNA target identification: Both computational and experimental approaches show that thousands of human genes are regulated by miRNAs [21, 22]. The functional characterization of miRNAs has become one of the most interesting research domains because of their critical roles in gene expression regulation. However, accurate target prediction is one of the major issue faced by miRNAs based research is the lack of computational tools. Various types of bioinformatics tools are developed to give insight into the molecular functionalities of microRNA gene regulatory network. One strategy for target prediction is to use machine learning approach but has not been applied to miRNA target prediction to a greater extent. An increasing number of experimentally validated microRNAs targets are now available that utilize this additional information in search of further targets. Which may help to improve the specificity of *in silico* based methods for target site prediction.

Structure and function of *GJB3*: The *GJB3* gene encoding the gap junction protein connexin 31 (Cx31) was initially mapped to chromosome 1p35- p33 and heterozygous mutations were shown to cause ADNSHL (Autosomal Dominant Non-Syndromic Hearing Loss) [23,24]. *GJB3* mutations have been reported to cause ARNSHL and a skin disorder called erythrokeratodermiavariabilis (MIM 133200). Biallelic *GJB3* mutations causing ARNSHL have been reported once in two families in which patients were compound heterozygote for two different *GJB3* mutations [25]. Digenic inheritance of non-syndromic deafness caused by mutations in *GJB2* and *GJB3* has recently been reported. Two different missense mutations (p.N166S and p.A194T) of *GJB3* were found in compound heterozygosity with the c.235delC and c.299delAT mutations of *GJB2* in three simplex families from China [26].

One gap junction consists of a cluster of closely packed pairs of trans-membrane channels, the connexins, through which materials of low molecular weight diffuse from one cell to a neighbouring cell. Involvement in disease, *GJB3* is a cause of erythrokeratodermiavariabilis (EKV) [MIM:133200]. EKV is a genodermatosis characterized by the appearance of two independent skin lesions: transient figurate erythematous patches and hyperkeratosis that is usually localized but occasionally

occurs in its generalized form. Clinical presentation varies significantly within a family and from one family to another. Palmoplantarkeratoderma is present in around 50% of all cases. Defects in *GJB3* are the cause of deafness autosomal dominant type 2B (DFNA2B) [MIM:612644]. DFNA2 is a form of sensorineural hearing loss. Sensorineural deafness results from damage to the neural receptors of the inner ear, the nerve pathways to the brain, or the area of the brain that receives sound information.

MATERIALS AND METHODS

List of genes associated with hearing loss identified in human were collected from literature and publicly available databases. All relevant publications were identified after searching PubMed (<http://www.ncbi.nlm.nih.gov/pubmed>) with key phrases, such as gene, genetics, hearing, candidate's genes for hearing, physiology of hearing impairments etc. Among the list of candidate genes, *GJB3* were selected for this study.

We used miRDB for potential miRNA target identification in *GJB3*. To briefly explain, miRDB is an online computational tool for miRNA target prediction and functional annotations [5]. All the targets were predicted by a bioinformatics tool MirTarget2, which was developed by analysing thousands of genes impacted by miRNAs with an SVM learning machine. Common features associated with miRNA target binding have been identified and used to predict miRNA targets. miRDB hosts predicted miRNA targets in five species: human, mouse, rat, dog and chicken.

RESULTS AND DISCUSSION

A large number of microRNAs have been identified across a variety of different species. Being as gene regulators, identification of microRNA targets has become an essential step toward understanding these regulatory mechanisms. *In silico* analysis of prediction presently remains the only source for rapid identification of a putative microRNA target because it helps in efficiently allocating experimental resources. *In silico* microRNA target prediction programs are based on specific parameters that can give slightly different results for the same target input. Such limitations can be partially compensated by predicting targets using more than one program. These approaches have been quite successful for a few top ranked results in different diseases' models. In the present study, an attempt was made to predict

target sites in the sequences of selected genes through advance bioinformatics approaches such as miRDB (online tool) were used.

Using this database, 11 potential miRNAs in the sequence of *GJB3* gene were identified, given in Fig. 2. This computer based approach for identification of target site in selected genes will provide base for experimental validation of these novel microRNAs in *GJB3*. If validated experimentally, these miRNAs may be used as novel biomarker for hearing impairment in human.

Salih *et al.* [27] carried out study on mutation in *GJB3* and *GJB4* genes involved in Deafness in two Sudanese families using next generation sequencing technique. Similarly a study was carried out in Kenyan and Sudanese children suffering from NSARD (non-syndromic autosomal recessive deafness) due to variants of *GJB2* gene. As compare to other areas and other ethnic groups, deafness-associated variants of the coding region of *GJB3* were rare in Kenya and Sudan, which clearly demonstrate a causative role of other genetic or epigenetic factors [28]. As compare to these studies our results also suggest that *GJB3* may be possibly responsible for hearing impairment.

GJB2 deafness-associated mutations are ethnicity-specific [30] and occur with a frequency of 30 - 50% in most Caucasian populations and 17% in Ghanaians, but they are remarkably rare among Kenyan and Sudanese populations [31]. Similarly the study of Lopez-Bigas *et al.*

[29] pointed out the strong association of E204A with deafness. From these studies [27-29] it can be hypothesized that E204A mutation has pathogenic effect on Connexins' activity which lead to hearing loss.

Trotta *et al.* [32] analysed the genotypic composition of *GJB2* gene and did not witness any significant difference between deaf and control group suggesting that *GJB2* gene or its genotypic combination with any other gene is not attributing to deafness in Northern Cameroon. Their results were different than those studies conducted at Kenya and Sudan. This also indicates that there are many factors contributing to deafness which vary within different populations such as presence of infections, malnutrition, poverty and poor access to health care. These factors may play a more predominant role than hereditary factors [33].

Details names of the candidate miRNAs, their sequence order, size, seed location, target score, target gene symbol and protein size of the target gene is given for each microRNA (Fig. 3 to Fig. 13).

Multiple alignment of selected miRNA sequences targeting *GJB3*: After identification of list of selected miRNAs (i.e: hsa-miR-466, hsa-miR-3613-3p, hsa-miR-16-1-3p, hsa-miR-144-3p, hsa-miR-561-5p, hsa-miR-326, hsa-miR-2110, hsa-miR-4510, hsa-miR-4419a, hsa-miR-3605-5p and hsa-miR-4738-3p), it was investigated, if there is any similarities among the sequences of these selected miRNAs. Multiple alignment of the mature sequences of

```

hsa-miR-466          auacacau-acacgca-----acacacau---
hsa-miR-3613-3p     -----caaaaaaaaagc---ccaacccuuc-
hsa-miR-16-1-3p     ---ccaguauuac--ugu--gcugcuga-----
hsa-miR-144-3p      -uacaguauagauaugu--acu-----
hsa-miR-561-5p      -----aucaaggauc-uuaaacuuugcc
hsa-miR-326         -----ccucugggcccuu--ccuccag--
hsa-miR-2110        -----uuggggaa-acggccgcugagug---
hsa-miR-4510        -----ugagggaguaggauuguu-----
hsa-miR-4419a       -----ugagggaggagacugca-----
hsa-miR-3605-5p     -----ugaggauaggau--agcaagggaagcc
hsa-miR-4738-3p     -----ugaaacuggag--cgccuggagga-
    
```

Fig. 1: Represents multiple sequence alignment of 11 selected miRNA genes.

Gene 2707 is predicted to be targeted by 11 miRNAs in miRDB

| Target Detail | Target Rank | Target Score | miRNA Name | Gene Symbol | Gene Description |
|-------------------------|-------------|--------------|-----------------|-------------|-------------------------------------|
| Details | 1 | 65 | hsa-miR-4510 | <i>GJB3</i> | gap junction protein, beta 3, 31kDa |
| Details | 2 | 58 | hsa-miR-4419a | <i>GJB3</i> | gap junction protein, beta 3, 31kDa |
| Details | 3 | 57 | hsa-miR-3605-5p | <i>GJB3</i> | gap junction protein, beta 3, 31kDa |
| Details | 4 | 56 | hsa-miR-16-1-3p | <i>GJB3</i> | gap junction protein, beta 3, 31kDa |
| Details | 5 | 55 | hsa-miR-144-3p | <i>GJB3</i> | gap junction protein, beta 3, 31kDa |
| Details | 6 | 55 | hsa-miR-3613-3p | <i>GJB3</i> | gap junction protein, beta 3, 31kDa |
| Details | 7 | 52 | hsa-miR-2110 | <i>GJB3</i> | gap junction protein, beta 3, 31kDa |
| Details | 8 | 51 | hsa-miR-561-5p | <i>GJB3</i> | gap junction protein, beta 3, 31kDa |
| Details | 9 | 51 | hsa-miR-4738-3p | <i>GJB3</i> | gap junction protein, beta 3, 31kDa |
| Details | 10 | 51 | hsa-miR-466 | <i>GJB3</i> | gap junction protein, beta 3, 31kDa |
| Details | 11 | 51 | hsa-miR-326 | <i>GJB3</i> | gap junction protein, beta 3, 31kDa |

Fig. 2: List of 11 miRNA predicted by online tool miRDB in the sequence of *GJB3* gene.

MicroRNA and Target Gene Description:

| | | | |
|-------------------------|-------------------------------------|--------------------------|------------------------------|
| miRNA Name | hsa-miR-4510 | miRNA Sequence | UGAGGGAGUAGGAUGUAUGUU |
| Target Score | 65 | Seed Location | 466, 630 |
| NCBI Gene ID | 2707 | GenBank Accession | NM_001005752 |
| Gene Symbol | GJB3 | 3' UTR Length | 806 |
| Gene Description | gap junction protein, beta 3, 31kDa | | |

3' UTR Sequence

```

1 CCACAGGGCA GGGTGGGGC AACATGCGG CTGCCAATG GACATGCAGG GCGGTGTGGC
61 AGGTGGAGAG GTCTACAGG GCGTGAGTGA CCCCACCTG AGTTCACTAA GTTATGCAAC
121 TTTCGTTTTG GCAGATATTT TTGACACTG GGAAGTGGG TGCTAGCCG GGTATAGSTA
181 ACCCACAGGC CCAGTGCCAG CCTCAAAGG ACATAGACTT TGAACACAGC GAATTAACATA
241 TCTACGCTGC CTGCAAGGGG CCACTTAGGG CACTGCTAGC AGGCCTTCAA CCAGGAAGGG
301 ATCAACCCAG GAAGGGATGA TCAGGAGAGG CTTCCCTGAG GACATAATGT GTAAGAGAGG
361 TGAGAAGTGC TCCCAAGCAG ACACAACAGC AGCACAGAGG TCTGGAGGCC ACACAAAAGG
421 TGATGCTCGC CTTGGGCTAG CCTCAGCAGA CCTAAGGCAT CTCTACTCCC TCCAGAGGAG
481 CCGCCAGAT TCCTGCAGTG GAGAGGAGGT CTTCCAGCAG CAGCAGGTCT GGAGGGCTGA
541 GAATGAACCT GACTAGAGGT TCTGGAGATA CCCAGAGGTC CCCCAGGTCA TCACCTTGGCT
601 CAGTGAAGC CCTCTTCCC CAAATCCTAC TCCCTCAGCC TCAGGCAGTG GTGCTCCCAT
661 CTTCTCTCCC ACAACTGTGC TCAGGCTGGT GCCAGCCTTT CAGACCCCTGC TCCCAGGGAC
721 TTGGGTGGAT GCGCTGATAG AACATCCTCA AGACAGTTTC CTTGAAATCA ATAAATACTG
781 TGTTTTATAC AAAAAAAAAA AAAAAA
    
```

Fig. 3: Illustrate the miRNA sequence (has-miR-4510), its seed sequence and it location in the sequence of target gene *GJB3* predicated by online tool miRDB.

MicroRNA and Target Gene Description:

| | | | |
|-------------------------|-------------------------------------|--------------------------|------------------------------|
| miRNA Name | hsa-miR-4419a | miRNA Sequence | UGAGGGAGGAGACUGCA |
| Target Score | 58 | Seed Location | 466, 630 |
| NCBI Gene ID | 2707 | GenBank Accession | NM_001005752 |
| Gene Symbol | GJB3 | 3' UTR Length | 806 |
| Gene Description | gap junction protein, beta 3, 31kDa | | |

3' UTR Sequence

```

1 CCACAGGGCA GGGTGGGGC AACATGCGG CTGCCAATG GACATGCAGG GCGGTGTGGC
61 AGGTGGAGAG GTCTACAGG GCGTGAGTGA CCCCACCTG AGTTCACTAA GTTATGCAAC
121 TTTCGTTTTG GCAGATATTT TTGACACTG GGAAGTGGG TGCTAGCCG GGTATAGSTA
181 ACCCACAGGC CCAGTGCCAG CCTCAAAGG ACATAGACTT TGAACACAGC GAATTAACATA
241 TCTACGCTGC CTGCAAGGGG CCACTTAGGG CACTGCTAGC AGGCCTTCAA CCAGGAAGGG
301 ATCAACCCAG GAAGGGATGA TCAGGAGAGG CTTCCCTGAG GACATAATGT GTAAGAGAGG
361 TGAGAAGTGC TCCCAAGCAG ACACAACAGC AGCACAGAGG TCTGGAGGCC ACACAAAAGG
421 TGATGCTCGC CTTGGGCTAG CCTCAGCAGA CCTAAGGCAT CTCTACTCCC TCCAGAGGAG
481 CCGCCAGAT TCCTGCAGTG GAGAGGAGGT CTTCCAGCAG CAGCAGGTCT GGAGGGCTGA
541 GAATGAACCT GACTAGAGGT TCTGGAGATA CCCAGAGGTC CCCCAGGTCA TCACCTTGGCT
601 CAGTGAAGC CCTCTTCCC CAAATCCTAC TCCCTCAGCC TCAGGCAGTG GTGCTCCCAT
661 CTTCTCTCCC ACAACTGTGC TCAGGCTGGT GCCAGCCTTT CAGACCCCTGC TCCCAGGGAC
721 TTGGGTGGAT GCGCTGATAG AACATCCTCA AGACAGTTTC CTTGAAATCA ATAAATACTG
781 TGTTTTATAC AAAAAAAAAA AAAAAA
    
```

Fig. 4: Illustrate the miRNA sequence (has-miR-4419a), its seed sequence and it location in the sequence of target gene *GJB3* predicated by online tool miRDB.

MicroRNA and Target Gene Description:

| | | | |
|-------------------------|-------------------------------------|--------------------------|------------------------------|
| miRNA Name | hsa-miR-3605-5p | miRNA Sequence | UGAGGAUGGAUAGCAAGGAAGCC |
| Target Score | 57 | Seed Location | 743 |
| NCBI Gene ID | 2707 | GenBank Accession | NM_001005752 |
| Gene Symbol | GJB3 | 3' UTR Length | 806 |
| Gene Description | gap junction protein, beta 3, 31kDa | | |

3' UTR Sequence

```

1 CCACAGGGCA GGGTGGGGC AACATGCGG CTGCCAATG GACATGCAGG GCGGTGTGGC
61 AGGTGGAGAG GTCTACAGG GCGTGAGTGA CCCCACCTG AGTTCACTAA GTTATGCAAC
121 TTTCGTTTTG GCAGATATTT TTGACACTG GGAAGTGGG TGCTAGCCG GGTATAGSTA
181 ACCCACAGGC CCAGTGCCAG CCTCAAAGG ACATAGACTT TGAACACAGC GAATTAACATA
241 TCTACGCTGC CTGCAAGGGG CCACTTAGGG CACTGCTAGC AGGCCTTCAA CCAGGAAGGG
301 ATCAACCCAG GAAGGGATGA TCAGGAGAGG CTTCCCTGAG GACATAATGT GTAAGAGAGG
361 TGAGAAGTGC TCCCAAGCAG ACACAACAGC AGCACAGAGG TCTGGAGGCC ACACAAAAGG
421 TGATGCTCGC CTTGGGCTAG CCTCAGCAGA CCTAAGGCAT CTCTACTCCC TCCAGAGGAG
481 CCGCCAGAT TCCTGCAGTG GAGAGGAGGT CTTCCAGCAG CAGCAGGTCT GGAGGGCTGA
541 GAATGAACCT GACTAGAGGT TCTGGAGATA CCCAGAGGTC CCCCAGGTCA TCACCTTGGCT
601 CAGTGAAGC CCTCTTCCC CAAATCCTAC TCCCTCAGCC TCAGGCAGTG GTGCTCCCAT
661 CTTCTCTCCC ACAACTGTGC TCAGGCTGGT GCCAGCCTTT CAGACCCCTGC TCCCAGGGAC
721 TTGGGTGGAT GCGCTGATAG AACATCCTCA AGACAGTTTC CTTGAAATCA ATAAATACTG
781 TGTTTTATAC AAAAAAAAAA AAAAAA
    
```

Fig. 5: Illustrate the miRNA sequence (has-miR-3605-5p), its seed sequence and it location in the sequence of target gene *GJB3* predicated by online tool miRDB.

MicroRNA and Target Gene Description:

| | | | |
|-------------------------|-------------------------------------|--------------------------|------------------------------|
| miRNA Name | hsa-miR-16-1-3p | miRNA Sequence | CCAGUAAUACUGUGUCUGUGA |
| Previous Name | hsa-miR-16-1* | | |
| Target Score | 56 | Seed Location | 774 |
| NCBI Gene ID | 2707 | GenBank Accession | NM_001005752 |
| Gene Symbol | GJB3 | 3' UTR Length | 806 |
| Gene Description | gap junction protein, beta 3, 31kDa | | |

3' UTR Sequence

```

1 CCACAGGGCA GGGGTGGGGC AACATGCGGG CTGCCAATGG GACATGCAGG GCGGTGTGGC
61 AGGTGGAGAG GTCCCTACAGG GGCTGAGTGA CCCCACTCTG AGTTCACATA GTTATGCAAC
121 TTTCGTTTTG GCAGATATTT TTTGACACTG GAACTGGGCG TGTCTAGCCG GGTATAGGTA
181 ACCCCAGAGC CCAAGTCCAG CCTCAAAGG ACATAGACTT TGAACAAGC GAATTAAGTA
241 TCTACGCTGC CTGCAAGGGG CCACTTAGGG CACTGCTAGC AGGGCTTCAA CCAGGAAGGG
301 ATCAACCCAG GAAGGGATGA TCAGGAGAGG CTTCCCTGAG GACATAATGT GTAAGAGAGG
361 TGAGAAGTGC TCCCAAGCAG ACACAACAGC AGCACAGAGG TCTGGAGGCC ACACAAAAGG
421 TGATGCTCGC CTTGGGCTAG CCTCAGCAGA CCTAAGGCAT CTCTACTCCC TCCAGAGGAG
481 CCGCCAGAT TCTGCAAGT GAGAGGAGGT CTTCCAGCAG CAGCAGGTCT GGAGGGCTGA
541 GAATGAACCT GACTAGAGGT TCTGGAGATA CCCAGAGGTC CCCAGGTCA TCACTTGGCT
601 CAGTGGAAAG CCTCTTTCCC CAAATCCTAC TCCCTCAGCC TCAGGCAGTG GTGCTCCCAT
661 CTTCTCCCCC ACAACTGTGC TCAGGCTGGT GCCAGCCTTT CAGACCCTGC TCCCAAGGAC
721 TTGGGTGGAT GCGCTGATAG AACATCCTCA AGACAGTTTC CTTGAAATCA ATRAATACTG
781 TGTITTTATC AAAAAAAAAA AAAAAA
    
```

Fig. 6: Illustrate the miRNA sequence (has-miR-16-1-3p), its seed sequence and it location in the sequence of target gene *GJB3* predicated by online tool miRDB.

MicroRNA and Target Gene Description:

| | | | |
|-------------------------|-------------------------------------|--------------------------|------------------------------|
| miRNA Name | hsa-miR-144-3p | miRNA Sequence | UACAGUUAUGAUGAUGUACU |
| Previous Name | hsa-miR-144 | | |
| Target Score | 55 | Seed Location | 775 |
| NCBI Gene ID | 2707 | GenBank Accession | NM_001005752 |
| Gene Symbol | GJB3 | 3' UTR Length | 806 |
| Gene Description | gap junction protein, beta 3, 31kDa | | |

3' UTR Sequence

```

1 CCACAGGGCA GGGGTGGGGC AACATGCGGG CTGCCAATGG GACATGCAGG GCGGTGTGGC
61 AGGTGGAGAG GTCCCTACAGG GGCTGAGTGA CCCCACTCTG AGTTCACATA GTTATGCAAC
121 TTTCGTTTTG GCAGATATTT TTTGACACTG GAACTGGGCG TGTCTAGCCG GGTATAGGTA
181 ACCCCAGAGC CCAAGTCCAG CCTCAAAGG ACATAGACTT TGAACAAGC GAATTAAGTA
241 TCTACGCTGC CTGCAAGGGG CCACTTAGGG CACTGCTAGC AGGGCTTCAA CCAGGAAGGG
301 ATCAACCCAG GAAGGGATGA TCAGGAGAGG CTTCCCTGAG GACATAATGT GTAAGAGAGG
361 TGAGAAGTGC TCCCAAGCAG ACACAACAGC AGCACAGAGG TCTGGAGGCC ACACAAAAGG
421 TGATGCTCGC CTTGGGCTAG CCTCAGCAGA CCTAAGGCAT CTCTACTCCC TCCAGAGGAG
481 CCGCCAGAT TCTGCAAGT GAGAGGAGGT CTTCCAGCAG CAGCAGGTCT GGAGGGCTGA
541 GAATGAACCT GACTAGAGGT TCTGGAGATA CCCAGAGGTC CCCAGGTCA TCACTTGGCT
601 CAGTGGAAAG CCTCTTTCCC CAAATCCTAC TCCCTCAGCC TCAGGCAGTG GTGCTCCCAT
661 CTTCTCCCCC ACAACTGTGC TCAGGCTGGT GCCAGCCTTT CAGACCCTGC TCCCAAGGAC
721 TTGGGTGGAT GCGCTGATAG AACATCCTCA AGACAGTTTC CTTGAAATCA ATRAATACTG
781 TGTITTTATC AAAAAAAAAA AAAAAA
    
```

Fig. 7: Illustrate themiRNA sequence (has-miR-144-3p), its seed sequence and it location in the sequence of target gene *GJB3* predicated by online tool miRDB.

MicroRNA and Target Gene Description:

| | | | |
|-------------------------|-------------------------------------|--------------------------|------------------------------|
| miRNA Name | hsa-miR-3613-3p | miRNA Sequence | ACAAAAAAAAGCCCAACCCUUC |
| Target Score | 55 | Seed Location | 138 |
| NCBI Gene ID | 2707 | GenBank Accession | NM_001005752 |
| Gene Symbol | GJB3 | 3' UTR Length | 806 |
| Gene Description | gap junction protein, beta 3, 31kDa | | |

3' UTR Sequence

```

1 CCACAGGGCA GGGGTGGGGC AACATGCGGG CTGCCAATGG GACATGCAGG GCGGTGTGGC
61 AGGTGGAGAG GTCCCTACAGG GGCTGAGTGA CCCCACTCTG AGTTCACATA GTTATGCAAC
121 TTTCGTTTTG GCAGATATTT TTTGACACTG GAACTGGGCG TGTCTAGCCG GGTATAGGTA
181 ACCCCAGAGC CCAAGTCCAG CCTCAAAGG ACATAGACTT TGAACAAGC GAATTAAGTA
241 TCTACGCTGC CTGCAAGGGG CCACTTAGGG CACTGCTAGC AGGGCTTCAA CCAGGAAGGG
301 ATCAACCCAG GAAGGGATGA TCAGGAGAGG CTTCCCTGAG GACATAATGT GTAAGAGAGG
361 TGAGAAGTGC TCCCAAGCAG ACACAACAGC AGCACAGAGG TCTGGAGGCC ACACAAAAGG
421 TGATGCTCGC CTTGGGCTAG CCTCAGCAGA CCTAAGGCAT CTCTACTCCC TCCAGAGGAG
481 CCGCCAGAT TCTGCAAGT GAGAGGAGGT CTTCCAGCAG CAGCAGGTCT GGAGGGCTGA
541 GAATGAACCT GACTAGAGGT TCTGGAGATA CCCAGAGGTC CCCAGGTCA TCACTTGGCT
601 CAGTGGAAAG CCTCTTTCCC CAAATCCTAC TCCCTCAGCC TCAGGCAGTG GTGCTCCCAT
661 CTTCTCCCCC ACAACTGTGC TCAGGCTGGT GCCAGCCTTT CAGACCCTGC TCCCAAGGAC
721 TTGGGTGGAT GCGCTGATAG AACATCCTCA AGACAGTTTC CTTGAAATCA ATRAATACTG
781 TGTITTTATC AAAAAAAAAA AAAAAA
    
```

Fig. 8: Illustrate the miRNA sequence (has-miR-3613-3p), its seed sequence and it location in the sequence of target gene *GJB3* predicated by online tool miRDB.

MicroRNA and Target Gene Description:

| | | | |
|-------------------------|-------------------------------------|--------------------------|------------------------------|
| miRNA Name | hsa-miR-2110 | miRNA Sequence | UUGGGGAAACGCCGCUGAGUG |
| Target Score | 52 | Seed Location | 616 |
| NCBI Gene ID | 2707 | GenBank Accession | NM_001005752 |
| Gene Symbol | GJB3 | 3' UTR Length | 806 |
| Gene Description | gap junction protein, beta 3, 31kDa | | |

3' UTR Sequence

```

1 CCACAGGCA GGGGTGGGC AACATGCGG CTGCCAATGG GACATGCAGG GCGGTGTGGC
61 AGGTGGAGAG GTCTACAGG GGCTGAGTGA CCCCACTCTG AGTTCACTAA GTTATGCAAC
121 TTTCGTTTTG SCAGATATTT TTTGACACTG GGAAGTGGGC TGTCTAGCCG GGTATAGGTA
181 ACCCACAGGC CCAGTGCCAG CCTTCAAGG ACATAGACTT TGAACAACGC GAATTAACCTA
241 TCTACGCTGC CTGCAAGGGG CCACTTAGGG CACTGCTAGC AGGGCTTCAA CCAGGAAGGG
301 ATCAACCCAG GAAGGGATGA TCAGGAGAGG CTTCCTGAG GACATAATGT GTAAGAGAGG
361 TGAGAAAGTGC TCCCAAGCAG ACACACACAGC AGCAGAGAGG TCTGGAGGCC ACACAAAAAG
421 TGATGCTCCG CCTGGGCTAG CCTCAGCAGA CCTAAGGCAT CTCTACTCCC TCCAGAGGAG
481 CCGCCAGAT TCTTGCAGTG GAGAGGAGGT CTTCAGCAG CAGCAGGCTC GGAGGGCTGA
541 GAATGAACCT GACTAGAGGT TCTGGAGATA CCCAGAGGTC CCCAGGTCA TCACTTGGCT
601 CAGTGGAAAG CCTCTTCCC CAATCTTAC TCCCTCAGCC TCAGGCAAGT GGTCTCCCAT
661 CTTCCTCCCC ACAACTGTGC TCAGGCTGGT GCCAGCCTTT CAGACCTGC TCCAGGGAC
721 TTGGGTGGAT GCGCTGATAG AACATCTCA AGACAGTTTC CTTGAAATCA ATAAATACTG
781 TGTTTTATAC AAAAAAAAAA AAAAAA
    
```

Fig. 9: Illustrate the miRNA sequence (has-miR-2110), its seed sequence and it location in the sequence of target gene *GJB3* predicated by online tool miRDB.

MicroRNA and Target Gene Description:

| | | | |
|-------------------------|-------------------------------------|--------------------------|------------------------------|
| miRNA Name | hsa-miR-561-5p | miRNA Sequence | AUCAAGGAUCUUAACUUUGCC |
| Target Score | 51 | Seed Location | 759 |
| NCBI Gene ID | 2707 | GenBank Accession | NM_001005752 |
| Gene Symbol | GJB3 | 3' UTR Length | 806 |
| Gene Description | gap junction protein, beta 3, 31kDa | | |

3' UTR Sequence

```

1 CCACAGGCA GGGGTGGGC AACATGCGG CTGCCAATGG GACATGCAGG GCGGTGTGGC
61 AGGTGGAGAG GTCTACAGG GGCTGAGTGA CCCCACTCTG AGTTCACTAA GTTATGCAAC
121 TTTCGTTTTG SCAGATATTT TTTGACACTG GGAAGTGGGC TGTCTAGCCG GGTATAGGTA
181 ACCCACAGGC CCAGTGCCAG CCTTCAAGG ACATAGACTT TGAACAACGC GAATTAACCTA
241 TCTACGCTGC CTGCAAGGGG CCACTTAGGG CACTGCTAGC AGGGCTTCAA CCAGGAAGGG
301 ATCAACCCAG GAAGGGATGA TCAGGAGAGG CTTCCTGAG GACATAATGT GTAAGAGAGG
361 TGAGAAAGTGC TCCCAAGCAG ACACACACAGC AGCAGAGAGG TCTGGAGGCC ACACAAAAAG
421 TGATGCTCCG CCTGGGCTAG CCTCAGCAGA CCTAAGGCAT CTCTACTCCC TCCAGAGGAG
481 CCGCCAGAT TCTTGCAGTG GAGAGGAGGT CTTCAGCAG CAGCAGGCTC GGAGGGCTGA
541 GAATGAACCT GACTAGAGGT TCTGGAGATA CCCAGAGGTC CCCAGGTCA TCACTTGGCT
601 CAGTGGAAAG CCTCTTCCC CAATCTTAC TCCCTCAGCC TCAGGCAAGT GGTCTCCCAT
661 CTTCCTCCCC ACAACTGTGC TCAGGCTGGT GCCAGCCTTT CAGACCTGC TCCAGGGAC
721 TTGGGTGGAT GCGCTGATAG AACATCTCA AGACAGTTTC CTTGAAATCA ATAAATACTG
781 TGTTTTATAC AAAAAAAAAA AAAAAA
    
```

Fig. 10: Illustrate the miRNA sequence (has-miR-561-5p), its seed sequence and it location in the sequence of target gene *GJB3* predicated by online tool miRDB.

MicroRNA and Target Gene Description:

| | | | |
|-------------------------|-------------------------------------|--------------------------|------------------------------|
| miRNA Name | hsa-miR-4738-3p | miRNA Sequence | UGAAACUGGAGCCUGGAGGA |
| Target Score | 51 | Seed Location | 754 |
| NCBI Gene ID | 2707 | GenBank Accession | NM_001005752 |
| Gene Symbol | GJB3 | 3' UTR Length | 806 |
| Gene Description | gap junction protein, beta 3, 31kDa | | |

3' UTR Sequence

```

1 CCACAGGCA GGGGTGGGC AACATGCGG CTGCCAATGG GACATGCAGG GCGGTGTGGC
61 AGGTGGAGAG GTCTACAGG GGCTGAGTGA CCCCACTCTG AGTTCACTAA GTTATGCAAC
121 TTTCGTTTTG SCAGATATTT TTTGACACTG GGAAGTGGGC TGTCTAGCCG GGTATAGGTA
181 ACCCACAGGC CCAGTGCCAG CCTTCAAGG ACATAGACTT TGAACAACGC GAATTAACCTA
241 TCTACGCTGC CTGCAAGGGG CCACTTAGGG CACTGCTAGC AGGGCTTCAA CCAGGAAGGG
301 ATCAACCCAG GAAGGGATGA TCAGGAGAGG CTTCCTGAG GACATAATGT GTAAGAGAGG
361 TGAGAAAGTGC TCCCAAGCAG ACACACACAGC AGCAGAGAGG TCTGGAGGCC ACACAAAAAG
421 TGATGCTCCG CCTGGGCTAG CCTCAGCAGA CCTAAGGCAT CTCTACTCCC TCCAGAGGAG
481 CCGCCAGAT TCTTGCAGTG GAGAGGAGGT CTTCAGCAG CAGCAGGCTC GGAGGGCTGA
541 GAATGAACCT GACTAGAGGT TCTGGAGATA CCCAGAGGTC CCCAGGTCA TCACTTGGCT
601 CAGTGGAAAG CCTCTTCCC CAATCTTAC TCCCTCAGCC TCAGGCAAGT GGTCTCCCAT
661 CTTCCTCCCC ACAACTGTGC TCAGGCTGGT GCCAGCCTTT CAGACCTGC TCCAGGGAC
721 TTGGGTGGAT GCGCTGATAG AACATCTCA AGACAGTTTC CTTGAAATCA ATAAATACTG
781 TGTTTTATAC AAAAAAAAAA AAAAAA
    
```

Fig. 11: Illustrate the miRNA sequence (has-miR-4738-3p), its seed sequence and it location in the sequence of target gene *GJB3* predicated by online tool miRDB.

| MicroRNA and Target Gene Description: | | | |
|---------------------------------------|-------------------------------------|-------------------|---|
| miRNA Name | hsa-miR-466 | miRNA Sequence | AUACACAUACCGCAACACACAU |
| Target Score | 51 | Seed Location | 347 |
| NCBI Gene ID | 2707 | GenBank Accession | NM_001005752 |
| Gene Symbol | GJB3 | 3' UTR Length | 806 |
| Gene Description | gap junction protein, beta 3, 31kDa | | |
| 3' UTR Sequence | | | |
| 1 | CCACAGGGCA | GGGGTGGGGC | AACATGCGGG CTGCCAATGG GACATGCAGG GCGGTGTGGC |
| 61 | AGGTGGAGAG | GTCTACAGG | GGCTGAGTGA CCCCCTCTG AGTTCACTAA GTTATGCAAC |
| 121 | TTTCGTTTTG | GCAGATATTT | TTTGACACTG GGAAGTGGGC TGCTAGCCG GGTATAGGTA |
| 181 | ACCCACAGGC | CCAGTGCCAG | CCTCAAAGG ACATAGACTT TGAACAACGC GAATTAAGTA |
| 241 | TCTACGCTGC | CTGCAAGGGG | CCACTTAGGG CACTGCTAGC AGGGCTTCAA CCAGGAGGGG |
| 301 | ATCAACCCAG | GAAGGGATGA | TCAGGAGAGG CTTCCCTGAG GACATAATGT GTAAGAGAGG |
| 361 | TGAGAAAGTG | TCCCAAGCAG | ACACACACAGC AGCACAGAGG TCTGGAGGCC ACACAAAAAG |
| 421 | TGATGCTCGC | CCTGGGCTAG | CCTCAGCAGA CCTAAGGCAT CTCTACTCCC TCCAGAGGAG |
| 481 | CCGCCAGAT | TCCTGCASTG | GAGAGGAGGT CTTCCAGCAG CAGCAGGCTC GGAGGGCTGA |
| 541 | GAATGAACTT | GACTAGAGGT | TCTGGAGATA CCCAGAGGTC CCCCAGGTCA TCACCTGGCT |
| 601 | CAGTGGAAAGC | CCTCTTTCCC | CAAATCCTAC TCCCTCAGCC TCAGGCAAGTGT GTGCTCCCAT |
| 661 | CTTCTCTCCC | ACAACTGTGC | TCAGGCTGGT GCCAGCCTTT CAGACCCCTGC TCCAGGGGAC |
| 721 | TTGGGTGGAT | GCGCTGATAG | AACATCCTCA AGACAGTTTC CTTGAAATCA ATAAATACTG |
| 781 | TGTTTTATAC | AAAAAAAAAA | AAAAAA |

Fig. 12: Illustrate themiRNA sequence (has-miR-466), its seed sequence and it location in the sequence of target gene *GJB3* predicated by online tool miRDB.

| MicroRNA and Target Gene Description: | | | |
|---------------------------------------|-------------------------------------|-------------------|---|
| miRNA Name | hsa-miR-326 | miRNA Sequence | CCUCUGGGCCUUCUCCAG |
| Target Score | 51 | Seed Location | 571 |
| NCBI Gene ID | 2707 | GenBank Accession | NM_001005752 |
| Gene Symbol | GJB3 | 3' UTR Length | 806 |
| Gene Description | gap junction protein, beta 3, 31kDa | | |
| 3' UTR Sequence | | | |
| 1 | CCACAGGGCA | GGGGTGGGGC | AACATGCGGG CTGCCAATGG GACATGCAGG GCGGTGTGGC |
| 61 | AGGTGGAGAG | GTCTACAGG | GGCTGAGTGA CCCCCTCTG AGTTCACTAA GTTATGCAAC |
| 121 | TTTCGTTTTG | GCAGATATTT | TTTGACACTG GGAAGTGGGC TGCTAGCCG GGTATAGGTA |
| 181 | ACCCACAGGC | CCAGTGCCAG | CCTCAAAGG ACATAGACTT TGAACAACGC GAATTAAGTA |
| 241 | TCTACGCTGC | CTGCAAGGGG | CCACTTAGGG CACTGCTAGC AGGGCTTCAA CCAGGAGGGG |
| 301 | ATCAACCCAG | GAAGGGATGA | TCAGGAGAGG CTTCCCTGAG GACATAATGT GTAAGAGAGG |
| 361 | TGAGAAAGTG | TCCCAAGCAG | ACACACACAGC AGCACAGAGG TCTGGAGGCC ACACAAAAAG |
| 421 | TGATGCTCGC | CCTGGGCTAG | CCTCAGCAGA CCTAAGGCAT CTCTACTCCC TCCAGAGGAG |
| 481 | CCGCCAGAT | TCCTGCASTG | GAGAGGAGGT CTTCCAGCAG CAGCAGGCTC GGAGGGCTGA |
| 541 | GAATGAACTT | GACTAGAGGT | TCTGGAGATA CCCAGAGGTC CCCCAGGTCA TCACCTGGCT |
| 601 | CAGTGGAAAGC | CCTCTTTCCC | CAAATCCTAC TCCCTCAGCC TCAGGCAAGTGT GTGCTCCCAT |
| 661 | CTTCTCTCCC | ACAACTGTGC | TCAGGCTGGT GCCAGCCTTT CAGACCCCTGC TCCAGGGGAC |
| 721 | TTGGGTGGAT | GCGCTGATAG | AACATCCTCA AGACAGTTTC CTTGAAATCA ATAAATACTG |
| 781 | TGTTTTATAC | AAAAAAAAAA | AAAAAA |

Fig. 13: Illustrate themiRNA sequence (has-miR-326), its seed sequence and it location in the sequence of target gene *GJB3* predicated by online tool miRDB.

these miRNAs were performed using Clustal Omega, which is maintained by EMBL-EBI (European Molecular Biology Laboratory- The European Bioinformatics Institute, i.e: part of the EMBL).

In our study the multiple alignment results showed that there is high level of sequences' variation among these selected miRNA except hsa-mir-4510, hsa-mir-4419a and hsa-miR-3605-5p where there is some sequence similarities, as shown in Fig. 1. As compare to the study of Mishra and Chandrasekharan [34], they used CLUSTAL-W alignment tool for alignment of all miRNA precursors in miRBase for selected species of hexapoda. They analysed conservation among *Apismellifera*, *Bombyxmori* and *Anopheles gambiae*. They concluded

that about 82% mature sequences fall under conserved region while 13% are outside the conserved region by 1-2 nucleotides.

CONCLUSION

It was concluded that there are 11 target sites (seed sequences) for miRNA on *GJB3* gene. These are non-identical. Every gene contains miRNA target sites that can be arranged using different bio informatics tools. This might help molecular biologists to diagnose genetic disorders such as unko gene expression rate, siRNA based drug designing and easy and short analysis of a genetically blemished pedigree for a specific

loci. In future amiRNA based DNA micro array technology can be developed for mutation's identification.

REFERENCES

1. Pasquinelli, A.E. and G. Ruvkun, 2002. Control of developmental timing by micromRNAs and their targets. *Annu. Rev. Cell Dev. Biol.*, 18: 495-513.
2. He, L. and G.J. Hannon, 2004. MicroRNAs: small RNAs with a big role in gene regulation. *Nat. Rev. Genet.*, 5(7): 522-31.
3. Sun, K. and E.C. Lai, 2013. Adult-specific functions of animal microRNAs. *Nat. Rev. Genet.*, 14(8): 535-548.
4. Kloosterman, W.P. and R.H. Plasterk, 2006. The diverse functions of microRNAs in animal development and disease. *Dev. Cell.*, 11(4): 441-450.
5. Lee, Y., K. Jeon, J.T. Lee, S. Kim and V.N. Kim, 2002. MicroRNA maturation: stepwise processing and subcellular localization. *Embryol. J.*, 21(17): 4663-4670.
6. Ambros, V., 2003. Micro RNA pathway in flies and worms: growth, death, fat, stress and timing. *Cell.*, 113(6): 673-676.
7. Tian, B., 2004. The double stranded-RNA-binding motifs: interference and much more. *Nat. Rev. Mol. Cell Biol.*, 5(12): 113-123.
8. Kim, V.N., J. Han and M.C. Siomi, 2009. Biogenesis of small RNAs in animals. *Nat. Rev. Mol. Cell Biol.*, 10(2): 126-139.
9. Moazed, D., 2009. Small RNAs in transcriptional gene silencing and genome defence. *Nature*. 457(7228): 413-420.
10. Han, J., 2004. The Drosha-DGCR8 complex in primary micro RNA processing. *Genes Dev.*, 18(24): 3016-3027.
11. Kohler, A. and E. Hurt, 2007. Exporting RNA from the nucleus to the cytoplasm. *Nat. Rev. Mol. Cell Biol.*, 8(10): 761-773.
12. Iwasaki, Y.W., K. Kiga, H. Kayo, Y. Fukuda-Yuzuwa, J. Weise, T. Inada, M. Tomita, Y. Ishihama and T. Fukao, 2013. Global microRNA elevation by inducible Exportin 5 regulates cell cycle entry. *RNA*. 19(4): 490-497.
13. Macfarlane, L.A. and P.R. Murphy, 2010. MicroRNA: Biogenesis, Function and Role in Cancer. *Current Genomics*. 11(7): 537-561.
14. Tomari, Y. and P.D. Zamore, 2005. Perspective: machines for RNAi. *Genes. Dev.*, 19(5): 517-529.
15. Ambros, V. and X. Chen, 2007. The regulation of genes and genomes by small RNAs. *Development*. 134(9): 1635-1641.
16. Pillai, R.S., 2005. MicroRNA function: multiple mechanisms for a tiny RNA?. *RNA*. 11(12): 1753-1761.
17. Thomson, D.W., C.P. Bracken and G.J. Goodall, 2011. Experimental strategies for microRNA target identification. *Nucleic Acids Res.*, 39(16): 6845-6853.
18. Siomi, H. and M.C. Siomi, 2010. Posttranscriptional regulation of microRNA biogenesis in animals. *Mol. Cell Biol.*, 38(3): 323-332.
19. Iorio, M.V. and C.M. Croce, 2012. Causes and consequences of microRNA dysregulation. *Cancer J.*, 18(3): 215-222.
20. Volinia, S., G.A. Calin, C.G. Liu, S. Ambs, A. Cimmino, F. Petrocca, R. Visone, M. Iorio, C. Roldo, M. Ferracin, R.L. Prueitt, N. Yanaihara, G. Lanza, A. Scarpa, A. Vecchione, M. Negrini, C.C. Harris and C.M. Croce, 2006. A microRNA expression signature of human solid tumours defines cancer gene targets. *Proc. Natl. Acad. Sci. USA*. 103(7): 2257-2261.
21. Jamieson, N.B., D.C. Morran, J.P. Morton, A. Ali, E.J. Dickson, C.R. Carter, O.J. Sansom, T.R. Evans, C.J. McKay and K.A. Oien, 2012. MicroRNA molecular profiles associated with diagnosis, clinicopathologic criteria and overall survival in patients with resectable pancreatic ductal adenocarcinoma. *Clin. Cancer Res.*, 18(2): 534-545.
22. Keller, A., P. Leidinger, J. Lange, A. Borries, H. Schroers, M. Scheffler, H.P. Lenhof, K. Ruprecht and E. Meese, 2009. Multiple sclerosis: microRNA expression profiles accurately differentiate patients with relapsing-remitting disease from healthy controls. *PLoS One*. 4(10): e7440.
23. Friedman, L.M., A.A. Dror, E. Mor, T. Tenne, G. Toren, T. Satoh, D.J. Biesemeier, N. Shomron, D.M. Fekete, E. Hornstein and K.B. Avraham, 2009. MicroRNAs are essential for development and function of inner ear hair cells in vertebrates. *Proc. Natl. Acad. Sci. USA*. 106(19): 7915-7920.
24. Lewis, B.P., C.B. Burge and D.P. Bartel, 2005. Conserved seed pairing, often flanked by adenosines, indicates that thousands of human genes are microRNA targets. *Cell*. 120(1): 15-20.
25. Lim, L.P., N.C. Lau, P. Garrett-Engele, A. Grimson, J.M. Schelter, J. Castle, D.P. Bartel, P.S. Linsley and J.M. Johnson, 2005. Microarray analysis shows that some microRNAs downregulate large numbers of target mRNAs. *Nature*. 433(7027): 769-773.

26. Miranda, K.C., T. Huynh, Y. Tay, Y.S. Ang, W.L. Tam, A.M. Thomson, B. Lim and I. Rigoutsos, 2006. A pattern-based method for the identification of MicroRNA binding sites and their corresponding heteroduplexes. *Cell*, 126(6): 1203-1217.
27. Salih, M.A., A.I. Dirar, N.O. Ibrahim, M.H. Osman, S.M.A. Sabeel, N.A. Yusif, A.A. Yusif, G.H. Abdelsamad and A.A Elagib, 2014. Mutation in (GJB3 and GJB4) Genes involved in Deafness in two Sudanese Families Using Next Generation Sequencing. *Int. J. Comp. Bioinfo. In Silico Model*. 3(3): 388-392.
28. Gasmelseed, N.M., M. Schmidt and M.M. Magzoub, 2004. Low frequency of deafness-associated GJB2 variants in Kenya and Sudan and novel GJB2 variants. *Hum Mutat.*, 23(2): 206-207.
29. Lopez-Bigas, N., S. Melchionda and P. Gasparini, 2002. A common frameshift mutation and other variants in GJB4 (connexin 30.3): Analysis of hearing impairment families. *Hum. Mutat.*, 19(4): 458.
30. Bonyadi, M., M. Esmaili, M. Abhari and A. Lotfi, 2009. Mutation analysis of familial GJB2-related deafness in Iranian Azeri Turkish patients. *Genet. Test Mol. Biomarkers*, 13: 689- 692.
31. Gasmelseed, N.M., M. Schmidt, M.M. Magzoub, M. Macharia and O.M. Elmustafa, 2004. Low frequency of deafness-associated GJB2 variants in Kenya and Sudan and novel GJB2 variants. *Hum Mutat.*, 23: 206-207.
32. Trotta, L., E. Lacona, P. Primignani, P. Castorina, C. Radaelli, L. Del Bo, D. Coviello and U. Ambrosetti. 2010. GJB2 and MTRNR1 contributions in children with hearing impairment from Northern Cameroon. *Int. J. Audiol.*, 33: 1-6.
33. Lasisi, O.A., J.K. Ayodele and G.T. Ijaduola, 2006. Challenges in management of childhood sensorineural hearing loss in sub-Saharan Africa, Nigeria. *Int. J. Pediatr. Otorhinolaryngol.*, 70: 625 - 629.
34. Mishra, A.K. and H. Chandrasekharan, 2014. Analytical study of Hexapoda miRNAs using phylogenetic methods. Unit of Simulation and Informatics, Indian Agricultural Research Institute, New Delhi, India. Available at: arxiv.org/ftp/arxiv/papers/1205/1205.5024.pdf