

Antioxidant and anticancer activities of *Brassica rapa*: a review

Abstract

Brassica rapa is one of the key *Brassica* species and used against different types of cancer. The different sub-species of *B. rapa* including Pak choi, Chinese cabbage, Sylvestris, Rapa and other vegetable types have both anticancer and antioxidant properties. Its various plant parts like root, leaves, fruit and seeds are full of some important anticancer and antioxidant compounds including glucosinolates, carotenoids, flavonoids, ketones, aldehydes, vitamin C, selenium, etc. are used against numerous types of cancer cell lines. However, the antioxidant and anticancer activity of *B. rapa* vary with type of genotype, changing dose concentration and associated time interval, type of solvent used for extraction, method of extraction, part of the plant used, type of cancer and cell line, gene expression of different transcription factors involved and other biotic or abiotic factors. The present review focuses on the antioxidant and anticancer activities of different *Brassica* genotypes. In addition the anti-mutagenic potentials of different *B. rapa* genotypes against various types of cancer cell lines have been discussed. However, further research is needed to identify and characterize useful bioactive compounds from diverse *B. rapa* sub-species or ecotypes against other lethal cancer types.

Keywords: anticancer, antioxidant, *brassica rapa*, cancer cell line, vegetable

Volume 3 Issue 5 - 2018

Sohail Ahmad Jan,^{1,2} Zabta Khan Shinwari,^{1,3} Maimoona Malik,⁴ Muhammad Ilyas⁵

¹Department of Biotechnology, Quaid-i-Azam University, Pakistan

²Department of Biotechnology, Hazara University, Pakistan

³Qarshi University, Lahore, Pakistan

⁴Riphah Institute of Pharmaceutical Sciences, Riphah International University Islamabad, Pakistan

⁵Department of Plant Breeding and Genetics, PMAS Arid Agriculture University Rawalpindi, Pakistan

Correspondence: Sohail Ahmad Jan, Department of Biotechnology, Hazara University, Mansehra, Pakistan, Email: sjan.parc@gmail.com

Received: July 30, 2018 | **Published:** October 03, 2018

Introduction

About 30-40% of cancer diseases are linked to imbalanced diet and other correlated factors.¹ Epidemiological reports give a direct relation between the intake of vegetables and fruits and lower mortality rate from cancer, heart and other degenerative diseases.²⁻⁵ Many vegetables of Cruciferous family are consumed in large quantities in all over the world.⁶ The important vegetable types of *Brassica oleracea* are kale, cabbage, broccoli, Brussels sprouts, cauliflower and *Brassica rapa* vegetable types are the turnip, Chinese cabbage and Pak choi.⁷ The *Brassica* vegetables i.e. broccoli, cabbage, kale, turnip, Brussels sprouts and cauliflower possess maximum antioxidant potentials.⁸ These vegetable species are full of many important secondary metabolites like glucosinolates, folic acid, vitamin C, selenium, carotenoids, and phenolics.⁹⁻¹³ and efficiently work against many types of cancer causing agents. These compounds work strongly against broad ranges of cancer types like ovary, colon, bladder, lung and breast.¹⁴⁻¹⁷ The anticancer activities of *B. rapa* against different types of cancer is given in Figure 1.




Figure 1 In-vitro and in-vivo anticancer activities of *B. rapa* against different types of cancer.

Important secondary metabolites of different *B. rapa* genotypes

Medicinal plants are used all over the world for control and treatment of many lethal diseases.¹⁸⁻²³ The mustard leaves possess many secondary metabolites having potent antioxidant activities.²⁴ The leaves of *B. campestris*/*B. rapa* show the protective activities against the chromosomal damage and oxidative stress.²⁵ Brassinosteroids are highly distributed in both pollen and seed of *B. campestris* and *B. napus* and its concentration varies with plant tissues used.²⁶ Brassinosteroids play vital role in controlling both prostate and breast cancers at micro molar concentration.²⁷⁻²⁹

The other sulphur and nitrogen based compounds like ketones, aldehyde, esters and norisoprenoids are also present.³⁰ Almost all parts (roots, both young and old leaves, flowers and stem) of *Brassica rapa* sub-species *pekinensis* contain carotenoids. However the higher concentrations of lutein and β -carotene are predominantly present in the flower and leaves.³¹ Chung et al.³² recorded high levels of glucosinolates, phenolic compounds and gene expression from the hairy root culture of *Brassica rapa* ssp. *rapa*. Aires and Carvalho³³ identified three important indole glucosinolates compounds (glucobrassicin, 4-methoxyglucobrassicin, and 1-methoxyglucobrassicin) from the hairy roots culture of *Brassica rapa* subsp. *pekinensis* by HPLC-DAD-UV/Vis method. Several other important antioxidant phenolic compounds were studied and identified from Chinese cabbage leaves³⁴⁻³⁸ Seo et al.³⁹ envisaged Glucosinolates (GSLs) amount and its expression analysis in 38 diverse *B. rapa* accessions from eight different sub-species. Twelve new GSLs were identified and were grouped into three different groups on the basis of GSL amount and its expression levels. However, they recorded varying GSLs amount and its BRMYB transcription factors (TFs) in all tested genotypes. The three BRMYB TFs (*BrMYB28.3*, *MYBB51.1* and *BrMYB122.2*) positively regulated the GSLs content in almost all genotypes. In addition high levels of gene expression of *BrMYB28s* and *BrMYB34* was observed in both biotic and abiotic stresses.

The antioxidant and anticancer activities of different *B. rapa* genotypes

The crude extract and different fractions of the fruit part of *B. rapa* shows strong antioxidant activity on glutathione peroxidase (GPx), superoxide dismutase (SOD) enzymes and total antioxidant status (TAS) in blood sample at concentration dependent manner.⁴⁰ The anticancer potential of vegetable *Brassica* species is directly linked to the presence of these bioactive substances. It enhances detoxification enzymes, minimizes the oxidative stress, evokes immune response, lower cancer risk and malignancy, inhibits the mutation and also lowers cancer cells proliferation. The presence of glucosinolates undergoes break down to other important bioactive compounds i.e. indoles and isothiocyanates with the help of important plant tissue enzyme called myrosinase. The resultant biologically active substances with the help of two phase enzymes (Phase I and II) of xenobiotic metabolism may lead the elimination of cancer causing factors; inhibit DNA methylation and cancer growth.⁴¹

The strongest antioxidant activities of the aqueous extract of both aerial parts and roots of *B. rapa* are also recorded by Beltagy.⁴² However more antioxidant activities were reported from root extract than turnip greens. The ethyl acetate fraction of *B. rapa* roots give maximum free radical scavenging, lipid peroxidation inhibition and reducing power activities due presence of highest amount of total phenolic compounds.⁴³ The antioxidant potential and phenolic content is higher in white cabbage (*B. oleracea*) than Chinese cabbage (*B. rapa*). But the antioxidant amount in cabbage was much higher in the first 8-12 weeks.⁴⁴ The choy sum genotype of *B. rapa* have higher antioxidant activity than three genotypes of *B. oleracea* genotypes (broccoli, cabbage and cauliflower). Among *B. oleracea* genotypes the broccoli shows maximum antioxidant activities followed by cabbage and cauliflower.⁴⁵ The antioxidant activities among different *Brassica* species vary with steaming, boiling and microwave processes. The antioxidant activity is higher in steamed followed by boiled and microwave respectively. In Chinese cabbage slight boiling leads to unchanged or minor decrease the antioxidant activity.⁴⁵ The nutrient rich soil increases the antioxidant capacity of green cabbage.⁴⁶ The sulphur availability leads to increased of antioxidant activity in two *B. rapa* sub-species sylvestris.⁴⁷

B. rapa has strong anticancer and moderate antioxidant activity against human lung cancer A-549 cell line (ATCC#CCL-185).^{48,49} The crude extract of *B. rapa* contain glucosinolates and it work as mutagenic agent in human lymphocytes.⁵⁰ The *in-vitro* cytotoxic activity of *B. rapa* roots was also recorded in three different types of cancer cell line including HeLa, Hep-2 and ANM-3. However, the cytotoxic potential of crude root extracts varies with type of dose concentrations and its timing and with type of cells. The maximum growth inhibition of 42 and 63% were recorded at 1250µg/ml concentration against Hep-2 and ANM-3 cell lines respectively. While, the optimum growth inhibition (64%) against HeLa cell was recorded at 10,000µg/ml concentration after exposure for 24hours exposure.⁵¹ The other important bioactive compounds of *B. rapa* i.e. brassica phenanthrene A, diarylheptanoid, 6-paradol and trans-6-shogaol, possessed high level of inhibitory activity against three human cancer cell lines i.e. HeLa, HCT-116 and MCF-7.⁵² The transgenic hairy roots of *Brassica rapa* ssp. *rapa* have more antioxidant potential than non-transgenic roots. Also the anticancer activities against MCF-7 and HT-29 cell lines were higher in hairy root culture than non-transformed roots.³² The juice of *Brassica* species provides protection to human hepatome cells against the genotoxic effect of the carcinogens.⁵³ Mayilsamy and

Krishnaswamy⁵⁴ studied the anti-tumor effect of methanolic extract and nanoparticles of *Brassica rapa* Chinensis at two different doses (800 mg/kg b.wt and 1mg/kg b.wt). They recorded strong antitumor effect against Dalton ascites lymphoma cells (DLA) in mice model experiment. They also reported that DLA tumor induced mice might survive long due to the presence of some important anti-tumor secondary metabolites.

Conclusion

Brassica rapa has rich diversity in the world including different sub-species. Its different plant parts show strong anticancer and antioxidant activities. In recent studies, its metabolites showed strong anti-cancerous potential against ovary, colon, bladder, lung, prostate, breast and many other types of cancer. The root and leaf parts of different vegetables sub-species are predominantly used against different types of cancer cell lines and showed strong inhibitory activities against cancer. But further studies are needed to identify other unknown secondary metabolites in other important *B. rapa* sub-species against broad ranges of cancer disease, as it is not possible to suggest one cultivar/variety as being the best in terms of antioxidant or anticancer properties. More nanotechnology and *Agrobacterium rhizogenes* hairy root culture techniques should be used to identify and characterize other important bioactive compounds of *B. rapa* against different types of cancer.

Acknowledgement

None

Conflict of interest

Author declare there is no conflict of interties.

References

1. Czapski J. Cancer preventing properties of cruciferous vegetables. *Veg Crops Res Bull.* 2009;70:5–18.
2. Art ICW, Hollmam PCH. Polyphenols and disease risk in epidemiologic studies. *Am J Clin Nutr.* 2005;81(1 Suppl):317–325.
3. Kaur C, Kapoor HC. Anti oxidants in fruits and vegetables—the millennium's health. *Int J Food Sci Technol.* 2001;36(7):703–725.
4. Scalbert A, Johnson IT, Saltmarsh M. Polyphenols:anti oxidants and beyond. *Am J Clin Nutr.* 2005;81(Suupl):215S–217.
5. Vita JA. Polyphenols and cardiovascular disease: effects on endothelial and platelet function. *Am J Clin Nutr.* 2005;81(suppl.):292S–297.
6. Fahey JW, Zalcmann AT, Talalay P. The chemical diversity and distribution of glucosinolates and isothiocyanates among plants. *Phytochem.* 2001;56(1):5–51.
7. Cartea ME, Francisco M, Soengas P, et al. Phenolic compounds in *Brassica* vegetables. *Molecules.* 2011;16(1):251–280.
8. Soengas Fernandez MDP, Sotelo Perez T, Velasco Pazos P, et al. Antioxidant properties of *Brassica* vegetables. *Functional Plant Sci Biotechnol.* 2011;5(2):43–55.
9. Howard LA, Jeffery EH, Wallig MA, et al. Retention of phytochemicals in fresh and processed broccoli. *J Food Sci.* 1997;62:1098–1100.
10. Femina A, Robertson JA, Waldron KW. Cauliflower (*Brassica oleracea* L.), globe artichoke (*Cynara scolymous* L.) and Chicory witloof (*Cichorium intybus*) processing by-products as source of dietary fibre. *J Sci Food Agric.* 1998;77:511–518.
11. Kushad MK, Brown AF, Kurillicn AC, et al. Variation in glucosinolates in

- vegetable crops of *Brassica oleracea*. *J Agric Food Chem*. 1999;47(4):1541–1548.
12. Conaway CC, Getachun SM, Liebes LL, et al. Disposition of glucosinolates and sulphoraphanes in human after ingestion of steam and fresh broccoli. *Nutr Cancer*. 2001;38(2):168–178.
 13. Fimognari C, Nusse M, Cesari R, et al. Growth inhibition, cell cycle arrest and apoptosis in human T-cell leukaemia by the isothiocyanates sulphoraphane. *Carcinogenesis*. 2002;23(4):581–586.
 14. Higdon JV, Delage B, Williams DE, et al. Cruciferous vegetable and human cancer risk: epidemiologic evidence and mechanistic basis. *Pharmacol Res*. 2007;55(3):224–236.
 15. Fowke JH. Head and neck cancer: a case for inhibition by isothiocyanates and indols from Cruciferous vegetables. *Eur J Cancer Prev*. 2007;16(4):348–356.
 16. Juge N, Mithen RF, Traka M. Molecular basis for chemoprevention by sulforaphane: a comprehensive review. *Cell Mol Life Sci*. 2007;64(9):1105–1127.
 17. Zhao H, Lin J, Grossman HB, et al. Dietary isothiocyanates GSTM 1, GSTT 1, NAT 2 polymorphism and bladder cancer risk. *Int J Cancer*. 2007;120(10):2208–2213.
 18. Khan I, Jan SA, Shinwari ZK, et al. Ethnobotany and medicinal uses of folklore medicinal plants belonging to family Acanthaceae: an updated review. *MOJ Biol Med*. 2017;1(2):00009.
 19. Shinwari ZK, Jan SA, Khalil AT, et al. Identification and phylogenetic analysis of selected medicinal plant species from Pakistan: DNA barcoding approach. *Pak J Bot*. 2018;50(2):553–560.
 20. Shinwari ZK, Ahmad N, Hussain J, et al. Antimicrobial evaluation and proximate profile of *Nepetale avigata*, *Nepeta kurramensis* and *Rhynchosia reniformis*. *Pak J Bot*. 2013;45(1):253–259.
 21. Ullah I, Wakeel A, Shinwari, et al. Antibacterial and antifungal activity of *Isatis tinctoria* L. using micro-plate method. *Pak J Bot*. 2017;49(5):1949–1957.
 22. Shamaki BU, Sandabe UK, Abdulrahman FI, et al. Toxicity studies and body weights changes in Wistar rats following oral administration of methanol extract from indigenous *Ganoderma* Sp. *Nigeria*. *MOJ Biol Med*. 2017;1(5):00030.
 23. Jan SA, Shinwari ZK, Zeb A, et al. Ethnobotany and research trends in *Trachyspermum ammi* L. (Ajowan) a popular folklore remedy. *Amer-Eurasian J Agric Environ Sci*. 2015;15(1):68–73.
 24. Kim HY, Yokozawa T, Cho EJ, et al. *In vitro* and *in vivo* antioxidant effects of mustard leaf (*Brassica juncea*). *Phytotherapy Res*. 2003;17(5):465–471.
 25. Tiku AB, Abraham SK, Kale RK. Protective effect of the Cruciferous vegetables mustard leaves (*Brassica campestris*) against *in vivo* chromosomal damage and oxidative stress induced by gamma radiation and genotoxic chemicals. *Environ Mol Mutagenesis*. 2008;49(5):335–342.
 26. Takatsuto S. Brassinosteroids: distribution in plants, bioassay and microanalysis by gas chromatography. *J Chromatogr*. 1994;658(1):3–5.
 27. Wachsman MB, Lopez EM, Ramirez JA, et al. Anti viral effect of brassinosteroids against herpes virus and arena viruses. *Antivir Chem Chemother*. 2000;11(1):71–77.
 28. Wachsman MB, Ramirez JA, Galagovsky LR, et al. Anti viral activity of brassinosteroids derivatives against measles virus in cell cultures. *Antivir Chem Chemother*. 2012;13(1):61–66.
 29. Malikova J, Swaczynova J, Kolar Z, et al. Anti cancer and anti-proliferative activity of natural brassinosteroids. *Photochem*. 2008;69(2):418–426.
 30. Taveira M, Fernandes F, de Pinho PG, et al. Evolution of *Brassica rapa* var. *rapa* L. volatile composition by HS-SPME and GC/IT-MS. *Microchem J*. 2009;93(2):140–146.
 31. Tuan PA, Kim JK, Lee J, et al. Analysis of carotenoid accumulation and expression of carotenoid biosynthesis genes in different organs of Chinese cabbage (*Brassica rapa* sub sp. *pekinensis*). *Excli J*. 2012;11:508–516.
 32. Chung IM, Rekha K, Rajakumar G, Thiruvengadam M (2016) Production of glucosinolates, phenolic compounds and associated gene expression profiles of hairy root cultures in turnip (*Brassica rapa* ssp. *rapa*). *3 Biotech*. 2016;6(2):175.
 33. Aires A, Carvalho R. Rapid Separation of Indole Glucosinolates in Roots of Chinese Cabbage (*Brassica rapa* Subsp. *Pekinensis*) by High-Performance Liquid Chromatography with Diode Array Detection. *Int J Anal Chem*. 2017;1–7.
 34. Seong GU, Hwang IW, Chung SK. Antioxidant capacities and polyphenolics of Chinese cabbage (*Brassica rapa* L. ssp. *Pekinensis*) leaves. *Food Chem*. 2016;199:612–618.
 35. Harbaum B, Hubbermann EM, Wolff C, et al. Identification of flavonoids and hydroxycinnamic acids in pak choi varieties (*Brassica campestris* L. ssp. *chinensis* var. *communis*) by HPLC-ESI-MSn and NMR and their quantification by HPLC-DAD. *J Agric Food Chem*. 2007;55(20):8251–8260.
 36. Kim DO, Padilla-Zakour OI, Griffiths PD. Flavonoids and antioxidant capacity of various cabbage genotypes at juvenile stage. *J Food Sci*. 2006;69(9):685–689.
 37. Miesan KH, Mohamed S. Flavonoid (myricetin, quercetin, kaempferol, luteolin, and apigenin) content of edible tropical plants. *J Agric Food Chem*. 2001;49(6):3106–3112.
 38. Roy MK, Takenaka M, Isobe S, et al. Antioxidant potential, antiproliferative activities, and phenolic content in water-soluble fractions of some commonly consumed vegetables: Effects of thermal treatment. *Food Chem*. 2007;103(1):106–114.
 39. Seo MS, Jin M, Sohn SH, et al. Expression profiles of Br MYB transcription factors related to glucosinolate biosynthesis and stress response in eight subspecies of *Brassica rapa*. *FEBS Open Bio*. 2017;7(11):1646–1659.
 40. Gul S, Ahmed S, Gul H, et al. The antioxidant potential of *Brassica rapa* L. on glutathione peroxidase, superoxide dismutase enzymes and total antioxidant status. *Rom Rev Lab Med*. 2013;21(2):161–169.
 41. Kapusta-Duch J, Kopec A, Piatkowska E, et al. The beneficial effects of *Brassica* vegetables on human health. *Panstw Zakl Hig*. 2012;63(4):389–395.
 42. Beltagy AM. Investigation of new antimicrobial and antioxidant activities of *Brassica rapa* L. *Int J Pharm Pharm Sci*. 2014;6(6): 84–88.
 43. Ryu JP, Kim DC, In MJ, et al. Antioxidant potential of ethanol extract of *Brassica rapa* L. root. *J Med Plants Res*. 2012;6(9): 1581–1584.
 44. Samec D, Piljac Zegarac J, Bogovic M, et al. Antioxidant potency of white (*Brassica oleracea* L. var. *capitata*) and Chinese (*Brassica rapa* L. var. *pekinensis* (Lour.)) cabbage: The influence of development stage, cultivar choice and seed selection. *Sci Hort*. 2011;128(2):78–83.
 45. Wachtel Galo S, Wong KW, Benzie IF. The effect of cooking on *Brassica* vegetables. *Food Chem*. 2008;110(3):706–710.
 46. Kim DO, Padilla Zakour OI, Griffiths PD. Flavonoids and antioxidant capacity of various cabbage genotypes at juvenile stage. *J Food Sci*. 2004;69(9):685–689.
 47. De Pascale S, Maggio A, Pernice R, et al. Sulphur fertilization may improve the nutritional value of *Brassica rapa* L. subsp. *sylvestris*. *Eur J Agron*. 2007;26(4):418–424.

48. Saeed M K, Anjum S, Ahmad I, et al. Nutritional facts and free radical scavenging activity of turnip (*Brassica rapa*) from Pakistan. *World Appl Sci J*. 2012;19(3):370–375.
49. Farag MA, Motaal AA. Sulforaphane composition, cytotoxic and antioxidant activity of crucifer vegetables. *J Adv Res*. 2010;1(1): 65–70.
50. Garcia EV, Lontok NN, Ramos AA. Micronucleus assay on crude petchay (*Brassica rapa chinensis*) extract: preliminary study on its cancer chemopreventive potential. *Proc Int Seminar Chem*. 2008;280–284.
51. Barakat NT, Obaid HH, Ali AM, et al. Cytotoxic effect of aqueous extract of *Brassica rapa* roots on cancer cell lines *in vitro*. *Iraqi J Sci*. 2010;51(4):550–560.
52. Wu Q, Cho JG, Yoo KH, et al. A new phenanthrene derivative and two diarylheptanoids from the roots of *Brassica rapa* ssp. *campestris* inhibit the growth of cancer cell lines and LDL-oxidation. *Arch Pharm Res*. 2013;36(4):423–429.
53. Steinkellner H, Rabot S, Freywald C, et al. Effects of cruciferous vegetables and their constituents on drug metabolizing enzymes involved in the bioactivation of DNA-reactive dietary carcinogens. *Mutat Res*. 2001;480(1):285–297.
54. Mayilsamy D, Krishnaswamy K. Evaluation of anticancer effect of *Brassica rapa chinensis* linn. using *in vivo* model. *Int J Pharm Sci Res*. 2016;7(11):4516–4522.