



# Photosynthetic Activity, Nutrient Concentration, Yield and Soil Properties Altered by Long Term INM Practices of Knol-khol (*Brassica oleracea* var. *gongylodes* L.) under Acidic Inceptisols

Suman G. Sahu <sup>a\*</sup>, Ashish Kumar Dash <sup>a</sup>, Narayan Panda <sup>a</sup>,  
Sushanta Kumar Pattanayak <sup>a</sup>, Meenakhi Prusty <sup>a</sup>, Ambika Prasad Mishra <sup>a</sup>,  
Kshitipati Padhan <sup>a</sup>, Sweta Nanda <sup>a</sup> and Bishnupriya Patra <sup>b</sup>

<sup>a</sup> Department of Soil Science and Agricultural Chemistry, College of Agriculture, Odisha University of Agriculture and Technology, Bhubaneswar, Odisha-751003, India.

<sup>b</sup> Department of Agronomy, College of Agriculture, Odisha University of Agriculture and Technology, Bhubaneswar, Odisha-751003, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/IJPSS/2022/v34i232558

### Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/93933>

Original Research Article

Received 20 September 2022  
Accepted 25 November 2022  
Published 26 November 2022

## ABSTRACT

A long-term field experiment was started in the campus of College of Agriculture, OUAT, Bhubaneswar, Odisha since *kharif* 2010 in a randomized block design with 08 treatments and three replications under a cereal-vegetable-pulse cropping system. Different observations on plant and soil parameters were taken during *kharif* 2018 to evaluate the performance of the test crop towards different package and practices followed. The test crop was Knol-khol, the 25th crop in the sequence. The treatments were T<sub>1</sub>: Control, T<sub>2</sub>: Soil Test Dose (STD), T<sub>3</sub>: STD + Farm Yard Manure (F), T<sub>4</sub>: STD + Vermicompost (VC), T<sub>5</sub>: STD + F + Biofertilizers (BFs), T<sub>6</sub>: STD + VC + BFs, T<sub>7</sub>: STD + F + L + BFs, and T<sub>8</sub>: STD + VC + L + BFs which consisted of different combinations of inorganic fertilizers, organic manures (FYM and vermicompost), ameliorant (lime) and biofertilizers. Knol-khol crop was grown in the experimental plot with different organic and inorganic fertilizers as per the treatment plans. Leaf samples from different treatments were collected at 45 days after

\*Corresponding author: E-mail: sumangsahu@gmail.com;

planting (DAP). The economic yield was highest (3.44 t ha<sup>-1</sup>) in the treatment supplied with inorganic and organic source of nutrient with lime (T<sub>7</sub>) which was at par with T<sub>8</sub> (3.34 t ha<sup>-1</sup>) whereas significantly higher than rest other treatments and lowest was obtained in STD (1.45 t ha<sup>-1</sup>). The highest values 2.68%, 1.11mg g<sup>-1</sup> of fresh leaves and 46.34 were recorded for leaf nitrogen content, chlorophyll content and SPAD value respectively were recorded with treatment T<sub>7</sub> whereas lowest value of 1.91%, 0.54 mg g<sup>-1</sup> of fresh of leaves and 21.81 were recorded for leaf nitrogen content, chlorophyll content and SPAD value respectively with T<sub>1</sub>. However, the soil properties after harvest of the crop were found to be better with T<sub>8</sub> where vermicompost was integrated with inorganic fertilizer and lime.

**Keywords:** Knol-khol; INM; photosynthetic activity; nutrient content; economic yield; soil properties.

## 1. INTRODUCTION

Indiscriminate use of chemical fertilizers has caused serious damage to the yield, soil health and ecology which also affects physico-chemical along with biological properties of soil. On the other hand, the organic sources of nutrients viz. FYM, poultry manure, neem cake etc., are gaining importance for sustainable crop production which should be integrated with chemical fertilizers for efficient utilisation of nutrients [1]. But these are not available in adequately due to increased mechanization and a decrease in farm animals. Thus, meeting the requirement of organic manures through composting vermicomposting is one of the best wastes management practices for recycling these substantial amounts of organic residues into a microbial decomposed product with better microbial activity and having plant growth regulators [2]. Inoculation of microbes as biofertilizer also enhances the bioavailability of essential nutrients to the plants in adverse condition like acidic condition [3,4] and drought condition by producing exopolysaccharides [5]. Organic matter plays an important role in improving the physico-chemical and biological properties of soil along with development of an eco-friendly system for mutual benefit of organisms for better yield of crop and soil health. The role of organic manure is more pronounced in vegetable production system than that of other cereal crops. Application of water-soluble fertilizers through foliar spray enhances the photosynthetic activity, yield, nutrient use efficiency and economics of rice [6].

Knol-khol (*Brassica oleracea* var. gongylodes L.) belongs to the family Brassicaceae that is the German name for cabbage turnip, also called kohlrabi which is generally used as a vegetable. It is high in minerals and vitamins A and C. It contains adequate amount of water (90.3 g), calories (29.0 g), protein (2.0 g), carbohydrate

(6.6 g), fiber (1.0 g) and ash (1.0 g) per 100 g of edible stem. It also contains satisfactory amount of calcium (41.0 mg), phosphorus (51.0 mg), iron (0.5 mg), sodium (8.0 mg), potassium (372.0 mg), vitamin A (20.0 mg), thiamin (0.06 mg), riboflavin (0.04 mg), niacin (0.03 mg), and vitamin C (66.0 mg) per 100 g of above ground stem and is rich in carbohydrates and minerals. It also contains the antioxidant, vitamin A, C, E, carotene and also a good source of dietary fiber. It also contains sulphoraphanes and other isothiocyanates which are believed to stimulate the production of protective enzyme in the body (Mishra et al. 2012). Nutrient requirement of Knol-khol is also high which can be managed both by combined application for organic and inorganic source of nutrients. Information regarding the effect of organic as well as inorganic source of nutrients on Knol-khol yield was scanty for Odisha soil for which present investigation was started in the campus of the College of Agriculture, OUAT, Bhubaneswar, Odisha since 2010 as a long-term field experiment where knol-khol was a test crop.

## 2. MATERIALS AND METHODS

The present investigation was conducted with the long-term experimental field laid out in 2010 at College of Agriculture, Bhubaneswar, Odisha. The experimental site is located at 20° 15' 86" north latitude and 85° 28' 68" east longitude and at an elevation of 25.9 m above mean sea level (MSL) and 60 kms west of the Bay of Bengal. The climate was hot and humid sub-tropical with dry season prevailing from October to June and wet season from July to September. The soil of the experimental site belongs to order *Inceptisols* having loamy sand texture and comes under sub-group *vertic ustochrept*. From the existing long term field experiment, the present observations were recorded during *kharif* 2018. It was designed in a randomized block design with 3 replications and 08 treatments such as T<sub>1</sub>;

Control, T<sub>2</sub>: Soil Test Dose (STD), T<sub>3</sub>: STD + Farm Yard Manure(F), T<sub>4</sub>: STD + Vermi Compost (VC), T<sub>5</sub>: STD + F + Biofertilisers (BFs), T<sub>6</sub>: STD + VC + BFs, T<sub>7</sub>: STD + F + L + BFs, and T<sub>8</sub>: STD + VC + L + BFs. The treatments were consisted of different combinations of inorganic fertilizers, organic manures (FYM and vermicompost), ameliorant (lime) and biofertilizers. As per the treatment plans Knol-khol crop was grown in the experimental plot with a spacing of 25 cm x 25 cm. Leaf samples were collected from different treatments for analysis of nitrogen, phosphorus and potassium estimated by following the standard methods as described by Page et al. [7] and Panda [8]. The chlorophyll content was estimated by ethanol extraction method as described by Panda [8]. Side by side SPAD value was recorded with the help of Soil Plant Analysis Development (SPAD) chlorophyll meter from different treatments at same growth stage of crop Knol-khol. The economic yield was also recorded at harvest of the crop.

Soil samples were collected from each experimental plot at a depth of 0-15 cm before planting the crop and after harvest of crop. Processed soil samples were preserved in polythene bags for analysis of different chemical parameters by using standard procedures [8]. The observed values of different parameters were analyzed statistically as per the procedure outlined by Gomez and Gomez [9].

### 3. RESULTS AND DISCUSSION

#### 3.1 Effect on Photosynthetic Activity and Nutrient Content

The nutrients like nitrogen (N), phosphorus (P), potassium (K) content in fresh leaves, chlorophyll content, SPAD value of fresh leaves of crop Knol-khol at 45 DAP and the economic yield are presented in Table 1. Nitrogen (N) content in fresh leaves varied from 1.91 to 2.68%. The highest N content of fresh leaves was recorded with T<sub>7</sub> (2.68%) which was significantly followed by 2.56 and 2.49 % with T<sub>8</sub> and T<sub>6</sub> respectively. The nitrogen content of 1.91% was estimated lowest in T<sub>1</sub> (control). Highest N content in fresh leaves with T<sub>7</sub> may be due to better translocation nutrients from soil to fresh leaves. The treatment which received only chemical fertilisers recorded low N content than that of other treatments receiving organic form of nutrients. The phosphorus content of fresh leaves in different treatments ranged from 0.11 to 0.44%. The

lowest was estimated in control (T<sub>1</sub>) and highest was in integrated packages (T<sub>7</sub>). There was an increase in of phosphorus content (30%) in limed package over un-limed package. Use of biofertilizers with organic manures increased 40-50% phosphorus content whereas organic manure increased 25-50% over STD with organic manure (FYM/VC). The potassium content in fresh leaves also varied between 1.07 and 1.74 %. The K content was recorded highest in treatment receiving STD+F+L+BFs (T<sub>7</sub>) and lowest in control (T<sub>1</sub>).

Photosynthetic activity in green leaves is measured by its chlorophyll content. The content varied in different package and practices which showed the influence of different source of inorganic and organic inputs in crop knol-khol. Proper integration of source of inorganic and organic (FYM or VC), inputs with biofertilizers and ameliorants (lime) significantly altered the chlorophyll content in fresh leaves of crop knol-khol. Chlorophyll content in fresh leaves of knol-khol varied from 0.54 to 1.11 mg g<sup>-1</sup> of fresh leaves in different treatments. The highest chlorophyll content of 1.11 mg g<sup>-1</sup> of fresh leaves was found with the treatment T<sub>7</sub> whereas lowest chlorophyll content of 0.54 mg g<sup>-1</sup> of fresh leaves was found with the control. The chlorophyll content of T<sub>3</sub> and T<sub>4</sub> in which chemical fertilisers added along with only organics were at par with each other. The treatments with organics and biofertilisers was recorded more chlorophyll content than that of other treatments without biofertilisers. The significant increase in amount of chlorophyll content might be due to application of organic manures which on decomposition releases magnesium and nitrogen for longer period and higher quantity. Nitrogen and magnesium are constituents of chlorophyll molecule and nitrogen is the main constituent of all amino acids and proteins which act as structural component of chloroplast. These results were in agreement with the findings of Nehra et al. [10] and Sanwal et al. [11]. The higher availability of nutrients might have increased chlorophyll content thereby SPAD value under these treatments. These results corroborated with the result of Saleh et al. [12] and Mishra et al. [13]. The SPAD value is also considered as an indicator of photosynthetic activities in green leaves. SPAD value of different treatments followed the same trend with that of chlorophyll content of fresh leaves. Significantly highest SPAD value of 46.34 was found with the treatment T<sub>7</sub> whereas the lowest value (21.81) was observed with T<sub>1</sub>.

Economic yield of knol-khol varied from 1.90 to 3.44 t ha<sup>-1</sup> in different treatments. Higher economic yield was recorded with the treatments getting integrated application of inorganic and organic source of nutrient with lime (T<sub>7</sub>) which were statistically and significantly higher than rest of the treatments. Enhanced economic yield of knol-khol due to application of integrated use nutrients might be resulted because of prolonged availability of the different macro and micro nutrients through decomposition of organics and mineralization of inorganic nutrients. This could be possible because of the influence of bioinoculants through use of biofertilizers. Use of bioinoculants infected the roots which resulted in better root proliferation, rhizosphere development, uptake of nutrients as well as water. This resulted into higher leaf area development along with higher rate of photosynthesis activity. The increase in average weight of Knol-khol knob due to INM has been also reported by Bahadur et al. [14], Maurya et al. [15] in broccoli, Padamwar and Dakore, [16]; Kumar et al. [17] in cauliflower; Upadhyay et al. [18] in cabbage; Swain et al. [1] in finger millate and Sahoo et al. [19] in a long term INM practice in cereal-vegetable-pulse cropping system and Sethi et al. [20] in production of *Acacia mangium* sapling.

### 3.2 Effect of INM on Soil Chemical Properties

Different soil parameters pertaining to present integrated study in knol knoll are presented in Table-2. The most important soil parameter pH varied from 4.33 to 5.77 in different treatments. Before planting the knol-khol the soil pH of different treatments ranged from 4.93 to 5.69. The soil was more acidic in the treatment receiving only soil test based chemical fertilizers (pH 4.33). The next less strong acidity developed in the treatments receiving STD + organics and BF's together (pH 5.37-5.49). Only organic integrated treatments recorded less acidity (pH 5.22 and 5.43) as compared to STD + organics + BF's. In control treatment, acidity development was less as compared to inorganic fertilizer treatments. Soil acidity was neutralised due to application of lime which raised the soil pH to a level of 5.77 in lime treated plots. During the initial year the soluble salt content was 0.18 dSm<sup>-1</sup>. Due to continuous cropping, content of soluble salt reduced to 0.05 dSm<sup>-1</sup> in control and 0.04-0.06 dSm<sup>-1</sup> in other treatments getting all the ingredients. The organic carbon in the soil ranged from 5.60 to 7.03 g kg<sup>-1</sup> soil in different

package of practices. Integrated treatments receiving chemical fertilizers along with farmyard manure exhibited a considerable increase in organic C content (7.03 g kg<sup>-1</sup>) over chemical fertilizer treatments might be due to addition of organic matter through FYM which stimulated higher root growth [21].

Maximum available N content (304 kg ha<sup>-1</sup>) was recorded in the treatment receiving organic manure in the form of vermicompost, inorganic manure, microbial inoculants and lime application. Nitrogen status of soil before planting of Knol-khol was increased in all the treatments except control and combined application of inorganic fertiliser with FYM and vermicompost as compared to initial status of soil (207 kg ha<sup>-1</sup>). Available N content was not increased significantly due to application of organic manure along with inorganic fertilizer than that of STD alone. However, the increase was significant due to Inclusion of biofertilizer. The increase in available N content with the incorporation of organic sources might be due to more N mineralisation [22]. Continuous removal by crops without external addition of fertilizers and FYM/VC over a period of time in control plot resulted in decline of soil available nitrogen. The result of the present study corroborated the result reported by Sharma et al. [23]. In INM treated with microbial consortia of *Azotobacter*, *Azospirillum* and PSB along with lime significantly increased the available nitrogen in soil, because of more nitrogen fixation, as well as mineralization which enhances release of plant growth regulating substances like IAA, NAA, GA, cytokinins along with production of antibiotics and biodegradation of organic matter [24].

Significantly highest (26.24 kg ha<sup>-1</sup>) amount of available P was recorded with the treatment which received all the ingredients whereas lowest (12.23 kg ha<sup>-1</sup>) value was recorded with control. Higher available phosphorus contents in soil were observed in the treated plot having combined application of NPK fertilizer with FYM or vermicompost than that of single application of NPK. Increase in available P with FYM application could be attributed more solubilization of the native P in the soil through release of various organic acids [25]. The organic anions and hydroxyl acids, such as tartaric, citric, malonic and malic acids liberated during the decomposition of organic matter might form complexes or chelates along with Fe, Al and Ca there by preventing these cations to react with

**Table 1. Nitrogen, Phosphorus, Potassium and chlorophyll content of Knol-khol under the influence of Integrated nutrient management practice**

SI. No.	INM practices	Nutrient content (%)			Total Chlorophyll content (mg g <sup>-1</sup> fresh leaves)	SPAD value	Economic Yield (t ha <sup>-1</sup> )
		N	P	K			
1	Control	1.91	0.11	1.07	0.54	21.81	1.90
2	STD	2.35	0.16	1.18	0.60	23.37	1.45
3	STD+F	2.13	0.20	1.26	0.77	28.65	2.02
4	STD+VC	2.31	0.25	1.29	0.79	30.02	2.03
5	STD+F+BFs	2.39	0.30	1.54	0.85	35.98	2.90
6	STD+VC+BFs	2.49	0.34	1.60	0.93	39.26	2.94
7	STD+F+L+BFs	2.68	0.44	1.74	1.11	46.34	3.44
8	STD+VC+L+BFs	2.56	0.40	1.68	1.08	43.68	3.34
LSD	(P=0.05)	0.24	0.077	0.276	0.15	2.51	0.06
CV	(%)	6	15	11	10	5	8

**Table 2. Effect of INM on available nutrients in soil before planting and after planting of Knol-khol**

SI. No.	INM practices	pH		EC		OC (g kg <sup>-1</sup> )		N (kg ha <sup>-1</sup> )		P (kg ha <sup>-1</sup> )		K (kg ha <sup>-1</sup> )	
		Initial	Final	Initial	Final	Initial	Final	Initial	Final	Initial	Final	Initial	Final
1	Control	5.47	5.57	0.03	0.05	5.40	5.60	193	245	11.7	12.23	86	130
2	STD	4.93	4.33	0.06	0.06	5.70	5.80	212	282	15.7	14.56	113	132
3	STD+F	5.24	5.43	0.05	0.04	6.10	6.40	206	291	17.5	16.50	136	143
4	STD+VC	5.46	5.22	0.04	0.05	6.03	6.10	203	288	18.3	18.38	139	141
5	STD+F+BFs	5.19	5.49	0.07	0.04	5.57	6.13	222	295	20.4	21.13	146	158
6	STD+VC+BFs	5.47	5.37	0.05	0.05	6.57	7.03	265	298	21.5	22.52	149	155
7	STD+F+L+BFs	5.35	5.38	0.05	0.05	5.67	6.70	287	302	21.8	25.89	150	172
8	STD+VC+L+BFs	5.69	5.77	0.05	0.05	6.07	6.90	304	317	24.6	26.24	155	191
<b>LSD (P=0.05)</b>		0.45	0.40	0.011	0.015	0.943	0.590	25	104	3.06	2.10	48.97	6.416
<b>CV (%)</b>		5	6	13	12	9	5	6	10	9	6	10	9
<b>Initial Year (2010)</b>		5.18		0.18		3.91		207		18.3		85	

phosphate. This result corroborated the result of [26]. Favourable effect of combined application of inorganic and organic source of nutrients in enhancing the availability of P was also reported by Prasad et al. [27].

Significantly highest amount (191 kg ha<sup>-1</sup>) of available K was observed with treatment T<sub>8</sub> which received all the organic and inorganic inputs whereas lowest value of 130 kg ha<sup>-1</sup> was recorded with control (T<sub>1</sub>). The treatments with STD + VC and DTS+FYM have lower available K (141 and 143 kg ha<sup>-1</sup>) than that of organics with biofertilizers (158 and 191 kg ha<sup>-1</sup>). This might be due to continuous addition of more amount of root through higher crop growth which on decomposition increases the total available P & K in the soil.

#### 4. CONCLUSION

From the present investigation, it was found that long term use of integrated package of practices taking inorganic nutrients with organics (FYM/VC), bioinoculants and lime altered the photosynthetic activity in green leaves of crop knol in terms of chlorophyll content (mg g<sup>-1</sup> fresh leaves) and SPAD value. The economic yield of Knol-khol was also recorded highest in package supplied with integrated doses of inorganic and organic source of nutrient with lime and biofertilizers. The highest leaf nutrient content (N, P and K) in fresh leaves were recorded in integrated package of practices (T<sub>7</sub>) whereas the lowest value was in control. The pH, organic carbon and other soil fertility parameters like available N, P and K were also elevated in integrated packages whereas these parameters declined critically in only inorganic package.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

- Swain P, Panda N and Pattanayak SK. Effect of long term integrated nutrient management practices on yield and nutrient uptake by finger millet (*Eleusine coracana* L.) in an acidic Inceptisols. *Annals of Plant and Soil Research*. 2021;23(4):473-476.
- Pandit L, Sethi D, Pattanayak SK and Nayak Y. Bioconversion of lignocellulosic organic wastes into nutrient rich vermicompost by *Eudrilus eugeniae*. *Bioresource Technology Reports*. 2020; 12:100580.
- Sethi D, Mohanty S, Pattanayak SK. Acid and salt tolerance behavior of *Rhizobium* isolates and their effect on microbial diversity in the rhizosphere of redgram (*Cajanus cajan* L.). *Indian Journal of Biochemistry & Biophysics*. 2019b;56:245-252.
- Khuntia D, Panda N, Mandal M, Swain P, Sahu SG, Pattanayak SK. Symbiotic effectiveness of acid tolerant nodulating rhizobia on growth, yield and nutrient uptake of pigeon pea (*Cajanus cajan* L.) in acidic Alfisols. *International Journal of Bio-resource and Stress Management*. 2022; 13(4):403-410.
- Sethi D, Mohanty S and Pattanayak SK. Effect of different carbon, nitrogen and vitamine sources on exopolysaccharide production of *Rhizobium* species isolated from root nodule of redgram. *Indian Journal of Biochemistry & Biophysics*. 2019a;56:86-93.
- Monica M, Dash AK, Panda N, Sahu SG, Prusty M, Pradhan PP. Photosynthetic activity, yield, nutrient use efficiency and economics of rice (*Oryza sativa* L.) as influenced by foliar supplementation of urea phosphate. *Journal of the Indian Society of Soil Science*. 2020;68(4): 423-430.
- Page AL, Millar RH, Keeney DR. *Methods of Soil Analysis: Part 2, Chemical and microbiological properties*, 2nd Ed., 1982, No. 9, In the series of Agronomy, American Society of Agronomy, Inc. Soil Science Society of America, Inc. Madison Wisconsin, USA; 1982.
- Panda N. *Soil, Plant, Water and Seed Testing- A Text Book*, Kalyani Publishers, New Delhi. 2019;141.
- Gomez KA and Gomez AA. *Statistical Procedure for Agriculture Research*. New York: John Wiley & Sons; 1983.
- Nehra AS, Hood IS and Singh KP. Effect of integrated nutrient management on growth and yield of wheat (*Triticum aestivum* L.). *Indian Journal of Agronomy*. 2001;112-117.

11. Sanwal SK, Lakinarayana K, Yadav RK, Rai N and Mousumi B. Effect of organic manures on soil fertility growth, physiology, yield and quality and turmeric. *Indian Journal of Horticulture*. 2007;64(4):444-449.
12. Saleh SA, ZakiMF, Nagwa, Hassan MK, Ezzo MI. Optimizing nitrogen sources and doses for optimum kohlrabi production in new reclaimed lands. *Journal of Applied Sciences Research*. 2013;9(3): 1642-1650.
13. Mishra PP, Das AK and Mishra N. Effect of integrated nutrient management on yield, quality and economics of knolkhol (*Brassica oleracea* L. cv. GONGYLODES). *Asian Journal of Horticulture*. 2014;9(2): 382-385.
14. Bahadur A, Singh J and Singh KP. Response of cabbage to organic manure and boifertilizers. *Indian Journal of Horticulture*. 2003;61(3):278-279.
15. Maurya AK, Singh MP, Srivastava BK, Singh YV, Singh DK, Singh S, Singh PK. Effect of organic manures and inorganic fertilizers on growth characters, yield and economics of sprouting broccoli cv. Fiesta. *Indian Journal of Horticulture*. 2008;65(1): 116- 118.
16. Padamwar SB and Dakore HG. Influence of organic fertilizers on morphological and nutritional parameters of cauliflower. *Bioinfolet*. 2009;6(2):158-160.
17. Kumar V, Kumar V, Tyagi AK, Singh B, Kumar N. Effect of vermicompost and VAM inoculation on growth and yield of cauliflower (*Brassica oleracea* var. Botrytis L.). *Progressive Agriculture*. 2010;10:197-199.
18. Upadhyay AK, Bahadur A, Singh J. Effect of organic manures and biofertilizers on yield, dry matter partitioning and quality traits of cabbage (*Brassicaoleracea* var. capitata L.). *Indian Journal of Agricultural Sciences*. 2012; 82(1):31-4.
19. Sahoo SK, Mishra KN, Panda N, Panda RK, Padhan K, Mohanty S, Kumar K and Sethi D. System productivity and nutrient recoveries as influenced by nine years of long term INM practices under acidic Inceptisols of India. *Biological Forum-An International Journal*. 2022;14(3):1036-1040.
20. Sethi D, Subudhi S, Rajput VD, Kusumavathi K, Sahoo TR, Dash S, Mangaraj S, Nayak DK, Pattanayak SK, Minkina T, Glinushkin AP and Kalinitchenko VP. Exploring the role of mycorrhizal and Rhizobium inoculation with organic and inorganic fertilizers on the nutrient uptake and growth of *Acacia mangium* saplings in acidic soil. *Forests*. 2021;12:1657.  
Available:<https://doi.org/10.3390/f12121657>
21. Banswasi R and Bajpai RK. Influence of organic and inorganic fertilizer sources on soil fertility, yield and nutrient uptake by wheat crop in a rice-wheat cropping system. *Journal of Soils and Crops*. 2006;16:300-304.
22. Sharma MP, Bali VS and Gupta DK.. Crop yield and properties of Inceptisols as influenced by residue management under rice-wheat cropping sequence. *Journal of Indian Society of Soil Science*. 2008a; 48:506-509.
23. Sharma RP, Datt N and Chander G. Effect of vermicompost, farmyard manure and chemical fertilizers on yield, nutrient uptake and soil fertility in okra- onion sequence in wet temperate zone of Himachal Pradesh. *Journal of the Indian Society of Soil Science*. 2008b;57:357-361.
24. Sinha RK, Valani D, Chauhan K, Agarwal S. Embarking on a second green revolution for sustainable agriculture by vermiculture biotechnology using earthworms: reviving the dreams of Sir Charles Darwin. *International Journal of Agriculture Health Safety*. 2014;1:50-64.
25. Tolanur SI and Badanur VP. Changes in organic carbon, available N, P and K under integrated use of organic manure, green manure and fertilizer on sustaining productivity of pearl millet-pigeonpea system and fertility of an inceptisol. *Journal of the Indian Society of Soil Science*. 2003;51(1):37-41.
26. Jagtap PB, Patil JD, Nimbalkar CA and Kadlag AD. Influence of integrated nutrient management on soil properties and release of nutrients in a saline-sodic soil. *Journal of the Indian Society of Soil Science*. 2007; 55:147-56.

27. Prasad PH, Bhunia P, Naik A, Thapa U. Response of nitrogen and phosphorus levels on the growth and yield of Chinese cabbage (*Brassica campestris* L. var. *pekinensis*) in the gangetic plains of West Bengal. Journal of Crop and Weed. 2009;5(2):75-77.

© 2022 Sahu et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*

<https://www.sdiarticle5.com/review-history/93933>