

Effects of exercise and physical activity on depression

P. C. Dinas · Y. Koutedakis · A. D. Flouris

Received: 14 December 2009 / Accepted: 26 October 2010 / Published online: 14 November 2010
© Royal Academy of Medicine in Ireland 2010

Abstract

Introduction Depression is a very prevalent mental disorder affecting 340 million people globally and is projected to become the leading cause of disability and the second leading contributor to the global burden of disease by the year 2020.

Aim In this paper, we review the evidence published to date in order to determine whether exercise and physical activity can be used as therapeutic means for acute and chronic depression. Topics covered include the definition, classification criteria and treatment of depression, the link between β -endorphin and exercise, the efficacy of exercise and physical activity as treatments for depression, properties of exercise stimuli used in intervention programs, as well as the efficacy of exercise and physical activity for treating depression in diseased individuals.

Conclusions The presented evidence suggests that exercise and physical activity have beneficial effects on depression symptoms that are comparable to those of antidepressant treatments.

Keywords Exercise · Depression · Physical activity · Mood · β -Endorphin

Introduction

Depression is a very prevalent mental disorder affecting 340 million people globally—independently of age, sex, and ethnic background—and is projected to become the leading cause of disability and the second leading contributor to the global burden of disease by the year 2020 [1, 2]. Current evidence also shows that 9% of children by age 14 have already experienced at least one episode of severe depression [3], while up to 10% of adolescents are affected by a major depressive disorder [4]. These statistics are a consequence of the multi-faceted effects of depression on health and well-being [1]. Indeed, individuals suffering from depression show low levels of mood, lack of interest or pleasure, disrupted appetite or sleep, low self-worth, feelings of guilt, deprived concentration and lack of energy. Very often these problems become chronic and lead to disability or, even worse to suicide attempts [1].

Recent reports conclude that depression can be reliably diagnosed in primary care and can be treated through pharmacological agents and psychotherapy with a 60–80% success [5]. However, only 10–25% of those affected by depression receive treatment for their condition, either due to lack of resources and/or trained providers or the social stigma associated with this condition [1, 5]. Therefore, it is very important to discover alternative therapeutic methods for depression. In this light, encouraging evidence has shown that exercise and physical activity have beneficial effects on depression symptoms that are often comparable to those of antidepressant treatments [6, 7]. Yet, to our knowledge, these new evidence linking exercise and physical activity with depression have not been critically evaluated. Therefore, in this paper, we review the evidence published to date in order to determine whether exercise and physical activity can be used as therapeutic means for

P. C. Dinas (✉) · Y. Koutedakis
Department of Sport and Exercise Science,
University of Thessaly, TEFAA Karies,
Trikala 42100, Greece
e-mail: pentinas@pe.uth.gr

A. D. Flouris
FAME Laboratory, Institute of Human Performance
and Rehabilitation, Centre for Research and Technology,
Thessaly, Greece

A. D. Flouris
Department of Research and Technology Development,
Biomnic Ltd., Trikala, Greece

acute and chronic depression. It is hoped that the information provided will be valuable not only to physicians, exercise psychologists and physiologists, but also to those interested in personal or public health, politics and economics.

Approach

A comprehensive search in PubMed and Science Direct was conducted using MeSH terms that are germane to acute and chronic depression (e.g., depression symptoms, mood, depressive disorder, psychotherapy) in conjunction with exercise and psychophysical mechanisms (e.g., exercise, physical activity, β -endorphin, opioids, serotonin, pituitary gland). The search also included the articles cited in the identified papers. All articles resulting from the above methods were considered in this review, and only those deemed irrelevant (i.e., not directly studying the effects of exercise and physical activity on depression) were omitted. The following pages cover information regarding exercise and physical activity as treatments for depression, the endorphin hypothesis and the hormonal effects of β -endorphins on depression, the effects and the required frequency and mode of exercise and physical activity on depression, as well as the influence of an individual's age on the successfulness of exercise and physical activity interventions for depression.

Definition, classification criteria and treatment of depression


Depression represents a major illness with both health and social consequences similar to chronic diseases such as diabetes, congestive heart failure and hypertension [5]. According to the World Health Organization, depression is an illness characterized by negative mood, decreased interest for pleasure, feelings of guilt, uneasy sleep, decreased appetite and energy, as well as poor brain concentration [1, 2]. These feelings can be either acute or chronic, resulting in a reduced interest for life which can lead to extreme actions such as suicide [2]. The criteria based on which classification of depression is accomplished are appointed by the International Classification of Diseases (ICD-10) for worldwide use [8] and the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV) developed in the United States [9]. Generally, these standardized criteria are used to separate “normal” depressed mood caused, for instance, by disappointment, from depressive disorders also known as “affective disorders” and “mood disorders” [8, 9].

Depression is normally treated with various pharmaceutical agents or psychotherapeutic interventions or a combination of these. Current evidence shows that patients seeing a primary care provider are more likely to have a failed treatment than patients seeing a psychiatrist [10]. This, however, does not suggest that pharmacological agents are unsuccessful in the treatment of depression. Indeed, a large number of studies show that different pharmacological treatments are successful in treating acute depressive episodes [5]. In this light, a successful cooperation between primary and specialty mental health sectors is crucial, since most patients with depression first seek help in primary care [11–13]. Therefore, consultative roles for mental health specialists have been recommended in some countries to support primary care physicians in the treatment of depression [14, 15]. Relatively, recent evidence suggests that an exercise science specialist may also need to complement the primary care physicians and mental health specialists in order to successfully tackle depression symptoms and episodes [5]. For instance, it is widely accepted that mood states are highly depended on endorphin secretion [16]. Moreover, according to the endorphins hypothesis, exercise augments endorphin secretion which, in turn, reduces anxiety and depression levels [17]. These notions are further discussed in the following section.

β -Endorphin and exercise

Endorphins are endogenous opioid polypeptide compounds produced by the pituitary gland and the hypothalamus in vertebrates during strenuous exercise, excitement, pain, and they resemble the opiates in their abilities to produce analgesia and a sense of well-being [16]. The opioid system plays a key role in mediating analgesia and social attachment and may also affect depression given the link between β -endorphins and depression symptoms [18–20]. To date, β -endorphin secretion has been used for the diagnosis of depression and it could be used as an agent in a therapeutic strategy [21]. Moreover, available data have shown that the μ -opioidergic system is considerably involved in the etiology of mental disorders, thus providing a rationale for the use of μ -opioid ligands such as β -endorphin in behavioral therapies [16]. A recent meta-analysis concluded that the mechanisms by which exercise may improve depression remain uncertain mainly due to methodological limitations of existing research [22]. Yet, based on recent biological evidence linking the opioid system with mood and depression [16, 19, 23], it seems reasonable to explore a relevant hypothesis that has received considerable attention. According to the ‘endorphins hypothesis’, exercise augments the secretion of endogenous opioid peptides in

Fig. 1 The endorphins hypothesis mechanism



the brain reducing pain and causing general euphoria (Fig. 1) [17, 24, 25]. In turn, the latter reduces anxiety and depression levels [17, 24]. These notions were confirmed by a recent study showing an exercise-induced beneficial effect of endorphins in mood, which suggests that further research is warranted on the endorphins hypothesis [25].

A recent study investigated the endorphins hypothesis in competitive long distance athletes (i.e., training > 4 h per week in the last 2 years) [26]. The participants underwent positron emission tomography “ligand activation” [27], toxicological examinations, and completed the visual analog mood scales [28] at rest and after 2 h of endurance running. Based on a large effect size, the results confirmed the endorphins hypothesis demonstrating that exercise leads to an increased secretion of endorphins which, in turn, improved mood states [26].

β -Endorphin, an endogenous μ -opioid receptor-selective ligand [29], has received much attention in the literature linking endorphins and depression or mood states. β -Endorphin is primarily synthesized in the anterior pituitary gland and cleaved from pro-opiomelanocortin, its larger precursor molecule [30]. It reaches the circulation either through secretion from the pituitary gland or via projection into specific brain regions through nerve fibers [30]. To date, clinical studies have shown an increased level of β -endorphin and μ -opioid receptors in plasma and brain of patients suffering from depression and schizophrenia [31–35]. Moreover, clinical trials on patients with different types of psychiatric disorders (i.e., depression, schizophrenia and neuroses) demonstrated an antidepressant effect of β -endorphin [36]. These results support, in part, the endorphins hypothesis given that exercise of sufficient intensity and duration can increase circulating β -endorphin levels [30, 37]. Indeed, β -endorphin levels of women with cardiologic problems (syndrome of prolapsed of maternal valve) show an increase following an exercise session to a degree that constitutes treatment [38]. Moreover, a recent study demonstrated that exercise and physical activity increased β -endorphin levels in plasma with positive effects on mood [39]. Interestingly, the researchers reported that, independently of sex and age, dynamic and anaerobic exercises increased β -endorphin, while resistance and aerobic exercises seem to have only small effects on β -endorphins [39]. In general, the evidence thus far

suggests that exercise-induced β -endorphin adaptations are depended on the type of exercise and population tested, and may differ in individuals with health problems. Furthermore, some of the proposed mechanisms inducing the exercise-induced β -endorphin changes include analgesia, lactate or base excess, and metabolic factors [30, 37]. Nevertheless, the social element of exercise and its effects on β -endorphin levels has not been examined. While individualized and group-based interventions appear to be equally beneficial for treating depression [40], we remain naïve as to the effects of these protocols on β -endorphin levels.

Exercise, physical activity and depression

It is known for some time that physical activity and exercise are inversely associated with depression levels. A relevant classic study was conducted by Paffenbarger and colleagues [41] who examined questionnaires of 31,000 Harvard College graduates in 1962, 1966, 1977, and 1988. Results showed that physically active individuals reported lower depression levels than their physically inactive peers. These results are supported by a more recent study that examined 5,877 individuals aged 15–54 years showing that physically active individuals report lower levels of depression than non-active individuals [42]. Moreover, there was an inverse linear relationship between physical activity participation and depression with individuals reporting as frequently active showing lower levels of depression, individuals reporting as occasionally active showing higher levels of depression and, finally, individuals reporting as inactive showing the highest levels of depression [42].

The beneficial effects of physical activity and exercise on depression symptoms and general mood have been confirmed in individuals of all ages. In children and adolescents, a small beneficial effect of exercise in reducing depression and anxiety scores has been recorded, yet the small number of studies included and the clinical diversity of participants, interventions and methods of measurement limit the ability to draw conclusions [43]. In younger adults, a previous investigation examined whether daily activity and the tendency to participate in exercise are

associated with mood states in a student cohort [44]. The results showed that mood tends to be higher in a day an individual exercises as well as that daily activity and exercise overall are strongly linked with mood states. In line with these findings, a recent study showed that exercise significantly improves mood states in non-exercisers, recreational exercisers, as well as marathon runners [45]. More importantly, the effects of exercise on mood were twofold in recreational exercisers and marathon runners [45].

In older adults, exercise interventions have shown very promising results for alleviating symptoms of major depression [46]. These findings are supported by a more recent study examining the effects of physical activity in a sample of depressed elderly patients showing that physical activity is associated with reduced concurrent depression [47]. Based on these findings, it was suggested that physical activity may be especially helpful in the context of medical problems and major life stressors, while encouraging depressed elderly patients to engage in physical activity is likely to have potential benefits on clinical depression symptoms [47]. Finally, elderly who remain physically active across time demonstrate lower levels of depression symptoms compared to those who adopt inactive lifestyles when they get older [48].

Exercise and physical activity as treatments for depression

A recent meta-analysis reported a large clinical effect of physical activity and exercise interventions on the symptoms of depression for adults of both sexes [22]. More specifically, aerobic exercise indicated a moderate clinical effect, while mixed and resistance exercise indicated large effect sizes. Moreover, when compared against other established treatments (i.e., cognitive behavior therapy and antidepressants), exercise appears to produce the same results [22]. Indeed, published evidence shows that exercise and physical activity interventions are generally successful in reducing depression symptoms. For instance, a recent study found that patients with major depression receiving aerobic exercise training performed either at home or in a supervised group setting achieved reductions in depression comparable to standard antidepressant medication (sertraline) and greater reductions in depression compared to placebo controls [7]. Interestingly, a different study reported that individuals with major depressive disorder undergoing an aerobic exercise program were as likely to be in remission as those taking standard antidepressant medication (sertraline) or medication and exercise combined [49]. These results are further supported by a study showing that a 4-month

aerobic exercise program was more successful in reducing depression symptoms than a placebo and a pharmacological treatment [50]. In line with these findings, a recent study demonstrated that an 8-month exercise program was successful in improving depression symptoms in 40–60 year-old depressed women, unlike a pharmacological treatment [51].

Properties of exercise stimulus

Methodological limitations in existing research preclude a clear identification of the optimum exercise properties for treating depression [22]. However, it is important to acknowledge that the majority of relevant published evidence suggests an inverted dose–response relationship between exercise frequency and depression symptoms. More specifically, a recent study examined the efficacy of low-frequency (one aerobic session/week) and high-frequency (3–5 aerobic exercise sessions/week) exercise interventions for 8 weeks showing that high-frequency exercise interventions are more efficacious in reducing depression symptoms [40]. These findings are supported by a Finnish study of 3,403 adults which demonstrated that exercise frequency is inversely associated with depression symptoms and overall well-being [52].

Based on evidence published to date, the duration of exercise intervention is not clearly related to outcome [22]. On the other hand, some—but not all [43]—evidence supports that more intense exercise regimes lead to larger improvements in mood [53–55], while aerobic + resistance training appears to generate stronger effects than aerobic + flexibility training [53]. These results mirror the finding that dynamic and anaerobic exercises have a more potent effect on β -endorphin than resistance and aerobic exercises [39]. Individualized and group-based interventions appear to be equally beneficial for treating depression [40]. Moreover, several sources of data—but not all [22, 40]—show that supervised physical activity programs may produce more potent effects on depression than non-supervised programs [7, 56, 57]. These findings may be explained by the fact that depressed patients in unsupervised programs fail to sustain a high-enough interest in order to complete the prescribed intervention, as shown in a previous study [58].

Exercise and physical activity for treating depression in diseased individuals

Two previous studies examining the efficacy of exercise interventions as a treatment for depression in sedentary individuals with hypertension reported significant

reduction of depression symptoms [57, 59]. Previous work in cardiovascular disease patients has shown that adding high-intensity resistance training to an outpatient cardiac rehabilitation aerobic exercise program is more efficacious in improving mood and depression levels compared to adding flexibility training [53]. Moreover, aerobic exercise appears to be beneficial in improving depression symptoms and quality of life in patients recovering from a heart attack [60]. Therefore, there is some support to the notion that the beneficial effects of exercise and physical activity on anxiety and depression can reduce the risk for death from cardiovascular illnesses, obesity and osteoporosis in cardiovascular disease patients [61]. However, given that classic depression treatments (i.e., not based on exercise/physical activity) do not reduce the associated morbidity/mortality in cardiac patients [62, 63], the beneficial effects of exercise and physical activity on cardiovascular morbidity/mortality may be independent of their effects on depression.

Discussion

Based on the presented evidence, it becomes clear that exercise and physical activity have beneficial effects on depression symptoms that are comparable to those of antidepressant treatments. Research thus far is showing that exercise and physical activity can be used as therapeutic means for acute and chronic depression in the general population as well as in hypertensive and cardiovascular disease patients. In this light, the presented evidence supports that an exercise science specialist may need to complement the primary care physicians and mental health specialists in order to successfully tackle depression symptoms and episodes.

Notwithstanding the attention on the effects of exercise and physical activity on depression and the excitement for the new discoveries in this area, our knowledge on the biophysical mechanisms (e.g., β -endorphins) involved in the exercise-induced decrease of depression symptoms is incomplete. Moreover, methodological limitations in existing research preclude a clear identification of the optimum exercise properties for treating depression, while the risks and cost effectiveness associated with relevant interventions remain unknown. These topics should be addressed by future research in order to improve and/or standardize the exercise prescription for individuals with depression symptoms.

Acknowledgments This work was supported in part by funding from the European Union 7th Framework Program (FP7-PEOPLE-IRG-2008 grant no. 239521).

References

- Murray CJL, Lopez AD (1996) The global burden of disease: a comprehensive assessment of mortality and disability from diseases, injuries and risk factors in 1990 and projected to 2020. Harvard University Press, Cambridge, MA
- Kruijshaar ME, Hoeymans N, Spijker J, Stouthard ME, Essink-Bot ML (2005) Has the burden of depression been overestimated? *Bull World Health Org* 83(6):443–448
- Lewinsohn PM, Rohde P, Seeley JR, Fischer SA (1993) Age-cohort changes in the lifetime occurrence of depression and other mental disorders. *J Abnorm Psychol* 102(1):110–120
- Escribá QR, Maestre MC, Amores LP, Pastor TA, Miralles ME, Escobar RF (2005) Depression prevalence in adolescents. *Actas Esp Psiquiatr* 33:298–302
- Möller HJ, Henkel V (2005) What are the most effective diagnostic and therapeutic strategies for the management of depression in specialist care? Health Evidence Network report. WHO Regional Office for Europe, Copenhagen. <http://www.euro.who.int/Document/E86602.pdf>, accessed 1/5/09
- Babyak M, Blumenthal JA, Herman S et al (2000) Exercise treatment for major depression: maintenance of therapeutic benefit at 10 months. *Psychosom Med* 62(5):633–638
- Blumenthal JA, Babyak MA, Doraiswamy PM et al (2007) Exercise and pharmacotherapy in the treatment of major depressive disorder. *Psychosom Med* 69(7):587–596
- American Psychiatric Association (1994) Diagnostic and statistical manual of mental disorders. 4th edn. American Psychiatric Association, Washington
- World Health Organization (1992) The ICD-10 Classification of mental and behavioural disorders: clinical descriptions and diagnostic guidelines. Royal College of Psychiatrists, London
- Powers RH, Knesner TJ, Croghan TW (2002) Psychotherapy and pharmacotherapy in depression. *J Ment Health Policy Econ* 5(4):153–161
- Klinkman MS (1997) Competing demands in psychosocial care. A model for the identification and treatment of depressive disorders in primary care. *Gen Hosp Psychiatry* 19(2):98–111
- Gilbody S, House A, Sheldon T (2002) Improving the recognition and management of depression in primary care. *Effective Health Care* 7(5):1–12
- Williams JW, Noël PH, Cordes JA (2002) Is this patient clinically depressed? *J Am Med Assoc* 287(9):1160–1170
- Von Korff M, Katon W, Bush T et al (1998) Treatment costs, cost offset, and cost-effectiveness of collaborative management of depression. *Psychosom Med* 60(2):143–149
- Oxman TE, Dietrich AJ, Schulberg HC (2003) The depression care manager and mental health specialist as collaborators within primary care. *Am J Geriatr Psychiatry* 11(5):507–516
- Fichna J, Janecka A, Costentin J, Do Rego JC (2007) The endomorphin system and its evolving neurophysiological role. *Pharmacol Rev* 59(1):88–123
- Cox RH (2002) Sport psychology: concepts and applications. McGraw-Hill Publishers, St. Louis, MO
- Navines R, Martin-Santos R, Gomez-Gil E, Martinez de Osaba MJ, Gasto C (2008) Interaction between serotonin 5-HT_{1A} receptors and beta-endorphins modulates antidepressant response. *Prog Neuropsychopharmacol Biol Psychiatry* 32(8):1804–1809
- Stein DJ, van Honk J, Ipsier J, Solms M, Panksepp J (2007) Opioids: from physical pain to the pain of social isolation. *CNS Spectr* 12 (9):669–670, 672–664
- Castilla-Cortazar I, Castilla A, Gurpegui M (1998) Opioid peptides and immunodysfunction in patients with major depression and anxiety disorders. *J Physiol Biochem* 54(4):203–215

21. Djurovic D, Milic-Askraic J, Majkic-Singh N (1998) Effect of fluvoxamine on the level of beta-endorphin in the sera and nervous tissue of rats. *Pharmazie* 53(2):143–144
22. Mead GE, Morley W, Campbell P, Greig CA, McMurdo M, Lawlor DA (2009) Exercise for depression. *Cochrane Database Syst Rev* 3:CD004366
23. Vaccarino AL, Olson GA, Olson RD, Kastin AJ (1999) Endogenous opiates: 1998. *Peptides* 20(12):1527–1574
24. Grisel JE, Bartels JL, Allen SA, Turgeon VL (2008) Influence of beta-endorphin on anxious behavior in mice: interaction with EtOH. *Psychopharmacology (Berl)* 200(1):105–115
25. Disham KR, O'Connor JP (2009) Lessons in exercise neurobiology: the case of endorphins. *Ment Health Phys Activity* (in press)
26. Boecker H, Sprenger T, Spilker ME et al (2008) The runner's high: opioidergic mechanisms in the human brain. *Cereb Cortex* 18(11):2523–2531
27. Wester HJ, Willoch F, Tolle TR et al (2000) 6-*O*-(2-[18F]Fluoroethyl)-6-*O*-desmethyldiprenorphine ([18F]DPN): synthesis, biologic evaluation, and comparison with [11C]DPN in humans. *J Nucl Med* 41(7):1279–1286
28. Stern RA, Arruba JE, Hooper CR, Wolfner GW, Morey CE (1997) Visual analogue mood scales to measure internal mood state in aphasic patients: description and initial validity evidence. *Aphasiology* 11:59–71
29. Finley JP, Bonet JF, Waxman MB (1979) Autonomic pathways responsible for bradycardia on facial immersion. *J Appl Physiol* 47(6):1218–1222
30. Goldfarb AH, Jamurtas AZ (1997) Beta-endorphin response to exercise. An update. *Sports Med* 24(1):8–16
31. Lindstrom LH, Widerlov E, Gunne LM, Wahlstrom A, Terenius L (1978) Endorphins in human cerebrospinal fluid: clinical correlations to some psychotic states. *Acta Psychiatr Scand* 57:153–164
32. Gross-Isseroff R, Dillon KA, Israeli M, Biegon A (1990) Regionally selective increases in μ -opioid receptor density in the brains of suicide victims. *Brain Res* 530:312–316
33. Scarone S, Gambini O, Calabrese G et al (1990) Asymmetrical distribution of β -endorphin in cerebral hemispheres of suicides: preliminary data. *Psychiatry Res* 32:159–166
34. Darko DF, Risch SC, Gillin JC, Golshan S (1992) Association of β -endorphin with specific clinical symptoms of depression. *Am J Psychiatry* 149:1162–1167
35. Gabilondo AM, Meana JJ, Garcia-Sevilla JA (1995) Increased density of μ -opioid receptors in the postmortem brain of suicide victims. *Brain Res* 682:245–250
36. Kline NS, Li CH, Lehman HL, Lajta A, Laski E, Cooper T (1977) β -Endorphin-induced changes in schizophrenics and depressed patients. *Arch Gen Psychiatry* 34:1111–1113
37. Jamurtas ZA, Fatouros JI (2004) The effects of exercise on the levels of beta-endorphin in blood. *Inq Sport Phys Educ* 2(1):93–102
38. Hung HF, Kao PF, Lin YS et al (2007) Changes of serum beta-endorphin by programmed exercise training are correlated with improvement of clinical symptoms and quality of life in female mitral valve prolapse syndrome. *Cardiology* 108(4):252–257
39. Bender T, Nagy G, Barna I, Tefner I, Kadas E, Geher P (2007) The effect of physical therapy on beta-endorphin levels. *Eur J Appl Physiol* 100(4):371–382
40. Legrand F, Heuze JP (2007) Antidepressant effects associated with different exercise conditions in participants with depression: a pilot study. *J Sport Exerc Psychol* 29(3):348–364
41. Paffenbarger RS Jr, Lee IM, Leung R (1994) Physical activity and personal characteristics associated with depression and suicide in American college men. *Acta Psychiatr Scand Suppl* 377:16–22
42. Goodwin DR (2003) Association between physical activity and mental disorders among adults in the United States. *Prev Med* 36(6):698–703
43. Larun L, Nordheim LV, Ekeland E, Hagen KB, Heian F (2006) Exercise in prevention and treatment of anxiety and depression among children and young people. *Cochrane Database Syst Rev* 3:CD004691
44. Giacobbi RP, Hausenblas AH, Frye N (2005) A naturalistic assessment of the relationship between personality, daily life events, leisure time exercise, and mood. *Psychol Sport Exerc* 6:67–81
45. Hoffman DM, Hoffman RD (2008) Exercisers achieve greater acute exercise-induced mood enhancement than non-exercisers. *Arch Phys Med Rehabil* 89:358–363
46. McNeil JK, LeBlanc EM, Joyner M (1991) The effect of exercise on depressive symptoms in the moderately depressed elderly. *Psychol Aging* 6(3):487–488
47. Harris AH, Cronkite R, Moos R (2006) Physical activity, exercise coping, and depression in a 10-year cohort study of depressed patients. *J Affect Disord* 93(1–3):79–85
48. Lampinen P, Heikkinen RL, Ruoppila I (2000) Changes in intensity of physical exercise as predictors of depressive symptoms among older adults: an eight-year follow-up. *Prev Med* 30(5):371–380
49. Herman S, Blumenthal JA, Babyak M et al (2002) Exercise therapy for depression in middle-aged and older adults: predictors of early dropout and treatment failure. *Health Psychol* 21(6):553–563
50. Hoffman MB, Babyak AM, Sherwood A et al (2009) Effects of aerobic exercise on sexual functioning in depressed adults. *Ment Health Phys Activity* (in press)
51. Carta MG, Hardoy MC, Pulu A et al (2008) Improving physical quality of life with group physical activity in the adjunctive treatment of major depressive disorder. *Clin Pract Epidemiol Ment Health* 4:1
52. Hassmen P, Koivula N, Uutela A (2000) Physical exercise and psychological well-being: a population study in Finland. *Prev Med* 30(1):17–25
53. Beniamini Y, Rubenstein JJ, Zaichkowsky LD, Crim MC (1997) Effects of high-intensity strength training on quality-of-life parameters in cardiac rehabilitation patients. *Am J Cardiol* 80(7):841–846
54. Dunn AL, Trivedi MH, Kampert JB, Clark CG, Chambliss HO (2002) The DOSE study: a clinical trial to examine efficacy and dose response of exercise as treatment for depression. *Control Clin Trials* 23(5):584–603
55. Singh NA, Stavrinou TM, Scarbek Y, Galambos G, Liber C, Fiatarone Singh MA (2005) A randomized controlled trial of high versus low intensity weight training versus general practitioner care for clinical depression in older adults. *J Gerontol A Biol Sci Med Sci* 60(6):768–776
56. National Institute for Health and Clinical Excellence (2007) Management of depression in primary and secondary care. <http://www.nice.org.uk/nicemedia/pdf/CG023fullguideline.pdf>. Accessed October 21 2009
57. Stanton JM, Arroll B (1996) The effect of moderate exercise on mood in mildly hypertensive volunteers: a randomized controlled trial. *J Psychosom Res* 40(6):637–642
58. Krawczynski M, Olszewski H (2000) Psychological well-being associated with a physical activity programme for persons over 60 years old. *Psychol Sport Exerc* 1(1):57–63
59. Smith PJ, Blumenthal JA, Babyak MA, Georgiades A, Hinderliter A, Sherwood A (2007) Effects of exercise and weight loss on depressive symptoms among men and women with hypertension. *J Psychosom Res* 63(5):463–469

-
60. Milani RV, Lavie CJ, Cassidy MM (1996) Effects of cardiac rehabilitation and exercise training programs on depression in patients after major coronary events. *Am Heart J* 132(4):726–732
 61. Fuchs R (2001) Physical activity and health. In: Smelser NJ, Baltes PB (eds) *International encyclopaedia of the social and behavioural sciences*, vol 17. Elsevier, New York, pp 11411–11415
 62. Berkman LF, Blumenthal J, Burg M et al (2003) Effects of treating depression and low perceived social support on clinical events after myocardial infarction: the Enhancing Recovery in Coronary Heart Disease Patients (ENRICHD) Randomized Trial. *JAMA* 289(23):3106–3116. doi:[10.1001/jama.289.23.3106](https://doi.org/10.1001/jama.289.23.3106)
 63. van Melle JP, de Jonge P, Honig A et al (2007) Effects of anti-depressant treatment following myocardial infarction. *Br J Psychiatry* 190:460–466. doi:[10.1192/bjp.bp.106.028647](https://doi.org/10.1192/bjp.bp.106.028647)