

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/271845779>

The phase diagram of solid hydrogen at high pressure: A challenge for first principles calculations

Conference Paper · March 2015

CITATIONS

0

READS

112

2 authors:



[Sam Azadi](#)

Imperial College London

44 PUBLICATIONS 631 CITATIONS

[SEE PROFILE](#)



[Matthew Foulkes](#)

Imperial College London

100 PUBLICATIONS 4,838 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Fast and Accurate Quantum Monte Carlo for Molecular Modeling [View project](#)



High pressure hydrogen [View project](#)

Abstract Submitted
for the MAR15 Meeting of
The American Physical Society

The phase diagram of solid hydrogen at high pressure: A challenge for first principles calculations¹ SAM AZADI, MATTHEW FOULKES , Imperial College London — We present comprehensive results for the high-pressure phase diagram of solid hydrogen. We focus on the energetically most favorable molecular and atomic crystal structures. To obtain the ground-state static enthalpy and phase diagram, we use semi-local and hybrid density functional theory (DFT) as well as diffusion quantum Monte Carlo (DMC) methods. The closure of the band gap with increasing pressure is investigated utilizing quasi-particle many-body calculations within the GW approximation. The dynamical phase diagram is calculated by adding proton zero-point energies (ZPE) to static enthalpies. Density functional perturbation theory is employed to calculate the proton ZPE and the infra-red and Raman spectra. Our results clearly demonstrate the failure of DFT-based methods to provide an accurate static phase diagram, especially when comparing insulating and metallic phases. Our dynamical phase diagram obtained using fully many-body DMC calculations shows that the molecular-to-atomic phase transition happens at the experimentally accessible pressure of 374 GPa. We claim that going beyond mean-field schemes to obtain derivatives of the total energy and optimize crystal structures at the many-body level is crucial.

¹This work was supported by the UK engineering and physics science research council under grant EP/I030190/1, and made use of computing facilities provided by HECTOR, and by the Imperial College London high performance computing centre.

Sam Azadi
Imperial College London

Date submitted: 13 Oct 2014

Electronic form version 1.4