

S_1 -PARACOMPACT SPACES

KHALID AL-ZOUBI, AMANI RAWASHDEH

ABSTRACT. In this paper we introduce a new class of spaces which will be called the class of S_1 -paracompact spaces. We characterize S_1 -paracompact spaces and study their basic properties. The relationships between S_1 -paracompact spaces and other well-known spaces are investigated.

2000 *Mathematics Subject Classification*: 54D20, 54C10, 54A10, 54G05.

1. INTRODUCTION AND PRELIMINARIES

In 1963, Levine [8] introduced and studied the concept of semi-open sets in topological spaces. In [1], Al-Zoubi used semi-open sets to define the class of s -expandable spaces. A space (X, T) is said to be s -expandable space if for every s -locally finite collection $\mathcal{F} = \{F_\alpha : \alpha \in I\}$ of subsets of X there exists a locally finite collection $\mathcal{G} = \{G_\alpha : \alpha \in I\}$ of open subsets of X such that $F_\alpha \subseteq G_\alpha$ for each $\alpha \in I$. In [1], Theorem 3.4, a space (X, T) is s -expandable if every semi-open cover of X has a locally finite open refinement.

In section 2 of this work we introduce and study a new class of spaces, namely S_1 -paracompact spaces, and we provide several characterizations of S_1 -paracompact spaces and investigate the relationship between S_1 -paracompact spaces and other well-known spaces such as paracompact spaces, s -expandable spaces, nearly paracompact spaces and semi-compact spaces. Finally, in section 3, we deal with some basic properties of S_1 -paracompact spaces, i.e. subspaces, sum, inverse image and product. Throughout this work a space will always mean a topological space on which no separation axioms are assumed unless explicitly stated. Let (X, T) be a space and A be a subset of X . The closure of A , the interior of A and the relative topology on A will be denoted by $cl(A)$, $int(A)$ and T_A respectively. A is called semi-open subset of (X, T) ([8]) if there exists an open set U of X such that $U \subseteq A \subseteq cl(U)$. The complement of a semi-open set is called a semi-closed set ([2]). The semiclosure of A ([2]), denoted by $scl(A)$, is the smallest semi-closed set that contains A .

A is called regular open if $A = \text{int}(\text{cl}(A))$. The family of all semi-open (resp. regular open) subsets of (X, T) is denoted by $SO(X, T)$ (resp. $RO(X, T)$).

Definition 1.1. A collection $\mathcal{F} = \{F_\alpha : \alpha \in I\}$ of subsets of a space (X, T) is said to be locally finite (resp. s -locally finite [1]), if for each $x \in X$, there exists $U \in T$ (resp. $U \in SO(X, T)$) containing x and U intersects at most finitely many members of \mathcal{F} .

Definition 1.2. A space (X, T) is said to be:

(a) semi-compact [3] if every semi-open cover of X has a finite subcover.
 (b) semi-regular [4] if for each semi-closed set F and each point $x \notin F$, there exist disjoint semi-open sets U and V such that $x \in U$ and $F \subseteq V$. This is equivalent to, for each $U \in SO(X, T)$ and for each $x \in U$, there exists $V \in SO(X, T)$ such that $x \in V \subseteq \text{scl}(V) \subseteq U$.

(c) extremally disconnected (briefly e.d.) if the closure of every open set in (X, T) is open.

Lemma 1.3 [10]. If (X, T) is e.d., then $\text{scl}(U) = \text{cl}(U)$ for each $U \in SO(X, T)$.

Proposition 1.4. Let (X, T) be an e.d. semi-regular space. Then:

- (a) $SO(X, T) = T$.
- (b) (X, T) is regular.

Proof. (a) Let $U \in SO(X, T)$ and $x \in U$. Since (X, T) is semi-regular, there exists $V \in SO(X, T)$ such that $x \in V \subseteq \text{scl}(V) \subseteq U$. Now, choose $W \in T$ such that $W \subseteq V \subseteq \text{cl}(W)$. But (X, T) is e.d., therefore, by Lemma 1.3, $\text{cl}(W) = \text{cl}(V) = \text{scl}(V)$ is an open set containing x such that $\text{cl}(W) \subseteq U$. Thus $U \in T$.

(c) Follows from part (a) and Lemma 1.3.

Lemma 1.5. ([2]). If A is an open set in (X, T) and $B \in SO(X, T)$ then $A \cap B \in SO(X, T)$.

2. S_1 -PARACOMPACT SPACES

Definition 2.1. A space (X, T) is said to be S_1 -paracompact space if every semi-open cover of X has a locally finite open refinement.

Every S_1 -paracompact space is s -expandable (Theorem 3.4 of [1]) but the converse is not true as may be seen from the following example.

Example 2.2. Let $X = \{1, 2, 3\}$ and $T = \{\emptyset, X, \{1\}, \{2\}, \{1, 2\}\}$. Then (X, T) is s -expandable (every finite space is s -expandable) but not S_1 -paracompact since $\{\{1, 2\}, \{2, 3\}\}$ is a semi-open cover of X which admits no locally finite open refinement. For any space (X, T) , we have $RO(X, T) \subseteq T \subseteq SO(X, T)$. Therefore the following implications are obvious. S_1 -paracompact \Rightarrow paracompact \Rightarrow nearly paracompact. Where a space (X, T) is said to be nearly paracompact if every regular open cover of X has a locally finite open refinement. Moreover we have the following theorem.

Theorem 2.3. *Let (X, T) be an e.d. semi-regular space. Then the following are equivalent:*

- (a) (X, T) is nearly paracompact.
- (b) (X, T) is paracompact.
- (c) (X, T) is S_1 -paracompact.

Proof. (a) \rightarrow (b): Follows from Proposition 1.4 and the fact that if a space (X, T) is regular then $T \subseteq RO(X, T)$.

(b) \rightarrow (c) : Follows from Proposition 1.4 (a).

The following examples show that the conditions “e.d. and semi-regular” on the space (X, T) in the above Theorem are essential.

Example 2.4. (a) Let $X = R$ be the set of the real numbers with the topology $T = \{\emptyset, X, \{1\}\}$. Then (X, T) is an e.d. paracompact space but not S_1 -paracompact since $\{\{1, x\} : x \in X\}$ is a semi-open cover of X which admits no locally finite open refinement.

(b) Consider the space (X, T) where $X = R$ and

$$T = \{U \subseteq R : 0 \notin U\} \cup \{U \subseteq R : 0 \in U \text{ and } R - U \text{ is finite}\}.$$

Then (X, T) is semi-regular and paracompact but not S_1 -paracompact since it is not s-expandable (Example 3.3,[1]).

(c) Let $X = R$ with the topology $T = \{U \subseteq R : 0 \in U\} \cup \{\emptyset\}$. Then (X, T) is an e.d. nearly paracompact space ($RO(X, T) = \{X, \emptyset\}$) but not paracompact since $\{\{0, x\} : x \in X\}$ is an open cover of X which admits no locally finite open refinement.

Theorem 2.5. *If (X, T) is an S_1 -paracompact T_1 -space, then $T = SO(X, T) = T^\alpha$.*

Proof. Let U be a semi-open in (X, T) . For each $y \notin U$, we choose an open set V_y containing y and $x \notin V_y$. Therefore the collection $\mathcal{V} = \{V_y : y \notin U\} \cup \{U\}$ is a semi-open cover of (X, T) and so it has a locally finite open refinement \mathcal{W} . Put $V = \cup\{W \in \mathcal{W} : x \in W\}$. Then V is an open set containing x and $V \subseteq U$. Thus U is open in (X, T) .

Corollary 2.6. *Let (X, T) be an S_1 -paracompact space.*

(a) *If (X, T) is T_1 then it is e.d.*

(b) *If (X, T) is T_2 then it is semi-regular.*

Proof. (a) Let $U \in T$. Then $cl(U) \in SO(X, T)$ and so $cl(U) \in T$ by Theorem 2.5.

(b) (X, T) is a paracompact T_2 -space. Therefore by Lemma 5.1.4 of ([6]), (X, T) is regular. Now let $U \in SO(X, T)$ and $x \in U$. By Theorem 2.5, $U \in T$ and so there exists an open set $V \in T$ such that $x \in V \subset cl(V) \subset U$. Thus $x \in V \subset scl(V) \subset$

$cl(V) \subset U$. It follows that (X, T) is semi-regular. Note that \mathbb{R} with the cofinite topology is S_1 –paracompact but not semi-regular. Recall that a space (X, T) is called semi-compact ([3]), if every semi-open cover of X has a finite subcover. We note that S_1 –paracompactness and semi-compactness are independent, since in Example 2.2, (X, T) is semi-compact (X is finite) but it is not S_1 –paracompact. On

the other hand, the space (X, T_{dis}) where X is an infinite set is S_1 –paracompact but it is not semi-compact.

Theorem 2.7. *Let (X, T) be a T_2 -space. Then the following are equivalent:*

- (a) (X, T) is semi-compact.
- (b) (X, T) is S_1 –paracompact and compact.

Proof. (a)→(b): Suppose that (X, T) is semi-compact. Then by Theorem 2.4 of ([3]), X is finite and so T is the discrete topology. Therefore (X, T) is S_1 –paracompact.

(b) →(a) : As (X, T) is compact T_2 -space, every locally finite family of open sets is finite.

For a space (X, T) , we denote by T_Ψ ([3]) the topology on X which has $SO(X, T)$ as a subbase.

It is clear that if a space (X, T) is e.d., then $T_\Psi = SO(X, T)$.

Note that in Example 2.4 part (a), the space (X, T) is paracompact and e.d while (X, T_Ψ) is not paracompact since $T_\Psi = \{U : 1 \in U\} \cup \{\phi\}$.

Proposition 2.8. *Let (X, T) be an e.d. space.*

- (a) *If (X, T) is S_1 –paracompact then (X, T_Ψ) is S_1 –paracompact.*
- (b) *(X, T_Ψ) is S_1 –paracompact if and only if (X, T) is paracompact.*

Proof. (a) Follows from the fact that for any space (X, T) , $T \subseteq T_\Psi$ and $SO(X, T_\Psi) \subseteq SO(X, T)$.

(b) Since (X, T) is e.d. then by Lemma 3.7 of ([1]), $SO(X, T_\Psi) = SO(X, T)$ and thus $T_\Psi = SO(X, T_\Psi)$,

Example 2.9. The converse of part (a) of Proposition 2.8 is not true in general. To see that, consider $X = \{1, 2, 3\}$ with $T = \{\phi, X, \{1\}\}$, then it is easy to see that (X, T) is e.d. and (X, T_Ψ) is S_1 –paracompact but (X, T) is not.

Corollary 2.10. *Let (X, T) be an e.d. semi-regular space. Then (X, T) is S_1 –paracompact if and only if (X, T_Ψ) is paracompact.*

Proof. Necessity follows from Proposition 2.8. For sufficiency, since (X, T) e.d. then $SO(X, T) = SO(X, T_\Psi)$ and so by Proposition 1.4 part (a), $T_\Psi = SO(X, T) = SO(X, T_\Psi) = T$. Recall that a subset A of a space (X, T) is said to be α –set if $A \subset int(cl(int(A)))$. The family of all α –sets of a space (X, T) , denote by T^α , forms a topology on X , finer than T .

Lemma 2.11. (a) *For any space (X, T) , $SO(X, T^\alpha) = SO(X, T)$ ([9]).*

(b) For a space (X, T) , if (X, T^α) is normal, then $T = T^\alpha$ ([5]).

Theorem 2.12. Let (X, T) be a T_2 -space. Then (X, T) is S_1 -paracompact space if and only if (X, T^α) is S_1 -paracompact space.

Proof. Necessity, follows from Lemma 2.11 and the fact that $T \subset T^\alpha$. For sufficiency, suppose that (X, T^α) is S_1 -paracompact. Then (X, T^α) is a paracompact T_2 -space and so it is normal ([6]). Therefore, by Lemma 2.11, $T = T^\alpha$. On the other hand $SO(X, T^\alpha) = SO(X, T)$ (Lemma 2.11) and so (X, T) is S_1 -paracompact. Note that, in Example 2.9, the space (X, T) is not S_1 -paracompact. However, (X, T^α) is S_1 -paracompact. Therefore the condition (X, T) is T_2 in Theorem 2.12, can not be dropped.

Theorem 2.13. If each semi-open cover of a space (X, T) has an open σ -locally finite refinement, then each semi-open cover of X has a locally finite refinement.

Proof. Let \mathcal{U} be a semi-open cover of X . Let $\mathcal{V} = \cup_{n \in N} \mathcal{V}_n$ be an open σ -locally finite refinement of \mathcal{U} , where \mathcal{V}_n is locally finite. For each $n \in N$ and each $V \in \mathcal{V}_n$, let $V'_n = V - \cup_{k < n} \mathcal{V}_k^*$ where $\mathcal{V}_k^* = \cup \{V : V \in \mathcal{V}_k\}$ and put $\mathcal{V}'_n = \{V'_n : V \in \mathcal{V}_n\}$. Now, put $\mathcal{W} = \{V'_n : n \in N, V \in \mathcal{V}_n\} = \cup \{\mathcal{V}'_n : n \in N\}$. We show \mathcal{W} is a locally finite refinement of \mathcal{U} . Let $x \in X$ and let n be the first positive integer such that $x \in \mathcal{V}_n^*$. Therefore $x \in V'$ for some $V' \in \mathcal{V}'_n$. Thus \mathcal{W} is a cover of X . To show \mathcal{W} is locally finite, let $x \in X$ and n be the first positive integer such that $x \in \mathcal{V}_n^*$. Then $x \in V$ for some $V \in \mathcal{V}_n$. Now, $V \cap V' = \emptyset$ for each $V' \in \mathcal{V}'_k$ and for each $k > n$. Therefore, V can intersect at most the elements of \mathcal{V}'_k for $k \leq n$. Since \mathcal{V}'_k is locally finite for each $k \leq n$, so we choose an open set $O_{x(k)}$ containing x such that $O_{x(k)}$ meets at most finitely many members of \mathcal{V}'_k . Finally, put $O_x = V \cap (\cap_{k=1}^n O_{x(k)})$. Then O_x is an open set containing x such that O_x meets at most finitely many members of \mathcal{W} .

Theorem 2.14. Let (X, T) be a semi-regular space. If each semi-open cover of a space X has a locally finite refinement, then each semi-open cover of X has a locally finite semi-closed refinement.

Proof. Let \mathcal{U} be a semi-open cover of X . For each $x \in X$, pick $U_x \in \mathcal{U}$ such that $x \in U_x$. Since (X, T) is semi-regular, then there exists $V_x \in SO(X, T)$ such that $x \in V_x \subset scl(V_x) \subset U_x$. The family $\mathcal{V} = \{V_x : x \in X\}$ is a semi-open cover of X and so, by assumption, has a locally finite refinement $\mathcal{W} = \{W_\alpha : \alpha \in I\}$. The collection $scl(\mathcal{W}) = \{scl(W_\alpha) : \alpha \in I\}$ is locally finite such that for each $\alpha \in I$, if $W_\alpha \subset V_x$, then $scl(W_\alpha) \subset U$ for some $U \in \mathcal{U}$. Thus $scl(\mathcal{W})$ is a semi-closed locally finite refinement of \mathcal{U} . For the next theorem we will use the statements:

- (a) (X, T) S_1 -paracompact.
- (b) Each semi-open cover of X has a σ -locally finite open refinement.
- (c) Each semi-open cover of X has a locally finite refinement.
- (d) Each semi-open cover of X has a locally finite semi-closed refinement.

Theorem 2.15. If (X, T) is a semi-regular space then (a) \rightarrow (b) \rightarrow (c) \rightarrow (d).

Proof. (a) \rightarrow (b) is obvious.

(b) \rightarrow (c) Follows from Theorem 2.13.

(c) \rightarrow (d) Follows from Theorem 2.14.

In Example 2.2, (X, T) is a semi-regular in which $\{1\}$, $\{2\}$ and $\{3\}$ are semi-closed sets. Therefore, (X, T) satisfies statement (d) of Theorem 2.15, but it is not S_1 -paracompact.

Note that if (X, T) is also e.d. in Theorem 2.15, then (d) \rightarrow (a).

3-PROPERTIES OF S_1 -PARACOMPACT SPACES

In this section we study some basic properties of S_1 -paracompact spaces such as subspaces, sums, products, images and inverse images under some types of functions.

Theorem 3.1. *Every regular open subspace of an S_1 -paracompact space is S_1 -paracompact.*

Proof. Let (X, T) be an S_1 -paracompact space and A be a regular open subspace of (X, T) . Let $\mathcal{U} = \{U_\alpha : \alpha \in I\}$ be a semi-open cover of A such that $U_\alpha \in SO(A, T_A)$ for each $\alpha \in I$. Since A is an open subset of X then $U_\alpha \in SO(X, T)$ for each $\alpha \in I$. Therefore the family $\mathcal{V} = \{U_\alpha : \alpha \in I\} \cup \{X - A\}$ is a semi-open cover of X . Let $\mathcal{W} = \{W_\beta : \beta \in B\}$ be a locally finite open refinement of \mathcal{U} in (X, T) . Then the family $\{W_\beta \cap A : \beta \in B\}$ is a locally finite open refinement of A in (A, T_A) . Thus (A, T_A) is S_1 -paracompact.

Corollary 3.2. *Every clopen subspace of an S_1 -paracompact space is S_1 -paracompact.*

Definition 3.3. [6] *Let $\{(X_\alpha, T_\alpha) : \alpha \in I\}$ be a collection of topological spaces such that $X_\alpha \cap X_\beta = \emptyset$ for each $\alpha \neq \beta$. Let $X = \cup_{\alpha \in I} X_\alpha$ be topologized by $T = \{G \subseteq X : G \cap X_\alpha \in T_\alpha, \alpha \in I\}$. Then (X, T) is called the sum of the spaces $\{(X_\alpha, T_\alpha) : \alpha \in I\}$ and we write $X = \oplus_{\alpha \in I} X_\alpha$.*

Theorem 3.4. *The topological sum $\oplus_{\alpha \in I} X_\alpha$ is S_1 -paracompact if and only if the space (X_α, T_α) is S_1 -paracompact for each $\alpha \in I$.*

Proof. Necessity follows from Corollary 3.2, since (X_α, T_α) is a clopen subspace of the space $\oplus_{\alpha \in I} X_\alpha$, for each $\alpha \in I$. To prove sufficiency, let \mathcal{U} be a semi-open cover of $\oplus_{\alpha \in I} X_\alpha$. For each $\alpha \in I$ the family $\mathcal{U}_\alpha = \{U \cap X_\alpha : U \in \mathcal{U}\}$ is a semi-open cover of the S_1 -paracompact space (X_α, T_α) . Therefore \mathcal{U}_α has a locally finite open refinement \mathcal{V}_α in (X_α, T_α) . Put $\mathcal{V} = \cup_{\alpha \in I} \mathcal{V}_\alpha$. It is clear that \mathcal{V} is a locally finite open refinement of \mathcal{U} . Thus $\oplus_{\alpha \in I} X_\alpha$ is S_1 -paracompact.

Recall that a function $f : (X, T) \rightarrow (Y, M)$ is said to be irresolute ([2]), if $f^{-1}(U) \in SO(X, T)$ for every $U \in SO(Y, M)$. It is well known that every continuous open surjective function is irresolute (Theorem 1.8 of [2]).

Theorem 3.5. *Let $f : (X, T) \rightarrow (Y, M)$ be a continuous, open, and closed surjective*

function such $f^{-1}(y)$ is compact for each $y \in Y$. If (X, T) is S_1 -paracompact, then so is (Y, M) .

Proof. Let $\mathcal{V} = \{V_\alpha : \alpha \in I\}$ be a semi-open cover of (Y, M) . Since f is irresolute, the collection $\mathcal{U} = f^{-1}(\mathcal{V}) = \{f^{-1}(V_\alpha) : \alpha \in I\}$ is a semi-open cover of the S_1 -paracompact (X, T) space and so it has a locally finite open refinement, say \mathcal{W} . The collection $f(\mathcal{W})$ is a locally finite open refinement of \mathcal{V} in (Y, M) .

Definition 3.6. A function $f : (X, T) \rightarrow (Y, M)$ is said to be semi-closed ([10]) if $f(A) \in SC(Y, M)$ for every closed subset A of X .

Proposition 3.7. [1] A function $f : (X, T) \rightarrow (Y, M)$ is semi-closed if and only if for every $y \in Y$ and every open set U in (X, T) which contains $f^{-1}(y)$, there exists $V \in SO(Y, M)$ such that $y \in V$ and $f^{-1}(V) \subseteq U$.

Theorem 3.8. Let $f : (X, T) \rightarrow (Y, M)$ be a continuous semi-closed surjection and $f^{-1}(y)$ is compact for each $y \in Y$. If (Y, M) is S_1 -paracompact space then (X, T) is paracompact.

Proof. Let $\mathcal{U} = \{U_\alpha, \alpha \in I\}$ be an open cover of X . For each $y \in Y$ and for each $x \in f^{-1}(y)$ choose $\alpha(x) \in I$ such that $x \in U_{\alpha(x)}$. Therefore the collection $\{U_{\alpha(x)} : x \in f^{-1}(y)\}$ is an open cover of $f^{-1}(y)$ and so there exists a finite subset $I(y)$ of I such that $f^{-1}(y) \subset \cup_{\alpha(x) \in I(y)} U_{\alpha(x)} = U_y$. But f is semi-closed so there exists a semi-open set V_y containing y and $f^{-1}(V_y) \subset U_y$. Thus $\mathcal{V} = \{V_y : y \in Y\}$ is a semi-open cover of Y and so it has a locally finite open refinement say $\mathcal{W} = \{W_\beta : \beta \in B\}$. Since f is continuous, then the family $\{f^{-1}(W_\beta) : \beta \in B\}$ is an open locally finite cover of X such that for each $\beta \in B$, $f^{-1}(W_\beta) \subset U_y$ for some $y \in Y$. Now, the family $\{f^{-1}(W_\beta) \cap U_y : \beta \in B, y \in \mathcal{V}\}$ is a locally finite open refinement of \mathcal{U} , where $f^{-1}(W_\beta) \cap U_y = \{f^{-1}(W_\beta) \cap U_y : \beta \in B, \alpha(x) \in I(y)\}$. Therefore (X, T) is paracompact. We finally study products of S_1 -paracompact spaces. Note that the space (X, T) where $X = \{1, 2\}$ and $T = \{\phi, X, \{1\}\}$ is an e.d. S_1 -paracompact (compact) space while $(X, T) \times (X, T)$ is not S_1 -paracompact since $\{(1, 1), (2, 2)\}, \{(1, 1), (1, 2), (2, 1)\}$ is a semi-open cover of $X \times X$ which admits no locally finite open refinement.

Corollary 3.9. If (X, T) is compact, (Y, M) is S_1 -paracompact and

$(X, T) \times (Y, M)$ is e.d. semi-reguler, then $(X, T) \times (Y, M)$ is S_1 -paracompact.

Proof. Since (Y, M) is paracompact then $(X, T) \times (Y, M)$ is paracompact (see[6]). Therefore $(X, T) \times (Y, M)$ is S_1 - paracompact by Theorem 2.3. In the above paragraph $(X, T) \times (X, T)$ is e.d. but not semi-reguler. Therefore the condition " semi-reguler " on $(X, T) \times (Y, M)$ in the above Corollary can not be dropped. On the other hand $(R, T) \times (R, T_{dis})$ (where (R, T) as in Example 2.4, part (b)) is semi-reguler but not S_1 -paracompact since it is not s-expandable (Example 3.14,[1]).

REFERENCES

- [1] K. Y. Al-Zoubi, *S-expandable spaces*, Acta Math. Hungar., 102 (2004), 203-212
- [2] S. G. Crossley and S. K. Hilderbrand, *Semi-topological properties*, Fund. Math., 74(1972), 233-254.
- [3] C. Dorsett, *Semi-compact R_1 and product spaces*, Bull. Malaysian Math., Soc. (2) 3 (1980), 15-19.
- [4] C. Dorsett, *Semi-regular spaces*, Soochow J. Math., 8(1982), 45-53.
- [5] J. Dontchev, *An answer to a question of Mrsevic and Reilly*, Q and A in Gen. Topology 12 (1994), no. 2, 205-207.
- [6] R. Engelking, *General Topology*, Helderman Verlag, (1989).
- [7] D. S. Jankovc, *A note on mappings of extremally disconnected spaces*, Acta Math. Hung. 46(1985), 83-92.
- [8] N. Levine, *Semi-open sets and semi-continuity in topological spaces*, Amer. Math. Monthly, 70(1963), 36-41.
- [9] O. Njastad, *On some classes of nearly open sets*, Pacific J. Math., 15(1965), 961-970.
- [10] T. Noiri, *Properties of S-closed spaces*, Acta Math. Hungar., 35 (1980), 431-436.
- [11] D. Sivaraj, *Semihomomorphism*, Acta Math. Hungar., 48(1986), 139-145.
- [12] M.K. Singal and S. P. Arya, *On nearly paracompact spaces*, Mat., Vesnik., 6 (21)(1969), 3-16.

Khalid Al-Zoubi
Department of Mathematics
Faculty of science, Yarmouk University
Irbid-Jordan
E-mail: *Khalidz@yu.edu.jo*.

Amani Rawashdeh
Department of Mathematics,
Faculty of science, Yarmouk University
Irbid-Jordan
E-mail: *a_rawashdeh85@yahoo.com*.