

Primary care

Randomised controlled trial of pelvic floor muscle training during pregnancy

Kjell Å Salvesen, Siv Mørkved

Department of
Obstetrics and
Gynaecology,
Trondheim
University Hospital
St. Olav, N-7006
Trondheim, Norway
Kjell Salvesen
*professor in obstetrics
and gynaecology*

Department of
Community
Medicine and
General Practice,
Norwegian
University of
Science and
Technology, N-7489
Trondheim, Norway
Siv Mørkved
physiotherapist

Correspondence to:
K Å Salvesen
pepes@medisin.
ntnu.no

BMJ 2004;329:378-80

Abstract

Objectives To examine a possible effect on labour of training the muscles of the pelvic floor during pregnancy.

Design Randomised controlled trial.

Setting Trondheim University Hospital and three outpatient physiotherapy clinics in a primary care setting.

Participants 301 healthy nulliparous women randomly allocated to a training group (148) or a control group (153).

Intervention A structured training programme with exercises for the pelvic floor muscles between the 20th and 36th week of pregnancy.

Main outcome measures Duration of the second stage of labour and number of deliveries lasting longer than 60 minutes of active pushing among women with spontaneous start of labour after 37 weeks of pregnancy with a singleton fetus in cephalic position.

Results Women randomised to pelvic floor muscle training had a lower rate of prolonged second stage labour (24%, 95% confidence interval 16% to 33%; 22 out of 105 women were at risk (undelivered) at 60 minutes in the survival analysis) than women allocated to no training (38% (37/109), 28% to 47%). The duration of the second stage was not significantly shorter (40 minutes *v* 45 minutes, *P* = 0.06).

Conclusions A structured training programme for the pelvic floor muscles is associated with fewer cases of active pushing in the second stage of labour lasting longer than 60 minutes.

Introduction

Training the pelvic floor muscles during pregnancy can prevent urinary incontinence,^{1,2} and pregnant women are encouraged to do exercises for these muscles. A myth prevails among birth attendants that strong pelvic floor muscles (for example, as a result of horse riding) may obstruct labour.³ However, training of the pelvic floor muscles may produce strong and well controlled muscles that will facilitate labour.⁴ A prospective study with 86 women found no effect of such training on the course of delivery,⁵ but possible effects of pelvic floor muscle training on labour have been sparsely scientifically documented.

The primary aim of this trial was to assess if training the muscles of the pelvic floor during pregnancy could prevent urinary incontinence. Women in the study group had stronger pelvic floor muscles and reported less urinary incontinence after the training period.² This report deals with secondary outcomes of the trial. We wanted to study any effect of pelvic floor muscle training on labour.

Methods

The population and methods are described elsewhere.² One group of women (*n* = 148) trained with a physiotherapist for 60 minutes once per week for a period of 12 weeks between the 20th and 36th week of pregnancy. In addition, the women were encouraged to perform eight to 12 intensive contractions of the pelvic floor muscle at home twice a day. Adherence to the training programme was 81% (120 women).² Women in the control group (*n* = 153) were not discouraged from doing pelvic floor muscle exercises on their own. Figure 1 shows the flow of participants through the trial.

We reviewed hospital records two to three years after delivery and recorded mode of delivery, epidural analgesia or oxytocin augmentation during labour, episiotomy, perineal tears, and neonatal outcomes. The reviewer (KÅS) was not involved in training the women and was blinded to group allocation while recording and plotting the data. The midwife in charge of labour judged the need for oxytocin augmentation or episiotomy. The obstetrician on call decided the need for operative delivery. Birth attendants were unaware of women's group status.

We recorded the lengths of the first and second stages of labour from partograms. We defined the onset of labour as the beginning of the active phase of the first stage of labour,⁶ or from the time of admission if the cervix was dilated more than 3 cm on arrival. Our definition of the second stage of labour was "active pushing time." We recorded the lengths of the first and second stages of labour in minutes. Most clinics have rules that limit the duration of the second stage.⁶ In Norway, the recommended "second stage rule" is one hour, with active pushing after complete dilatation of the cervix.⁷ In this study we defined prolonged second stage as active pushing for longer than 60 minutes.

This article was posted on *bmj.com* on 14 July 2004: <http://bmj.com/cgi/doi/10.1136/bmj.38163.724306.3A>

We undertook our analysis by intention to treat. It was restricted to 111 women in the training group and 113 women in the control group (fig 1). They had spontaneous start of labour after 37 weeks of pregnancy with a singleton fetus in cephalic position.

We used a Kaplan-Meier survival analysis to test for differences between groups in proportions of women with prolonged second stage and the duration of labour. We censored operative deliveries and deliveries with a prolonged second stage. We also performed a Cox regression analysis with possible confounding variables and appropriate statistical tests for categorical and normally distributed variables. We considered P values <0.05 significant.

Results

Women randomised to pelvic floor muscle training had a lower rate of prolonged second stage labour than women allocated to no training (table). The number needed to treat (NNT) to benefit was 8. In a survival analysis the difference between groups was 24% (95% confidence interval 16% to 33%) for the training group versus 38% (28% to 47%) for the control group.

The duration of the second stage of labour was not statistically different between groups (40 min *v* 45 min, P=0.06). Figure 2 shows the time to delivery from the start of active pushing up to 60 minutes. The infants in the training group were slightly younger and smaller (table). A Cox regression analysis with gestational length, birth weight, and head circumference as possible confounders did not change the estimates materially.

Fewer women had breech presentations (fig 1; 1 *v* 9, P=0.01). The rates of operative delivery for prolonged second stage did not differ between the two groups (table 1). Fewer women had episiotomies (51% *v* 64%, odds ratio 0.59, 0.35 to 1.00; NNT 7), but we found no other significant differences in outcomes related to labour. Apgar scores and umbilical artery pH did not differ between groups (data not shown).

Discussion

Pelvic floor muscle training during pregnancy results in improved muscle control and strong flexible




Fig 1 Flow of participants through the trial

muscles. The effect may be on the central nervous system and the muscles, and training seems to facilitate rather than obstruct labour.

Possible limitation of the study

Since this report deals with secondary outcomes and the differences reached borderline significance, the results should be viewed with caution. However, this was a randomised controlled trial with blinding technique, few withdrawals, and high adherence to the training protocol.

The second stage of labour begins when cervical dilatation is complete and ends with fetal expulsion.⁶ We defined the second stage as “active pushing time.” This definition of the second stage is suitable for clinical

Mean outcome variables (with standard deviations) among 224 women with a singleton fetus in cephalic position with spontaneous start of labour after 37 weeks of pregnancy. Values are numbers (percentages) of women unless otherwise indicated

	Study group (n=111)	Control group (n=113)	P value
Median duration of first stage of labour in minutes (95% CI)	260 (195 to 325)	259 (215 to 303)	0.44*
Acute caesarean section in the first stage	5 (5)	3 (3)	0.46†
Duration of second stage of labour in minutes (95% CI)	40 (33 to 47)	45 (38 to 52)	0.06*
Deliveries with prolonged second stage (≥60 minutes)	22 (21)	37 (34)	0.03†
Vaginal operative delivery in the second stage	15 (15)	19 (17)	0.57†
Episiotomy	56 (51)	72 (64)	0.05†
Third or fourth degree tears	7 (6)	9 (8)	0.64†
Epidural analgesia	47 (42)	49 (43)	0.88†
Oxytocin augmentation	69 (63)	74 (66)	0.67†
Characteristics of infants at birth:			
Gestational age in days	280 (8.6)	283 (7.1)	0.04‡
Weight (g)	3514 (391)	3622 (431)	0.05‡
Length (cm)	50.4 (1.9)	50.8 (1.7)	0.11‡
Head circumference (cm)	35.3 (1.4)	35.9 (1.6)	0.002‡

*Log rank test.

† χ^2 test or Fisher's exact test.

‡t test.




Fig 2 Survival plot of the duration of the second stage of labour for women in the training group (n=105) and the control group (n=109). Discrepancies in numbers are due to some missing data in each group (six in the training group and four in the control group). Operative deliveries for fetal distress (n=9) and slow progress (n=3) during the first hour, and all deliveries lasting longer than 60 minutes (n=59) were censored. Log rank test, P=0.06 for comparison of the two survival plots in the Kaplan-Meier analysis

research, since the start of active pushing is easily identified from partograms, and it is related to a second stage rule.⁷ The risk for bias should be small since the reviewer of the partograms was blinded to group status.

Chance finding

The difference in breech presentations should be interpreted as a possible chance finding. The women trained in different positions, but there is insufficient evidence from well controlled trials to support the use of postural management of breech presentations.⁸

Role of body mass index or exercise

We found no differences in body mass index or self reported regular physical exercise after the training period (data not shown). This argues for an effect of increased strength and better control of pelvic floor muscles rather than a general effect of physical training during pregnancy. New trials from other populations are needed.

Contributors: KÅS and SM were involved in designing and conducting the study, analysing the data, and writing the report. Kari Bø contributed to the design of the study. The physiothera-

What is already known about this topic

Exercising the pelvic floor muscles prevents urinary incontinence in about one in six women during pregnancy and in one in eight women after pregnancy

It increases the strength of the pelvic floor muscles

What this study adds

Intensive training of the pelvic floor muscles during pregnancy seems to facilitate rather than to obstruct labour.

It could prevent a prolonged second stage in one in eight women

pists Hildegunn Børsting, Trude Hoff Leirvik, Bente Olsen, Monica U Tøndel, and Bjørg Vada led the group training sessions. Pål Romundstad gave statistical advice. KÅS is the guarantor.

Funding: Norwegian Fund for Postgraduate Training in Physiotherapy and Norwegian Women's Public Health Association.

Competing interests: None declared.

Ethical approval: Regional medical ethics committee.

- 1 Reilly ETC, Freeman RM, Waterfield MR, Waterfield AE, Steggle P, Pedlar F. Prevention of postpartum stress incontinence in primigravidae with increased bladder neck mobility: a randomised controlled trial of antenatal pelvic floor exercises. *Br J Obstet Gynaecol* 2002;109:68-76.
- 2 Mørkved S, Bø K, Schei B, Salvesen KÅ. Pelvic floor muscle training during pregnancy to prevent urinary incontinence: A single-blind randomized controlled trial. *Obstet Gynecol* 2003;101:313-9.
- 3 UK Midwifery Archives. Hannah, midwife. Does horse-riding affect the pelvic floor? www.radmid.demon.co.uk/pelvicfloor.htm (accessed 25 Jun 2004).
- 4 Baby Centre. How do pelvic floor exercises help me? www.babycentre.co.uk/refcap/536339 (accessed 25 Jun 2004).
- 5 Nielsen CA, Sigsgaard I, Olsen M, Tolstrup M, Danneskiold-Samsøe B, Bock JE. Trainability of the pelvic floor. A prospective study during pregnancy and after delivery. *Acta Obstet Gynecol Scand* 1988;67:437-40.
- 6 Cunningham FG, MacDonald PC, Gant NF, Leveno KJ, Gilstrap LC, Hankins GDV, *Williams obstetrics 20th edition*. Stamford, CT: Appleton and Lange, 1997.
- 7 Bergsjø P, Maltau JM, Molne K, Nesheim B-L. *Obstetrikk* [in Norwegian]. Oslo: Universitetsforlaget, 1987.
- 8 Hofmeyr GJ, Kulier R. Cephalic version by postural management for breech presentation *Cochrane Database Syst Rev* 2000;(3):CD000051. (Accepted 25 May 2004)

doi 10.1136/bmj.38163.724306.3A

For the pleasure of their company

It was a normal surgery, nothing exciting, the usual complaints. Patients came and went. On looking back, however, I find that three patients stand out.

One patient gave me a hug, another told me that I was the best doctor in the world, and the third played a song for me with great care and feeling. Such encounters are not common; in fact, they are unique. The hug was given before I had time to anticipate it; the compliment was well meant without any hidden agenda; and the song—well, that was quite a treat on a grey winter morning.

It so happened that all three men had Down's syndrome, and any concern about having a too familiar relationship with their doctor wouldn't cross their minds. What they also had in common was their total dedication to and enjoyment of the moment, of an encounter with me. I know they will bring the same quality of human awareness and directness to the next person they meet, and the next doctor will be the best ever. These three men, with ages from 22 to 60, are all healthy, have a good

lifestyle, don't have many worries, and take pleasure in family and social life. In short, they are a delight to meet.

On reflection, I feel troubled. These are exactly the people whom we medics try with all our skills to prevent being born. What is so awful, so dreadful about their destiny that it is not worth living?

The likelihood that one of them will commit a crime or become a drug addict is quite slim. All three are in need of guidance, which will cost society money, but is such money less well spent than on the military or the new parliament building in Scotland? And I even refuse to believe they give their family so much heartache, after the initial shock at birth.

I am in the unusual situation of seeing patients with Down's syndrome quite regularly in my surgery, as more than 3% of our practice population have this condition. And I can tell you, I don't mind a bit.

Marga Hogenboom *general practitioner, Camphill Medical Practice, Aberdeen*