



Review Article

PRAGMATIC USAGE OF HARITAKI (*TERMINALIA CHEBULA* RETZ): AN AYURVEDIC PERSPECTIVE VIS-À-VIS CURRENT PRACTICE

Thrigulla Saketh Ram^{1*}, Bandari Srinivasulu², Ala Narayana³

^{1*}Research Officer (Ayu.), ²Consultant (Ayurveda), ³Director, National Institute of Indian Medical Heritage (NIIMH), Sy.No. 314, Revenue Board Colony, Gaddiannaram, Dilsukhnagar, Hyderabad, Andhra Pradesh, India.

ABSTRACT

Haritaki (Chebulic myrobalan-*Terminalia chebula* Retz (Combretaceae Family) denotes “a fruit having dark greenish yellow colour, which drives away diseases” The widely acclaimed Ayurvedic drug is the best wholesome substance and safe for bodily passages (*haritaki pathyanam*). *Haritaki* enjoys wide patronage in Tibetan medicine also which is known as A-ru-ra and praised with the adjective Sman-mchog-rgyal-pa (the king of best medicines). *Haritaki* in day to day life by judicious combinations of different adjuvants will be a boon for the person who consumes it. Though the Ayurvedic texts provide decent options such as *Rituharitaki* (Seasonal use of *Haritaki* with selective adjuvant), specific indications and contra indications, may practitioners as well as others do not pay much attention to them. This has led to the underutilization of this wonderful fruit drug. It is the time to take fresh look at the Ayurvedic perspective of pragmatic usage of *Haritaki* in the light of current therapeutic evidence, pharmaceutical practices.

INTRODUCTION

Haritaki consists of the pericarp of mature fruits of *Terminalia chebula* Retz. (Fam. Combretaceae), a moderate sized or large tree found throughout India, chiefly in deciduous forests and areas of light rainfall, but occasionally also in slightly moist forests, upto about 1500 m elevation, throughout India, flowers appear from April, August and fruits ripen from October-January.¹ It is one of the three fruits of *Triphala* and a universally acclaimed Ayurvedic medicine world over. The very first quartet of chapters of *Carakasamhita*, *Cikitsasthana* begins with the title “*Abhayamalakiyarasayanapada*”² where the term *Abhaya* means *Haritaki*, apart from this almost all the famous *Dravyaguna Nighantu* (Lexicons) have a chapter titled “*Haritakyadivarga*” indicating the therapeutic importance of it among Ayurvedic faculty. About seven varieties of this wonderful fruit along with instructions of how to use are widely discussed under these sections. The drug is the best wholesome substance and safe for bodily passages³. *Haritaki* enjoys wide patronage in Tibetan medicine and is known as A-ru-ra and praised with the adjective Sman-mchog-rgyal-pa (the king of best medicines)⁴.

The term *Haritaki* denotes “a fruit having dark greenish yellow colour, which drives away diseases”. It is also known as *Abhaya* (Very safe to use), *Avyatha* (cures pain), *Pathya* (safe for bodily passages), *Kayastha* (which provides strength to body), *Putana* (eliminates waste), *Haimavati* (grows on Himalayas), *Cetaki* (Helps to attain clear mind by cleansing channels), *Shreyasi* (Takes care of the person who ingests it) and *Shiva* (Provides all good things) in Sanskrit⁵.

Vernacular names⁶

Sanskrit	:	<i>Abhaya, Kayastha, shiva, Pathya, Vijaya</i> (Not Bhanga)
Asamese	:	<i>Shilikha</i>
Bengali	:	Haritaki
English	:	Myrobalan
Gujrati	:	Hirido, Himaja, Pulo-harda
Hindi	:	Harre, Harad, Harar
Kannada	:	Alalekai
Kashmiri	:	Halela
Malayalam	:	Katukka
Marathi	:	Hirda, Haritaki, Harda, Hireda
Oriya	:	Harida
Punjabi	:	Halela, Harar
Tamil	:	Kadukkai

Commentary on the Tibetan name of *Haritaki* A-ru-ra provides an interesting insight. The first syllable A denotes that it is the best of medicines to handle the deranged Doshas, Ru suggests that fruit consist of flesh (pericarp), bone (stone) and skin (epicarp) and ra is a similie comparing the shape of *Haritaki* fruit with that of Rhinoceros and has its effect on all the seven Dhatu (tissues)⁷.

Terminalia = proceeding from the extremity at the end; chebula= distorted from of the word kabul⁸.

The drug is equated with nectar drops fallen from heaven while it was consumed by *Indra*. Bhavaprakasha narrates this story in reverence as a dialogue between Ashivini kumara and *Daksha*.

Haritaki mentioned in Samhitas various groups, *Kushtagna dashaimani*⁹, *Virecanopag dashaimani*¹⁰, *Hikkanigraha dashaimani*¹¹, *Kasahara dashaimani*¹², *Jvarahara dashaimani*¹³, *Prajasthapana dashaimani*¹⁴, *Vayahsthapana*

*dashaimani*¹⁵, *Mustadi gana*¹⁶, *Triphala*¹⁷, *Amalakyadi gana*¹⁸ *Virecanopaga*¹⁹

Haritaki fruit comes bundled with five rasa viz., rind – *Madhura* (sweet), ridges - *Amla* (sour), stalk - *Tikta* (bitter), membrane - *Katu* (pungent), seed/stone - *Kashaya* (astringent). Owing to this formidable combination of five Rasas the whole fruit is capable of managing all the three *Doshas*. It has *Ruksha* (dryness producing), guru (causing heaviness) properties. The *Virya* is *Ushna* (in contrast to *Amalaki* - *Embelica officinalis*) and *Vipaka* (taste at the end of digestion) is *Madhura* (sweet).

Synonyms

Abhaya, Amruta, Avyatha, Bhashagvara, Cetaki, Cetanika, Devi, Divya, Haimavati, Jaya, Jivanika, Jivanti, Jivapriya, Jivya, Kayastha, Pathya, Pranada, Prapathya, Putana, Rohini, Shiva, Shreyasi, Vayastha, Vijaya^{20, 21}.

Important Karma (therapeutic activities)²² vis-à-vis current research leads

Karma (therapeutic activity)	Research lead
Rasayana	<p>Antioxidant and free radical scavenging activity</p> <ul style="list-style-type: none"> ➤ The leaves, bark and fruit of <i>T. chebula</i> demonstrated high antioxidant activity due to presence of phenolics²³ ➤ <i>T. chebula</i> (Aqueous extract) exhibited xanthine/xanthine oxidase inhibition, 2,2-diphenyl-1-picrylhydrazyl (DPPH) radicals scavenging activity²⁴ ➤ <i>T. chebula</i> (aqueous extract) Inhibits free radical induced hemolysis, nitric oxide release from lipopolysaccharide stimulated murine macrophages²⁵ ➤ <i>T. Chebula</i> (aqueous extract) has strong antioxidant activity due to inhibition of radiation induced lipid peroxidation, which is observed in rat liver microsomes at different doses²⁶ ➤ <i>T. Chebula</i> (methanolic extract) found to inhibit lipid peroxide formation and to scavenge hydroxyl and superoxide radicals <i>in vitro</i>²⁷ ➤ <i>T. Chebula</i> (Acetone extract) has stronger antioxidant activity than alpha²⁸ ➤ <i>T. chebula</i> (Gallic acid and chebulagic acid, isolated from fruit extract) blocked cytotoxic T lymphocyte (CTL)-mediated cytotoxicity. Granule exocytosis in response to anti-CD3 stimulation was also blocked by the above phytochemicals at the equivalent concentrations²⁹ ➤ <i>T. chebula</i> (ethanol extract) inhibited oxidative stress and the age-dependent shortening of the telomeric DNA length.³⁰ ➤ <i>T. chebula</i> gall were tested for antioxidative and tyrosinase inhibition activities as well as for proliferative and MMP-2 inhibition activities on early aging human skin fibroblasts to evaluate <i>in vitro</i> anti-aging activity. The cold water extract of <i>T. chebula</i> gall indicated the highest stimulation index (SI) on normal human fibroblast proliferation. The extract also demonstrated MMP-2 inhibition on fibroblasts 1.37 times more potent than ascorbic acid. The

	<p>study confirmed the traditional use of <i>T. chebula</i> gall in many Thai medicinal plant recipes for longevity³¹</p> <p>Cytoprotective activity</p> <ul style="list-style-type: none"> ➤ <i>T. chebula</i> extract on the age dependent shortening of the telomere length as shown by the Southern Blots of the terminal restriction fragments of DNA extracted from sub-culture passages³² ➤ It exhibited the development of duodenal ulcers and appeared to exert a cytoprotective effect on the gastric mucosa <i>in vitro</i>³³ ➤ oxidative stress and inhibitory effect on cellular aging of its fruits have also been documented³⁴ <p>Anticarcinogenic activity</p> <ul style="list-style-type: none"> ➤ Crude extract of <i>T. chebula</i> fruit have chebulinic acid, tannic acid and ellagic acid, which are found to be the most growth inhibitory phenolics of <i>T. chebula</i>³⁵ ➤ <i>T. chebula</i> fruit (Ethanol extract) inhibited cell proliferation and induced cell death in a dose dependent manner in several malignant cell lines including human (MCF-7) and mouse (S115) breast cancer cell line, human osteosarcoma cell line (HOS-1), human prostate cancer cell (PC-3) and a non-tumorigenic immortalized human prostate cell line (PNT1A)³⁶ ➤ <i>T. chebula</i> bark and fruit powder (acetone extract) has constituents with promising anticarcinogenic activity³⁷ <p>Antimutagenic, radioprotective and chemopreventive activity</p> <ul style="list-style-type: none"> ➤ <i>T. chebula</i> (aqueous extract and hydrolyzable tannins) demonstrated Antimutagenic activity in <i>Salmonella typhimurium</i>.³⁸ ➤ <i>T. chebula</i> (aqueous extract) breaks formati Gamma radiation induced strand in plasmid PBR322 DNA³⁹ ➤ <i>T. chebula</i> reduces irradiation effects in mice, human lymphocytes while undergoing the gamma radiation- <i>in vitro</i>⁴⁰ ➤ <i>T. chebula</i> showed chemopreventive effect on nickel chloride -induced renal oxidative stress, toxicity and cell proliferation response in male Wistar rats⁴¹
Cakshushya	<ul style="list-style-type: none"> ➤ <i>T. chebula</i> seeds (chloroform extract) in a quantity of 100, 200 and 300 mg/kg body weight produced dose-dependent reduction in blood glucose of diabetic rats in both short term and long term study (300 mg/kg body weight for 8 weeks). Further, remarkable renoprotective activity was also observed in <i>T. chebula</i> treated rats⁴²
Dipana / Anulomana, Vibandha Aruci, Udavarta, Gulma, Arsha	<p>Purgative property</p> <ul style="list-style-type: none"> ➤ Purgative action of an oil fraction from <i>T. chebula</i> has been reported⁴³ ➤ <i>T. chebula</i> increases the stools and has got property of evacuating the bowel completely demonstrated in a short term clinical trials carried out on patients with simple constipation⁴⁴ <p>Gastrointestinal motility improving and anti-ulcerogenic activity</p> <ul style="list-style-type: none"> ➤ <i>T. chebula</i> fruit has been shown to increase gastric emptying time⁴⁵ ➤ Antiulcerogenic activity of <i>T. chebula</i> appeared to be balanced with a protective effect on the gastrointestinal mucosa, with the improvement in the secretory status of Brunner's gland involved in the protection against duodenal ulcer⁴⁶ <p>Antispasmodic activity</p> <ul style="list-style-type: none"> ➤ <i>In vitvo</i> and <i>in vitro</i> studies on <i>T. chebula</i> demonstrated its 'anti-vata' or 'anti-spasmodic' properties by the reduction of abnormal blood pressure as well as intestinal spasms. This confirm its traditional usefulness for spastic colon

	<p>and other intestinal disorders⁴⁷</p> <p>Antiulcerogenic activity</p> <ul style="list-style-type: none"> ➤ Animals pretreated at 200 and 500 mg/kg body weight with hydroalcoholic extract of <i>T. chebula</i> showed reduction in lesion index, total affected area and percentage of lesion in comparison with control groups in the aspirin, ethanol and cold restraint stress-induced ulcer models. The <i>T. chebula</i> extract increased mucus production in aspirin and ethanol-induced ulcer models and showed antisecretory activity in pylorus ligated model leading to a reduction in the gastric juice volume, free acidity, total acidity, and significantly increased gastric pH⁴⁸
<i>Udararoga</i>	<p>Hepatoprotective activity</p> <ul style="list-style-type: none"> ➤ <i>T. chebula</i> fruits showed strong hepatoprotective activity through antioxidant effect in isolated rat hepatocytes⁴⁹ ➤ <i>T. chebula</i> was found to prevent the hepatotoxicity caused by the administration of rifampicin, isoniazid and pyrazinamide (combination) in sub-chronic model (12 weeks)⁵⁰ ➤ <i>T. chebula</i> fruit (95% ethanolic extract) showed hepatoprotective activity against anti-tuberculosis (anti-TB) drug-induced toxicity which could be attributed to its prominent anti-oxidative and membrane stabilizing activities⁵¹ ➤ Protective effects of an aqueous extract of <i>T. chebula</i> fruit on the tert-butyl hydroperoxide-induced oxidative injury was observed in cultured rat primary hepatocytes and rat liver^{52, 53}
<i>Hrudya /Pandu/ Hrudroga</i>	<p>Cardioprotective activity</p> <ul style="list-style-type: none"> ➤ <i>T. chebula</i> demonstrated cardioprotective activity in isolated frog heart model⁵⁴ ➤ <i>T. chebula</i> fruits (500 mg/kg body weight) was investigated in isoproterenol induced myocardial damage in rats. It was reported that the pretreatment with <i>T. chebula</i> extract had cardioprotective effect due to the lysosomal membrane stabilization preventing myocardial necrosis and inhibition of alterations in the heart mitochondrial ultrastructure and function in the experimental rats^{55,56,57} <p>Hypolipidemic and hypocholesterolemic activity</p> <ul style="list-style-type: none"> ➤ Hypolipidemic activity of <i>T. chebula</i> extract against experimentally induced atherosclerosis have been recorded⁵⁸ ➤ Combination of aqueous extract of <i>Terminalia chebula</i> and Gaumutra possessed hypocholesterolemic activity against cholesterol-induced hypercholesterolemia and atherosclerosis in rabbits⁵⁹ ➤ <i>T. chebula</i> extract administration showed hypolipidaemic activity against experimentally induced atherosclerosis and hypocholesterolemic activity against cholesterol-induced hypercholesterolemia and atherosclerosis⁶⁰ ➤ Triphala formulation was found to have hypolipidaemic effects on the experimentally induced hypercholesterolemic rats⁶¹
<i>Shotha</i>	<p>Anti-inflammatory and anti-arthritic activity</p> <ul style="list-style-type: none"> ➤ Terminalia chebula hydroalcoholic extract (TCHE) demonstrated Anti-arthritic and disease modifying activity of <i>Terminalia chebula</i> Retz. in experimental models⁶² <p>Wound healing activity</p> <ul style="list-style-type: none"> ➤ Topical administration of an alcoholic extract of <i>T. chebula</i> leaves on the healing of rat dermal wounds showed that <i>T. chebula</i> treated wounds healed

	<p>faster as indicated by improved rates of contraction and decreased period of epithelialization⁶³</p> <ul style="list-style-type: none"> ➤ <i>T. chebula</i> (ethanol extract) against the indomethacin-induced stomach ulceration was reported⁶⁴ ➤ Tannins extracted from immature fruits of <i>T. chebula</i> inhibited <i>Staphylococcus aureus</i> and <i>Klebsiella Pneumonia in vitro</i> and promoted cutaneous wound healing in rats due to a powerful anti-bacterial and angiogenic activity of the extract⁶⁵ ➤ The wound healing activity of ethanolic extract of fruits of <i>T. chebula</i> in the form of an ointment with two concentrations (5% and 10% w/w ointment of bark extract in simple ointment base) showed significant response in excision and incision models in albino rats compared to controls⁶⁶
Vishamajvara/ Jirnajvara	<p>anti plasmodial activity against <i>Plasmodium falciparum</i>⁶⁷</p> <ul style="list-style-type: none"> ➤ <i>T. chebula</i> (water extract) showed antiplasmodial activity <i>in vitro</i> by its ability to inhibit the uptake of [3H] hypoxanthine into the <i>Plasmodium falciparum</i> K1 multidrug-resistant strain and <i>in vivo</i>⁶⁸ ➤ Acetone seed extract of <i>T. chebula</i> was also found to have good antiplasmodial activity in a study⁶⁹ <p>Antiviral activity</p> <ul style="list-style-type: none"> ➤ It protects epithelial cells against influenza A virus, supporting its traditional use for aiding in recovery from acute respiratory infections⁷⁰
Prameha	<p>Antidiabetic and renoprotective activity</p> <ul style="list-style-type: none"> ➤ <i>T. chebula</i> fruit and seeds exhibited dose dependent reduction in blood glucose of streptozotocin induced diabetic rats both in short term and long term study and also had renoprotective activity^{71, 72} ➤ Oral administration of 75% methanolic extract of <i>T. chebula</i> (100 mg/kg body weight) reduced the blood sugar level in normal and alloxan diabetic rats significantly within 4 h. Continued daily administration of the drug produced a sustained effect⁷³ ➤ Oral administration of ethanolic extract of fruits of <i>T. chebula</i> (200 mg/kg body weight for 30 days) reduced the levels of blood glucose and glycosylated hemoglobin in streptozotocin (STZ)-induced experimental diabetic rats⁷⁴ ➤ <i>T. chebula</i> (200 mg/kg body weight for two months) reduced the elevated blood glucose and increase in glycosylated hemoglobin. The same dose also showed a marked improvement in controlling the elevated blood lipids as well as decreased serum insulin levels. The <i>in vitro</i> studies with pancreatic islets showed that the insulin release was nearly two times more than that in untreated diabetic animals. The treatment did not have any unfavorable effect on liver and kidney function tests⁷⁵
Shiroroga (dantaroga)	<p>Anticaries activity</p> <ul style="list-style-type: none"> ➤ <i>T. chebula</i> (aqueous extract) strongly inhibited the growth, sucrose induced adherence and glucan induced aggregation of <i>Streptococcus mutans</i>. Mouth rinsing with a 10% solution of the extract inhibited the salivary bacterial count and glycolysis of salivary bacteria for upto 90 min post rinsing^{76, 77} ➤ <i>T. chebula</i> demonstrated preventive action on dental caries⁷⁸

Important formulations⁷⁹

Name of the formulation	Indication(s)	Dose
<i>Abhayarishtha</i> ⁸⁰ (Bhaishajyaratnavali, Arshorogadhikara: 105-	<i>Arsha</i> (Haemorrhoids), <i>Udara</i> (Diseases of abdomen / enlargement of abdomen), <i>Mutra Vibandha</i> (Retention of urine), <i>Agnimandya</i> (Digestive impairment),	12 to 24 ml

110)	<i>Varcovibandha</i> (Impaction due to Hard stools)	
<i>Agastya Haritaki</i> Rasayana ⁸¹ (Ashtangahrudaya, Cikitsasthana, Adhyaya 3: 125-130)	<i>Hikka</i> (Hiccup), <i>Kasa</i> (Cough), <i>Shvasa</i> (Dyspnoea/Asthma), <i>Kshaya</i> (Pthisis), <i>Vishamajvara</i> (Intermittent fever), <i>Vali Palita</i> (Wrinkles in skin and graying of hair), Used as <i>Rasayana</i> (Nutrient to body and mind with adapto-immuno-neuro-endocrino-modulator properties)	6 to 12 g with Warm water or Milk
<i>Citraka Haritaki</i> ⁸² (Bhaishajyaratnavali, Nasarogadhikara: 31-33)	<i>Gulma</i> (Abdominal lump), <i>Udavarta</i> (Condition in which there is upward movement of <i>vayu</i>), <i>Pinasa</i> (Chronic rhinitis/sinusitis), <i>Kasa</i> (Cough), <i>Shvasa</i> (Dyspnoea/Asthma), <i>Arsha</i> (Haemorrhoids), <i>Agnimandya</i> (Digestive impairment), <i>Kshaya</i> (Pthisis), <i>Krumi</i> (Helminthiasis/Worm infestation)	6 - 12 g with milk.
<i>Danti Haritaki</i> ⁸³ (Ashtangahrudaya, Cikitsasthana, Adhyaya14: 92-96)	<i>Grahani</i> (Malabsorption syndrome), <i>Gulma</i> (Abdominal lump), <i>Arocaka</i> (Tastelessness), <i>Anaha</i> (Distension of abdomen due to obstruction to passage of urine and stools), <i>Pliharoga</i> (Splenic disease), <i>Vishamajvara</i> (Intermittent fever), <i>Kamala</i> (Jaundice), <i>Hrudroga</i> (Heart disease), <i>Arsha</i> (Haemorrhoids), <i>Shopha</i> (Oedema), <i>Gara</i> (Slow / accumulated poison), <i>Udara</i> (Diseases of abdomen / enlargement of abdomen), <i>Kushtha</i> (Diseases of skin), <i>Utklesha</i> (Nausea), <i>Pandu</i> (Anaemia), Used as <i>Virecaka</i> (Purgative)	1/2 to 1 Haritaki and 12 to 24g. of Leha
<i>Dashamula Haritaki</i> ⁸⁴ (Ashtangahrudaya, Cikitsasthana, Adhyaya 17: 14 -16)	<i>Shopha</i> (Oedema), <i>Arocaka</i> (Tastelessness), <i>Gara-Udara</i> (Abdominal disorder due to slow / accumulated poison), <i>Gulma</i> (Abdominal lump), <i>Pliha Roga</i> (Splenic disease), <i>Vaivarnya</i> (Discolouration), <i>Mutrakrucchra</i> (Dysuria), <i>Shukra Dosh</i> (Vitiation of semen), <i>Shvasa</i> (Dyspnoea/Asthma), <i>Jvara</i> (Fever), <i>Meha</i> (Excessive flow of urine), <i>Karshya</i> (Emaciation), <i>Raktapitta</i> (Bleeding disorder), <i>Amavata</i> (Rheumatism)	6 to 12 g with Water, Milk
<i>Brahma Rasayana</i> ⁸⁵ (Ashtangahrudaya, Uttarasthana, Adhyaya 39 : 15-23)	<i>Tandra</i> (Lassitude), <i>Shrama</i> (Fatigue/lethargy), <i>Klama</i> (Tiredness without exertion/Langour), <i>Manodaurbalya</i> (Mental weakness), <i>Jara</i> (Senility/Progeriasis), <i>Vali Palita</i> (Wrinkles in skin and graying of hair), <i>Smrutibhrama</i> (Impairment of memory)	12g with Milk
<i>Triphala Cur,</i> ^{a86} (Bhavaprakasha, Haritakyadi Varga : 41- 42)	<i>Anaha</i> (Distension of abdomen due to obstruction to passage of urine and stools), <i>Prameha</i> (Urinary disorders), <i>Netraroga</i> (Eye disorder), <i>Kaphapittaroga</i> (Disease due to <i>Kapha dosha</i> and <i>Pitta dosha</i>), <i>Kushtha</i> (Diseases of skin), <i>Mandagni</i> (Impaired digestive fire), <i>Aruci</i> (Tastelessness), <i>Vishamajvara</i> (Intermittent fever).	3 to 6 g
<i>Triphaladi Taila</i> ⁸⁷ (Sahasrayoga, Tailaprakarana : 44)	<i>Shiroroga</i> (Disease of head), <i>Pratishyaya</i> (Coryza), <i>Khalitya</i> (Alopecia), <i>Palitya</i> (Graying of hair, particularly of scalp), <i>Keshashata</i> (Loss of hair), <i>Urdhvajatrugata Roga</i> (Disorders of body parts above clavicle)	Used externally for Abhyanga
<i>Abhaya Lavana</i> ⁸⁸ (Bhaishajyaratnavali, Plihayakrudrogadhikara: 35-43)	<i>Anaha</i> (Distension of abdomen due to obstruction to passage of urine and stools), <i>Gulma</i> (Abdominal lump), <i>Yakrutplihodara</i> (Disease of liver and spleen), <i>Hrudroga</i> (Heart disease), <i>Ashthila</i> (Prostatic Hyperplasia),	1 to 2 g

	<i>Mandagni</i> (Impaired digestive fire), <i>Shirahshula</i> (Headache), <i>Sharkara</i> (Gravel in urine), <i>Ashmari</i> (Calculus)	
<i>Pathyadi Lepa</i> ⁸⁹ (Yogaratanakara, Kushthadhikara : Page 682)	<i>Kushtha</i> (Diseases of skin)	External application

Best quality of *Haritaki* dry fruit

Best variety of *Haritaki* has smooth texture, bulky, heavy and drowns in water. Roughly it weighs 20 grams equivalent to two *Karsha*. This type of dry *Haritaki* is considered as best for medicinal usage⁹⁰.

Usage

The drug is significantly prescribed in therapeutic as well as preventive use and process of restorative or alterative indication (*Rasayana karma*), and it is classically recommended as *Ritu haritaki* for using the drug during the course of different seasons viz., *Varsha* (Rains) – *Saindhavalavana* (Rock salt), *Sharad Ritu* (Autumn)- *Sarkara* (sugar), *Hemanta* (Cold/winter)-*Shunti* (Dry Ginger), *Sishira* (late winter early spring)- *Pippali* (long pepper), *Vasanta* (Spring)- *Madhu* (honey) and *Grishma* (Summer)- *Guda* (Jaggery)⁹¹.

The drug is stomochic (*Agnivardhana*) on chewing (*Carvita*), it is laxative (*Malashodhana*) on licking (*Peshita*) astringent (*Sangrahi*) on boiling (*Swinna*) and alleviating three *Doshas* (*Tridoshanut*) on frying (*Bhrishta*). Taken along with food it enhances mental ability, alleviates the ill effects of aggravated *Pitta*, *Kapha* and *Vata* and restores the free flow of urine, faeces and sweat. Taking the fruit after food ensure removal of defects due to faulty food intake⁹².

The fruit consumed along with *Lavana* (salt) mitigates *Kapha*, with sugar-*Pitta* and with ghee *Vata*. The combination of *Haritaki* with Jaggery is considered best as it takes care of all the three disturbed *Doshas*⁹³.

1. 2-3 g of *Haritaki* powder mixed with equal amount of jiggery before consuming food is very good for Persons suffering with *Arshas* (Hemorrhoids).
2. Chronic colitis *Haritaki* powder with hot water is advised.

3. Licking *Haritaki* powder mixed with honey is good to prevent or control motion sickness and vomitings.
4. Prescribing *Haritaki* powder with Jaggery is good to manage *Vatarakta* (Descriptive arthropathies).
5. *Haritaki* Seed powder mixed with hot milk is good for treating Urinary Calculi.
6. *Haritaki* acts as General Tonic promoting strength and vigor when used with Ghee. The best method to achieve this is to first fry *Haritaki* Fruits in Ghee followed by powdering of the same. The leftover ghee after frying has to be kept aside (not to be discarded). The *Haritaki* powder prepared after frying in ghee is to be mixed with the left over ghee and to be taken twice a day.
7. *Haritaki* decoction mixed with boiled milk is very useful to manage *madatyaya* (Hangover).
8. *Haritaki* powder mixed with equal quantity of Ghee is prescribed for managing Colicky abdominal pain due to *Pittaja shula* (Acid peptic disorders). The same results can be achieved with combination of *Draksha* (Dry Grapes) and *Haritaki* powder.
9. Use of *Erandataila* (castor oil) with *Haritaki* powder is recommended for all types of rheumatological conditions. (*Shodhala Nighantu*).
10. Applying the paste of Fine *Haritaki* powder mixed with fresh turmeric rhizome juice manages Chippa (Nail bed infection, Whitlow).

CONTRAINDICATIONS

The fruit is not recommended for regular use for people who are suffering with fatigue, tired, weight loss, dry parched skin, not having physical strength (weak). *Haritaki* is contraindicated during *Pitta prakopa* state, *Garbhini* (pregnancy), during acute fever, person who has undergone *Raktamokshana* (blood letting)⁹⁴.

In summary use of *Haritaki* in day to day life by judicious combinations of different

adjuvants will be a boon for the person who consumes it. All Ayurveda physicians and people who follow Ayurvedic way of life must exploit the advantages offered by this wonderful drug. The mantra for sure success in Ayurveda practice lies in making the best out of well established, genuine drugs like *Haritaki* than trying to go after very rarely available ones. The author wishes to remind you the words from *Carakasamhita* which state that use of *Haritaki* results in *Shivam* i.e everything good and in the interest of the user.

REFERENCES

- 1 . Department of ISM, MoHFW, Government of India. (2001 (Reprint)). *The Ayurvedic Pharmacopoeia of India*. New Delhi: NISCOM (CSIR).p.47.
- 2 . Charaka Samhita, Ram Karan Sharma & Vaidya Bhagwan Dash, Chowkambha Sanskrit Series office, Varanasi. Reprint: 2008; Vol- 3; pp.3
- 3 . Ibid; Vol- 1; pp.429
- 4 . Vd. Bhagwan Dash, Kailash, Volume IV, 1976, Number 11, p. 5 at http://himalaya.socanth.cam.ac.uk/collections/journals/kailash/pdf/kailash_04_01_01.pdf and Vd. Bhagwan Dash, Haritaki – A Comparative Study of Literature in Ayurveda and Tibetan Medicine, BIIMH, Vo. IV NO.1 p. 1-8
- 5 . Bhavaprakasha of Bhavamishra, Commentary by Dr. B.Sitaram, Chaukhambha Orientalia, Varanasi 2006
- 6 . Department of ISM, MoHFW, Government of India. (2001 (Reprint)). *The Ayurvedic Pharmacopoeia of India*. New Delhi: NISCOM (CSIR).p.47.
- 7 . Vd. Bhagwan Dash, Kailash, Volume IV, 1976, Number 11, p. 6 at http://himalaya.socanth.cam.ac.uk/collections/journals/kailash/pdf/kailash_04_01_01.pdf and Vd. Bhagwan Dash, Haritaki – A Comparative Study of Literature in Ayurveda and Tibetan Medicine, BIIMH, Vo. IV NO.1 p. 1-8
8. Vaidya Vishnu Mahadev Gogte. Ayurvedic Pharmacology and Therapeutic Uses of Medicinal Plants (Dravyagunavignyan). Shri S. Ramakrishnan, Executive Secretary, Bharatiya Vidya Bhavan, Kulapti Munshi Marg, Mumbai. Edition: 2000; pp.515-516
- 9 Charaka Samhita, Ram Karan Sharma & Vaidya Bhagwan Dash, Chowkambha Sanskrit Series office, Varanasi. Reprint: 2008; Vol- 1; pp.91
- 10 Ibid. pp. 94
- 11 Ibid. pp. 95-96
- 12 Ibid. pp. 97
- 13 Ibid. pp. 98
- 14 Ibid. pp. 100-101
- 15 Ibid. pp. 101
- 16 G.D. Singhal & Colleagues, *Susruta-Samhita Ancient Indian Surgery*, Chaukhamba Sanskrit Pratishthan, Delhi. Reprint: 2007: Vol: 3; pp.89
- sush. Vol.1 pp.316
- 17 Ibid pp. 316-317
- 18 Ibid pp. 317
- 19 Ibid pp. 322
- 20 Bhavaprakasa of Bhavamisra, Commentary by Bulusu Sitaram, Chaukhambha Orientalia, Varanasi. Vol-1, Edition: 2006; pp. 131
- 21 Rajanighantu of Pandit Narahari by Dr.Indradev Tripathi, Chowkhambha Krishnadas Academy, Varanasi. 2006; pp.384
- 22 . Department of ISM, MoHFW, Government of India. (2001 (Reprint)). *The Ayurvedic Pharmacopoeia of India*. New Delhi: NISCOM (CSIR).p.47.
- 23 Chang CL, Lin CS. Development of antioxidant activity and pattern recognition of *Terminalia chebula* Retzius extracts and its fermented products. *HungKuang J.* 2010;61:115–129.
- 24 Naik GH, Priyadarsini KI, Naik DB, Gangabhairathi R, Mohan H. Studies on the aqueous extract of *Terminalia chebula* as a potent antioxidant and a probable radioprotector. *Phytomedicine.* 2004;11(6):530–538.
- 25 Mahesh R, Bhuvana S, Begum VM. Effect of *Terminalia chebula* aqueous extract on oxidative stress and antioxidant status in the liver and kidney of young and aged rats. *Cell Biochem Funct.* 2009;27(6):358–363.
- 26 Lee HS, Won NH, Kim KH, Lee H, Jun W, Lee KW. Antioxidant effects of aqueous extract of *Terminalia chebula* *in vivo* and *in vitro*. *Biol Pharm Bull.* 2005;28(9):1639–1644.
- 27 Lee HS, Jung SH, Yun BS, Lee KW. Isolation of chebulic acid from *Terminalia chebula* Retz. and its antioxidant effect in isolated rat hepatocytes. *Arch Toxicol.* 2007;81(3):211–218.
- 28 Chen X, Sun F, Ma L, Wang J, Qin H, Du G. *In vitro* evaluation on the antioxidant capacity of triethylchebulate, an aglycone from *Terminalia chebula* Retz fruit. *Indian J Pharmacol.* 2011;43(3):320–323
- 29 Hamada S, Kataoka T, Woo JT, Yamada A, Yoshida T, Nishimura T, Otake N, Nagai K. Immunosuppressive effects of gallic acid and chebulagic acid on CTL-mediated cytotoxicity. *Biological & Pharmaceutical Bulletin* 1997;20(9):1017-19.
- 30 Na MK, Bae KH, Kang SS, Min BS, Yoo JK, Kamiryo Y, Senoo YI, Yokoo S, Miwa N. Cytoprotective effect on

oxidative stress and inhibitory effect on cellular aging of *Terminalia chebula* fruit. *Phytotherapy Res* 2004;18(9) 737-41.

31 Manosroi A, Jantrawut P, Akihisa T, Manosroi W, Manosroi J. *In vitro* anti-aging activities of *Terminalia chebula* gall extract. *Pharm Biol.* 2010;48(4):469-81.

32 Minkyun NA, Wan BAE, Kang SS, Min BS, Yoo JK, Yuk OK, et al. et al. Cytoprotective effect on oxidative stress and inhibitory effect on cellular aging of *Terminalia chebula* fruit. *Phytother Res.* 2004;18:737-741.

33 Lee HS, Koo YC, Suh HJ, Kim KY, Lee KW. Preventive effects of chebulic acid isolated from *Terminalia chebula* on advanced glycation endproduct-induced endothelial cell dysfunction. *J Ethnopharmacol.* 2010;131(3):567-574.

34 Na M, Bae M, Keng SS, Min BS, Yoo JK, Kamiryo Y, et al. et al. Cytoprotective effect on oxidative stress and inhibitory effect on cellular aging of *Terminalia chebula* fruit. *Phytother Res.* 2004;18(9):737-741.

35 Saleem M, Hushum P, Harkonen K, Pihlaja Inhibition of cancer cell growth by crude extract and phenolics of *Terminalia chebula* fruit. *J Ethnopharmacol.* 2002;81:327-336.

36 Reddy DB, Reddy TC, Jyotsna G, Sharan S, Priya N, Lakshmipathi V, et al. et al. Chebulagic acid, a COX-LOX dual inhibitor isolated from the fruits of *Terminalia chebula* Retz., induces apoptosis in COLO-205 cell line. *J Ethnopharmacol.* 2009;124(3):506-512.

37 Reddy DB, Reddy TC, Jyotsna G, Sharan S, Priya N, Lakshmipathi V, et al. et al. Chebulagic acid, a COX-LOX dual inhibitor isolated from the fruits of *Terminalia chebula* Retz., induces apoptosis in COLO-205 cell line. *J Ethnopharmacol.* 2009;124(3):506-512.

38 Grover IS, Bala S. Antimutagenic activity of *Terminalia chebula* (myroblan) in *Salmonella typhimurium*. *Indian J Exp Biol.* 1992;30(4):339-341

39 Naik GH, Priyadarsini KI, Naik DB, Gangabhairathi R, Mohan H. Studies on the aqueous extract of *Terminalia chebula* as a potent antioxidant and a probable radioprotector. *Phytomedicine.* 2004;11(6):530-538.

40 Gandhi NM, Nayar CKK. Radiation protection by *Terminalia chebula* some mechanistic aspects. *Mol Cell Biochem.* 2005;277(1-2):43-48

41 Prasad L, Husain Khan T, Jahengir T, Sultana S. Chemomodulatory effect of *Terminalia chebula* against nickel chloride-induced oxidative stress and tumor promotion response in male Wistar rats. *J Trace Elem Med Biol.* 2006;20(4):233-239

42 Rao NK, Nammi S. Antidiabetic and renoprotective effects of the chloroform extract of *Terminalia chebula* Retz. seeds in streptozotocin-induced diabetic rats. *BMC Complementary and Alternative Medicine* 2006;6:17.

43 Vani T, Rajani M, Sarkar S, Shishoo CJ. Antioxidant properties of ayurvedic formulation triphala and its constituents. *Int J Pharmacog.* 1997;35:313-317.

44 Mukherjee P, Roy S, Bhattacharyya S, Debnath PK, Jan U, Pandit S, et al. et al. Clinical study of Triphala- a well known phytomedicine from India. *Indian J Pharmcol Ther.* 2006;5:51-54.

45 Tamhane MD, Thorate SP, Rege NN, Dahanukar SA. Effect of oral administration of *Terminalia chebula* on gastric emptying: An experimental study. *J Postgrad Med.* 1997;43(1):12-13.

46 Sharma P, Prakash T, Kotresha D, Ansari MA, Sahrm UR, Kumar B, et al. et al. Antiulcerogenic activity of *Terminalia chebula* fruit in experimentally induced ulcer in rats. *Pharm Biol.* 2011;49(3):262-268.

47 Seyyed AM, Ali V, Mohammad KGN, Peyman M. Spasmogenic activity of the seed of *Terminalia chebula* Retz in rat small intestine: *In vitvo* and *in vitro* studies. *Malays J Med Sci.* 2011;18(3):18-26.

48 Sharma P, Prakash T, Kotresha D, Ansari MA, Sahrm UR, Kumar B, Debnath J, Goli D. Antiulcerogenic activity of *Terminalia chebula* fruit in experimentally induced ulcer in rats. *Pharm Biol.* 2011;49(3):262-68.

49 Lee HS, Jung SH, Yun BS, Lee KW. Isolation of chebulic acid from *Terminalia chebula* Retz. and its antioxidant effect in isolated rat hepatocytes. *Arch Toxicol.* 2007;31(3):211-218

50 Tasduq SS, Singh AK, Salti NK, Gupta DK, Suri K. *Terminalia chebula* fruits prevent liver toxicity caused by sub-chronic administration of rifampicin, isoniazid and pyrazinamide (PZA) in combination. *Hum Exp Toxicol.* 2006;25(3):11-18.

51 Tasduq SA, Singh K, Satti NK, Gupta DK, Suri KA. *Terminalia chebula* (fruit) prevents liver toxicity caused by sub-chronic administration of rifampicin, isoniazid and pyrazinamide in combination. *Hum Exp Toxicol* 2006;25:111-18.

52 Lee HS, Won NH, Kim KH, Lee H, Jun W, Lee KW. Antioxidant effects of aqueous extract of *Terminalia chebula* *in vitvo* and *in vitro*. *Biol Pharm Bull.* 2005;28(9):1639-1644.

53 Lee HS, Jung SH, Yun BS, Lee KW. Isolation of chebulic acid from *Terminalia chebula* Retz. and its antioxidant effect in isolated rat hepatocytes. *Arch Toxicol.* 2007;81(3):211-218.

54 Reddy VRC. Cardioprotective activity of the fruit of *Terminalia chebula*. *Fitoterapia.* 1990;61:517-525

55 Suchalatha S, Shyamala Devi CS. Protective effect of *Terminalia chebula* against experimental myocardial injury induced by isoproterenol. *Indian J Exp Biol* 2004;42(2):174-78.

56 Suchalatha S, Shyamala Devi CS. Protective effect of *Terminalia chebula* against lysosomal enzyme

- alterations in isoproterenol-induced cardiac damage in rats. *Exp Clin Cardiol* 2005;10(2):91-95.
- 57 Suchalatha S, Srinivasan P, Shyamala Devi CS. Effect of *T. chebula* on mitochondrial alterations in experimental myocardial injury. *Chemico-Biological Interactions* 2007;169:145-53.
- 58 Maruthappan V, Shree KS. Hypolipidemic activity of Haritaki (*Terminalia chebula*) in atherogenic diet induced hyperlipidemic rats. *J Adv Pharm Tech Res.* 2010;1:229-235
- 59 Israni DA, Patel KV, Gandhi TR. Anti-hyperlipidemic activity of aqueous extract of *Terminalia chebula* and Gaumutra in high cholesterol diet fed rats. *Int J Pharm Sci.* 2010;1(1):48-59.
- 60 Shaila HP, Udupa SL, Udupa AL. Hypolipidemic activity of three indigenous drugs in experimentally induced atherosclerosis. *Int J Cardiol* 1998;67(2):119-24.
- 61 Saravanan S, Shrikumar R, Manikandan S, Jayaparthasarathy N, Sheela Devi R. Hypolipidemic effect of Triphala in experimentally induced hypercholesteremic rats. *Yakugaku Zasshi* 2007;127:2385-388.
- 62 Nair V, Singh S, Gupta YK. Anti-arthritis and disease modifying activity of *Terminalia chebula* Retz. in experimental models. *J Pharm Pharmacol.* 2010;62(12):1801-1806.
- 63 Li K, Diao Y, Zhang H, Wang S, Zhang Z, Yu B, et al. et al. Tannin extracts from immature fruits of *Terminalia chebula* Fructus Retz. promote cutaneous wound healing in rats. *BMC Comp Alter Med.* 2011;11:1-9.
- 64 Bhattacharya S, Chaudhuri SR, Chattopadhyay S, Bandyopadhyay SK. Healing properties of some Indian medicinal plants against indomethacin-induced gastric ulceration of rats. *J Clin Biochem and Nutrition* 2007;41(2):106-14.
- 65 Li K, Diao Y, Zhang H, Wang S, Zhang Z, Yu B, Huang S, Yang H. Tannin extracts from immature fruits of *Terminalia chebula* Fructus Retz. promote cutaneous wound healing in rats. *BMC Complementary and Alternative Med* 2011;11:8.
- 66 Choudhary GP. Wound healing activity of ethanolic extract of *Terminalia chebula* Retz. *Int J Pharma and Bio Sci* 2011;2(1):48-52.
- 67 Bagavan A, Rahuman AI, Kamaraj C, Kaushik NK, Mohanakrishnan D, Sahal D. Antiplasmodial activity of botanical extracts against *Plasmodium falciparum*. *Parasitol Res.* 2011;108(5):1099-1109.
- 68 Pinmai K, Hiriote W, Soonthornchareonnon N, Jongsakul K, Sireeratawong S, Tor-Udom S. *In vitro* and *in vivo* antiplasmodial activity and cytotoxicity of water extracts of *Phyllanthus emblica*, *Terminalia chebula*, and *Terminalia bellerica*. *J Med Assoc Thai.* 2010;93(7):120-26.
- 69 Bagavan A, Rahuman AA, Kamaraj C, Kaushik NK, Mohanakrishnan D, Sahal D. Antiplasmodial activity of botanical extracts against *Plasmodium falciparum*. *Parasitol Res* 2011;108:1099-109.
- 70 Badmaev V, Nowakowski M. Protection of epithelial cells against influenza A virus by plant derived biological response modifier Ledretan-96. *Phytother Res.* 2000;44(4):245-249.
- 71 Kannan VR, Rajasekar GS, Rajesh P, Balasubramanian V, Ramesh N, Solomon EK, et al. et al. Anti-diabetic activity on ethanolic extracts of fruits of *Terminalia chebula* Retz. Alloxan induced diabetic rats. *Am J Drug Discov Dev.* 2012;2:135-142.
- 72 Senthilkumar GP, Subramanian SP. Biochemical studies on the effect of *Terminalia chebula* on the levels of glycoproteins in streptozotocin-induced experimental diabetes in rats. *J Appl Biomed.* 2008;6:105-115.
- 73 Sabu MC, Kuttan R. Anti-diabetic activity of medicinal plants and its relationship with their antioxidant property. *J Ethnopharmacol* 2002;81:155-60.
- 74 Kumar GPS, Arulselvan P, Kumar DS, Subramanian SP. Anti-diabetic activity of fruits of *Terminalia chebula* on streptozotocin induced diabetic rats. *J Health Sci* 2006;52(3):283-91.
- 75 Murali YK, Anand P, Tandon V, Singh R, Chandra R, Murthy PS. Long-term effects of *Terminalia chebula* Retz. on hyperglycemia and associated hyperlipidemia, tissue glycogen content and *in vitro* release of insulin in streptozotocin induced diabetic rats. *Exp Clin Endocrinol Diabetes.* 2007;115(10):641-46.
- 76 Aneja KR, Joshi R. Evaluation of antimicrobial properties of fruit extracts of *Terminalia chebula* against dental caries pathogens. *Jundishapur J Microbiol.* 2009;2(3):105-111.
- 77 Carounanidy U, Satyanarayanan R, Velmurugan A. Use of an aqueous extract of *Terminalia chebula* as an anticaries agent: a clinical study. *Indian J Dent Res.* 2007;18(4):152-156.
- 78 Aneja KR, Joshi R. Evaluation of antimicrobial properties of fruit extracts of *Terminalia chebula* against dental caries pathogens. *Jundishapur J Microbiol.* 2009;2(3):105-111.
79. Department of ISM, MoHFW, Government of India. (2001 (Reprint)). *The Ayurvedic Pharmacopoeia of India*. New Delhi: NISCOM (CSIR).p.47.
- 80 . Department of ISM, MoHFW, Government of India. (2003). *The Ayurvedic Formulary of India, Part-1*. New Delhi: NISCOM(CSIR) p. 6.

- 81 . Ibid p. 6 & 31
82 . Ibid 6 & 37
83 . Ibid p. 6 & 39
84 . Ibid p. 6 & 40
85 . Ibid p. 6 & 43
86 . Ibid p. 6 & 110
87 . Ibid p. 136
88 . Ibid p. 164
89 . Ibid p. 173
90 . Bulusu Sitaram. (2006). *Bhavaprakasa of Bhava Misra Vol. 1*. Varanasi: Chaukhamba Orientalia. p.133
91 . B Sitaram. (2010). *Bhavaprakasa of Bhava Misra Vol. 2*. Varanasi: Chaukhamba Orientalia. P.737
91 . Bulusu Sitaram. (2006). *Bhavaprakasa of Bhava Misra Vol. 1*. Varanasi: Chaukhamba Orientalia.
92 . Ibid, p.133
93 . Ibid.
94 . Ibid

Cite this article as:

Thrigulla Saketh Ram, Bandari Srinivasulu, Ala Narayana.
Pragmatic Usage of Haritaki (*Terminalia Chebula Retz*): An
Ayurvedic Perspective Vis- à -vis Current Practice. Int. J.
Ayur. Pharma Research 2013; 1 (3): 72-82

Source of support: Nil, Conflict of interest: None Declared

***Address for correspondence**

Dr. Thrigulla Saketh Ram
Research officer (Ayu.)
National Institute of Indian Medical
Heritage, Survey No. 314
Revenue Board Colony
Gaddiannaram, Hyderabad - 500036
Andhra Pradesh, India.
Cell: +919849995719
Email: dr.saketram@gmail.com