WATER RESEARCH 40 (2006) 851–852

Available at www.sciencedirect.com

SCIENCE DIRECT®

journal homepage: www.elsevier.com/locate/watres

Erratum

Erratum to "Modeling monochloramine loss in the presence of natural organic matter" [Water Research 39(14), 3418–3431]

*Stephen Edward Duirk**

United States Environmental Protection Agency, 960 College Station Road, Athens, GA 30605, USA

Page 3422, Fig. 1—Hypochlorous acid in the monochloramine reaction pathway schematic should be HOCl and not HOCl3.

Page 3426, Fig. 5b—Figs. 5(a) and (b) are the same figure. Fig. 5b should show ammonia formation due to monochloramine loss in the presence of IRWC.



Fig. 1 – Schematic of monochloramine reaction pathways in the presence of NOM. DOC_{ox} represents oxidized NOM.

DOI of original article: 10.1016/j.watres.2005.06.003

*Tel.: +1 319 335 5653; fax: +1 319 355 5660.
E-mail address: duirk.stephen@epa.gov.

0043-1354/$ - see front matter © 2006 Elsevier Ltd. All rights reserved.

doi:10.1016/j.watres.2005.12.022



Fig. 5 – (b) Ammonia formation as a function of pH in IRWC. $[NH_2Cl]_o = 0.05$ mM, $[TOTCO_3] = 4.0$ mM, $\mu = 6.0$ mM, temperature = 25 °C, Cl/N = 0.7 mol/mol and [TOC] = 3.3 mg C/L. Lines represent model results.