

## Biologic response of sperm and seminal plasma to transient testicular heating

Zhen-Ya Fang<sup>1</sup>, Wei Xiao<sup>2</sup>, Su-Ren Chen<sup>3</sup>, Mei-Hua Zhang<sup>1</sup>, Yi Qiu<sup>1</sup>, Yi-Xun Liu<sup>3, 4</sup>

<sup>1</sup>Key Laboratory of Birth Regulation and Control Technology of National Health and Family Planning Commission of China, Key Laboratory for Improving Birth Outcome Technique, Maternal and Child Health Care Hospital of Shandong Province, Jinan, Shandong 250014, China, <sup>2</sup>Shandong Provincial Hospital Affiliated to Shandong University, <sup>3</sup>State Key Laboratory of Stem Cell and Reproductive Biology, Institute of Zoology, Chinese Academy of Sciences, Beijing, 100101, China, <sup>4</sup>University of Chinese Academy of Sciences, Beijing, 100049, China

### TABLE OF CONTENTS

1. Abstract
2. Introduction
3. Research strategy, experimental animals, subjects, and materials for TTH as a contraceptive method
  - 3.1. Research strategy
  - 3.2. TTH method
4. Evidence of alteration of spermatogenesis and sperm parameters under TTH
5. Effects of TTH on the testis, histology and germ cells
6. Sperm DNA damage, oxidative stress and apoptosis in TTH treatment phase
  - 6.1. Sperm DNA damage, oxidative stress and reactive oxygen species (ROS) after TTH treatment
  - 6.2. Effect of TTH on sperm chromosome and acrosin activity
  - 6.3. Effect of TTH on caspase
7. Effects of TTH on sperm protein expression, autophagy and molecular changes
  - 7.1. Effect of TTH on sperm protein expression
  - 7.2. Autophagy and apoptosis of sperm or germ cells in TTH treatment phase
  - 7.3. Molecular changes in TTH
8. Effect of TTH on epididymis and seminal biochemistry
  - 8.1. Effect of TTH on epididymal proteins
  - 8.2. Effect of TTH on epididymal sperm
  - 8.3. Effect of TTH on NO and NOS
  - 8.4. Effect of TTH on MIF
  - 8.5. Changes in seminal NAG, L-carnitine and epidermal growth factor
9. Conclusions
10. Acknowledgment
11. References

### 1. ABSTRACT

Currently, there are few male contraceptive methods that are purely based on prevention of the entry of the sperm into the female reproductive tract. An alternative approach for designing reversible male contraceptive is achieved by transient testicular heating (TTH). This treatment, through massive germ cell apoptosis, causes reversible oligospermia or azoospermia. Here, we describe as how TTH causes DNA damage, oxidative stress, apoptosis, autophagy, sperm protein expression, and alters the

biochemical components of seminal plasma. Further understanding of TTH will help design safe and reversible male contraception.

### 2. INTRODUCTION

Ideal birth control methods for males must reduce the number of fertile sperm to reliably prevent fertilization, and such methods must be relatively long-lasting, reversible, easy to use and have no

effect on the hormone secretion (1-5). Unfortunately, among currently available male contraceptive options, the reversible methods are not reliable, while the reliable methods are not reversible. Although female contraceptive methods (such as sterilization, intrauterine devices, or hormonal contraception) are very effective in preventing unintended pregnancy, some women cannot use them due to health conditions or side effects. In addition, many men wish to take active responsibility for family planning. As early as 1959, testicular heating is considered as a potential contraceptive method for men (2, 6). It is well known that functions of mammalian testes are very sensitive to temperature. Elevated testicular temperature detrimentally affects the spermatogenesis, leading to reversible oligospermia or azoospermia in animals and humans via germ cell apoptosis (7-10). The scrotum of most adult mammalian species, including humans, is a local thermo-regulator, and spermatogenesis optimally occurs at temperatures slightly lower than the body temperature (at least 2°C–8°C below core body temperature) (7, 8, 11-14). In general, the temperature of hot spring and transient testicular heating (TTH) is higher than the scrotal temperature (from 37°C to 45°C). Therefore, TTH exposure can reduce testis size, lead to spermatogenic damage, decrease sperm production, cause apoptosis of sperm and impair semen quality if the scrotal temperature is increased over time (15, 16). An elevated testicular temperature damages the seminiferous epithelium, resulting in germ cell loss as well as morphological and functional alternations in Sertoli cells (SCs) (17, 18). TTH can impair protein and RNA biosynthesis, cause oxidative stress and apoptosis, damage spermatocyte DNA, and disrupt chromatin packing in sperm nucleus (10, 12, 19). The mechanisms underlying molecular response of germ cells to TTH may involve heat-induced germ cell damage, including apoptosis, DNA damage and autophagy (13). TTH mainly leads to heat stress and affects the metabolism and apoptosis of spermatogenic cells, resulting in compromised sperm quality and impaired fertility of many animal models (20-22). These findings provide an important theoretical basis for designing male contraceptive methods (5).

The biochemical components of seminal plasma, such as fructose, neutral  $\alpha$ -glucosidase (NAG), carnitine, nitric oxide (NO), nitric oxide synthase (NOS), macrophage migration inhibitory factor (MIF) and epidermal growth factor (EGF), are important for sperm motility, morphology and capacitation (20, 21). TTH may affect the testis and epididymis, thus reducing the semen quality, damaging the sperm membrane and nucleus, and interfering with sperm maturation.

In the present review, we aimed to give an update on the TTH-based male contraceptive methods, including the materials, devices and technologies as

well as the molecular determinants of sperm quality and biochemical components in seminal plasma.

### 3. RESEARCH STRATEGY, EXPERIMENTAL ANIMALS, SUBJECTS, AND MATERIALS FOR TTH AS A CONTRACEPTIVE METHOD

#### 3.1. Research strategy

We conducted a systematic review of the literature concerning subjecting individuals to TTH, and its effect on semen quality, sperm protein and RNA biosynthesis, spermatocyte DNA, biochemical components of seminal plasma, and fertility or reproductive health (Figure 1). Regarding scrotal heating, we used comprehensive terms in order to optimize the search. Regarding the sperm quality, we selected a strategy that combined comprehensive terms and scrotal heat stress or hyperthermia in order not to omit any articles.

#### 3.2. TTH method

At present, only few approaches other than condoms and vasectomy have been adequately developed for male contraception (5, 10, 23-26). Condoms and vasectomy cannot satisfy all users (27, 28). The major limitations of condoms for contraception are relatively high failure rates and interference with sexuality (29). As a quick, simple, highly effective and convenient method of permanent sterilization, vasectomy has been identified as one of the indicators of male involvement in family planning. However, the application of vasectomy has been limited due to fear of surgery and postoperative complications, such as pain, infection, sperm granuloma, epididymal and testicular damage, and its limited reversibility. Since male hormonal contraception may lead to other long-term health issues in men, it is not a preferred approach, resulting in limited experimental research and restricted clinical applications (24). A quarter of the world's adult men are willing to share responsibility for contraception (30). Therefore, it is necessary to develop more effective and reversible methods for males in order to fulfill such greater participation by men in sharing the burdens and benefits of effective family planning. In recent years, as a potential male contraceptive method, TTH has been carried out in a large number of animal experiments and human trials, such as mice (14, 28), rats (18), rabbits (30, 31), sheep or goats (32), dogs (33), monkeys (5, 34) and humans (7, 20, 21, 35). Each animal is anesthetized, and the lower third of the body (hind legs, tail and scrotum) is submerged in a water bath at 40°C or 43°C for 30 min (36-38). It is the most precise way to expose testes to an increased temperature for a short time by immersing the scrotum in a water bath at the desired temperature. This technique has been used in rats and monkeys (Figure 2) (18, 34). Local scrotal exposure to 42°C for




Figure 1. Review framework.





Figure 2. TTH for animals and humans. Left, water bath for monkeys. Right, designed sketch of scrotal heat stress device (heating underpants). No. 1 and 7 show the fixed belts, No. 2 shows the water bag, No. 3 shows the electric heating pad, No. 4 shows the protective layer of the water bag, No. 5 shows the thermal insulation layer and No. 6 shows the temperature probe which links with a digital display. (a) vertical section map of scrotal heat stress device, (b) cross-section map of scrotal heat stress device. This Figure is modified with permission from (21).

approximately 1 h with a home-made electric blanket is used in boars (39). In rats, scrotal temperature is altered within the range of 13-43°C by a thermode (40). Insulated bag (with layers of nylon, cotton batting, mylar, and then canvas) covers the scrotum

and inguinal region, and the temperature ranges from 34.9 ± 0.3°C to 38.0 ± 0.1°C. In the scrotum of ram, an insulating bag is placed for 288 consecutive hours (32, 41, 42). TTH for men includes water bag and electric heating system (Figure 2). The electric warming bag or



**Figure 3.** Changes in sperm parameters, DNA fragmentation and chromosomes under TTH. In addition to mice, primates such as monkey and humans are used to assess the role of TTH on spermatogenesis and male fertility. TTH leads to changes in sperm quality (e.g., sperm concentration, motility, morphology, and membrane characteristics), sperm DNA (e.g., chromosome structure, caspase activity, and acrosin), and several important biochemical substances in seminal plasma. (A) and (B) Sperm concentration (left) and motility (right) before, during and after scrotal heating. This Figure is modified with permission from (22). (C) DFI from SCSA under TTH. (D) Sperm FISH assay. (a) and (b), sperm chromosome X centromere (green), chromosome Y centromere (red) and chromosome 18 centromere (blue) specific probes. (a) Before TTH, spermatozoa with one chromosome Y (3 spermatozoa) or one chromosome X (one spermatozoon) and one chromosome 18. (b) during TTH 3 months, a spermatozoon with 2 chromosomes Y and one chromosome 18 (the right). (c) and (d), sperm chromosome 13 centromere (green) and chromosome 21 centromere (red) specific probes. (c) before TTH, spermatozoa with one chromosome 13 and one chromosome 21. (d) during TTH 3 months, a spermatozoon with one chromosome 13 and one chromosome 21 (the top) and a spermatozoon with two chromosomes 13 (the bottom left) and a spermatozoon with one chromosome 21.

water bag can be attached to the underpants, and the temperature is controlled (20-22). For water bath, the scrotum is immersed in a water bathtub at 43°C for 10 times (30 min each time) (20-22, 35).

#### 4. EVIDENCE OF ALTERATION OF SPERMATOGENESIS AND SPERM PARAMETERS UNDER TTH

TTH can affect spermatogenesis and reduce semen quality. A large number of experiments on TTH have reported that local testicular heat treatment in water at 43°C can induce reversible oligospermia or azoospermia in rodents and monkeys (5, 18, 34, 43-45). Short-term exposure of the testes to heat results in selective, but reversible damage to the seminiferous

epithelium (18, 46). TTH may damage spermatogenesis and semen quality, leading to declined fertility of many animal models (18, 44-49). Exposure of the testes to TTH, cryptorchidism or varicocele can increase the death of germ cells. Accumulated evidence indicates that a single exposure of the rat (43 °C for 15 min) or monkey (43 °C for 30 min) testes to heat can result in selective and reversible damage to morphology, motility and concentration of sperm (43, 47, 50). When adult male mice are exposed to whole-body heat of 37-38°C for 8 h per day for 3 consecutive days, the sperm motility is reduced, and damage in sperm membrane is increased (51). In boars, a single heat exposure results in the decreased number of spermatozoa as well as the impaired motility and reduced proportion of normal spermatozoa (50, 52, 53). The motility and

proportion of normal spermatozoa may be recovered in approximately 6 weeks after TTH in boars (53, 54, 55) and 4-8 weeks in human males (7, 20-22, 35).

TTH consequently changes the concentration, motility, morphology and membrane of sperm (Figure 3). The sperm count and motility can be affected by a sauna exposure from 37 to 45 °C (19). Ahmad *et al.* (56) have reported that TTH reduces the proportion of motile and viable spermatozoa and total sperm count of men. Oligozoospermic men with a varicocele have significantly higher scrotal temperatures than normozoospermic men (15). Galil *et al.* (57) have found that exposure of one or both testes of rats to heat at 43°C for 30 min results in a significant reduction in blood flow per testis. Testicular blood flow decreases along with testicular weight loss. Parameters of rabbit spermatozoa, such as metabolic activity and motility, are largely modified after a short exposure of spermatozoa to an environment of 42°C for only 3 h (31). Experimental TTH has also been demonstrated to severely suppress spermatogenesis in several studies on rats (58), mice (59), monkeys (60) and humans (7, 20-22, 35). Therefore, TTH may be used as the potential target of contraceptive development for men (10, 12, 61-63). Spermatogenesis and parameters of sperm concentration, morphology and motility in semen samples are adversely affected by TTH. In human, the minimum sperm concentration and total sperm count can be achieved during 6 or 8 weeks of TTH, and a good reversibility is observed in the recovery period (20, 35).

### 5. EFFECTS OF TTH ON THE TESTIS, HISTOLOGY AND GERM CELLS

Testicular thermoregulation is important. Shortly after heat exposure, a loss in testicular weight occurs. This reduction in testicular weight can be ascribed to loss of germ cells. Although the testicular weight may be partially regained several weeks after heat exposure in the rats, the testis remains lighter than before heat exposure (64). Spermatogenesis is temperature-sensitive, and an increase of 1°C entails a 14% drop in spermatogenesis, consequently leading to poorer sperm production (65). However, elevation of testicular temperature causes testicular germinal atrophy and spermatogenic arrest in male subfertility (66). Exposure of testes to TTH can cause severe damage in the seminiferous tubules. Selective and reversible damage to the seminiferous epithelium has been found in mouse (10), rat (12), monkey (34, 65) and human testes (20-23) after exposure to 43°C. Liu *et al.* (14) have reported that after a 42°C heat shock for 30 min in mice, acidophilic, coagulated germ cells are found in the seminiferous epithelium, most spermatocytes are affected, and full of abnormal cells appear in the corpus and caput of the epididymis. Degeneration, vacuolation, disorganization,

multinucleated giant cells, destructive spermatocytes and spermatids, and hyperplasia of Leydig cells are observed in a large proportion of seminiferous tubules in the atrophic testes in mice (67). In addition, 7 days after an exposure to a 43°C water bath for 20 min or after a 2-h 42°C heat treatment for 8 days, low testicular weight and severe vacuolization of seminiferous tubules along with loss of spermatogenic cells are observed in mice, and appearance of multinucleated giant cells can be found (58, 68-70). Hamilton *et al.* (42) have reported that in rams treated with insulating bag, initial testicular degeneration as well as mild and multifocal vacuolization of SCs are observed, exhibiting a small to moderate proportion of desquamated cells containing three or more nuclei (giant cells) inside the seminiferous tubules, desquamation of germ cells and mild multifocal interstitial fibrosis. In rhesus monkeys, after TTH at 43°C for 30 min daily for 6 consecutive days, the disrupted structure of seminiferous epithelium is found on day 8 after hyperthermia with some germ cells sloughing from SCs, and most germ cells are lost from the seminiferous epithelium at days 15 and 30 after treatment (68, 71). The structure of seminiferous epithelium starts to recover between days 45 and 75, and complete recovery is found on day 90 (71). In addition, spermatogenesis and seminiferous tubule diameter are decreased after the testosterone undecanoate plus heating in fertile males (7). Among the germ cells, the types that are most vulnerable to heat are the pachytene and diplotene spermatocytes as well as the early round spermatids in humans (72). Therefore, the effects of TTH on testicular weight, seminiferous epithelium, germ cells and spermatogenesis are evident. However, these alterations can be recovered after termination of TTH. Liu *et al.* (14) have reported that after 1-week recovery period of TTH, new spermatocytes can be found in injured tubules which do not contain spermatid cells, and spermatogenesis and the morphology of all seminiferous epithelium can be completely recovered at 6 weeks after termination of TTH.

### 6. SPERM DNA DAMAGE, OXIDATIVE STRESS AND APOPTOSIS IN TTH TREATMENT PHASE

#### 6.1. Sperm DNA damage, oxidative stress and reactive oxygen species (ROS) after TTH treatment

The integrity of sperm and oocyte DNA plays a fundamental role in the development and quality of embryos. Spermatozoa deliver DNA to the oocyte at fertilization. Sperm DNA fragmentation (sDF) is the most common DNA abnormality presenting in the infertile males (71) and TTH treatment procedures (11-13, 20-23). Single and double DNA strand breaks in the sperm nucleus are the main types of sDF. Indeed, there is evidence that sperm DNA breaks may

originate in the testis and epididymis during transit in the ejaculatory ducts, and following ejaculation. Pathways of DNA breaks may be the beginning of sperm apoptosis, leading to endonuclease activation and a direct attack to DNA by free radicals which produce both base oxidation and strand breaks. The apoptotic process mainly occurs during spermatogenesis by affecting the testicular function and derailment of the chromatin condensation. The germ cell damage by TTH includes sperm apoptosis (8, 17, 18), DNA damage (73, 74) and generation of ROS (75-77). Under oxidative stress, ROS can cause severe damage to cellular components and increase sDF. TTH has been associated with an increase in testicular oxidative stress (35). Several studies have reported the mechanisms underlying the heat-induced alteration of semen parameters and loss of germ cell DNA integrity (7, 8, 11-13, 23). TTH to temperatures higher than the normal physiological temperature can change the structure of sperm chromatin, resulting in the disruption of spermatogenesis (78, 79). In several rodent models, testicular heat stress, including TTH and surgical induction of cryptorchidism, results in reduced testicular weights, germ cell loss and increased rate of sperm apoptosis (12, 80-82). Oxidative stress directly or indirectly causes sDF. Excessive levels of ROS can damage normal spermatozoa by inducing lipid peroxidation and DNA damage. Some studies have found a clear relationship between heat and oxidative stress (83, 84). Other studies on rams have reported a reduced integrity of sperm DNA under heat stress conditions induced by testicular insulation (32, 42). Oxidative stress is described as the main cause of sDF and may be related to protamine deficiency and sperm apoptosis (85). As a major cause of impaired spermatogenesis, TTH is greatly associated with oxidative stress and germ cell apoptosis (86). Early studies of TTH have found that DNA fragmentation remains high in cryptorchid mouse testis, suggesting that male infertility is likely caused by high sDF (17, 87). An abnormal increase in ROS levels can overwhelm local antioxidants and lead to oxidative stress (11, 88). Kanter *et al.* (38) and Park *et al.* (68) have reported a significantly decreased expression of proliferating cell nuclear antigen and an elevated activity of terminal deoxynucleotidyl transferase dUTP nick end labeling (TUNEL) (TUNEL-positive apoptotic cells) after TTH in rats and mice. Another study has indicated that TTH can cause azoospermia and oligozoospermia in monkeys by increasing the germ cell apoptosis (44). A previous study has reported elevated sperm mitochondrial ROS generation, increased sperm membrane fluidity and DNA damage, and an activated oxidative stress cascade in adult male mice exposed to an elevated ambient temperature of 35°C for 24 h (88). A slight increase in scrotal temperature caused by occupational exposure, lifestyle or cryptorchidism

can disrupt spermatogenesis and ultimately cause problems with fertility (11, 19).

### 6.2. Effect of TTH on sperm chromosome and acrosin activity

TTH may contribute to sperm chromosomal changes by reducing antioxidant power and increasing DNA strand breaks. The study by Sailer *et al.* (78) has suggested that scrotal exposure to 40°C in mice decreases the number of testicular haploid cells but increases the diploid germ cells, and some thermal ranges above normal allow production of sperm with compromised nuclear chromatin structure. Heat shock to spermatozoa can distort the sex chromosome (X and Y) ratio (78). The proportion of sperm aneuploidy is correlated with that of apoptotic sperm and high frequency of abnormal sperm morphology (89-91). TTH can affect sperm chromosome (22) (Figure 3) and increase in the incidence of X and Y univalents at metaphase I (92). Chao *et al.* (93) have reported that when mouse sperms are heated in a water bath at 50-95°C for 30 min, the aberrant chromosome rate is increased from 16.3% to 100%. TTH at 42°C results in the reduction of pregnancy rate, decreased placental weight and litter size, and abnormal embryonic development (10).

TTH can cause a rapid and transient suppression of spermatogenesis, and damage to the sperm DNA and acrosin. As an acrosomal protease, acrosin is synthesized as a pro-enzyme and activated into beta-acrosin during acrosin release. TTH may decrease acrosin activity in sperm (22, 35). Oxidative stress can be deleterious for DNA fragmentation and acrosin release. Acrosin plays a key role in the fertilizing capacity of human spermatozoa, and its activity directly affects the fertilization rate (94, 95). Several human studies have found that poor semen quality of infertile patients is related to a high scrotal temperature, and thermoregulatory failure-triggered heat stress can damage acrosin and increase the risk of infertility (96-98). Acrosin may act as a matrix-degrading proteinase and play an important role in sperm penetration of the zona pellucida (99, 100). In many cell types, hypoxia and oxidative stress trigger apoptosis and cell death. Heat stress may induce oxidative damage to sperm plasma membrane, sperm parameters, germ cells, the cell division of SCs, and acrosin activity (20, 101-103).

### 6.3. Effect of TTH on caspase

Hyperthermia has detrimental effects on the testis, including DNA damage in germ cells and mature sperms, poor-quality spermatozoa, and a lengthy recovery period of the testis from TTH (10, 12). Some studies have provided evidence that the mitochondria-

dependent apoptotic pathway is involved in the activation of the initiator caspase-9 and executioner caspase-3 in heat-induced germ cell death (51, 104-106). During apoptosis, caspase-3 cleaves the inhibitor, thereby allowing the nuclease to cut the chromatin. Caspase-3 is activated by cleavage of its proenzyme (32 kDa) to yield two active subunits p18 and p12, and the activation of caspase-3 is used as an important index of apoptosis (104, 105). In fertile males, during TTH treatment, the sperm motility, number and normal morphology are decreased, whereas the rates of sDF, abnormal chromatin condensation and the caspase-3 activity are increased, and these indicators are restored to the normal levels before the TTH in recovery period (20-22). The testis displays severe damage, and the number of cleaved caspase-3-positive germ cells per tubule is dramatically increased when mice are exposed to a single TTH treatment (42°C for 25 min) (10, 105). Wang *et al.* (105) have reported that after TTH (40°C for 2 h), the enzyme activities of caspase 3, 8 and 9 in newt testis are significantly elevated. In non-apoptotic cells, caspase-activated deoxyribonuclease is present as an inactive complex with inhibitor of caspase-activated deoxyribonuclease. Following a single mild TTH exposure (40°C or 42°C for 30 min), an increased expression of cleaved caspase-3 and a decreased expression of inhibitor of caspase-activated deoxyribonuclease are found in mouse testis. The reduced expression of inhibitor of caspase-activated deoxyribonuclease contributes to the increased activity of caspase-activated deoxyribonuclease, and this finding is consistent with the increased rate of DNA fragmentation (19). When 8-week-old mice are exposed to a single TTH (42°C for 25 min), the testes display severe damage with multinucleated giant cells, and the number of cleaved caspase-3-positive germ cells per tubule is dramatically increased (10, 12). After TTH (42 °C, 30 min) in male Balb/c mice, the number of TUNEL-positive cells (apoptotic index of spermatogenic cells) and expressions of caspase 3, 8, 9 are increased, whereas the expression of Bcl-2 AKT at the mRNA level is decreased (105). When adult mice are exposed to 37-38°C for 8 h, considerably more caspase-3-positive germ cells are found in heat-treated mice at 16 h after heat exposure compared with the controls left at 23-24°C (51). In mice treated with TTH at 43 °C for five cycles, the levels of cleaved caspase-3 and TUNEL-positive apoptotic spermatocytes are increased (12, 106, 107). However, histological examination of excised testes and epididymides for apoptotic (TUNEL and activated caspase-3) and proliferating cells (Ki-67 antigen) indicates only marginal effects of scrotal insulation on tissue morphology in adult beagle dogs (insulated with a self-made scrotal support for 48 h, 9 weeks later). It suggests that a mild TTH in dogs does not cause substantial changes in sperm quantity and quality, and canine testes and epididymides may have a higher competence to compensate such thermal stress (33).

## 7. EFFECTS OF TTH ON SPERM PROTEIN EXPRESSION, AUTOPHAGY AND MOLECULAR CHANGES

Previous studies have reported the changes in sperm parameters (Figure 3), sperm apoptosis, and molecular mechanisms of TTH exposure (18, 31, 34, 108, 109). The protein expression and autophagy of heat-induced spermatogenic impairment play important and complex roles in maintaining a series of cellular functions. Within the testis, spermatogenesis is a strictly regulated, protracted and complex process, and it must, in part, be dictated by networks of many genes (110). These processes are particularly vulnerable to perturbation by genetic and environmental factors.

### 7.1. Effect of TTH on sperm protein expression

During the sperm development, correct timing of gene expression plays a fundamental role within biological processes. Gene expression during spermatogenesis is highly orchestrated and strictly regulated at the transcriptional and posttranscriptional levels, and the basis of male germ cell differentiation and male fertility depends on the correct gene expression patterns (111). Serial analysis of gene expression and microarray profiling of rodent testis samples and enriched cell populations have identified a large number of important genes for spermatogenesis (111-113). Spermatogenesis related factor-2 is one of the cloned genes involved in division and differentiation of germ cells (112). The occludin expression at the protein and mRNA levels is decreased in boars under local scrotal exposure to 42°C (39). Chihara *et al.* (109) have found that heat-induced testicular damage through the MRL/MpJ-derived locus on Chr 1 may play a pivotal role in the recovery from heat-induced testicular damage, especially via the inhibition of calcification. The microRNAs (miRNAs) are small, non-coding RNAs with approximately 17-25 nucleotides in length, which act as potent posttranscriptional regulators of gene expression, and they are involved in a variety of biological processes, including cell proliferation, apoptosis and tumorigenesis. Recent studies have reported the role of miRNAs in heat stress-induced spermatogenic impairment. TTH at 43 °C for 30 min down-regulates the relative expression of cold-inducible RNA binding protein at the mRNA level but up-regulates the expressions of transforming growth factor  $\beta$ 2 and transforming growth factor  $\beta$ 3 in mice (114), and some identified miRNAs, including miR-449a-3p, miR-92a-1-5p, miR-423-3p and miR-128-3p, have been closely correlated with apoptosis of germ cells (115). Heat shock proteins (HSPs) can protect cells from harmful stress and regulate cell apoptosis (8). Most proteins in spermatozoa are formed before spermatid DNA condensation, and mature spermatozoa exhibit almost no protein synthesis. Therefore, the alteration of protein levels in spermatozoa can largely reflect

the effect of hyperthermia on spermatogenesis, not only on spermatozoa itself. These proteins modulate the correct folding of stress-induced misfolded proteins and protect germ cells and spermatozoa from apoptosis (8, 19). A sufficient increase in blood flow may not support an increased metabolism in the testes after heat stress, and the germ cells may be under relatively hypoxic conditions (12). Zhang *et al.* (73) have demonstrated that besides initiating apoptotic pathways, heat also induces autophagic pathways in germ cells. The expression of autophagy-related gene 7 is positively correlated with the level of autophagy in germ cells. TTH induces lipid droplet accumulation and up-regulates the expression of adipose differentiation-related protein (14). The expressions of Bax and Bcl-2 at the protein and mRNA levels can be increased after local testicular heating (42°C, 1 h) in boar testis (116). TTH can cause the apoptosis of germ cells by regulating the expressions of Bax and Bcl-2 at the protein level. In the mouse SC line, the expression of RNA binding motif protein 3 is increased or decreased within 12 h after temperature shift from 37 °C to 32 °C or 39 °C, respectively (117).

### 7.2. Autophagy and apoptosis of sperm or germ cells in TTH treatment phase

As a survival, lysosomal degradation and tightly regulated pathway for the clearance and recycling of damaged cytoplasmic organelles or cellular components, autophagy plays fundamental roles in cellular homeostasis (118-122). ROS can induce various cellular responses, including apoptosis, programmed necrosis and autophagy, depending on the cellular setting. During the response to sublethal stress and autophagy, cells undergo rapid changes to adapt their metabolism and protect themselves against potential damage. This is orchestrated through a multifaceted cellular program and evolutionarily conserved physiological process, and it can serve as a cell survival pathway, suppressing apoptosis (120-123). Autophagy has been shown to engage in a complex interplay with apoptosis and to promote cell survival because nucleotides, amino acids and free fatty acids can be generated during the degradation of cellular contents and then recycled and reused for macromolecular synthesis and ATP generation (73, 123). In the testis, autophagic pathway can be stimulated by multiple forms of cellular stress, including nutrient or growth factor deprivation, hypoxia, ROS, DNA damage, protein aggregates, chemical insults, withdrawal of gonadotropin or testosterone, heat, radiation, damaged organelles, or intracellular pathogens (69, 73, 123). Hyperthermia causes germ cell death by oxidative stress in testes, triggering antioxidant signals, including autophagy and nuclear factor erythroid 2-related factor 2. Under a 2-h heat treatment at 42°C for 8 days, heat exposure increases the expression of erythroid 2-related factor 2 at the

protein and mRNA levels in male mice, and the testes may be protected by the autophagy inducer (69). Varicocele is in chronic state of heat stress and can lead to impaired spermatogenesis by apoptotic and autophagic pathways. Moreover, apoptotic markers (active caspases 3/7 and DNA fragmentation) and autophagic markers (autophagy-related gene 7 and light chain 3 proteins) are significantly up-regulated in infertile men with varicocele compared with fertile individuals (124). In heat-treated (water bath at 42°C for 15 min) mice, besides initiating apoptotic pathways, autophagic pathways are also induced in germ cells, and TTH results in several specific features of the autophagic process, including autophagosome formation and the conversion of light chain 3-I to light chain 3-II (73). Recently, Chen *et al.* (125) have reported that supplementation of methylmercury may induce germ cell apoptosis of rats through oxidative stress and autophagy. Horibe *et al.* (119) have observed high levels of autophagy in stages VII–VIII of the spermatogenic cycle, and autophagy is specifically up-regulated in SCs of adult rats under stress of ethanol treatment. The up-regulation of autophagy is a specific stage, and it may be related to androgen receptor suppression, mitochondrial damage, lipid accumulation, and phagocytosis of apoptotic cells. Autophagy and endocytosis can efficiently transport transcription factors, adhesion molecules, or secreted factors. Mitochondrial-ER contact sites are important in autophagosome formation and have been recently recognized as a major mechanism in the regulation of self-renewal and differentiation of stem cells as well as a regulator of cell migration (119). Scrotal insulation in boars can damage meiotic germ cells due to changes in the integrity of the blood-testis barrier (BTB), which induce apoptosis, autophagy and DNA damage in the germ cells (126).

### 7.3. Molecular changes in TTH

Zhu *et al.* have reported the cellular and molecular mechanisms of TTH in murine testis (127), showing that 108 proteins are differentially expressed between the heat shock tissues and the control mouse testes, and 36 proteins are identified by comparing with the control reference map. Some of the proteins may be early molecular targets responsible for suppression of spermatogenesis induced by TTH in rhesus monkeys and humans (127). Another study has also investigated the molecular mechanisms underlying the heat-induced suppression of spermatogenesis in human testis, and found 32 and 26 differentially expressed proteins at week 2 and week 9 after TTH treatment, respectively (128). Heterogeneous nuclear ribonucleoprotein H1 is an anti-apoptotic protein that can regulate the expressions of other heat-induced proteins (128). After TTH, mice exhibit an increased expression of hypoxia-inducible factor 1 alpha (Hif1 $\alpha$ ) at the mRNA level and a translocation



of Hif1 $\alpha$  protein to the germ cell nucleus (12). TTH alters the expression levels of various genes in a complex manner, accompanied with post-translational modification and protein localization, and altered gene expressions will lead to changes in the protein composition of spermatocytes. However, another study has found an increased Hif1 $\alpha$  expression at the mRNA level in spermatozoa after sauna exposure (19). Hif1 $\alpha$  is a transcription factor, its target gene is involved in vasodilation, angiogenesis and glycolysis, and it can induce the expression of vascular endothelial growth factor. Hikim *et al.* (129) have found that the initiation of apoptosis in wild-type mice is preceded by a redistribution of Bax from a cytoplasmic to paranuclear localization in heat-susceptible germ cells. The relocation of Bax is accompanied by sequestration of ultra-condensed mitochondria into paranuclear areas of apoptotic germ cells, and cytosolic translocation of mitochondrial cytochrome C and Bax is associated with activation of the initiator caspase-9 and the executioner caspase-3 (129).

Sertoli-Sertoli and Sertoli-germ cells are the specialized junctions at contact sites of seminiferous epithelium, and they play pivotal roles in spermatogenesis. Cai *et al.* (102) have reported that after a TSH (43°C for 30 min) treatment for 24-48 h, the expressions of tight junction components of occludin, claudin-3 and zonula occludens-1 (ZO-1) are decreased, whereas the claudin-11 expression at the mRNA level is increased, and the protein localization of occludin and ZO-1 is lost from the BTB site. A transient induction of transforming growth factor- $\beta$ 2, -3 and p38 mitogen-activated protein kinase activation may lead to the reversible BTB disruption by TTH. In humans, HSPs (such as HSP70 and HSF1) and mitochondria functions are up-regulated, and Bcl-2, mitofusin-2 and spermatozoa-associated antigen 6 are down-regulated in TTH treatment (130). Chen *et al.* (34) have reported that the expression of androgen receptor in primary monkey SCs is dramatically decreased during TTH. The expressions of adherens junction-associated molecules of N-cadherin and beta-catenin, and tight junction-associated molecule ZO-1 are decreased in 24-48 h by TTH treatment, and the intermediate filament vimentin is up-regulated in 6-48 h. The expressions of androgen receptor and Wilms' tumor gene 1 are dramatically decreased during TTH. Both proteins completely disappear immediately after termination of TTH treatment and begin to recover after 6 h (34). The androgen receptor plays a crucial role in the heat-induced reversible change in BTB via the partitioning-defective protein polarity complex, which may be the mechanism of TTH underlying a reversible disruption of the BTB (59). When the androgen receptor is over-expressed in SCs using an adenovirus, the heat stress-induced down-regulation of BTB-associated proteins of ZO-

1, N-cadherin, E-cadherin,  $\alpha$ -catenin and  $\beta$ -catenin is partially rescued. Ablation of androgen receptor by RNAi or flutamide treatment (an androgen receptor antagonist) in SCs can inhibit the recovery of BTB-associated protein expression after TTH treatment of 43°C. The recovery of BTB permeability induced by temporal heat stress may be regulated by the androgen receptor in mice (59). TTH can change the expressions of some protein components in the spliceosome and result in a reversible dysfunction of RNA splicing in its target pre-mRNAs in animal and human males.

As stress-responsive proteins, HSPs are the most prominent group of proteins involved in folding and unfolding of other proteins. The most important event in stressed cells is the synthesis of a highly conserved family of proteins, such as HSPs. HSPs play an important role in the homeostasis, apoptosis regulation and the maintenance of various physiological processes. HSPs are considered to be essential factors in animal reproduction and required for spermatogenesis, and they also protect cells from environmental hazards, such as heat, radiation, and chemicals (131, 132). The expressions of HSPs are regulated by heat shock transcription factors (HSFs) for their cytoprotective functions during cellular stress. Proper spermatogenesis can be maintained by cooperation of different HSFs. For example, knockout cooperation of HSF1 and HSF2 results in meiosis arrest and spermatocyte apoptosis (131, 132). Elevated testicular temperatures can induce activation of HSF1. Activation of HSF1 by heat shock initiates apoptosis of spermatogenic cells, leading to infertility of males. In the male germ cells (spermatocytes), caspase-3-dependent apoptosis is induced upon HSF1 activation, and spermatogenic cells are actively eliminated (133, 134). In addition to the up-regulation of HSF1, which in turn increases HSP production, other up-regulated genes include those responsible for apoptosis, cell adhesion and signal transduction during TTH. Rockett *et al.* (135) have used DNA microarrays to interrogate the expressions of 2, 208 genes and thousands more expressed sequence tags (ESTs) in mouse testis. Of these genes, 27 are up-regulated and 151 are down-regulated after TTH (135). TTH can reduce the expressions of antioxidant-related enzymes, spermatogenesis-related proteins and sex hormone receptors (androgen receptor, luteinizing hormone receptor and follicle-stimulating hormone receptor) in rats (136). Heat increases oxidative stress, affects mitochondria and results in DNA damage, eventually leading to spermatozoal apoptosis. The expressions of steroidogenic acute regulatory protein and HspB10 genes are decreased by long-term heat stress via the expressions of steroidogenic pathway and small HSPs in rat testis at 38°C for 9 weeks (137). TTH may adversely induce effects on sperm parameters and gene expression. TTH has effects on expressions of Bax,

Bcl-2 and sperm parameters in male mice. Bcl-2 family members may induce sperm apoptosis under heat stress (138). Fas/FasL is another important cell death pathway. When Fas/FasL is activated and caspase-3 is cleaved, the cell apoptotic rate is decreased (139).

### 8. EFFECT OF TTH ON EPIDIDYMIS AND SEMINAL BIOCHEMISTRY

Spermatozoa are produced in the testis, which acquire the maturation ability in the epididymis. The caput and corpus of the epididymis as well as the epididymal fluids play a critical role in sperm maturation (140, 141). During the epididymal transit, spermatozoa undergo critical changes in structure and function. Some of biochemical substances and proteins in epididymal fluid, such as fructose, NAG, L-carnitine, MIF, EGF and dipeptidyl HSP, are the key factors for maturation and function of epididymal sperm (22, 23, 142-146).

#### 8.1. Effect of TTH on epididymal proteins

Numerous studies have investigated the effect of TTH on spermatogenesis and sperm maturation (143, 144, 146). In epididymis, strong staining signal of HSP10 is detected in pseudostratified columnar epithelium (144). A total of 270 proteins may participate in epididymal sperm maturation, and 34 epididymal milieu proteins contribute to the composition of seminal fluid proteomes (146). Wang *et al.* (143) have identified the key heat stress-associated sperm maturation proteins in the rat caput epididymal fluids responsible for sperm maturation, and demonstrated 21 proteins corresponding to 29 differential protein spots, including 10 down-regulated and 11 up-regulated proteins in heat-treated rats. In ram epididymis, 17 seminal plasma proteins are affected by treatment, actin, albumin, HSP 70 kDa, protein DJ-1 and so on in seminal plasma are up-regulated, and some of proteins, such as dipeptidyl peptidase 3 and isoforms of HSP 90 kDa, are down-regulated after scrotal insulation. Expressions of most such proteins return to normal levels after scrotal insulation for 113 days (146). Changes in seminal plasma proteome by heat challenge may be correlated with variations in semen parameters, such as sperm protection, maturation and fertilization.

#### 8.2. Effect of TTH on epididymal sperm

TSH can affect epididymal spermatozoa. The TTH-treated epididymal sperm has lower sperm-zona pellucida binding and oocyte penetrating capacity. These effects are first seen at 1 week, and then they become more prominent at 2 weeks after heat exposure (147). In another study, mice exposed to whole-body heat show lower motility and membrane changes of sperm obtained from the cauda epididymis (148). In a previous study, when TTH is applied to male

mice at 42°C for 30 min, the sperm count, motility and viability are reduced, and a sex ratio distortion occurs in epididymal spermatozoa (149).

#### 8.3. Effect of TTH on NO and NOS

When the TTH causes sperm apoptosis and changes in the testis, the epididymal microenvironment is also destroyed in a short time, thereby altering the contents of semen biochemical substances (20). ROS are metabolites of NO, playing a role in many physiological functions of sperm. NO is a free radical produced by most cells, including human male and female reproductive tracts, and it is an important regulator in many physiologic processes and a well-known oxidative stress agent that directly inhibits mitochondrial respiration and DNA synthesis. Shiraishi and Naito (150) and Carreira *et al.* (151) have both reported that NO is synthesized through the enzymatic conversion of L-arginine to L-citrulline by NOS. Many previous studies have shown that mature human spermatozoa synthesize NO by up-regulating NOS which exists in at least three distinct isoforms, neural NOS (nNOS), endothelial NOS (eNOS) and inducible NOS (iNOS). Three distinct isoforms of NOS are localized in the sperm head and midpiece (152, 153). Cellular injury and apoptosis occur in various cell systems through the up-regulation of iNOS to increase NO synthesis (154-157). It has been reported that various physiologic activities via different signaling pathways, including the cyclic guanosine monophosphate (cGMP)-protein kinase G and mitogen-activated protein kinase pathway, can be regulated by lower concentrations of NOS (158, 159). At low concentrations, NO stimulates guanylate cyclase activity and triggers the formation of cyclic guanosine monophosphate. Higher concentrations of NO produced by iNOS interact with thiol groups (158). At higher concentrations of NOS, the molecule can promote the germ cell apoptosis through peroxynitrite, which is generated by a reaction between NO and superoxide (160). High concentrations of NO have injurious effects on sperm properties, such as motility, morphology, and DNA stability, which can increase apoptosis and block all sperm functions (161-163). Intracellular NO, a free radical, is a factor that physiologically regulates spermatozoa function and pathophysiology of male reproductive system (164, 165). Previous study has well documented the role of NO in multiple signal transduction pathways of male germ cells, such as capacitation and acrosome reaction (AR), and activation of apoptosis signaling is negatively correlated with NO production (166). The signaling for iNOS expression may be similar to that induced by cellular injury and apoptosis. This concept is supported by the evidence that a significant loss of germ cells is caused by up-regulation of testicular iNOS and the seminiferous epithelium damage after treatment with endotoxin lipopolysaccharides

(167-169). iNOS may play a functional role in spermatogenesis via apoptosis, leading to reduced sperm count in monkeys (170). Low concentrations of NO improve the motility of mouse (171), hamster (172) and human (173, 174) spermatozoa, as well as the zona pellucida-binding ability of human spermatozoa. Maturity of human spermatozoa has been correlated with active NO production by NOS and inactivated apoptosis signaling, indicating a rather anti-apoptotic effect of NO. NO may be enough to affect the fecundity potential *in vivo* (171, 175). A previous study by our group (21) has shown that the levels of NO and NOS in semen are negatively correlated with sperm concentration, motile and progressive motile sperm and normal morphological sperm, but positively correlated with sDF and caspase-3 activity during TTH in human.

### 8.4. Effect of TTH on MIF

NO, NOS and MIF are important regulators in many physiologic processes (21, 22). MIF is a well-known pro-inflammatory mediator with important functions in human reproduction and prostatic physiology. As a T-cell cytokine, MIF is present in large quantities in human semen and it is transferred to spermatozoa during the epididymal transit, which has been correlated with sperm maturation and stability (176-178). A higher level of MIF is found in epididymis and associated with spermatozoa, prostasomes as well as the soluble fraction (179). Eickhoff *et al.* (180) have suggested that MIF plays an important role in the maturation process of rat sperm during epididymal transit. Some studies have indicated that MIF is secreted by the Leydig cells of the testis, and it is a constituent of seminal fluid (177, 181, 182). Interestingly, the expression of sperm-associated MIF has been negatively correlated with the proportion of motility in different semen samples (183). Komori *et al.* (184) have reported that MIF in seminal plasma is higher in patients with abnormal AR. Therefore, MIF in seminal plasma may negatively affect sperm function. Our group has reported (21, 22) that the MIF level is negatively correlated with conventional sperm parameters and NAG in seminal plasma but positively correlated with sDF index.

### 8.5. Changes in seminal NAG, L-carnitine and EGF

Apart from proteins, seminal plasma also contains several small molecular substances, which are shown to play active role in sperm maturation/protection processes. Seminal NAG, L-carnitine and fructose are regarded as the reliable biochemical parameters to examine epididymal patency, ejaculatory duct obstruction, and prostate and seminal vesicle function or its hypoplasia (184-188). They are used for the diagnosis of sperm function and obstructive azoospermia.

Fructose in seminal plasma is synthesized and secreted by the seminal vesicles and accessory sex glands. It is the major carbohydrate source in seminal plasma and essential for normal sperm motility (189, 190). Rao *et al.* (35) have reported that fructose and zinc concentrations in TTH procedures show no significant differences compared with their baseline levels. Such finding may be explained by that during TTH, only the scrotum is immersed into the warm water, therefore the seminal vesicles are not affected by TTH.

Epididymis is the main source of NAG in semen. The neutral form of alpha-glucosidase is the main active ingredient. NAG is correlated with the location of epididymal obstruction in men with azoospermia, which helps to locate obstructive positions (22, 130, 186, 191). The NAG level remains very low in seminal plasma of epididymal duct obstruction or ejaculatory duct occlusion. NAG is considered as the best index of epididymal function, and it is correlated with human sperm concentration and motility (184, 192). In infertile oligoasthenozoospermic men with varicocele, the seminal alpha-1, 4-glucosidase activity is correlated with sperm count, sperm motility and serum testosterone (193). Varicocele causes a reduction in NAG activity by the epididymis, and the NAG level is correlated with the quality of the sperm membrane and nucleus (194). Watanabe *et al.* (195) have suggested that sDF index is correlated with neutral NAG activity, and the determination of NAG activity is a valuable tool in the diagnosis and would also aid in the prognosis.

Spermatozoa undergo post-gonadal modifications in the epididymis to acquire fertilizing ability. In epididymal plasma, high-molecular-weight proteins and small molecules, such as free-L-carnitine, convert the gametes into functional cells (196). As a small polar molecule present in high concentrations in mammalian seminal plasma, L-carnitine plays roles (only L-isomer) in acyl chain translocation and regulation of  $\beta$ -oxidation of fatty acid chains in mitochondria (197). More than 94% of the free carnitine originates from the epididymis, and free carnitine does not originate from seminal vesicles, but rather flows into the seminal fluid via the ductus deferens (198). The L-carnitine level in seminal plasma plays an essential role in maintaining male fertility, and it is correlated with sperm count, total motility and normal morphology, which can be taken as a biochemical index for the clinical treatment of male infertility as well as for the study on the mechanisms of male reproduction (198, 199). L-carnitine has effective superoxide anion radical scavenging, hydrogen peroxide scavenging, total reducing power and metal chelating activities on ferrous ions (200). Carnitine is effective in infertility treatment, and it improves male reproductive capacity in adulthood (201).

Our group (22, 142) has reported that during TTH, the NAG activity and L-carnitine level in the semen are decreased, whereas the levels of MIF and DNA fragmentation are increased. These findings indicate that TTH may affect the testis and epididymis, thereby reducing the seminal quality measures and impairing the quality of the sperm membrane and nucleus. An experimental trial consisting of 20 male volunteers reveals no significant variation in the three biochemical markers, seminal NAG, fructose and zinc concentration, before and after TTH (35). This finding may be attributed to that the functions of epididymis and accessory sex gland are not severely affected by TTH, or the number of the subjects in the clinical trial is small. Seminal plasma zinc, fructose and NAG are the functional indexes of prostate, seminal vesicle and epididymis, respectively. During TTH, prostate and seminal vesicles are farther away from heating bag or warm bath than testes and epididymis. The heat has less influence on the first two organs. Therefore, their functions are less affected.

EGF is a low-molecular-weight polypeptide that is mainly produced by submandibular and parotid glands (142, 202). It is known that EGF affects both male and female reproduction and may play some roles in testicular steroidogenesis (199). Some authors have observed that EGF has either no effect or an inhibitory/negative effect on the human sperm cell capacitation and/or AR, especially at higher concentrations (203, 204). In the semen, EGF is mainly produced by prostate (205, 206). Zhang *et al.* (142) have suggested that the content of seminal EGF is not changed in the TTH process, and it may be because TSH for men is placed to the scrotum and not to the prostate. Therefore, it is necessary to assess changes in EGF in semen upon TTH treatment in future studies.

## 9. CONCLUSIONS

As a male contraceptive method, TTH has potential advantages and feasibility. Obvious changes are observed in testis, epididymis, sperm DNA, chromosomes, mRNA and protein expression, autophagy, oxidative stress and apoptosis of germ cells, and seminal plasma biochemistry after TTH. Sperm concentration, motility, normal morphology and acrosin activity during TTH are decreased, whereas sDF and chromosomal aneuploidy are increased. TTH is correlated with the generation of oxidative stress and ROS. The rate of sperm chromosomal aneuploidy is correlated with sperm concentration, motility, morphology, membrane, and DNA integrity, caspase-3 and acrosin activities. The alteration of sperm DNA, chromosomes, mRNA and protein expression, autophagy, oxidative stress, apoptosis and molecular biology after TTH may be caused by increased levels of NO, NOS and MIF or reduced levels of NAG and L-carnitin in seminal plasma. Given that TTH is a

reversible process, and further studies are needed to design contraceptive methods based on TTH.

## 10. ACKNOWLEDGMENT

Zhen-Ya Fang, Wei Xiao and Su-Ren Chen contributed equally to this work. This study was approved by the national “12th Five-year Plan” of Science and Technology Support (No: 2012BAI31B08). The authors would like to thank Doctor Zhi-Da Shi for their help in English language of this paper.

## 11. REFERENCES

1. B. P. Setchell: Possible physiological bases for contraceptive techniques in the male. *Hum Reprod*, 9(6), 1081-7 (1994)  
DOI: 10.1093/oxfordjournals.humrep.a138637
2. R. Mieusset and L. Bujan: The potential of mild testicular heating as a safe, effective and reversible contraceptive method for men. *Int J Androl*, 17(4), 186-91 (1994)  
DOI: 10.1111/j.1365-2605.1994.tb01241.x
3. S. T. Page, J. K. Amory and W. J. Bremner: Advances in male contraception. *Endocr Rev*, 29(4), 465-93 (2008) DOI: 10.1210/er.2007-0041
4. E. A. McLaughlin and R. J. Aitken: Is there a role for immunocontraception? *Mol Cell Endocrinol*, 335(1), 78-88 (2011)  
DOI: 10.1016/j.mce.2010.04.004
5. Y. X. Liu: Temperature control of spermatogenesis and prospect of male contraception. *Front Biosci (Schol Ed)*, 2, 730-55 (2010)  
DOI: 10.2741/s97
6. A. M. Henken, G. R. Kleinhout and W. van der Hel: The effect of environmental temperature on immune response and metabolism of the young chicken. 1. Effect of intramuscular injection on heat production. *Poult Sci*, 61(3), 503-9 (1982)  
DOI: 10.3382/ps.0610503
7. C. Wang, Y. G. Cui, X. H. Wang, Y. Jia, A. Sinha Hikim, Y. H. Lue, J. S. Tong, L. X. Qian, J. H. Sha, Z. M. Zhou, L. Hull, A. Leung and R. S. Swerdloff: transient scrotal hyperthermia and levonorgestrel enhance testosterone-induced spermatogenesis suppression in men through increased germ cell apoptosis. *J Clin Endocrinol Metab*, 92(8), 3292-304 (2007)  
DOI: 10.1210/jc.2007-0367

8. Y. H. Lue, B. L. Lasley, L. S. Laughlin, R. S. Swerdloff, A. P. Hikim, A. Leung, J. W. Overstreet and C. Wang: Mild testicular hyperthermia induces profound transitional spermatogenic suppression through increased germ cell apoptosis in adult cynomolgus monkeys (*Macaca fascicularis*). *J Androl*, 23(6), 799-805 (2002)  
DOI: 10.1002/j.1939-4640.2002.tb02336.x
9. B. Kim, H. J. Cooke and K. Rhee: DAZL is essential for stress granule formation implicated in germ cell survival upon heat stress. *Development*, 139(3), 568-78 (2012)  
DOI: 10.1242/dev.075846
10. C. Paul, A. A. Murray, N. Spears and P. T. Saunders: A single, mild, transient scrotal heat stress causes DNA damage, subfertility and impairs formation of blastocysts in mice. *Reproduction*, 136(1), 73-84 (2008)  
DOI: 10.1530/REP-08-0036
11. S. Banks, S. A. King, D. S. Irvine and P. T. Saunders: Impact of a mild scrotal heat stress on DNA integrity in murine spermatozoa. *Reproduction*, 129(4), 505-14 (2005)  
DOI: 10.1530/rep.1.00531
12. C. Paul, S. Teng and P. T. Saunders: A single, mild, transient scrotal heat stress causes hypoxia and oxidative stress in mouse testes, which induces germ cell death. *Biol Reprod*, 80(5), 913-9 (2009)  
DOI: 10.1095/biolreprod.108.071779
13. D. Durairajanayagam, A. Agarwal and C. Ong: Causes, effects and molecular mechanisms of testicular heat stress. *Reprod Biomed Online*, 30(1), 14-27 (2015)  
DOI: 10.1016/j.rbmo.2014.09.018
14. M. Liu, L. Qi, Y. Zeng, Y. Yang, Y. Bi, X. Shi, H. Zhu, Z. Zhou and J. Sha: Transient scrotal hyperthermia induces lipid droplet accumulation and reveals a different ADFP expression pattern between the testes and liver in mice. *PLoS One*, 7(10), e45694 (2012)  
DOI: 10.1371/journal.pone.0045694
15. A. Jung and H. C. Schuppe: Influence of genital heat stress on semen quality in humans. *Andrologia*, 39(6), 203-15 (2007)  
DOI: 10.1111/j.1439-0272.2007.00794.x
16. A. Jung, F. Leonhardt, W. B. Schill and H. C. Schuppe: Influence of the type of undertrousers and physical activity on scrotal temperature. *Hum Reprod*, 20(4), 1022-7 (2005)  
DOI: 10.1093/humrep/deh697
17. Y. Yin, K. L. Hawkins, W. C. DeWolf and A. Morgentaler: Heat stress causes testicular germ cell apoptosis in adult mice. *J Androl*, 18(2), 159-65 (1997)  
DOI: 10.1002/j.1939-4640.1997.tb01896.x
18. Y. H. Lue, A. P. Hikim, R. S. Swerdloff, P. Im, K. S. Taing, T. Bui, A. Leung and C. Wang: Single exposure to heat induces stage-specific germ cell apoptosis in rats: role of intratesticular testosterone on stage specificity. *Endocrinology*, 140(4), 1709-17 (1999)  
DOI: 10.1210/endo.140.4.6.629
19. A. Garolla, M. Torino, B. Sartini, I. Cosci, C. Patassini, U. Carraro and C. Foresta: Seminal and molecular evidence that sauna exposure affects human spermatogenesis. *Hum Reprod*, 28(4), 877-85 (2013)  
DOI: 10.1093/humrep/det020
20. M. H. Zhang, Z. D. Shi, J. C. Yu, Y. P. Zhang, L. G. Wang and Y. Qiu: Scrotal heat stress causes sperm chromatin damage and cysteinyl aspartate-specific proteinases 3 changes in fertile men. *J Assist Reprod Genet*, 32(5), 747-55 (2015)  
DOI: 10.1007/s10815-015-0451-0
21. M. H. Zhang, A. D. Zhang, Z. D. Shi, L. G. Wang and Y. Qiu: Changes in levels of seminal nitric oxide synthase, macrophage migration inhibitory factor, sperm DNA integrity and Caspase-3 in fertile men after scrotal heat stress. *PLoS One*, 10(10), e0141320 (2015)  
DOI: 10.1371/journal.pone.0141320
22. M. H. Zhang, L. P. Zhai, Z. Y. Fang, A. N. Li, Y. Qiu and Y. X. Liu: Impact of a mild scrotal heating on sperm chromosomal abnormality, acrosin activity and seminal alpha-glucosidase in human fertile males. *Andrologia* (2018)  
DOI: 10.1111/and.12985
23. Y. X. Liu: Control of spermatogenesis in primate and prospect of male contraception. *Arch Androl*, 51(2), 77-92 (2005)  
DOI: 10.1080/01485010490485768
24. S. R. Chen, A. Batool, Y. Q. Wang, X. X. Hao, C. S. Chang, C. Y. Cheng and Y. X. Liu: The

- control of male fertility by spermatid-specific factors: searching for contraceptive targets from spermatozoon's head to tail. *Cell Death Dis*, 7(11), e2472 (2016)  
DOI: 10.1038/cddis.2016.344
25. E. Nieschlag and A. Henke: Hopes for male contraception. *Lancet*, 365(9459), 554-6 (2005)  
DOI: 10.1016/S0140-6736(05)17924-0
  26. Q. Yu, X. Q. Mei, X. F. Ding, T. T. Dong, W. W. Dong and H. G. Li: Construction of a catsper1 DNA vaccine and its antifertility effect on male mice. *PLoS One*, 10(5), e0127508 (2015)  
DOI: 10.1371/journal.pone.0127508
  27. J. Stover, J. E. Rosen, M. N. Carvalho, E. L. Korenromp, H. S. Friedman, M. Cogan and B. Deperthes: The case for investing in the male condom. *PLoS One*, 12(5), e0177108 (2017)  
DOI: 10.1371/journal.pone.0177108
  28. L. Li, J. Shao and X. Wang: Percutaneous no-scalpel vasectomy via one puncture in China. *Urol J*, 11(2), 1452-6 (2014)  
DOI: 10.22037/uj.v11i2.2053
  29. M. Mullinax, S. Sanders, B. Dennis, J. Higgins, J. D. Fortenberry and M. Reece: How condom discontinuation occurs: interviews with emerging adult women. *J Sex Res*, 54(4-5), 642-650 (2017)  
DOI: 10.1080/00224499.2016.1143440
  30. D. Waller, D. Bolick, E. Lissner, C. Premanandan and G. Gamerman: Reversibility of Vasalgel male contraceptive in a rabbit model. *Basic Clin Androl*, 27, 8 (2017)  
DOI: 10.1186/s12610-017-0051-1
  31. M. Sabes-Alsina, O. Tallo-Parra, M. T. Mogas, J. M. Morrell and M. Lopez-Bejar: Heat stress has an effect on motility and metabolic activity of rabbit spermatozoa. *Anim Reprod Sci*, 173, 18-23 (2016)  
DOI: 10.1016/j.anireprosci.2016.08.004
  32. J. S. Fleming, F. Yu, R. M. McDonald, S. A. Meyers, G. W. Montgomery, J. F. Smith and H. D. Nicholson: Effects of scrotal heating on sperm surface protein PH-20 expression in sheep. *Mol Reprod Dev*, 68(1), 103-14 (2004)  
DOI: 10.1002/mrd.20049
  33. H. Henning, C. Masal, A. Herr, K. Wolf, C. Urhausen, A. Beineke, M. Beyerbach, S. Kramer and A. R. Gunzel-Apel: Effect of short-term scrotal hyperthermia on spermatological parameters, testicular blood flow and gonadal tissue in dogs. *Reprod Domest Anim*, 49(1), 145-57 (2014)  
DOI: 10.1111/rda.12244
  34. M. Chen, H. Cai, J. L. Yang, C. L. Lu, T. Liu, W. Yang, J. Guo, X. Q. Hu, C. H. Fan, Z. Y. Hu, F. Gao and Y. X. Liu: Effect of heat stress on expression of junction-associated molecules and upstream factors androgen receptor and Wilms' tumor 1 in monkey sertoli cells. *Endocrinology*. 149(10):4871-82 (2008)
  35. M. Rao, X. L. Zhao, J. Yang, S. F. Hu, H. Lei, W. Xia and C. H. Zhu: Effect of transient scrotal hyperthermia on sperm parameters, seminal plasma biochemical markers, and oxidative stress in men. *Asian J Androl*, 17(4), 668-75 (2015)  
DOI: 10.4103/1008-682X.146967
  36. D. S. Hwang, H. G. Kim, S. Park, N. D. Hong, J. H. Ryu and M. S. Oh: Capsaicin attenuates spermatogenic cell death induced by scrotal hyperthermia through its antioxidative and anti-apoptotic activities. *Andrologia*. 49(5) (2017)  
DOI: 10.1111/and.12656
  37. A. Kheradmand, O. Dezfoulian and M. Alirezaei: Ghrelin regulates Bax and PCNA but not Bcl-2 expressions followingscrotalhyperthermia in the rat. *Tissue Cell*, 44(5), 308-15 (2012)  
DOI: 10.1016/j.tice.2012.04.009
  38. M. Kanter, C. Aktas and M. Erboğa: Heat stress decreases testicular germ cell proliferation and increases apoptosis in short term: an immunohistochemical and ultrastructural study. *Toxicol Ind Health*, 29(2), 99-113 (2013)  
DOI: 10.1177/0748233711425082
  39. Z. Zhang, X. Fan, H. Xi, R. Ji, H. Shen, A. Shi and J. He: Effect of local scrotal heating on the expression of tight junction-associated molecule Occludin in boar testes. *Reprod Domest Anim*, 53(2), 458-62 (2018)  
DOI: 10.1111/rda.13131
  40. R. F. Hellon and N. K. Misra: Neurones in the dorsal horn of the rat responding to

- scrotal skin temperature changes. *J Physiol*, 232(2), 375-88 (1973)
41. T. R. Hamilton, C. M. Mendes, L. S. de Castro, P. M. de Assis, A. F. Siqueira, J. C. de Delgado, M. D. Goissis, T. Muiño-Blanco, J. Á. Cebrián-Pérez, M. Nichi, J. A. Visintin and M. E. Assumpção: Evaluation of lasting effects of heat stress on sperm profile and oxidative status of ram semen and epididymal sperm. *Oxid Med Cell Longevity*, 2016, 12 (2016)  
DOI: 10.1155/2016/1687657
  42. T. R. D. S. Hamilton, A. F. P. Siqueira, L. S. de Castro, C. M. Mendes, J. C. Delgado, P. M. de Assis, L. P. Mesquita, P. C. Maiorka, M. Nichi, M. D. Goissis, J. A. Visintin and M. E. O. D. Á. Assumpção: Effect of Heat Stress on Sperm DNA: Protamine Assessment in Ram Spermatozoa and Testicle. *Oxid Med Cell Longev*, 2018, 5413056 (2018)  
DOI: 10.1155/2018/5413056
  43. Y. Lue, A. P. Hikim, C. Wang, M. Im, A. Leung and R. S. Swerdloff: Testicular heat exposure enhances the suppression of spermatogenesis by testosterone in rats: the “two-hit” approach to male contraceptive development. *Endocrinology*, 141(4), 1414-24 (2000)  
DOI: 10.1210/endo.141.4.7.416
  44. Y. Lue, C. Wang, Y. X. Liu, A. P. Hikim, X. S. Zhang, C. M. Ng, Z. Y. Hu, Y. C. Li, A. Leung and R. S. Swerdloff: Transient testicular warming enhances the suppressive effect of testosterone on spermatogenesis in adult cynomolgus monkeys (*Macaca fascicularis*). *J Clin Endocrinol Metab*, 91(2), 539-45 (2006)  
DOI: 10.1210/jc.2005-1808
  45. X. S. Zhang, Y. H. Lue, S. H. Guo, J. X. Yuan, Z. Y. Hu, C. S. Han, A. P. Hikim, R. S. Swerdloff, C. Wang and Y. X. Liu: Expression of HSP105 and HSP60 during germ cell apoptosis in the heat-treated testes of adult cynomolgus monkeys (*macaca fascicularis*). *Front Biosci*, 10, 3110-21 (2005)  
DOI: 10.2741/1767
  46. B. P. Setchell: The Parkes Lecture. Heat and the testis. *J Reprod Fertil*, 114(2), 179-94 (1998)  
DOI: 10.1530/jrf.0.1140179
  47. F. R. Kandeel and R. S. Swerdloff: Role of temperature in regulation of spermatogenesis and the use of heating as a method for contraception. *Fertil Steril*. 1988;49:1-23.
  48. J. I. McNitt and N. L. First: Effects of 72-hour heat stress on semen quality in boars. *Int J Biometeorol*, 14(4), 373-80 (1970)  
DOI: 10.1007/BF01462914
  49. M. Spinaci, S. Volpe, C. Bernardini, M. de Ambrogi, C. Tamanini, E. Seren and G. Galeati: Sperm sorting procedure induces a redistribution of Hsp70 but not Hsp60 and Hsp90 in boar spermatozoa. *J Androl*, 27(6):899-907 (2006)  
DOI: 10.2164/jandrol.106.001008
  50. H. Shen, X. Fan, Z. Zhang, H. Xi, R. Ji, Y. Liu, M. Yue, Q. Li and J. He: Effects of elevated ambient temperature and local testicular heating on the expressions of heat shock protein 70 and androgen receptor in boar testes. *Acta Histochem*, (2019)  
DOI: 10.1016/j.acthis.2019.01.009
  51. H. Wechalekar, B. P. Setchell, K. R. Pilkington, C. Leigh, W. G. Breed and E. Peirce: Effects of whole-body heat on male germ cell development and sperm motility in the laboratory mouse. *Reprod Fertil Dev*, 28(5):545-55 (2016)  
DOI: 10.1071/RD13395
  52. Y. Li, A. Wang, K. Taya and C. Li: Declining semen quality and steadying seminal plasma ions in heat-stressed boar model. *Reprod Med Biol*, 14(4):171-7 (2015)  
DOI: 10.1007/s12522-015-0205-9
  53. B. A. Stone: Thermal characteristics of the testis and epididymis of the boar. *J Reprod Fertil*, 63(2), 551-7 (1981)  
DOI: 10.1530/jrf.0.0630551
  54. K. Larsson and S. Einarsson: Seminal changes in boars after heat stress. *Acta Vet Scand*, 25(1), 57-66 (1984)
  55. L. Malmgren and K. Larsson: Semen quality and fertility after heat stress in boars. *Acta Vet Scand*, 25(3), 425-35 (1984)
  56. G. Ahmad, N. Moïnard, C. Esquerré-Lamare, R. Mieusset and L. Bujan: Mild induced testicular and epididymal hyperthermia alters sperm chromatin integrity in men. *Fertil Steril*, 97(3), 546-53 (2012)  
DOI: 10.1016/j.fertnstert.2011.12.025

57. K. A. Galil and B. P. Setchell: Effects of local heating of the testis on testicular blood flow and testosterone secretion in the rat. *Int J Androl*, 11(1), 73-85 (1988)  
DOI: 10.1111/j.1365-2605.1988.tb01218.x
58. J. Guo, S. X. Tao, M. Chen, Y. Q. Shi, Z. Q. Zhang, Y. C. Li, X. S. Zhang, Z. Y. Hu and Y. X. Liu: Heat treatment induces liver receptor homolog-1 expression in monkey and rat sertoli cells. *Endocrinology*, 148(3), 1255-65 (2007)  
DOI: 10.1210/en.2006-1004
59. X. X. Li, S. R. Chen, B. Shen, J. L. Yang, S. Y. Ji, Q. Wen, Q. S. Zheng, L. Li, J. Zhang, Z. Y. Hu, X. X. Huang and Y. X. Liu: The heat-induced reversible change in the blood-testis barrier (BTB) is regulated by the androgen receptor (AR) via the partitioning-defective protein (Par) polarity complex in the mouse. *Biol Reprod*, 89(1), 1-10 (2013)  
DOI: 10.1095/biolreprod.113.109405
60. X. S. Zhang, Z. H. Zhang, X. Jin, P. Wei, X. Q. Hu, M. Chen, C. L. Lu, Y. H. Lue, Z. Y. Hu, A. P. Sinha Hikim, R. S. Swerdloff, C. Wang and Y. X. Liu: Dedifferentiation of adult monkey Sertoli cells through activation of extracellularly regulated kinase 1/2 induced by heat treatment. *Endocrinology*, 147(3), 1237-45 (2006)  
DOI: 10.1210/en.2005-0981
61. G. R. Dohle, G. M. Colpi, T. B. Hargreave, G. K. Papp, A. Jungwirth, W. Weidner and E. A. U. W. G. o. M. Infertility: EAU guidelines on male infertility. *Eur Urol*, 48(5), 703-11 (2005)  
DOI: 10.1016/j.eururo.2005.06.002
62. E. Geva, J. B. Lessing, L. Lerner-Geva and A. Amit: Free radicals, antioxidants and human spermatozoa: clinical implications. *Hum Reprod*, 13(6), 1422-4 (1998)  
DOI: 10.1093/oxfordjournals.humrep.a019709
63. M. Heidary, S. Vahhabi, J. Reza Nejadi, B. Delfan, M. Birjandi, H. Kaviani and S. Givrad: Effect of saffron on semen parameters of infertile men. *Urol J*, 5(4), 255-9 (2008)
64. B. P. Setchell, L. Plöen and E. M. Ritzen: Effect of local heating of rat testes after suppression of spermatogenesis by pretreatment with a GnRH agonist and an anti-androgen. *Reproduction*, 124(1), 133-40 (2002)  
DOI: 10.1530/rep.0.1240133
65. C. Wang, V. McDonald, A. Leung, L. Superlano, N. Berman, L. Hull and R. S. Swerdloff: Effect of increased scrotal temperature on sperm production in normal men. *Fertil Steril*, 68(2), 334-9 (1997)  
DOI: 10.1016/S0022-5347(01)62981-1
66. R. Munkelwitz and B. R. Gilbert: Are boxer shorts really better? A critical analysis of the role of underwear type in male subfertility. *J Urol*, 160(4), 1329-33 (1998)  
DOI: 10.1097/00005392-199810000-00030
67. C. Lin, Y. S. Choi, S. G. Park, L. W. Gwon, J. G. Lee, J. M. Yon, I. J. Baek, B. J. Lee, Y. W. Yun and S. Y. Nam: Enhanced protective effects of combined treatment with  $\beta$ -carotene and curcumin against hyperthermic spermatogenic disorders in mice. *Biomed Res Int*, 2016, 2572073 (2016)  
DOI: 10.1155/2016/2572073
68. S. G. Park, J. M. Yon, C. Lin, L. W. Gwon, J. G. Lee, I. J. Baek, B. J. Lee, Y. W. Yun and S. Y. Nam: Capsaicin attenuates spermatogenic cell death induced by scrotal hyperthermia through its antioxidative and anti-apoptotic activities. *Andrologia*. 2016 Jul 12.  
DOI: 10.1111/and.12656.
69. Z. Li, Y. Li, X. Zhou, P. Dai and C. Li: Autophagy involved in the activation of the Nrf2-antioxidant system in testes of heat-exposed mice. *J Therm Biol*, 71(2), 142-52 (2018)  
DOI: 10.1016/j.jtherbio.2017.11.006
70. C. Lin, D. G. Shin, S. G. Park, S. B. Chu, L. W. Gwon, J. G. Lee, J. M. Yon, I. J. Baek and S. Y. Nam: Curcumin dose-dependently improves spermatogenic disorders induced by scrotal heat stress in mice. *Food Funct*, 6(12), 3770-7 (2015)  
DOI: 10.1039/c5fo00726g
71. M. Rao, S. Ma, S. Hu, H. Lei, Y. Wu, Y. Zhou, W. Xia and C. Zhu: Isobaric tags for relative and absolute quantification-based proteomic analysis of testis biopsies in rhesus monkeys treated with transient scrotal hyperthermia. *Oncotarget*, 8(49), 85909-25 (2017)  
DOI: 10.18632/oncotarget.20719
72. E. Carlsen, A. M. Andersson, J. H. Petersen and N. E. Skakkebaek: History of febrile illness and variation in semen quality. *Hum Reprod*, 18, 2089-92 (2003)  
DOI: 10.1093/humrep/deg412



73. A. Eisenberg-Lerner, S. Bialik, H. U. Simon and A. Kimchi: Life and death partners: apoptosis, autophagy and the cross-talk between them. *Cell Death Differ*, 16, 966-75 (2009)  
DOI: 10.1038/cdd.2009.33
74. M. Zhang, M. Jiang, Y. Bi, H. Zhu, Z. Zhou and J. Sha: Autophagy and apoptosis act as partners to induce germ cell death after heat stress in mice. *PLoS ONE*, 7, e41412 (2012)  
DOI: 10.1371/journal.pone.0041412
75. M. Ahotupa and I. Huhtaniemi: Impaired detoxification of reactive oxygen and consequent oxidative stress in experimentally cryptorchid rat testis. *Biol Reprod*, 46(6), 1114-8 (1992)  
DOI: 10.1095/biolreprod46.6.1.114
76. M. Ikeda, H. Kodama, J. Fukuda, Y. Shimizu, M. Murata, J. Kumagai and T. Tanaka: Role of radical oxygen species in rat testicular germ cell apoptosis induced by heat stress. *Biol Reprod*, 61(2), 393-9 (1999)  
DOI: 10.1095/biolreprod61.2.3.93
77. V. Peltola, I. Huhtaniemi and M. Ahotupa: Abdominal position of the rat testis is associated with high level of lipid peroxidation. *Biol Reprod*, 53(5), 1146-50 (1995)  
DOI: 10.1095/biolreprod53.5.1.146
78. C.C. Love and R.M. Kenney: Scrotal heat stress induces altered sperm chromatin structure associated with a decrease in protamine disulfide bonding in the stallion. *Biol Reprod*. 60:615-620 (1999)
79. B. L. Sailer, L. J. Sarkar, J. A. Bjordahl, L. K. Jost and D. P. Evenson: Effects of heat stress on mouse testicular cells and sperm chromatin structure. *J Androl*, 18(3), 294-301 (1997)  
DOI: 10.1002/j.1939-4640.1997.tb01922.x
80. B. K. Zhu and B. P. Setchell: Effects of paternal heat stress on the *in vivo* development of preimplantation embryos in the mouse. *Reprod Nutr Dev*, 44(6), 617-29 (2004)  
DOI: 10.1051/rnd:2004064
81. B. Zhu, S. K. Walker, H. Oakey, B. P. Setchell and S. Maddocks: Effect of paternal heat stress on the development *in vitro* of preimplantation embryos in the mouse. *Andrologia*, 36(6), 384-94 (2004)  
DOI: 10.1111/j.1439-0272.2004.00635.x
82. K. M. Cammack, H. Mesa and W. R. Lamberson: Genetic variation in fertility of heat-stressed male mice. *Theriogenology*, 66(9), 2195-201 (2006)  
DOI: 10.1016/j.theriogenology.2006.06.011
83. M. Nichi, P. E. Bols, R. M. Züge, V. H. Barnabe, I. G. Goovaerts, R. C. Barnabe and C. N. Cortada: Seasonal variation in semen quality in *Bos indicus* and *Bos taurus* bulls raised under tropical conditions. *Theriogenology*, 66(4), 822-8 (2006)  
DOI: 10.1016/j.theriogenology.2006.01.056
84. C. G. Blumer, R. M. Fariello, A. E. Restelli, D. M. Spaine, R. P. Bertolla and A. P. Cedenho: Sperm nuclear DNA fragmentation and mitochondrial activity in men with varicocele. *Fertil Steril*, 90(5), 1716-22 (2008)  
DOI: 10.1016/j.fertnstert.2007.09.007
85. D. Sakkas and J. G. Alvarez: Sperm DNA fragmentation: mechanisms of origin, impact on reproductive outcome, and analysis. *Fertil Steril*, 93(4), 1027-36 (2010)  
DOI: 10.1016/j.fertnstert.2009.10.046
86. K. Shiraishi, H. Takihara and H. Matsuyama: Elevated scrotal temperature, but not varicocele grade, reflects testicular oxidative stress-mediated apoptosis. *World J Urol*, 28(3), 359-64 (2010)  
DOI: 10.1007/s00345-009-0462-5
87. T. Shikone, H. Billig and A.J. Hsueh: Experimentally induced cryptorchidism increases apoptosis in rat testis. *Biol Reprod*, 51(5), 865-72 (1994)  
DOI: 10.1095/biolreprod51.5.8.65
88. B. J. Houston, B. Nixon, J. H. Martin, G. N. De Iuliis, N. A. Trigg, E. G. Bromfield, K. E. McEwan and R. J. Aitken: Heat exposure induces oxidative stress and DNA damage in the male germ line. *Biol Reprod*, 98(4), 593-606 (2018)  
DOI: 10.1093/biolre/i0y009
89. R. Ramasamy, S. Besada and D. J. Lamb: Fluorescent *in situ* hybridization of human sperm: diagnostics, indications, and therapeutic implications. *Fertil Steril*, 102(6), 1534-9 (2014)  
DOI: 10.1016/j.fertnstert.2014.09.013
90. Z. Sarrate, F. Vidal and J. Blanco: Role of sperm fluorescent *in situ* hybridization studies in infertile patients: indications,

- study approach, and clinical relevance. *Fertil Steril*, 93(6), 1892-902 (2010)  
DOI: 10.1016/j.fertnstert.2008.12.139
91. W. J. Huang, D. J. Lamb, E. D. Kim, J. de Lara, W. W. Lin, L. I. Lipshultz and F. Z. Bischoff: Germ-cell nondisjunction in testes biopsies of men with idiopathic infertility. *Am J Hum Genet*, 64(6), 1638-45 (1999)  
DOI: 10.1086/302402
  92. B. P. Carreira, C. M. Carvalho and I. M. Araujo: Regulation of injury-induced neurogenesis by nitric oxide. *Stem Cells Int*. 2012; 895659. (2012)  
DOI: 10.1155/2012/895659
  93. S. B. Chao, L. Guo, X. H. Ou, S. M. Luo, Z. B. Wang, H. Schatten, G. L. Gao and Q. Y. Sun: Heated spermatozoa: effects on embryonic development and epigenetics. *Hum Reprod*, 27(4), 1016-24 (2012)  
DOI: 10.1093/humrep/des005
  94. R. Oko and P. Sutovsky: Biogenesis of sperm perinuclear theca and its role in sperm functional competence and fertilization. *J Reprod Immunol*, 83(1-2), 2-7 (2009)  
DOI: 10.1016/j.jri.2009.05.008
  95. Y. Yu, J. Vanhorne and R. Oko: The origin and assembly of a zona pellucida binding protein, IAM38, during spermiogenesis. *Microsc Res Tech*, 72(8), 558-65 (2009)  
DOI: 10.1002/jemt.20696
  96. N. H. Hjollund, L. Storgaard, E. Ernst, J. P. Bonde and J. Olsen: Impact of diurnal scrotal temperature on semen quality. *Reprod Toxicol*, 16(3), 215-21 (2002)  
DOI: 10.1016/S0890-6238(02)00025-4
  97. A. Ilacqua, G. Izzo, G.P. Emerenziani, C. Baldari and A. Aversa: Lifestyle and fertility: the influence of stress and quality of life on male fertility. *Reprod Biol Endocrinol*. 16(1):115. (2018)  
DOI: 10.1186/s12958-018-0436-9.143
  98. Y. Qiu, L. G. Wang, L. H. Zhang, J. Li, A. D. Zhang and M. H. Zhang: Sperm chromosomal aneuploidy and DNA integrity of infertile men with anejaculation. *J Assist Reprod Genet*. 2012;29(2):185-94.  
DOI: 10.1007/s10815-011-9688-4
  99. T. Planchenault, D. Cechová and V. Keil-Dlouha: Matrix degrading properties of sperm serine proteinase, acrosin. *FEBS Lett*. 294(3):279-281. (1991)
  100. R.T. Richardson, B.S. Nikolajczyk, L.H. Abdullah, J.C. Beavers and M.G. O'Rand: Localization of rabbit sperm acrosin during the acrosome reaction induced by immobilized zona matrix. *Biol Reprod*. 45(1):20-26. (1991)
  101. Z. H. Zhang, Z. Y. Hu, X. X. Song, L. J. Xiao, R. J. Zou, C. S. Han and Y. X. Liu: Disrupted expression of intermediate filaments in the testis of rhesus monkey after experimental cryptorchidism. *Int J Androl*, 27(4), 234-9 (2004)  
DOI: 10.1111/j.1365-2605.2004.00477.x
  102. H. Cai, Y. Ren, X. X. Li, J. L. Yang, C. P. Zhang, M. Chen, C. H. Fan, X. Q. Hu, Z. Y. Hu, F. Gao and Y. X. Liu: Scrotal heat stress causes a transient alteration in tight junctions and induction of TGF-beta expression. *Int J Androl*, 34(4), 352-62 (2011)  
DOI: 10.1111/j.1365-2605.2010.01089.x
  103. M. Chen, J. X. Yuan, Y. Q. Shi, X. S. Zhang, Z. Y. Hu, F. Gao and Y. X. Liu: Effect of 43 degrees treatment on expression of heat shock proteins 105, 70 and 60 in cultured monkey Sertoli cells. *Asian J Androl*, 10(3), 474-85 (2008)  
DOI: 10.1111/j.1745-7262.2008.00391.x
  104. D. H. Wang, J. R. Hu, L. Y. Wang, Y. J. Hu, F. Q. Tan, H. Zhou, J. Z. Shao and W. X. Yang: The apoptotic function analysis of p53, Apaf1, Caspase3 and Caspase7 during the spermatogenesis of the Chinese fire-bellied newt *Cynops orientalis*. *PLoS One*, 7(6), e39920 (2012)  
DOI: 10.1371/journal.pone.0039920
  105. S. Kaur and M. P. Bansal: Protective role of dietary-supplemented selenium and vitamin E in heat-induced apoptosis and oxidative stress in mice testes. *Andrologia*, 47(10), 1109-19 (2015)  
DOI: 10.1111/and.12390
  106. J. H. Kim, S. J. Park, T. S. Kim, H. J. Park, J. Park, B. K. Kim, G. R. Kim, J. M. Kim, S. M. Huang, J. I. Chae, C. K. Park and D. S. Lee: Testicular hyperthermia induces Unfolded Protein Response signaling activation in spermatocyte. *Biochem Biophys Res Commun*, 434(4), 861-6 (2013)  
DOI: 10.1016/j.bbrc.2013.04.032
  107. Y. Vera, M. Diaz-Romero, S. Rodriguez, Y. Lue, C. Wang, R. S. Swerdloff and A. P. Sinha Hikim: Mitochondria-dependent pathway is involved in heat-induced male

- germ cell death: lessons from mutant mice. *Biol Reprod*, 70(5), 1534-40 (2004)  
DOI: 10.1095/biolreprod.103.024661
108. S. G. Park, J. M. Yon, C. Lin, L. W. Gwon, J. G. Lee, I. J. Baek, B. J. Lee, Y. W. Yun and S. Y. Nam: Capsaicin attenuates spermatogenic cell death induced by scrotal hyperthermia through its antioxidative and anti-apoptotic activities. *Andrologia*, 49(5) (2017)  
DOI: 10.1111/and.12656
109. M. Chihara, T. Nakamura, N. Sakakibara, S. Otsuka, O. Ichii and Y. Kon: The onset of heat-induced testicular calcification in mice: involvement of the telomeric locus on chromosome 1. *Am J Pathol*, 184(9), 2480-92 (2014)  
DOI: 10.1016/j.ajpath.2014.06.004
110. M. M. Matzuk and D. J. Lamb: The biology of infertility: research advances and clinical challenges. *Nat Med*, 14(11), 1197-213 (2008)  
DOI: 10.1038/nm.f.1895
111. A. Laiho, N. Kotaja, A. Gyenesei and A. Sironen: Transcriptome profiling of the murine testis during the first wave of spermatogenesis. *PLoS One*, 8(4), e61558 (2013)  
DOI: 10.1371/journal.pone.0061558
112. F. Rossi, G. Bernardini, P. Bonfanti, A. Colombo, M. Prati and R. Gornati: Effects of TCDD on spermatogenesis related factor-2 (SRF-2): gene expression in *Xenopus*. *Toxicol Lett*, 191(2-3), 189-94 (2009)  
DOI: 10.1016/j.toxlet.2009.08.022
113. S. M. Wu, V. Baxendale, Y. Chen, A. L. Pang, T. Stitely, P. J. Munson, M. Y. Leung, N. Ravindranath, M. Dym, O. M. Rennert, and W. Y. Chan: Analysis of mouse germ-cell transcriptome at different stages of spermatogenesis by SAGE: Biological significance. *Genomics*, 84(6), 971-81 (2004)  
DOI: 10.1016/j.ygeno.2004.08.018
114. M. Rao, D. Ke, G. Cheng, S. Hu, Y. Wu, Y. Wang, F. Zhou., H. Liu, C. Zhu and W. Xia: The regulation of CIRBP by transforming growth factor beta during heat shock-induced testicular injury. *Andrology*, (2018).  
DOI: 10.1111/andr.12566
115. M. Rao, Z. Y. Zeng, L. Tang, G. P. Cheng, W. Xia and C. H. Zhu: Next-generation sequencing-based microRNA profiling of mice testis subjected to transient heat stress. *Oncotarget*. 8(67):111672–82. (2017)  
DOI: 10.18632/oncotarget.22900
116. H. Xi, X. Fan, Z. Zhang, Y. Liang, Q. Li and J. He: Bax and Bcl-2 are involved in the apoptosis induced by local testicular heating in the boar testis. *Reprod Domest Anim*, 52(3), 359-65 (2017)  
DOI: 10.1111/rda.12904
117. S. Danno, K. Itoh, T. Matsuda and J. Fujita: Decreased expression of mouse Rbm3, a cold-shock protein, in Sertoli cells of cryptorchid testis. *Am J Pathol*, 156(5), 1685-92 (2000)  
DOI: 10.1016/S0002-9440(10)65039-0
118. A. Eisenberg-Lerner and A. Kimchi: PKD at the crossroads of necrosis and autophagy. *Autophagy*. 8(3):433-34. (2012)  
DOI: 10.4161/auto.19288
119. A. Horibe, N. Eid, Y. Ito, Y. Otsuki and Y. Kondo: Ethanol-induced autophagy in Sertoli cells is specifically marked at androgen-dependent stages of the spermatogenic cycle: potential mechanisms and implications. *Int J Mol Sci*. 20(1). pii: E184. (2019)  
DOI: 10.3390/ijms20010184
120. M. Hamasaki, N. Furuta, A. Matsuda, A. Nezu, A. Yamamoto, N. Fujita, et al: Autophagosomes form at ER-mitochondria contact sites. *Nature*, 495(7441):389-93. (2013)  
DOI: 10.1038/nature11910
121. T. D. Oberley, J. M. Swanlund, H. J. Zhang and K. C. Kregel: Aging results in increased autophagy of mitochondria and protein nitration in rat hepatocytes following heat stress. *J Histochem Cytochem*. 56:615–27. (2008)  
DOI: 10.1369/jhc.2008.950873
122. B. Levine and D. J. Klionsky: Development by self-digestion: molecular mechanisms and biological functions of autophagy. *Dev Cell*. 2004;6:463–77.
123. G. Kroemer, G. Mariño and B. Levine: Autophagy and the integrated stress response. *Mol Cell*. 40(2):280-93. (2010)  
DOI: 10.1016/j.molcel.2010.09.023

124. S. Foroozan-Broojeni, M. Tavalaei, R. A. Lockshin, Z. Zakeri, H. Abbasi and M. H. Nasr-Esfahani: Comparison of main molecular markers involved in autophagy and apoptosis pathways between spermatozoa of infertile men with varicocele and fertile individuals. *Andrologia*. Oct 23:e13177. (2018)  
DOI: 10.1111/and.13177
125. N. Chen, M. Lin, N. Liu, S. Wang and X. Xiao: Methylmercury-induced testis damage is associated with activation of oxidative stress and germ cell autophagy. *J Inorg Biochem*, 190:67-74. (2019)  
DOI: 10.1016/j.jinorgbio.2018.10.007
126. J. J. Parrish, K.L. Willenburg, K. M. Gibbs, K.B. Yagoda, M. M. Krautkramer, T. M. Loether, F.C.S.A. Melo: Scrotal insulation and sperm production in the boar. *Mol Reprod Dev*. 84(9):969-78. (2017)  
DOI: 10.1002/mrd.22841
127. Y. F. Zhu, Y. G. Cui, X. J. Guo, L. Wang, Y. Bi, Y. Q. Hu, X. Zhao, Q. Liu, R. Huo, M. Lin, Z. M. Zhou and J. H. Sha: Proteomic analysis of effect of hyperthermia on spermatogenesis in adult male mice. *J Proteome Res*, 5(9), 2217-25 (2006)  
DOI: 10.1021/pr0600733
128. H. Zhu, Y. Cui, J. Xie, L. Chen, X. Chen, X. Guo, Y. Zhu, X. Wang, J. Tong, Z. Zhou, Y. Jia, Y. H. Lue, A. S. Hikim, C. Wang, R. S. Swerdloff and J. Sha: Proteomic analysis of testis biopsies in men treated with transient scrotal hyperthermia reveals the potential targets for contraceptive development. *Proteomics*, 10(19), 3480-93 (2010)  
DOI: 10.1002/pmic.201000281
129. A. P. Hikim, Y. Lue, C. M. Yamamoto, Y. Vera, S. Rodriguez, P. H. Yen, K. Soeng, C. Wang and R. S. Swerdloff: Key apoptotic pathways for heat-induced programmed germ cell death in the testis. *Endocrinology*, 144(7), 3167-75 (2003)  
DOI: 10.1210/en.2003-0175
130. M. Rao, W. Xia, J. Yang, L. X. Hu, S. F. Hu, H. Lei, Y. Q. Wu and C. H. Zhu: Transient scrotal hyperthermia affects human sperm DNA integrity, spermapoptosis, and spermprotein expression. *Andrology*, 4(6), 1054-63 (2016)  
DOI: 10.1111/andr.12228
131. W. Widlak and N. Vydra: The Role of heat shock factors in mammalian spermatogenesis. *Adv Anat Embryol Cell Biol*. 222:45-65. (2017)
132. K. Purandhar, P.K. Jena, B. Prajapati, P. Rajput and S. Seshadri: Understanding the role of heat shock protein Isoforms in male fertility, aging and apoptosis. *World J Mens Health*. 32(3):123-32. (2014)
133. M. Kus-Liśkiewicz, J. Polańska, J. Korfanty, M. Olbryt, N. Vydra, A. Toma and W. Widlak: Impact of heat shock transcription factor 1 on global gene expression profiles in cells which induce either cytoprotective or pro-apoptotic response following hyperthermia. *BMC Genomics*, 14, 456 (2013)  
DOI: 10.1186/1471-2164-14-456
134. J. Korfanty, T. Stokowy, P. Widlak, A. Gogler-Pigłowska, L. Handschuh, J. Podkowiński, N. Vydra, A. Naumowicz, A. Toma-Jonik and W. Widlak: Crosstalk between HSF1 and HSF2 during the heat shock response in mouse testes. *Int J Biochem Cell Biol*, 57, 76-83 (2014)  
DOI: 10.1016/j.biocel.2014.10.006
135. J. C. Rockett, F. L. Mapp, J. B. Garges, J. C. Luft, C. Mori and D. J. Dix: Effects of hyperthermia on spermatogenesis, apoptosis, gene expression, and fertility in adult male mice. *Biol Reprod*, 65(1), 229-39 (2001)  
DOI: 10.1095/biolreprod65.1.2.29
136. M. K. Kim, K. M. Cha, S. Y. Hwang, U. K. Park, S. K. Seo, S. H. Lee, M. S. Jeong, S. Cho, S. R. Kopalli and S. K. Kim: Pectinase-treated Panax ginseng protects heat stress-induced testicular damage in rats. *Reproduction*, 153(6), 737-47 (2017)  
DOI: 10.1530/REP-16-0560
137. F. Bozkaya, M. O. Atli, A. Guzeloglu, S. A. Kayis, M. E. Yildirim, E. Kurar, R. Yilmaz and N. Aydilek: Effects of long-term heat stress and dietary restriction on the expression of genes of steroidogenic pathway and small heat-shock proteins in rat testicular tissue. *Andrologia*. 49(6). (2017)  
DOI: 10.1111/and.12668
138. F.A. Gohari, B. Saranjam, M. Asgari, L. Omid, H. Ekrami and S.A. Moussavi-Najarkola: An experimental study of the effects of combined exposure to microwave and heat on gene expression and sperm parameters in mice. *J Hum Reprod Sci*. 10(2):128-34. (2017)  
DOI: 10.4103/jhrs.JHRS\_136\_16

139. X. Guo, S. Chi, X. Cong, H. Li, Z. Jiang, R. Cao and W. Tian: Baicalin protects sertoli cells from heat stress-induced apoptosis via activation of the Fas/FasL pathway and Hsp72 expression. *Reprod Toxicol.* 57:196-203. (2015)
140. S. M. Downs: Nutrient pathways regulating the nuclear maturation of mammalian oocytes. *Reprod Fertil Dev*, 27(4), 572-82 (2015)  
DOI: 10.1071/RD14343
141. J. L. Gatti, S. Castella, F. Dacheux, H. Ecroyd, S. Métayer, V. Thimon and J. L. Dacheux: Post-testicular sperm environment and fertility. *Anim Reprod Sci*, 82-83, 321-39 (2004)  
DOI: 10.1016/j.anireprosci.2004.05.011
142. M. H. Zhang, L. P. Zhai, Z. Y. Fang, A. N. Li, W. Xiao and Y. Qiu: Effect of scrotal heating on sperm quality, seminal biochemical substances, and reproductive hormones in human fertile men. *J Cell Biochem*, 119(12), 10228-38 (2018)  
DOI: 10.1002/jcb.27365
143. X. Wang, F. Liu, X. Gao, X. Liu, X. Kong, H. Wang and J. Li: Comparative proteomic analysis of heat stress proteins associated with rat sperm maturation. *Mol Med Rep*, 13(4), 3547-52 (2016)  
DOI: 10.3892/mmr.2016.4958
144. F. J. Liu and X. F. Shen: Comparative analysis of human reproductive proteomes identifies candidate proteins of sperm maturation. *Mol Biol Rep*, 39(12), 10257-63 (2012)  
DOI: 10.1007/s11033-012-1902-7
145. W. Xun, L. Shi, T. Cao, C. Zhao, P. Yu, D. Wang, G. Hou and H. Zhou: Dual functions in response to heat stress and spermatogenesis: Characterization of expression profile of small heat shock proteins 9 and 10 in goat testis. *Biomed Res Int*, 2015, 686239 (2015)  
DOI: 10.1155/2015/686239
146. D. R. Rocha, J. A. Martins, M. F. van Tilburg, R. V. Oliveira, F. B. Moreno, A. C. Monteiro-Moreira, R. A. Moreira, A. A. Araújo and A. A. Moura: Effect of increased testicular temperature on seminal plasma proteome of the ram. *Theriogenology*, 84(8), 1291-305 (2015)  
DOI: 10.1016/j.theriogenology.2015.07.008
147. J. Yaeram, B. P. Setchell and S. Maddocks: Effect of heat stress on the fertility of male mice *in vivo* and *in vitro*. *Reprod Fertil Dev*, 18(6), 647-53 (2006)  
DOI: 10.1071/RD05022
148. H. Wechalekar, B. P. Setchell, E. J. Peirce, M. Ricci, C. Leigh and W. G. Breed: Whole-body heat exposure induces membrane changes in spermatozoa from the cauda epididymidis of laboratory mice. *Asian J Androl*, 12(4), 591-8 (2010)  
DOI: 10.1038/aja.2010.41
149. M. Pérez-Crespo, B. Pintado and A. Gutiérrez-Adán: Scrotal heat stress effects on sperm viability, sperm DNA integrity, and the offspring sex ratio in mice. *Mol Reprod Dev*, 75(1), 40-7 (2008)  
DOI: 10.1002/mrd.20759
150. K. Shiraishi and K. Naito: Nitric oxide produced in the testis is involved in dilatation of the internal spermatic vein that compromises spermatogenesis in infertile men with varicocele. *BJU Int*, 99(5), 1086-90 (2007)  
DOI: 10.1111/j.1464-410x.2007.06800.x
151. B. P. Carreira, C. M. Carvalho and I. M. Araújo: Regulation of injury-induced neurogenesis by nitric oxide. *Stem Cells Int*, 2012, 895659 (2012)  
DOI: 10.1155/2012/895659
152. H. Meiser and R. Schulz: Detection and localization of two constitutive NOS isoforms in bull spermatozoa. *Anat Histol Embryol*, 32(6), 321-5 (2003)  
DOI: 10.1111/j.1439-0264.2003.00459.x
153. S. E. Lewis, E. T. Donnelly, E. S. Sterling, M. S. Kennedy, W. Thompson and U. Chakravarthy: Nitric oxide synthase and nitrite production in human spermatozoa: evidence that endogenous nitric oxide is beneficial to sperm motility. *Mol Hum Reprod*, 2(11), 873-8 (1996)  
DOI: 10.1093/molehr/2.11.873
154. S. S. du Plessis, D. A. McAllister, A. Luu, J. Savia, A. Agarwal and F. Lampiao: Effects of H<sub>2</sub>O<sub>2</sub> exposure on human sperm motility parameters, reactive oxygen species levels and nitric oxide levels. *Andrologia*, 42(3), 206-10 (2010) <http://med.wanfangdata.com.cn/Periodical/N2007EPST0000007>  
DOI: 10.1111/j.1439-0272.2009.00980.x

155. S. S. du Plessis, K. Hagenaar and F. Lampiao: The *in vitro* effects of melatonin on human sperm function and its scavenging activities on NO and ROS. *Andrologia*, 42(2), 112-6 (2010)  
DOI: 10.1111/j.1439-0272.2009.00964.x
156. P. C. Rodriguez, L. B. Valdez, T. Zaobornyj, A. Boveris and M. T. Beconi: Nitric oxide and superoxide anion production during heparin-induced capacitation in cryopreserved bovine spermatozoa. *Reprod Domest Anim*, 46(1), 74-81 (2011)  
DOI: 10.1111/j.1439-0531.2010.01583.x
157. J. M. Moran, L. Madejón, C. Ortega Ferrusola and F. J. Peña: Nitric oxide induces caspase activity in boar spermatozoa. *Theriogenology*, 70(1), 91-6 (2008)  
DOI: 10.1016/j.theriogenology.2008.02.010
158. K. F. Beck, W. Eberhardt, S. Frank, A. Huwiler, U. K. Mebmer, H. Muhl and J. Pfeilschifter: Inducible NO synthase: role in cellular signaling. *J Exp Biol*, 202(Pt 6), 645-53 (1999)
159. L. E. Shao, T. Tanaka, R. Gribi and J. Yu: Thioredoxin-related regulation of NO/NOS activities. *Ann N Y Acad Sci*, 962, 140-50 (2002)  
DOI: 10.1111/j.1749-6632.2002.tb04064.x
160. E. Bonfoco, D. Krainc, M. Ankarcona, P. Nicotera and S. A. Lipton: Apoptosis and necrosis: two distinct events induced, respectively, by mild and intense insults with N-methyl-D-aspartate or nitric oxide/superoxide in cortical cell cultures. *Proc Natl Acad Sci U S A*, 92(16), 7162-6 (1995)  
DOI: 10.1073/pnas.92.16.7162
161. T. Ramya, M. M. Misro, D. Sinha, D. Nandan and S. Mithal: Altered levels of seminal nitric oxide, nitric oxide synthase, and enzymatic antioxidants and their association with sperm function in infertile subjects. *Fertil Steril*, 95(1), 135-40 (2011)  
DOI: 10.1016/j.fertnstert.2010.06.044
162. Y. Sakamoto, T. Ishikawa, Y. Kondo, K. Yamaguchi and M. Fujisawa: The assessment of oxidative stress in infertile patients with varicocele. *BJU Int*, 101(12), 1547-52 (2008)  
DOI: 10.1016/S0022-5347(08)61860-1
163. M. Salter and R. G. Knowles: Assay of NOS activity by the measurement of conversion of oxyhemoglobin to methemoglobin by NO. *Methods Mol Biol*, 100, 61-5 (1998)  
DOI: 10.1385/1-59259-749-1:61
164. P. Ghosh, S. Mukherjee, A. Bhoumik and S. R. Dungdung: A novel epididymal quiescence factor inhibits sperm motility by modulating NOS activity and intracellular NO-cGMP pathway. *J Cell Physiol*, 233(5):4345-59 (2018)
165. S. S. Gross and M. S. Wolin: Nitric oxide: pathophysiological mechanisms. *Annu Rev Physiol*, 57, 737-69 (1995)
166. C. Romeo, R. Ientile, G. Santoro, P. Impellizzeri, N. Turiaco, P. Impalà, S. Cifalà, G. Cutroneo, F. Trimarchi and C. Gentile: Nitric oxide production is increased in the spermatic veins of adolescents with left idiopathic varicocele. *J Pediatr Surg*, 36(2), 389-93 (2001)  
DOI: 10.1053/jpsu.2001.20724
167. C. Roessner, U. Paasch, H. J. Glander and S. Grunewald: Activity of nitric oxide synthase in mature and immature human spermatozoa. *Andrologia*, 42(2), 132-7 (2010)  
DOI: 10.1111/j.1439-0272.2009.01005.x
168. M. K. O'Bryan, S. Schlatt, O. Gerdprasert, D. J. Phillips, D. M. de Kretser and M. P. Hedger: Inducible nitric oxide synthase in the rat testis: evidence for potential roles in both normal function and inflammation mediated infertility. *Biol Reprod*, 63(5), 1285-93 (2000)  
DOI: 10.1095/biolreprod63.5.1.285
169. Y. Lue, A. P. S. Hikim, C. Wang, A. Leung and R. S. Swerdloff: Functional role of inducible nitric oxide synthase in the induction of male germ cell apoptosis, regulation of sperm number, and determination of testes size: evidence from null mutant mice. *Endocrinology*, 144(7), 3092-100 (2003)  
DOI: 10.1210/en.2002-0142
170. J. Guo, Y. Jia, S. X. Tao, Y. C. Li, X. S. Zhang, Z. Y. Hu, N. Chiang, Y. H. Lue, A. P. Hikim, R. S. Swerdloff, C. Wang and Y. X. Liu: Expression of nitric oxide synthase during germ cell apoptosis in testis of cynomolgus monkey after testosterone and heat treatment. *J Androl*, 30(2), 190-9 (2009)  
DOI: 10.2164/jandrol.108.005538
171. M. B. Herrero, E. Cebral, M. Boquet, J. M. Viggiano, A. Vitullo and M. A. Gimeno:

- Effect of nitric oxide on mouse sperm hyperactivation. *Acta Physiol Pharmacol Ther Latinoam*, 44(3), 65-9 (1994)
172. R. R. Yeoman, W. D. Jones and B. M. Rizk: Evidence for nitric oxide regulation of hamster sperm hyperactivation. *J Androl*, 19(1), 58-64 (1998)  
DOI: 10.1002/j.1939-4640.1998.tb02470.x
173. W. J. Hellstrom, M. Bell, R. Wang and S. C. Sikka: Effect of sodium nitroprusside on sperm motility, viability, and lipid peroxidation. *Fertil Steril*, 61(6), 1117-22 (1994)  
DOI: 10.1016/s0015-0282(16)56766-1
174. E. Miraglia, F. De Angelis, E. Gazzano, H. Hassanpour, A. Bertagna, E. Aldieri, A. Revelli and D. Ghigo: Nitric oxide stimulates human sperm motility via activation of the cyclic GMP/protein kinase G signaling pathway. *Reproduction*, 141(1), 47-54 (2011)  
DOI: 10.1530/REP-10-0151
175. B. S. Joo, S. H. Park, S. J. Park, H. S. Kang, H. S. Moon and H. D. Kim: The effect of nitric oxide on sperm cell function and embryo development. *Am J Reprod Immunol*, 42(6), 327-34 (1999)  
DOI: 10.1111/j.1600-0897.1999.tb00109.x
176. B. Aljabari, A. E. Calogero, A. Perdichizzi, E. Vicari, R. Karaki, T. Lahloub, R. Zadari, K. El-Abed, F. Nicoletti, E. J. Miller, V. A. Pavlov and Y. Al-Abed: Imbalance in seminal fluid MIF indicates male infertility. *Mol Med*, 13(3-4), 199-202 (2007)  
DOI: 10.2119/2006-00114.Aljabari
177. G. Frenette, R. R. Tremblay, J. Y. Dube, C. Lazure and M. Lemay: High concentrations of the macrophage migration inhibitory factor in human seminal plasma and prostatic tissues. *Arch Androl*, 41(3), 185-93 (1998)  
DOI: 10.3109/01485019808994890
178. G. Frenette, C. Lessard, E. Madore, M. A. Fortier and R. Sullivan: Aldose reductase and macrophage migration inhibitory factor are associated with epididymosomes and spermatozoa in the bovine epididymis. *Biol Reprod*, 69(5), 1586-92 (2003)  
DOI: 10.1095/biolreprod.103.019216
179. G. Frenette, C. Légaré, F. Saez and R. Sullivan: Macrophage migration inhibitory factor in the human epididymis and semen. *Mol Hum Reprod*, 11(8), 575-82 (2005)  
DOI: 10.1093/molehr/gah197
180. R. Eickhoff, C. Baldauf, H. W. Koyro, G. Wennemuth, Y. Suga, J. Seitz, R. Henkel and A. Meinhardt: Influence of macrophage migration inhibitory factor (MIF) on the zinc content and redox state of protein-bound sulphhydryl groups in ratsperm: indications for a new role of MIF in spermmaturation. *Mol Hum Reprod*, 10(8), 605-11 (2004)  
DOI: 10.1093/molehr/gah075
181. A. Meinhardt, M. Bacher, J. R. McFarlane, C. N. Metz, J. Seitz, M. P. Hedger, D. M. de Kretser and R. Bucala: Macrophage migration inhibitory factor production by Leydig cells: evidence for a role in the regulation of testicular function. *Endocrinology*, 137(11), 5090-5 (1996)  
DOI: 10.1210/endo.137.11.8895383
182. A. G. Utleg, E. C. Yi, T. Xie, P. Shannon, J. T. White, D. R. Goodlett, L. Hood and B. Lin: Proteomic analysis of human prostasomes. *Prostate*, 56(2), 150-61 (2003)  
DOI: 10.1002/pros.10255
183. K. Komori, A. Tsujimura, Y. Okamoto, Y. Matsuoka, T. Takao, Y. Miyagawa, S. Takada, N. Nonomura and A. Okuyama: Relationship between substances in seminal plasma and Acrobeads Test results. *Fertil Steril*, 91(1), 179-84 (2009)  
DOI: 10.1016/j.fertnstert.2007.11.003
184. T. G. Cooper, C. H. Yeung, D. Nashan and E. Nieschlag: Epididymal markers in human infertility. *J Androl*, 9(2), 91-101 (1988)  
DOI: 10.1002/j.1939-4640.1988.tb01016.x
185. R. Henkel, G. Maass, H. C. Schuppe, A. Jung, J. Schubert and W. B. Schill: Seasonal changes of neutral alpha-glucosidase activity in human semen. *J Androl*, 27(1), 34-9 (2006)  
DOI: 10.2164/jandrol.05064
186. P. Pena, J. Risopatron, J. Villegas, W. Miska, W. B. Schill and R. Sanchez: Alpha-glucosidase in the human epididymis: topographic distribution and clinical application. *Andrologia*, 36(5), 315-20 (2004)  
DOI: 10.1111/j.1439-0272.2004.00625.x
187. B. Kret, M. Milad and R. S. Jeyendran: New discriminatory level for glucosidase activity to diagnose epididymal obstruction

- or dysfunction. *Arch Androl*, 35(1), 29-33 (1995)  
DOI: 10.3109/01485019508987850
188. Y. Qui, K. Tian, D. Yang, Q. Zhang, Z. Wang, Z. Huang, M. Sun and S. Mingzhang: The change of carnitine content in seminal plasma after reversible injection occlusion of vas deferens. *Contraception*, 51(4), 261-3 (1995)  
DOI: 10.1016/0010-7824(95)00043-A
189. W. M. Buckett and D. I. Lewis-Jones: Fructose concentrations in seminal plasma from men with nonobstructive azoospermia. *Arch Androl*. 48(1):23-7 (2002)
190. M. K. Ahmad, A. A. Mahdi, K. K. Shukla, N. Islam, S. P. Jaiswar and S. Ahmad: Effect of *Mucuna pruriens* on semen profile and biochemical parameters in seminal plasma of infertile men. *Fertil Steril*, 90(3):627-35 (2008)  
DOI: 10.1016/j.fertnstert.2007.07.1314
191. C. H. Yeung, T. G. Cooper and T. Senge: Histochemical localization and quantification of alpha-glucosidase in the epididymis of men and laboratory animals. *Biol Reprod*, 42(4), 669-76 (1990)  
DOI: 10.1095/biolreprod42.4.6.69
192. M. H. Viljoen, M. S. Bornman, M. P. van der Merwe and D. J. du Plessis: Alpha-glucosidase activity and sperm motility. *Andrologia*, 22(3), 205-8 (1990)  
DOI: 10.1111/j.1439-0272.1990.tb01967.x
193. M. M. Roaiah, T. Mostafa, D. Salem, A. R. El-Nashar, I. I. Kamel and M. S. El-Kashlan: alpha-1, 4-Glucosidase activity in infertile oligoasthenozoospermic men with and without varicocele. *Andrologia*, 39(1), 28-32 (2007)  
DOI: 10.1111/j.1439-0272.2006.00756.x
194. G. Vivas-Acevedo, R. Lozano-Hernández and M. I. Camejo: Varicocele decreases epididymal neutral  $\alpha$ -glucosidase and is associated with alteration of nuclear DNA and plasma membrane in spermatozoa. *BJU Int*, 113(4), 642-9 (2014)  
DOI: 10.1111/bju.12523
195. M. Watanabe, R. Roussev, P. Ahlering, R. Sauer, C. Coulam and R. S. Jeyendran: Correlation between neutral alpha-glucosidase activity and sperm DNA fragmentation. *Andrologia*, 41(5), 316-8 (2009)  
DOI: 10.1111/j.1439-0272.2009.00950.x
196. C. Sudheer Kumar and M. J. Swamy: Modulation of chaperone-like and membranolytic activities of major horse seminal plasma protein HSP-1/2 by L-carnitine. *J Biosci*, 42(3), 469-79 (2017)  
DOI: 10.1007/s12038-017-9693-6
197. S. D. Ahmed, K. A. Karira, Jagdesh and S. Ahsan: Role of L-carnitine in male infertility. *J Pak Med Assoc*, 61(8), 732-6 (2011)  
DOI: 10.1016/j.trsl.2010.12.017
198. U. Wetterauer and H. J. Heite: Carnitine in seminal fluid as parameter for the epididymal function. *Andrologia*, 10(3), 203-10 (1978)
199. K. Li, W. Li and Y. Huang: Determination of free L-carnitine in human seminal plasma by high performance liquid chromatography with pre-column ultraviolet derivatization and its clinical application in male infertility. *Clin Chim Acta*, 378(1-2), 159-63 (2007)  
DOI: 10.1016/j.cca.2006.11.008
200. I. Gülçin: Antioxidant and antiradical activities of L-carnitine. *Life Sci*, 78(8), 803-11 (2006)  
DOI: 10.1016/j.lfs.2005.05.103
201. R. E. L. Cabral, T. B. Mendes, V. Vendramini and S. M. Miraglia: Carnitine partially improves oxidative stress, acrosome integrity, and reproductive competence in doxorubicin-treated rats. *Andrology*, 6(1), 236-46 (2018)  
DOI: 10.1111/andr.12426
202. E. Spaleková, A. V. Makarevich and N. Lukáč: Ram sperm motility parameters under the influence of epidermal growth factor. *Vet Med Int*, 2011, 642931 (2011)  
DOI: 10.4061/2011/642931
203. S. C. Stubbs, T.B. Hargreave and F. K. Habib: Localization and characterization of epidermal growth factor receptors on human testicular tissue by biochemical and immunohistochemical techniques. *J Endocrinol*. 125(3):485-92 (1990)
204. K. Zitta, M. Albrecht, S. Weidinger, A. Mayerhofer and F. Köhn: Protease activated receptor 2 and epidermal growth factor receptor are involved in the regulation of



**TSH on sperm and bio-components**

human sperm motility. *Asian J Androl*, 9(5), 690-6 (2007)  
DOI: 10.1111/j.1745-7262.2007.00289.x

205. R. K. Naz and P. Kaplan: Effects of epidermal growth factor on human sperm cell function. *J Androl*, 14(4), 240-7 (1993)  
DOI: 10.1002/j.1939-4640.1993.tb03361.x

206. H. Fuse, M. Sakamoto, M. Okumura and T. Katayama: Epidermal growth factor contents in seminal plasma as a marker of prostatic function. *Arch Androl*, 29(1), 79-8 (1992)  
DOI: 10.3109/01485019208987712

**Key Words:** Transient scrotal heating, TSH, Autophagy, Apoptosis, Sperm protein, Bio-components, Review

**Send correspondence to:** Yi Qiu, Maternal and Child Health Care Hospital of Shandong Province, 238 East Road of Jingshi, Jinan, Shandong 250014, China, Tel: 86-531-68795969, Fax: 86-531-68795005, E-mail: qiuyi987@sina.com