

A Review on Prescription and Non-Prescription Appetite Suppressants and Evidence Based Method to Treat Overweight and Obesity

SURUCHI SINGH¹, PANKAJ BHATT², SATISH K SHARMA³

Glocal School Of Pharmacy, Glocal University, Saharanpur, U.P, India, 247121

Received: 04.01.20, Revised: 10.02.20, Accepted: 21.03.20

ABSTRACT

Shedding pounds isn't simply a question of fitting into a smaller size, it's about your wellbeing. Additional pounds raise your danger of creating interminable and once in a while destructive infections, including stroke, coronary illness, type 2 diabetes, and hypertension. At the point when you're hefty or overweight, you're additionally at a more serious danger of medical problems that meddle with your personal satisfaction, for example, joint pain, sleep apnea, and depression. Appetite suppressants are prescription medications that assist you with controlling your craving, what's more, accordingly your food consumption. Contingent upon the formulation utilized, they may assist you with feeling full sooner. Longings and issues with craving are basic reasons individuals neglect to adhere to a low-calorie weight reduction diet. Appetite suppressants help control your yearning so you can concentrate on giving your body the nutritious food it needs to work ideally as you get in shape — and feel better while doing it. Ordinarily, appetite suppressants are utilized alongside a decreased calorie diet. You get backing and direction with respect to both your medications and your dinner plan at all times. More than 1 of every 3 American grown-ups are overweight or obese. At Weight No More, we need to help decrease this measurement by helping you accomplish a healthy size and a healthy life.

Keywords: review, evidence, overweight.

INTRODUCTION

Health care suppliers utilize the Body Mass Index (BMI), which is a proportion of your weight according to your height, to characterize overweight and obesity. Individuals who have a BMI somewhere in the range of 25 and 30 are viewed as overweight. Obesity is characterized as having a BMI of 30 or more noteworthy.^{1,2} You can calculate your BMI in the event that you are overweight or obese. Being overweight or large may build the danger of health issues. Your health care supplier can evaluate your individual hazard because of your weight.³

Obesity is a chronic condition that influences more than one of every three adults in the United States. Another in three adults is overweight. In the event that you are battling with your weight, you may locate that a good dieting arrangement and standard physical movement assist you with getting more fit⁴ and keep it off over the long haul. On the off chance that these way of life changes are insufficient to assist you with getting in shape or keep up your weight reduction, your primary care physician may prescribe medications as a feature of your weight-control program.^{5,6} Appetite suppressants are drugs that stunt the body into trusting it isn't ravenous. Medications delegated appetite suppressants follow up on the

body's central nervous system, fooling the body into accepting that it isn't ravenous. A few instances of remedy appetite suppressants include: benzphetamine, diethylpropion, mazindol and phentermine. These medications for the most part come as tablets or expanded delivery capsules. Appetite suppressants can be endorsed or bought over-the-counter.⁶

Appetite suppressants are utilized as a transient treatment for patients with weight.

It's not unexpected to feel hungry. It's your body's method of revealing to you it's an ideal opportunity to fuel up. Be that as it may, on the off chance that you end up wanting food despite the fact that you just ate, there's a decent possibility you'll put on weight.⁷

Classifications Of Appetite Suppressants

1. Prescription Appetite Suppressants
2. Non Prescription Appetite Suppressants

Prescription Appetite Suppressants

Appetite suppressants are drugs intended to control hunger and, thusly, assist you with getting in shape.^{9,11} The FDA has endorsed these medications that you can just get with a Doctor's solution:

Liraglutide

Also called Saxenda. You take this as an injection. It was additionally showcased as a diabetes treatment under the brand name Victoza. The medication hoses hunger by following up on a hormone in the gut.⁸

Naltrexone

Also called Contrave. It contains two medications and may influence the reward framework in your brain, so eating certain nourishments that would regularly cause you to feel great do not do anymore.¹⁰ It additionally works on the hypothalamus, the part of your brain that manages appetite, temperature, and different capacities.¹²

Phentermine

Also called Qsymia. It's a combo of two medications. Phentermine is a stimulant that causes you to feel less hungry.¹³ Topiramate is a medication utilized for seizures and migraine, however as a component of a combo with phentermine may cause you to feel not so much hungry but rather more full.

There are likewise some different choices - like phentermine, benzphetamine, diethylpropion, and phendimetrazine - yet those must be utilized for as long as 12 weeks.¹⁴

Non Prescription Appetite Suppressants

Hunger suppressants check your hunger and prevent you from gorging. Here are 6 characteristic hunger suppressants that you can remember for your eating regimen to remain on target.

Fenugreek

Fenugreek is a spice that has a place with legumes family. The little yellow seeds, which are generally utilized in Indian food for including flavor are additionally known for its medicinal properties. The seeds contain 45% fiber, which is for the most part insoluble. The fiber hinders the procedure of carb and fat absorption in the body,¹⁵ saving your fuller for a more drawn out time. Having fenugreek seeds are viewed as valuable in bringing down blood pressure.^{16,17}

How to have it:

You can straightforwardly chew one teaspoon of these seeds two times every day or absorb 1 teaspoon of the seeds a glass of water for the time being and drink it the morning.

Green tea extract

Green tea extract is an incredible fat eliminator and appetite suppressant. It contains two significant mixes, which helps in the weight

reduction process-caffeine and catechins. Studies recommend that catechins additionally help to support digestion, which is urgent when you plan to shed kilos.¹⁸

How to have it: Have 200-250 ml of green tea twice day by day for compelling outcomes.

Almonds

Nuts are constantly considered excellent tidbits to control your mid-afternoon hunger aches. In any case, almonds specifically ought to be your preferred choice when you need to dodge binge eating.¹⁹ Loaded with fiber, almonds are a rich source of antioxidants, vitamin E, and magnesium, which helps to suppress your untimely hunger aches. They increase the feeling of fullness.²⁰

How to have it: Have a bunch of almonds when you feel hungry.

Ginger

We as a whole love the kind of ginger in our tea and dishes, however this herb makes them flabbergast medicinal properties. It stimulates your digestive system, can ease nausea and can likewise suppress your appetite.²¹ According to a little 2012 investigation, men who expended ginger at breakfast felt more full for at any rate 3 hours after their dinner.²²

How to have it: You can have ginger tea or can even have ginger enhancements subsequent to counseling your primary care physician.

Yerba mate

Yerba mate is a native plant of South America. It is regularly used to make a drink called mate. The beverage is known for its energy stimulating and appetite stifling properties. It hinders the procedure of stomach purging, which keeps you from careless eating.

How to have it: You can either take yerba mate in powdered shape or have supplements or have yerba mate tea twice day by day.²³

Coffee

Because of high caffeine content, coffee isn't viewed as useful for wellbeing. In any case, when we talk about weight reduction it very well may be very useful. Coffee diminishes your hunger, expands calorie consume and fat breakdown. These things can be gainful when attempting to shed kilos.²⁴ Yet, by coffee, we mean dark coffee, not the standard one weighed down with sugar and milk.²⁵

How to have it: Have 200-250 ml of coffee

twice daily. Due to high caffeine content, coffee isn't viewed as useful for wellbeing. Yet, when we talk about weight reduction it very well may be very advantageous. Coffee diminishes your hunger, expands calorie consume and fat breakdown. These things can be helpful when attempting to shed kilos.²⁶

Mechanism Of Action

Centrally acting appetite suppressant drugs utilized in the treatment of weight loss into 2 wide pharmacological classifications; those which demonstration through cerebrum catecholamine pathways and those which demonstration by means of serotonin pathways.²⁷ Of the previous gathering, amphetamine and phenmetrazine are no longer suggested in view of their energizer properties and addictive potential. The rest of the medications in this class incorporate amfepramone (diethylpropion),²⁸ phentermine, mazindol and phenylpropanolamine. All have been appeared to decrease craving and lower food consumption, subsequently helping obese patients more effectively keep to a low-calorie diet and get more fit.²⁹

Do Appetite Suppressant Works?

Indeed, however likely not as much as you would trust. A review of studies on five significant FDA-approved prescription medications for obesity, including orlistat, shows that any of them work better than a placebo for helping individuals lose at any rate 5% of their body weight through the span of a year. Phentermine-topiramate and liraglutide had the most noteworthy chances of getting that going.³⁰

It's additionally imperative to comprehend that these medications don't take a shot at their own. Studies show that weight reduction medication, including appetite suppressants, work best when you're additionally rolling out solid improvements to your eating and exercise propensities simultaneously.³¹

When Appetite Suppressants Are Included In Weight Loss Plan?

In the event that you've just battled with weight reduction for a considerable length of time, appetite suppressants might be a piece of your underlying arrangement. Be that as it may, if this is your first endeavor at weight reduction, we'll sit back and watch how much weight you lose over a couple of long periods of following an eating routine and exercise plan.

There are likewise clinical rules overseeing the utilization of appetite suppressants. These proposals depend on your weight file (BMI), which

mirrors the measure of muscle versus fat you convey. On the off chance that your BMI is 25-29.9, you're overweight. Any worth that is at least 30 is viewed as obese.³²

Prescription appetite suppressants are suggested for patients who meet two rules. You're a decent applicant in the event that you have a BMI of 27 or higher and you're determined to have at any rate one weight-related ailment, for example, hypertension or Type 2 diabetes.³³ You can likewise get the medication if your BMI is more than 30, with or without other medical issues.

Settling on a choice about utilizing an appetite suppressant isn't simply founded on your BMI. We'll likewise think about your general wellbeing and check whether there's any explanation the medication may not be ok for you. For instance, appetite suppressants may interface with different medications you take. You additionally shouldn't take them in case you're pregnant or plan to get pregnant.³⁴

Benefits Of Using Prescription Medication To Lose Weight

At the point when joined with changes to behavior, including eating and physical movement propensities, prescription medications may assist some with people to lose weight. Weight loss of 5 to 10 percent of your beginning body weight may help improve your wellbeing by bringing down glucose, pulse, and fatty substances. Getting in shape additionally can improve some other medical issues identified with overweight and obese, for example, joint disease or sleep apnea. Most weight loss happens inside the initial a half year of beginning the medication.³⁵

Tips For Taking Weight Loss Medication

1. Adhere to your doctor's instructions about weight reduction medications.
2. Purchase your medication from a pharmacy or web distributor endorsed by your doctor.
3. Take weight reduction medication to help your healthy eating and physical activity program.³⁶
4. Know the side effects and warnings for taking any medicine.
5. Inquire as to whether you should quit taking your medication in the event that you are not getting thinner following 12 weeks.
6. Talk about different medications, including supplements and vitamins, you are taking with your doctor when considering weight-loss medications.
7. Abstain from assuming weight-loss medications during pregnancy or on the off chance that you are planning a pregnancy.³⁷

Evidence-Based Methods To Suppress Appetite **Eat more protein and fortifying fats**

Not all nourishments fulfill hunger similarly. Contrasted with sugars, proteins and certain fats are more powerful for fulfilling craving and keeping individuals feeling full for more. An individual can supplant a few wellsprings of sugar with proteins and invigorating fats to help monitor their craving. The accompanying high-protein nourishments are lean meats, eggs, beans and peas, soy items, Greek yogurt.

Drink water before each entrée

Drinking a huge glass of water legitimately before eating has been found to cause an individual to feel more full, more fulfilled, and less ravenous after the supper. Another investigation, which took a gander at hunger in 50 overweight females, indicated that drinking 1.5 liters of water a day for about two months caused a decrease in craving and weight, and furthermore prompted more noteworthy fat misfortune. A soup starter may likewise extinguish the appetite.³⁸ Research from 2007 indicated that individuals revealed feeling more full following the supper on the off chance that they had a fluid starter.³⁹

Eat all the more high-fiber nourishments

Fiber doesn't separate like different nourishments, so it remains in the body for more. This hinders assimilation and keeps individuals feeling full for the duration of the day. Exploration proposes that fiber can be a compelling appetite suppressant. High-fiber eats less carbs are likewise connected with lower corpulence rates. Then again, another audit found that bringing additional fiber into the eating regimen was viable in under portion of the investigations they took a gander at. More exploration is expected to recognize which wellsprings of fiber are the best for smothering appetite.⁴⁰ Healthful high-fiber nourishments incorporate entire grains, beans and heartbeats, apples and avocados, almonds, chia seeds, vegetables.⁴¹

Exercise before a meal

Exercise is another sound and viable hunger suppressant. An audit dependent on 20 distinct examinations found that hunger hormones are stifled following activity, particularly high-force workouts.⁴² They discovered lower levels of ghrelin in the body, a hormone that makes us hungry, and more significant levels of "completion hormones, for example, PYY and GLP-1."^{43,44}

Drink Yerba Maté tea

Examination shows that a tea called Yerba Maté, which originates from the *Ilex paraguariensis*

plant, can diminish hunger and improve mind-set when joined with high-power work out. Yerba Maté is accessible for buy on the web.

Switch to dark chocolate

Dark chocolate has been shown to suppresses appetite compared to milk chocolate. One study showed that people ate less during their next meal after snacking on dark instead of milk chocolate.⁴⁵

Eat some ginger

Consuming a small amount of ginger powder has been shown to reduce appetite and increase fullness, possibly because of its stimulating effect on the digestive system. This was a small-scale study,⁴⁶ so more research is needed to confirm this effect. Ginger powder is available for purchase online.⁴⁷

Eat bulky, low-calorie foods

Reducing general food intake while dieting can leave people with a ravenous appetite. This can cause a relapse into binge eating. However, dieting does not have to mean going hungry. Some foods are high in nutrients and energy, but low in calories. These include vegetables, fruits, beans, and whole grains. Eating a large volume of these foods will stop the stomach from growling and still allow a person to burn more calories than they consume.⁴⁸

Stress less

Comfort eating due to stress, anger, or sadness is different from physical hunger. Research has linked stress with an increased desire to eat, binge eating, and eating non-nutritious food. Mindfulness practices and mindful eating may reduce stress-related binge eating and comfort eating, according to one research. Regular sleep, social contact, and time spent relaxing can also help tackle stress.⁴⁹

Mindful eating

The brain is a major player in deciding what and when a person eats. If a person pays attention to the food they are eating instead of watching TV during a meal, they may consume less. Research published in the journal *Appetite* found that eating a huge meal in the dark led people to consume 36 percent more. Paying attention to food during meals can help a person reduce overeating. Another article showed that mindfulness might reduce binge eating and comfort eating, which are two significant factors that influence obesity. The National Institute of Health recommend using mind and body-based techniques, such as meditation and yoga, to curb

appetite.^{49,48}

CONCLUSION

Researchers are as of now studying a few new medications and combination of medications in animals and individuals. Researchers are attempting to recognize more secure and more viable medications to help individuals who are overweight or corpulent get thinner and keep up a healthy weight for quite a while.

Future medications may utilize new methodologies, for example, to combined medications that influence appetite and those that influence enslavement craving, stimulate gut hormones that lessen appetite, shrink the blood vessels that feed fat cells in the body, along these lines keeping them from developing, target genes that influence body weight, change microscopic organisms in the gut to control weight.

REFERENCES

1. Alberts HI, Thewissen R, Raes L. Dealing with problematic eating behaviour. The effects of a mindfulness-based intervention on eating behaviour, food cravings, dichotomous thinking and body image concern. *Appetite*. 2012;58(3):847–851.
2. Allen GS. A double-blind trial of diethylpropion hydrochloride, mazindol and placebo in the treatment of exogenous obesity. *Current Therapy Research* 22: 678–683, 1977
3. Altschuler S, Conte A, Sebok M, Marlin R, Winick C. Three controlled trials of weight loss with phenylpropanolamine. *International Journal of Obesity* 6: 549–556, 1982
4. Altschuler S, Frazer DL. Double-blind clinical evaluation of the anorectic activity of phenylpropanolamine hydrochloride drops and placebo drops in the treatment of exogenous obesity. *Current Therapy Research* 40: 211–217, 1986
5. American Medical Association. Drugs used in obesity. *Drug Evaluations Subscriptions* 3(6): 1–616, 1990
6. Angel I. Central receptors and recognition sites mediating the effects of monoamines and anorectic drugs on feeding behaviour. *Clinical Neuropharmacology* 13: 361–391, 1990
7. Ashwell M. A survey of patients' views on doctors' treatment of obesity. *Practitioner* 211: 653–661, 1973
8. Atkinson RL, Greenway FL, Bray GA, Dahms WT, Molitch ME. Treatment of obesity: comparison of physician and non-physician therapists using placebo and anorectic drugs in a double-blind trial. *International Journal of Obesity* 1: 113–120, 1977
9. Baer RA, Fischer S, Huss DB. Mindfulness and acceptance in the treatment of disordered eating. *Journal of Rational-Emotive & Cognitive-Behavior Therapy*. 2005;23(4):281–300.
10. Baer RA, Smith GT, Allen KB. Assessment of mindfulness by self-report the Kentucky Inventory of Mindfulness Skills. *Assessment*. 2004;11(3):191–206.
11. Blackburn G, Morgan J, Lavin P, Noble R, Funder Burk F. Determinants of the pressor effect of phenylpropanolamine in healthy subjects. *Journal of the American Medical Association* 261: 3267–3272, 1989
12. Blundell IE, Latham EM, McArthur RA. Structural analysis of the actions of amphetamine and fenfluramine on food intake and feeding behaviour in animals and in man. *Current Medical Research and Opinion* 6 (Suppl. 1): 34–54, 1979
13. Bolding OT. Diethylpropion hydrochloride: an effective appetite suppressant. *Current Therapy Research* 16: 40–43, 1974
14. British National Formulary. Appetite suppressants. British Medical Association and the Royal Pharmaceutical Society of Great Britain, 1991
15. Bunnell DW, Shenker IR, Nussbaum MP, Jacobson MS, Cooper P. Subclinical versus formal eating disorders: differentiating psychological features. *International Journal of Eating Disorders*. 2006;9(3):357–362.
16. Burton P, Smit HJ, Lightowler HJ. The influence of restrained and external eating patterns on overeating. *Appetite*. 2007;49(1):191–197.
17. Cosentino G, Conrad AO, Uwaifo GIPhentermine and topiramate for the management of obesity: a review..*Drug Des Devel Ther*. 2011 Apr 5;7:267-78.
18. Farello G, Ferrara P, Antenucci A, Basti C, Verrotti A. The link between obesity and migraine in childhood: a systematic review. *Ital J Pediatr*. 2017 Mar 7;43(1):27
19. fernstrom ID, Choi S. The development of tolerance to drugs that suppress food intake.*Pharmacol Ther*. 2008 Jan;117(1):105-22.
20. Flegal KM, Carroll MD, Ogden CL, Curtin LR. Prevalence and trends in obesity among US adults, 1999–2008. *JAMA*. 2010;303(3):235–241.
21. Hainer V, Kabrnova K, Aldhoon B, Kunesova M, Wagenknecht M. Serotonin and norepinephrine reuptake inhibition and eating behavior. *Ann N Y Acad Sci*. 2006 Nov;1083:252-69. doi: 10.1196/annals.1367.017.
22. Hudson JI, Hiripi E, Pope HG, Kessler RC. The prevalence and correlates of eating disorders in the National Comorbidity Survey Replication. *Biological psychiatry*. 2007;61(3):348–358.
23. Hurt RT, Edakkanambeth Varayil J, Ebbert JONew pharmacological treatments for the management of obesity..*Curr Gastroenterol Rep*. 2014;16(6):394.

24. Kristeller JL, Wolever RQ. Mindfulness-based eating awareness training for treating binge eating disorder: the conceptual foundation. *Eat Disord*. 2011;19(1):49–61.
25. Macht M. How emotions affect eating: a five-way model. *Appetite*. 2008;50(1):1–11.
26. National Heart, Lung, and Blood Institute: Overweight and Obesity
27. National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK): Prescription Medications for the Treatment of Obesity
28. Nonogaki K, Nozue K, Kuboki T, Oka Y. Milnacipran, a serotonin and norepinephrine reuptake inhibitor, induces appetite-suppressing effects without inducing hypothalamic stress responses in mice. *Am J Physiol Regul Integr Comp Physiol*. 2007 May;292(5):R1775–81.
29. Ogden CL, Carroll MD. Prevalence of overweight, obesity, and extreme obesity among adults: United States, trends 1960–1962 through 2007–2008. *NCHS Health and Stats*. 2010.
30. Piet I, Hougaard E. The effect of mindfulness-based cognitive therapy for prevention of relapse in recurrent major depressive disorder: a systematic review and meta-analysis. *Clin Psychol Rev*. 2011;31(6):1032–1040.
31. Shapiro SL, Carlson LE, Astin IA, Freedman B. Mechanisms of mindfulness. *Journal of clinical psychology*. 2005;62(3):373–386.
32. Singh J, Kumar R. Phentermine-topiramate: First combination drug for obesity. *Int J Appl Basic Med Res*. 2015 May-Aug;5(2):157–8.
33. Smith BW, Shelley BM, Leahigh L, Vanleit B. A preliminary study of the effects of a modified mindfulness intervention on binge eating. *Complementary Health Practice Review*. 2006;11(3):133–143.
34. Torres SJ, Nowson CA. Relationship between stress, eating behavior, and obesity. *Nutrition*. 2007;23(11):887–894.
35. Treanor M. The potential impact of mindfulness on exposure and extinction learning in anxiety disorders. *Clin Psychol Rev*. 2011;31(4):617–625.
36. Wang Y, Beydoun MA, Liang L, Caballero B, Kumanyika SK. Will all Americans become overweight or obese? Estimating the progression and cost of the US obesity epidemic. *Obesity*. 2008;16(10):2323–2330.
37. Weltzin TE, Weisensel N, Franczyk D, Burnett K, Klitz C, Bean P. Eating disorders in men: Update. *The journal of men's health & gender*. 2005;2(2):186–193.
38. Woolhouse H, Knowles A, Crafti N. Adding mindfulness to CBT programs for binge eating: a mixed-methods evaluation. *Eat Disord*. 2012;20(4):321–339.
39. Yanovski SZ, Yanovski JA. *JAMA*. Long-term drug treatment for obesity: A systematic and clinical review. 2014; 311(1):74–86.
40. Zgierska A, Rabago D, Chawla N, Kushner K, Koehler R, Marlatt A. Mindfulness meditation for substance use disorders: a systematic review. *Subst Abus*. 2009;30(4):266–294.
41. Godsey J. The role of mindfulness based interventions in the treatment of obesity and eating disorders: An integrative review. *Complementary therapies in medicine*. 2013
42. Epel ES, Tomiyama AJ, Dallman M. Stress and reward neural networks, eating, and obesity. In: Brownell K, Gold M, editors. *Handbook of Food Addiction*. Oxford; Oxford University Press; in press.
43. Dallman MF, Pecoraro NC, la Fluer SE. Chronic stress and comfort foods: Self-medication and abdominal obesity. *Brain, Behavior, and Immunity*. 2005;19:275–280.
44. Ouwens MA, van Strien T, van der Staak CP. Tendency toward overeating and restraint as predictors of food consumption. *Appetite*. 2003;40:291–8.
45. Polivy I, Coleman I, Herman CP. The effect of deprivation on food cravings and eating behavior in restrained and unrestrained eaters. *International Journal of Eating Disorders*. 2005;38:301–309.
46. Rutters F, Nieuwenhuizen AG, Lemmens SG, Born JM, Westerterp-Plantenga MS. Acute stress-related changes in eating in the absence of hunger. *Obesity*. 2009;17:72–77.
47. Westenhoefer J. Dietary restraint and disinhibition: Is restraint a homogenous construct? *Appetite*. 1991;16:45–55.
48. Arnsten AFT. Stress signaling pathways that impair prefrontal cortex structure and function. *Nature Reviews Neuroscience*. 2009;10:410–422.
49. Chandler-Laney PC, Castaneda E, Viana IB, Oswald KD, Maldonado CR, Boggiano MM. A history of human-like dieting alters serotonergic control of feeding and neurochemical balance in a rat model of binge-eating. *International Journal of Eating Disorders*. 1997;40:136–142.