

Differential Roles of M1 and M2 Microglia in Neurodegenerative Diseases

Yu Tang · Weidong Le

Received: 2 November 2014 / Accepted: 29 December 2014
© Springer Science+Business Media New York 2015

Abstract One of the most striking hallmarks shared by various neurodegenerative diseases, including Parkinson's disease, Alzheimer's disease (AD), and amyotrophic lateral sclerosis, is microglia-mediated neuroinflammation. Increasing evidence indicates that microglial activation in the central nervous system is heterogeneous, which can be categorized into two opposite types: M1 phenotype and M2 phenotype. Depending on the phenotypes activated, microglia can produce either cytotoxic or neuroprotective effects. In this review, we focus on the potential role of M1 and M2 microglia and the dynamic changes of M1/M2 phenotypes that are critically associated with the neurodegenerative diseases. Generally, M1 microglia predominate at the injury site at the end stage of disease, when the immunoresolution and repair process of M2 microglia are dampened. This phenotype transformation is very complicated in AD due to the phagocytosis of regionally distributed β -amyloid ($A\beta$) plaque and tangles that are released into the extracellular space. The endogenous stimuli including aggregated α -synuclein, mutated superoxide dismutase, $A\beta$, and tau oligomers exist in the milieu that may persistently activate M1 pro-inflammatory responses and finally lead to irreversible neuron loss. The changes of microglial phenotypes depend on the disease stages and severity; mastering the stage-specific switching of M1/M2 phenotypes within appropriate time windows may provide better therapeutic benefit.

Keywords Neurodegenerative diseases · Microglial phenotypes · Classical activation · Alternative activation · M2 microglia · M1/M2 switching

Abbreviations

AD	Alzheimer's disease
ALS	Amyotrophic lateral sclerosis
APP	Amyloid precursor protein
Arg1	Arginase 1
$A\beta$	β -Amyloid
BDNF	Brain-derived neurotrophic factor
CD206	Mannose receptor
Chi3l3	Chitinase-3-Like-3
CNS	Central nervous system
DA	Dopaminergic
ECM	Extracellular matrix
FIZZ1	Found in inflammatory zone 1
IFN- γ	Interferon- γ
IGF-I	Insulin-like growth factor 1
IL	Interleukin
iNOS	Induced nitric oxide synthase
LBs	Lewy bodies
LPS	Lipopolysaccharide
MHC	Major histocompatibility complex
MPTP	1-Methyl-4-phenyl-1,2,3,6-tetrahydropyridine
mSOD1	Mutated superoxide dismutase
NO	Nitric oxide
PD	Parkinson's disease
PET	Positron emission tomography
PS1	Presenilin-1
RELM	Resistin-like molecules
ROS	Reactive oxygen species
SN	Substantia nigra
SRA	Scavenger receptors
TAM	Tumor-associated macrophages
TDP-43	TAR DNA-binding protein 43

Y. Tang
Key Laboratory of Stem Cell Biology, Institute of Health Sciences,
Shanghai Institutes for Biological Sciences, Chinese Academy of
Sciences/Shanghai JiaoTong University School of Medicine,
200025 Shanghai, China

W. Le (✉)
Center for Translational Research of Neurology Disease,
1st Affiliated Hospital, Dalian Medical University,
116011 Dalian, China
e-mail: wdle@sibs.ac.cn

TGF- β	Transforming growth factor- β
TLRs	Toll-like receptors
TNF- α	Tumor necrosis factor- α
TTBK	Tau-tubulin kinase

Introduction

Neuroinflammation is a prominent feature shared by various neurodegenerative diseases such as Parkinson's disease (PD), Alzheimer's disease (AD), and amyotrophic lateral sclerosis (ALS) [1–3]. Microglial activation is the principal component of neuroinflammation in the central nervous system (CNS), which provides the first line of defense whenever injury or disease occurs [2, 4]. The molecular and clinical evidences from postmortem analysis and positron emission tomography (PET) imaging have shown an increase of microglial activation and an increasing accumulated inflammatory mediators during the pathogenesis of neurodegenerative diseases [5–7].

Neuroinflammation is now considered as a double-edged sword that executes both detrimental and beneficial effects on the neurons [8, 9]. Many evidences point to the neurotoxic nature of microglia [6], whereas others indicate that neuroinflammation is actually beneficial in certain circumstances to stimulate myelin repair, remove toxic aggregated proteins and cell debris from the CNS, as well as secretion of neurotrophic factors to prevent neural injury [10–12]. Immune cells within the CNS milieu such as microglia appear to be heterogeneous with diverse functional phenotypes that range from pro-inflammatory M1 phenotypes to immunosuppressive M2 phenotypes. In recent years, M1/M2 paradigm of microglial activation has been increasingly studied in several neurodegenerative diseases in attempt to uncover the mechanisms of immunopathogenesis. In this review, we focus on the roles of microglial phenotypes and their switch in multiple neurodegenerative diseases.

Microglial Phenotypes

M1/M2 Paradigm

Inside the human body, immune cells including peripheral macrophages and CNS microglia cells often communicate with the resident functional cells in the milieu. In the normal condition, the immune responses are fine-regulated in the process of either initiation or resolution, so as to keep tissue homeostasis. In the pathological condition, however, the immune responses are uncontrolled and skew to either extreme of the immune balance that highly integrates with cell loss or cell dysfunction occurred within the inflammatory processes.

The M1/M2 paradigm is a simplified model to decipher the two polars of the inflammatory responses, which resembles the “Ying and Yang” principles in the nature. M1 and M2 macrophages were extensively studied to differentially affect the functional cells in the inflammation-induced pathologies of several human diseases. For example, transition of tissue-resident macrophages (TAM) from the M2 to the M1 phenotype is tightly associated with obesity, insulin resistance, and type 2 diabetes that is also believed to be a chronic inflammatory disease [13–16]. Obesity induces the accumulation of newly recruited M1 macrophages overwhelming M2 macrophages in the adipose tissue which secrete pro-inflammatory mediators, thereby inducing an insulin-resistant state in the adipose tissue that is a strong risk factor for the development of type 2 diabetes mellitus [16, 13–15]. In spinal cord injury, an M1 macrophage response is rapidly induced and then maintained at sites of traumatic injury, and this action overwhelms a comparatively smaller and transient M2 macrophage response, which promotes a regenerative growth in adult sensory axons [17].

The already depicted “M1/M2 paradigm” in insulin resistance and spinal cord injury shed light on the research of microglial activation states in CNS. The category of M1 and M2 microglia is a common category shared by various neurodegenerative diseases. Depending on the milieu in which they become activated or the factors by which they are stimulated, microglia possess states of “classical activation,” “alternative activation,” and “acquired deactivation” [18, 19]. Classical activation is associated with the production of pro-inflammatory cytokines such as tumor necrosis factor- α (TNF- α), interleukin-1 β (IL-1 β), superoxide, nitric oxide (NO), reactive oxygen species (ROS), and proteases that have already been studied for a long time [6, 20, 21]. Microglia in this state are also termed “M1 microglia,” whereas “M2 microglia” is used to include the states of both alternative activation and acquired deactivation (Fig. 1). Alternative activation is limited to the activation state treated by IL-4 or IL-13 and is closely associated with M2 genes that promote anti-inflammation, tissue repair, and extracellular matrix (ECM) reconstruction [19, 22]. Acquired deactivation is another state to alleviate acute inflammation and is induced primarily by uptake of apoptotic cells or exposure to the insult of anti-inflammatory cytokines such as IL-10 and transforming growth factor- β (TGF- β) [18, 19, 23].

It is still not clear whether there are any morphology differences between the two phenotypes or whether they can coexist. Nevertheless, the two phenotypes could be transited into each other in different context that may contribute to pathogenic forms of inflammation in neurodegenerative diseases.

Resolution Mechanisms of M2 Microglia

M1 microglia originally respond to the injury and infection, and generally act in the first line to defense tissue and promote




Fig. 1 M1 and M2 microglia. Microglia possess states of “classical activation,” “alternative activation,” and “acquired deactivation,” depending on the milieu in which they become activated and the factors they are stimulated. Microglia in classical activation state are also termed M1 microglia, which induce iNOS and NF-κB pathways and produce various pro-inflammatory cytokines such as TNF- α , IL-1 β , and IL-6, as well as superoxide, ROS and NO. M2 microglia include the states of both alternative activation and acquired deactivation, which are induced by IL-4/IL-13 and IL-10/TGF- β , respectively. M2 microglia facilitate phagocytosis of cell debris and misfolded proteins, promote ECM reconstruction and tissue repair, and support neuron survival by neurotrophic factors. M2 microglia are driven by the coordinated regulation of various anti-inflammatory factors and antagonize the M1 pro-inflammatory responses that finally results in immunosuppression and neuron protection

the destruction of invading pathogens. However, they also induce neurotoxicity due to the release of pro-inflammatory factors and various neurotoxic mediators and often setup a vicious cycle between dying neurons and acute inflammation [6, 24]. After the onset of classical activation, an anti-inflammatory and repair phase is rapidly initiated that leads to wound healing and brings back tissue homeostasis. M2 microglia are the major effector cells with the potential to dampen pro-inflammatory immune responses and promote the repair genes expression (Fig. 1).

Four major anti-inflammatory cytokines including IL-4, IL-13, IL-10, and TGF- β are employed by M2 microglia to antagonize the pro-inflammatory responses [25, 26]. IL-4 and IL-13 are well-described anti-inflammatory cytokines, which could suppress the production of pro-inflammatory cytokines such as IL-8, IL-6, and TNF- α , and reduce NO release, which collectively protect against lipopolysaccharide (LPS)-induced neuron injury both in vitro and in vivo [25, 27–29]. TGF- β is

a pleiotropic cytokine with diverse functions on angiogenesis, ECM deposition, and participates in suppressing microglial responses [30, 31]. Upregulation of scavenger receptors (SRA) such as CD163 are observed in acquired deactivation to promote debris clearance [32–34]. Besides that, M2 microglia can also enhance neurotrophic factors, such as insulin-like growth factor 1 (IGF-I) release to assist inflammation resolution and promote neuron survival [35]. Anti-inflammatory factors are also released when apoptotic cells or myelin debris are removed [36], accompanied by the induction of typical M2 markers, such as Arginase 1 (Arg1), Mannose receptor (CD206), Found in inflammatory zone 1 (FIZZ1), and Chitinase-3-Like-3 (Chi3l3) to help tissue reconstruction [37, 22].

Notably, these neuroprotective signals are orchestrating each other in a coordinated way against the pro-inflammation responses (Fig. 1). For example, TGF- β can enhance IL-4-induced M2 microglia by increasing the expression of Arg1 and Ym1; whereas in turn, treatment with IL-4 can increase the level of TGF- β 2, suggesting that TGF- β and IL-4 signals are working together in promoting protective effects [26].

Functions of Typical M2 Markers

The resolution states are critically important in chronic neuro-inflammation; thus, the genes specifically associated with these states are of great importance. Although the exact functions of many of those genes are not clear in the CNS, some alternative activation genes such as Arg1 have been well studied in peripheral macrophages, which provide a link between M2 microglia and immunosuppression or repair processes.

Arg1

Arginine metabolism varies in tissues throughout the body including the brain due to the differential activation of multiple enzymatic pathways [38]. Arg1 is a typical marker for M2 macrophage/microglia activation that participates in arginine metabolism. Specifically, both Arg1 and induced nitric oxide synthase (iNOS) expressed in the CNS utilize arginine as the sole substrate for biosynthetic pathways [37, 39]. On one hand, arginine is catalyzed by iNOS to produce citrulline and NO. On the other hand, Arg1 metabolizes arginine into urea and ornithine, which are further metabolized into hydroxyproline and proline, and polyamine. The enzymatic products in the Arg1 pathway contribute greatly to the tissue repair [40]. For instance, hydroxyproline and proline are important sources of collagen synthesis, or by large, ECM synthesis that helps to physically strengthen the tissue and are also used for repair at the sites of injury [41, 42]. Polyamines such as spermines are multivalent cations required for cell proliferation and differentiation [43], and could help protect neurons

from injury by pro-inflammatory cytokines [44]. Arg1 is induced by IL-4 or IL-13 insult and produced anti-inflammatory effects by competing utilization of the common substrate arginine to suppress NO production [45]. The maintenance of high Arg1 expression directs arginine metabolism toward the production of proline or polyamines and keep NO production at a low level, which therefore contributes to the neuroprotection [45].

FIZZ1

FIZZ1 (also known as RELM- α) encodes a 9.4-kDa cysteine-rich protein that is induced by IL-4 and IL-13 and increases collagen expression and myofibroblast differentiation [46, 47]. FIZZ1 belongs to the resistin-like molecule (RELM) family of secreted mammalian proteins, the members of which are upregulated in several infectious and inflammatory settings, including helminth infection, allergic airway inflammation, and colitis [48–50]. FIZZ1 limited Th2 cytokine-dependent inflammatory responses in lung through immunoregulatory effects on CD4⁺ T cell responses [47]. After challenge with *Schistosoma mansoni* eggs, FIZZ1-deficient mice develop exacerbated lung inflammation [47]. FIZZ1 promotes the activation of innate immune cells in the intestine, including macrophages and eosinophils, in the chemically induced colitis [49, 50]. FIZZ1 may also contribute to insulin resistance linking with the effects of promotion of angiogenesis, stimulation of collagen synthesis, and inhibition of apoptosis [49].

Chi313

Chi313 (also known as Ym1) is a secretory protein of 45 kDa synthesized by activated peripheral macrophages that binds saccharides and heparin sulfate on cell surfaces [51], and helps protect the ECM scaffold at the injury sites [52, 53]. Heparin sulfate serves as a docking site for growth factors in the ECM and is degraded by heparinases during inflammation. Ym1 thus acts by binding to heparin so as to slow the loss of growth factors which may be required for the tissue reconstruction [52, 54]. Ym1 is induced by IL-4 or IL-13 stimulation by a STAT6-dependent mechanism [55] and is essential for alternative activation of the microglia/macrophages that are antagonized by LPS and interferon- γ (IFN- γ) [22, 56, 54, 57].

CD206

CD206 is a transmembrane glycoprotein in the macrophage/microglia or dendrite cells and is a member of the C-type lectin family [58]. The N-terminal

cysteine-rich domain of CD206 plays an important role in recognizing terminal mannose, sulfated glycoproteins as well as fucose residues on glycans attached to proteins on the surface of several microorganisms and then helps their clearance from the circulation [59]. To ensure inflammatory agents are removed, CD206 is expressed at low levels during inflammation and at high levels during the resolution of inflammation [60]. In general, CD206 initiates phagocytosis of its ligand and activates immunosuppressive pathways that results in decreased TNF- α and IL-12, whereas increased the expression levels of anti-inflammatory factors such as IL-10 and IL-1R α [61, 62]. In line with the immunosuppressive effects, CD206 is a characteristic of the alternative activated state that could promote CNS repair in the spinal cord injury while limiting secondary inflammatory-mediated injury [63, 64].

Microglial Phenotypes in Neurodegenerative Diseases

Microglial Phenotypes in PD

PD is clinically manifested by progressive loss of dopaminergic (DA) neurons in the substantia nigra (SN) of the midbrain and pathologically characterized by the accumulation of protein aggregates called Lewy bodies (LBs) in the remaining DA neurons [65, 66]. Microglia-mediated neuroinflammation is an important component in PD pathogenesis that shows inversely correlated with the DA neuron survival in patients [5]. In general, activated microglia are prominent and surround DA neurons exhibiting classically activated M1 phenotypes. Among those neurodegenerative diseases, the role and function of M2 microglia specifically in PD are not well studied.

In PD, microglial activation might be initiated either directly or indirectly by the misfolded proteins, pathogens or environmental toxins. α -Synuclein is one of the most prevalent pathological genes altered in familial PD, and it is originally acts as an intracellular component localized at presynaptic terminals [67–69]. Generally, the mutated forms of α -synuclein are released and aggregated, nitrated, or oxidized, which constitute the major components of LBs [67, 70]. Numerous studies have shown that aggregated α -synuclein released into the extracellular space from dying or dead DA neurons can directly induce microglia towards M1 phenotype with the activation of NADPH oxidase, increasing production of ROS and pro-inflammatory cytokines [71–75]. Overexpression of mutant α -synuclein solely in microglia switches microglia into a more reactive M1 phenotype characterized by

elevated levels of pro-inflammatory cytokines including TNF- α and NO [76]. However, a contradictory conclusion comes from a study to characterize the microglial phenotype differences caused by lack of α -synuclein expression [77]. Deficient of α -synuclein in microglia impairs the phagocytic ability and enhances the secretion of TNF- α and IL-6 after LPS stimulation [77]. Those studies complicate the role of α -synuclein in microglia but indeed hint an autonomous microglial reaction in the α -synuclein transgenic model (Fig. 2).

1-Methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP) is an environmental toxin that causes parkinsonism and is used to establish the animal model of PD [65, 78]. MPTP induces DA neuron injury by blocking the electron transport chain of the mitochondria, and thereby indirectly activates microglia [65]. Similarly, typical characteristics of M1 phenotype including the activation of NADPH oxidase and NF- κ B pathways, as well as the release of various pro-inflammatory mediators were observed in the MPTP-intoxicated models [79–81]. LPS as a classical ligand of toll-like receptors (TLRs) is definitely evoking

M1 microglial activation. And therefore, administration of LPS causes extensive DA neuron death both in vitro and in vivo [20, 21, 82].

However, little is known about the activation of the M2 phenotype in the PD pathogenesis. To evaluate the possible link of alternative activation and α -synuclein, Theodore et al. established a mouse model overexpressing human α -synuclein by a recombinant adeno-associated virus vector (AAV2-SYN) [83]. However, the results come out that the expression of cytokines IL-4 and IL-13 as well as M2 marker Arg1 in the SN of mice treated with AAV2-SYN was not significantly changed after 2 or 4 weeks [83]. Considering that M2 microglia generally execute immunoresolution at a later phase, the detected endpoints in the early time may not be convincing. It will be interesting to examine at later time points when neurodegeneration becomes apparent.

As the two microglia phenotypes can transit each other, it might be available to make microglia protective by switching their phenotypes. For instance, treatment with LPS primed microglia into the M1 phenotype in both BV2 cells and primary microglia, while the addition of fasudil, one type of Rho




Fig. 2 Microglial phenotypes in neurodegenerative diseases. Microglial activation states are closely associated with the aggregated misfolded proteins seen in various neurodegenerative diseases including PD, AD, and ALS. Mutant α -synuclein in microglia is potential to switch microglia to either M1 or M2 phenotype. Microglia harboring excessive mSOD1 or TDP-43 are easily induced to be more M1-like phenotypes. Aggregated α -synuclein, mSOD1, or A β /tau oligomers released into the extracellular space from neurons can directly induce microglia towards M1 phenotypes. At the stage of disease onset, M2 microglia might be predominated to phagocytize cell debris, enhance tissue reconstruction,

and produce anti-inflammatory factors, in attempt to quench pro-inflammation and keep tissue homeostasis. However, the endogenous stimuli including aggregated α -synuclein, mSOD1, A β plaques, and tau oligomers, as well as environmental toxins persistently exist in the milieu that skew microglia into M1 phenotypes and compromise the immunoresolution process at the later stage of disease progression, which eventually leads to irreversible neuron loss. Notably, the regional distributed A β plaque are surrounded by M2 microglia that is observed till the end stage of pathology, which may complicate the changing M1 microglial phenotypes during the disease development

kinase inhibitor, skews M1 toward M2 microglia characterized by lower NF- κ B activity and pro-inflammatory cytokines (IL-1 β , IL-6 and TNF- α) as well as increases anti-inflammatory cytokines (IL-10) [84]. Administration of fasudil in the LPS model increases Arg1⁺/CD11b⁺ M2 microglia while decreases iNOS⁺/CD11b⁺ M1 microglia [84]. Inhibition of NADPH oxidase or genetic deletion of its functional p47^{phox} subunit switches microglial activation from M1 to M2 in response to LPS challenge [57]. Apart from drug treatment, an epigenetic machinery is employed to modulate M2 microglial phenotype. The histone H3K27me3 demethylase Jmjd3 was demonstrated to be essential for alternative microglial activation [45]. Suppression of Jmjd3 inhibits the expression of Arg1 and CD206 in IL-4-treated microglia, whereas enhances the production of pro-inflammatory cytokines and NO, which eventually causes more DA neuron death [45]. Since targeting M1/M2 microglia is promising to halt the PD development, the dynamic changes of microglial activation states should be explored in depth, whatever in MPTP or α -synuclein induced animal models, or most importantly in PD patients.

Microglial Phenotypes in AD

AD is the most common form of dementia, the hallmarks of which are abundant numbers of “plaques” and “tangles” by postmortem examination [85]. Senile plaques are extracellular deposits of fibrils and aggregates of β -amyloid (A β), which are derived from altered proteolytic processing of the amyloid precursor protein (APP) with the help of presenilin-1 (PS1) [85, 86]. Amyloid plaques can attract and stimulate microglia cells *in vivo* [87], and A β peptides are found to induce activation of primary microglia and stimulate the NO production *in vitro* [88, 89]. However, the activated microglia could adopt different phenotypes, which is much complicated by the presence of extracellular A β peptides and amyloid fibrils.

Microglia can be neuroprotective by degrading A β plaques as a reaction against A β accumulation [90]. However, an age-dependent increase in both the number and size of A β plaques in AD might reflect a diminution in the microglial phagocytic capability [91, 92]. Microglia surrounding the plaques for A β phagocytosis generally manifest M2 activation phenotype as labeled by Ym1 [92]. The phagocytic activity of microglia is attenuated by pro-inflammatory cytokines such as IFN- γ , IL-1 β , and TNF- α , which most likely shifts microglia into the pro-inflammatory M1 state [93].

A β peptides have the capacity to self-assemble, transforming from monomeric to oligomeric, and then to insoluble heavy aggregates. Oligomeric A β , with a molecular weight of around 56 kDa, has the most markable toxic effects on the neuronal functions. Interestingly, different forms of A β peptides may induce different inflammatory profiles of microglia. The oligomeric A β appears to be a stronger M1-inductor than the fibrillar form [94]. Moreover, a cytokine-

induced anti-inflammatory environment reduces the microglial reactivity towards oligomeric A β [94].

In the hippocampus of PS1^{M146L}/APP^{751SL} double transgenic AD mice, an age-dependent phenotypic change of microglial activation is an active processing [92]. M2 microglia with A β phagocytic capabilities in AD mice at 6 months of age can switch into M1 phenotypes at 18 months of age, which is coincident with accumulated levels of soluble A β oligomers [92]. The YM1 positive and TNF- α negative microglia cells are exclusively located surrounding and infiltrating the A β plaques (Fig. 2). Furthermore, this M2 phenotype seems to maintain even at relative old ages, suggesting that activated microglia surrounding A β plaques adopted an alternative phenotype, regardless of the age [92]. In the mice of 18 months old, microglial activation is expanded into hippocampal areas free of plaques, showing that classic M1 phenotypes producing cytotoxic effects to neurons [92].

Pro-inflammatory factors IL-1 β and IFN- γ as well as LPS suppress the microglial phagocytosis of fibrillar A β peptides, which are antagonized by anti-inflammatory cytokines including IL-4, IL-13, TGF- β , and IL-10 both *in vitro* and *in vivo* [25, 95–98]. Activation of M1 microglia results in an increase of iNOS expression. Ablation of iNOS in the APP/PS1 mice can protect the mice from plaque formation and premature mortality [99]. However, in other reports, microglial activation by acute LPS treatment reduces A β load in APP transgenic mice, which is prevented by co-treatment with dexamethasone [100–102]. APPswDI mice deficient in iNOS displayed extensive tau pathology associated with regions of dense microvascular amyloid deposition [103]. Similar effects with different pro-inflammatory mediators such as IFN- γ , TNF- α , and IL-6 were observed in the TgCRND8 mice [104–106]. Thus in the closed *in vivo* system, the role of inflammatory cytokines impairing A β clearance is still controversial, as seen not all consistent with the *in vitro* results.

Anti-inflammatory factors were believed to be promising molecules in AD therapy. Intra-cerebral injection of IL-4 and IL-13 reduced A β plaque load in APP23 mice, accompanied with improved cognition and upregulation of Arg1 and YM1 positive M2 cells [107]. There was a reduction in A β plaques and an improvement in spatial memory of APP/PS1 mice 5 months after the intrahippocampally injected AAV2 carrying IL-4 while not IL-10 [108, 109]. In another study, however, Chakrabarty et al. utilized rAAV2/1 to overexpress murine IL-4 expression in the hippocampus of TgCRND8 mice with preexisting amyloid plaques, which resulted in establishment of an “M2-like” phenotype in the brain but exacerbated amyloid deposition after 6 weeks [110]. They showed that IL-4 treatment attenuated soluble A β 40 uptake by microglia but does not affect aggregated A β 42 internalization by microglia or soluble A β 40 internalization by astrocytes [110]. This short-term focal IL-4 expression led to reduced glia phagocytosis and acute suppression of glial clearance mechanisms. It

appears that A β clearance in AD might be primed by moderate level of M1 microglial activation and maintained by M2 microglia polarization, since amyloid deposition is associated with high expression of alternative activation and acquires deactivation genes. The acute incorporation of either pro-inflammatory or anti-inflammatory factors might cause unwarranted effects.

In most cases, microglia in AD patients may exhibit mixed activation phenotypes. Colton et al. probed cortical tissue from two transgenic mouse models and from AD patients for evidence of alternative activation [46]. Interestingly, the results are always correlated in those animal models and AD patients. Cortical tissue from the Tg2576 mouse and individuals with AD demonstrate a mixed profile of alternative activation and classical activation genes, while the Tg-SwDI mouse primarily demonstrates classical activation [46]. Alternative activation genes such as Arg1, CD206, and YM1 were greatly increased in the Tg2576 mice while TNF- α and iNOS, respectively, increase or have no change. In AD patients, TNF- α , Arg1, CD206, Chi311, and Chi3112 are found significantly increased while iNOS and IL-1 β mRNA levels are unchanged [46]. Microarray analysis on brain samples from AD subjects shows up-regulation of apoptotic and pro-inflammatory signaling represented by major histocompatibility complex (MHC) class II, IFN- γ , and IL-1 β elevation [111, 112]. The discrepancies of gene expression patterns in different models and in different stages of disease suggest a very complicated cytokine environment in the brain and its role in modifying microglial responses to A β plaques. Microglia might exhibit specific dominant phenotypes during a chronic neuroinflammatory process. Hence, understanding the sequence and timing of the alterations in M1/M2 phenotypes in AD is important.

As another hallmark of AD, misfolded tau protein plays a crucial role in the formation of intracellular tangles, which is a driving force of neurofibrillary degeneration in AD or other diseases. Oligomeric tau can also be released into the extracellular space and spread throughout the brain. Activated microglia have been frequently present in the proximity of neurofibrillary tangles and tangle-bearing neurons in the hippocampus of AD patients, as well as various tau transgenic models at early and late stages of tangle formation, indicating a close relationship between inflammatory response and tau neurofibrillary lesions [113–117]. Tau oligomers and fibrils which were induced by arachidonic acid *in vitro* can augment the production of nitrites and pro-inflammatory cytokine IL-6 in microglia cells [118].

In early stages of tau structural metamorphosis, some pro-inflammatory cytokines such as IL-1, IL-6 and TNF- α , as well as fractalkine (CX3CL1) can modify tau phosphorylation patterns and alter the structure and function of tau proteins [119]. Enhanced pro-inflammatory activation of microglia by disruption of CX3CR1, an anti-inflammatory CX3CL1 receptor,

accelerates tangle formation [120]. Inflammatory changes were shown to be tau-dependent in the Tg4510 mice, which overexpress a mutant form of human tau, as they were reversed by the suppression of tau expression [121]. While treatment with immunosuppressant drug FK506 can attenuate microglial activation and delay the tau-related neuropathology in P301S tau mice [122].

In the cortical area of transgenic mice expressing P301L mutant tau (JNPL3) [123], immunoreactivity of phosphorylated tau (AT8) was closely associated with the number of Iba1-positive microglia [117]. Tau-associated microglia showed lower expression of MHC class II antigen and SRA. Interestingly, in the affected spinal cord region, JNPL3 mutant mice showed accumulation of alternatively activated microglia [124]. Tau-tubulin kinase 1 (TTBK1) expression is significantly elevated in AD brains, and directly phosphorylates tau protein, especially at Ser422 [125]. By crossing the JNPL3 mice with TTBK1 transgenic mice, there was a striking switch in activation phenotypes in the anterior horn of the spinal cord from alternative activated microglia (M2 state) in P301L tau mutant mice to pro-inflammatory infiltrating monocytes (M1 state), showing that tau phosphorylation is responsible for mediating M1-activated microglia-induced neurotoxicity [124].

Microglial Phenotypes in ALS

ALS, commonly known as Lou Gehrig's disease, is an adult-onset progressive disorder that selectively affects upper and lower motoneurons [126, 127]. The common hallmark shared by familial and sporadic ALS patients is neuroinflammation, characterized by microglial/astroglial activation and infiltration of peripheral T cells. Accumulation of misfolded human mutated forms of Cu, Zn-superoxide dismutase (mSOD1), or TAR DNA-binding protein 43 (TDP-43) seen in inherited ALS is critically contributing to the neurotoxic M1 inflammatory responses [126, 128] (Fig. 2). Mutant TDP-43^{M337V} evoked robust microglial activation around diseased motoneurons in the ventral horns [129]. Microglia expressing higher amounts of TDP-43 produced more pro-inflammatory cytokines and neurotoxic mediators after stimulation with LPS or ROS, due to the activation of p65 subunit of NF- κ B that interacts with TDP-43 as a coactivator [128]. Compared to TDP-43, SOD1 is extensively studied in the immunopathogenesis of ALS. Intracellular and extracellular mSOD1 employs different pathways to enhance the production of ROS and exaggerates the pro-inflammatory signaling in microglia [130, 131]. Whereas treatment with IL-4 suppresses M1 microglial activation by reducing the release of ROS and promotes an M2 phenotype by enhancing IGF-I secretion that improves motoneuron survival [28].

Microglia harboring excessive mSOD1 are easily induced to be more M1-like phenotypes. Following activation with

LPS, primary microglia isolated from mSOD1^{G93A} transgenic mice are more neurotoxic than activated wild-type microglia, due to the increased production of superoxide, NO, and pro-inflammatory cytokines IL-1 β and TNF- α , as well as the less release of IGF-I [132]. A lack or reduction of mSOD1 expression in microglia could slow disease progression and prolong the survival of mSOD1^{G93A} or mSOD1^{G37R} mice [133, 134].

Microglia in mSOD1^{G93A} mouse model appear to switch from an M2 microglial phenotype observed at the beginning of pathology to an M1 phenotype as disease progresses with increasing expressions of CD86, iNOS, and the NADPH oxidase isoform NOX2 and pro-inflammatory cytokines, including TNF- α , IL-1 β , and IL-6 [135, 136]. Microglia isolated from mSOD1^{G93A} mice at disease onset express higher levels of Ym1, CD163, brain-derived neurotrophic factor (BDNF), and IGF-I, and lower levels of NOX2 compared with those isolated from mSOD1^{G93A} mice at end-stage [137, 138]. This may indicate the diminished function of neuroprotective microglia in the late disease stage. What's more, end-stage M1 microglia are toxic to cocultured motoneurons, whereas those M2-like microglia at disease onset are neuroprotective to promote motoneuron survival [137].

Deletion of the pro-inflammatory factors such as NOX2 rescues almost three times motoneuron death in the spinal cords of mSOD1^{G93A} mice [139]. NF- κ B is upregulated in spinal cords of ALS patients and mSOD1^{G93A} mice. Ablation of NF- κ B signaling in microglia rescues motoneurons in vitro and extends survival in mSOD1^{G93A} mice [140]. The number of iNOS⁺/Iba1⁺ cells and CD86⁺/Iba1⁺ cells is significantly decreased in mice with microglial NF- κ B inhibition [140], while the levels of M2 markers as CD206, Arg1, or CD204 assessed by immunostaining are not altered, suggesting that NF- κ B inhibition does not promote an M2 phenotype [140]. Similarly, administration of minocycline delays the pathogenesis of mSOD1^{G93A} mice by selectively attenuating the induction of M1 microglia markers during the progressive phase, without affecting the transient enhancement of expression of M2 microglia markers at the early onset stage [141].

The Arg1/iNOS balance, however, seems to be in a more complicated condition. By immunostaining, Arg1-positive and iNOS-positive microglia were found to increase 18-fold and 7-fold, respectively, between 10 and 25 weeks of age in the lumbar spinal cord of mSOD1^{G93A} mice compared to the control mice [142]. This appears to contradict the notion that M2 microglia activation is dampened in ALS progression, which hints the complexity of Arg1/iNOS unbalance in ALS development.

Collectively, microglia have both neuroprotective and cytotoxic functions in ALS. In the first response, M2 microglia may augment a neuroprotective effect, and then following sustained neuronal stress and signaling, a transformation of microglial phenotypes happens. During the disease progression, M1 microglia predominate in the milieu, which may be promoted and

amplified by misfolded and aggregated SOD1 proteins, and eventually exacerbate motoneuron injury (Fig. 2).

Complexity of Microglial Phenotypes

Different models and different stages of disease underlie the complexity of the cytokine environment and its role in modifying microglial activation states. Thus, there exist various controversial results that link M1/M2 microglia switch and disease development. Notably, the status of M1/M2 microglia in AD is much more complicated than other two diseases, probably because of different triggers. The major trigger in AD is the extracellular A β oligomers, whereas the synuclein or SOD1 aggregation is mostly intracellular, which may play a greatest role after their release from injured or dead cells. Secondly, the regionally distributed A β plaques also make the unevenly distribution of M1/M2 microglia in the lesion area, which is different in PD and ALS. Thirdly, infiltrated peripheral macrophages or monocytes in AD are more often to penetrate into the CNS to help eliminate the extracellular oligomeric A β or tau antigens [143]. Those peripheral macrophages are also undergoing phenotypic switch that may compromise the role of microglia. Therefore, it is not easy to make a quick conclusion regarding the role of M1/M2 microglia in AD, considering the molecular mechanisms of M2 microglia are still less studied in this disease. The extra complexity arises from that M2 microglia might be also activated slightly in some models while not suppressed by M1 microglia along the disease pathogenesis. So far, the majority of studies on microglial phenotypes are focused on the various animal models, the in-depth functions of microglial phenotype switch in neurodegenerative diseases especially in AD are still very open. Establishment of an animal model showing pathology that truly represents those neurodegenerative diseases especially progressive neuronal loss in human body may partially mitigate those discrepancies.

Aging and Microglial Phenotypes

Neurodegenerative diseases are usually seen in the middle-age to elderly people. In the aged brain, many microglia cells undergo various molecular and cellular changes and even morphological features indicative of senescence, such as fragmented cytoplasmic processes, rendering them lose the ability to protect the brain [144, 145]. It is also hypothesized as "microglia dysfunction or dystrophic" that provides initial evidence for the age-associated changes in microglia. More importantly, aged microglia are also manifested by altered inflammatory profiles [146]. Normal aging in the brain is accompanied by increasing number of pro-inflammatory mediators such as IL-1 β and IL-6 while compromising IL-10 level [147–150, 45]. The DNA-binding activity of NF- κ B is increased in aged brain compared to adult and juvenile brain that

contributes to more IL-6 expression [150]. Additionally, treatment with MPTP in aged mice can cause severer DA neuron loss and greater microglial activation in the SN [151]. As the classical activation in the CNS is enhanced along aging, alternative activation appears to be mitigated, which is manifested by the reduction in the IL-4/IL-13 signaling pathway [152]. Therefore, the age-associated inflammation profiles might switch microglia phenotypes to be more M1-like, which renders aging brains are more easily affected by genetic or environmental insults during the onset of neurodegenerative diseases.

Therapeutic Perspectives


Switching of M1/M2 Phenotypes

Neurodegenerative diseases commonly implicate the failure in the resolution of neuroinflammation and presence of a defective repair process. Clinical therapy of neurodegenerative diseases faces numerous challenges with respect to timing, efficacy, and safety. Targeting any one of the vast pro-inflammatory mediators or pathways may not be efficacious, and the anti-inflammatory strategy varies in different diseases. Early stage intervention with drugs targeting dysregulated pro-inflammatory cytokine production might be therapeutically beneficial. A small molecule named MW-151 has been

tested during two distinct but overlapping therapeutic time windows at the early pathological stage in APP/PS1 transgenic mice. MW-151 treatment attenuates the microglial activation and the production of pro-inflammatory cytokines in the cortex, which protects from synaptic dysfunction implicated in learning and memory [153].

As stated above, it might be interesting to manipulate microglia phenotypes from cytotoxic to neuroprotective by drug treatment or genetic modification. To be clinically effective, targeting M1/M2 balance also depends on the time window, since the timing, stages, and severity of diseases are critically associated with the changing microglial activation states (Fig. 3). Anti-inflammatory therapies will have to gain access to the CNS; it is not realistic that cytokines are used to modulate microglia polarization because most of the cytokines cannot enter brain tissue. The chemicals as fasudil [84] and minocycline [141], which have the superior ability to cross the blood–brain barrier, have demonstrated to enhance M2 microglial anti-inflammatory responses, and importantly manipulating the demethylase Jmjd3 is capable to skew microglia towards beneficial M2 phenotype [45]. Therefore, it is necessary to further delve into the effects of this manipulation within appropriate time windows. In addition, as the cell replacement therapy emerges in recent years to replace damaged neurons with fresh ones derived from embryonic stem cells or induced pluripotent stem cells, the immunosuppressive milieu modified through M1/M2 switching might help achieve a beneficial clinical outcome.

Fig. 3 Therapeutic perspectives on switching of M1/M2 phenotypes. In the progression of neurodegenerative diseases, it might be possible to switch microglial phenotypes from cytotoxic to neuroprotective by drug treatment or genetic modification, so as to alleviate pro-inflammation and attenuate neuron loss. To be clinically effective, targeting M1/M2 balance greatly depends on the optimal therapeutic time windows, since the timing, stages, and severity of diseases are critically associated with the changing microglial phenotypes. The treatment before or after the stage of disease onset may produce different therapeutic outcomes and may also vary in different neurodegenerative diseases. *Dash lines* indicate the neuron survival and microglial activation after M1/M2 switch therapy



Too Much or too Little?

After learning the optimal time window to manipulate M2 microglia in different diseases, there is still a long way to go to understand to which extent this balance of M1/M2 may skew, neither too much nor too little. M2 microglia might resemble M2 macrophages to participate in the immunosuppressive and repair process. A long-term repair phase after a rapid pro-inflammatory response that is driven principally by M2 macrophages results in fibrosis and other aberrant repair. For example, alternatively activated alveolar macrophages contribute to the fibrotic lesion in idiopathic pulmonary fibrosis and in the liver fibrosis associated with *S. mansoni* [48, 46, 154]. Elevated arginase activity shifts the metabolism of L-arginine dramatically to produce increased ornithine and proline that stimulate cell division leading to hyperplasia and fibrosis, and meanwhile an uncoupling decrease in NO production results in endothelial dysfunction [154, 155]. Moreover, loss of iNOS and NO also dampens the effectiveness of the innate immune response against bacteria and virus, since NO-mediated modification of bacterial proteins is an efficient way to kill bacteria [156]. Considering the M1/M2 microglia switching requires fine regulation, more in-depth investigations are urgently needed.

Conclusion

The demonstrated role of microglial phenotypes may boost the research of M1/M2 paradigm in the human body. The balance of M1 and M2 microglial activation is broken down during the chronic inflammation progress in neurodegenerative diseases, with the highest complexity in AD. Setting a standard for measuring M1/M2 ratio might be critical, since M2 microglia also increases to a certain extent on some occasions. The endogenous stimuli including aggregated α -synuclein, mSOD1, and A β plaques persistently exist in the milieu that compromise the immunoresolution process and finally lead to irreversible neuron loss. Thus, stage-specific switching of the M1/M2 microglial phenotypes within appropriate time windows may produce therapeutic benefits.

Acknowledgments This work was supported by grants from the National Natural Sciences Foundation of China (No. 81171201) and the National Basic Research Program of China (No. 2011CB510003).

Conflict of Interest The authors declare no conflicts of interest.

References

- Gao HM, Hong JS (2008) Why neurodegenerative diseases are progressive: uncontrolled inflammation drives disease progression. *Trends Immunol* 29(8):357–365
- Glass CK, Saijo K, Winner B, Marchetto MC, Gage FH (2010) Mechanisms underlying inflammation in neurodegeneration. *Cell* 140(6):918–934
- Minghetti L (2005) Role of inflammation in neurodegenerative diseases. *Curr Opin Neurol* 18(3):315–321
- Block ML, Hong JS (2005) Microglial and inflammation-mediated neurodegeneration: multiple triggers with a common mechanism. *Prog Neurobiol* 76(2):77–98
- Ouchi Y, Yoshikawa E, Sekine Y, Futatsubashi M, Kanno T, Ogusu T et al (2005) Microglial activation and dopamine terminal loss in early Parkinson's disease. *Ann Neurol* 57(2):168–175
- Block ML, Zecca L, Hong JS (2007) Microglia-mediated neurotoxicity: uncovering the molecular mechanisms. *Nat Rev Neurosci* 8(1):57–69
- Gerhard A, Pavese N, Hotton G, Turkheimer F, Es M, Hammers A et al (2006) In vivo imaging of microglial activation with [¹¹C](R)-PK11195 PET in idiopathic Parkinson's disease. *Neurobiol Dis* 21(2):404–412
- Hanisch UK, Kettenmann H (2007) Microglia: active sensor and versatile effector cells in the normal and pathologic brain. *Nat Neurosci* 10(11):1387–1394
- Tang Y, Le W (2014) "Good" and "bad" microglia in Parkinson's disease: an understanding of homeostatic mechanisms in immunomodulation. In: Thomas M (ed) *Inflammation in Parkinson's disease*. Springer, New York, pp 105–126
- Glezer I, Simard AR, Rivest S (2007) Neuroprotective role of the innate immune system by microglia. *Neuroscience* 147(4):867–883
- Simard AR, Soulet D, Gowing G, Julien JP, Rivest S (2006) Bone marrow-derived microglia play a critical role in restricting senile plaque formation in Alzheimer's disease. *Neuron* 49(4):489–502
- Ding YM, Jaumotte JD, Signore AP, Zigmond MJ (2004) Effects of 6-hydroxydopamine on primary cultures of substantia nigra: specific damage to dopamine neurons and the impact of glial cell line-derived neurotrophic factor. *J Neurochem* 89(3):776–787
- Lumeng CN, Bodzin JL, Saltiel AR (2007) Obesity induces a phenotypic switch in adipose tissue macrophage polarization. *J Clin Invest* 117(1):175–184
- Lumeng CN, Deyoung SM, Bodzin JL, Saltiel AR (2007) Increased inflammatory properties of adipose tissue macrophages recruited during diet-induced obesity. *Diabetes* 56(1):16–23
- Odegaard JI, Ricardo-Gonzalez RR, Goforth MH, Morel CR, Subramanian V, Mukundan L et al (2007) Macrophage-specific PPAR γ controls alternative activation and improves insulin resistance. *Nature* 447(7148):1116–1120
- Hevener AL, Olefsky JM, Reichart D, Nguyen MT, Bandyopadhyay G, Leung HY et al (2007) Macrophage PPAR γ is required for normal skeletal muscle and hepatic insulin sensitivity and full antidiabetic effects of thiazolidinediones. *J Clin Invest* 117(6):1658–1669
- Kigerl KA, Gensel JC, Ankeny DP, Alexander JK, Donnelly DJ, Popovich PG (2009) Identification of two distinct macrophage subsets with divergent effects causing either neurotoxicity or regeneration in the injured mouse spinal cord. *J Neurosci* 29(43):13435–13444
- Colton C, Wilcock DM (2010) Assessing activation states in microglia. *CNS Neurol Disord Drug Targets* 9(2):174–191
- Colton CA (2009) Heterogeneity of microglial activation in the innate immune response in the brain. *J Neuroimmune Pharmacol* 4(4):399–418

20. Le W, Rowe D, Xie W, Ortiz I, He Y, Appel SH (2001) Microglial activation and dopaminergic cell injury: an in vitro model relevant to Parkinson's disease. *J Neurosci* 21(21):8447–8455
21. Li R, Huang YG, Fang D, Le WD (2004) (-)-Epigallocatechin gallate inhibits lipopolysaccharide-induced microglial activation and protects against inflammation-mediated dopaminergic neuronal injury. *J Neurosci Res* 78(5):723–731
22. Ponomarev ED, Maresz K, Tan Y, Dittel BN (2007) CNS-derived interleukin-4 is essential for the regulation of autoimmune inflammation and induces a state of alternative activation in microglial cells. *J Neurosci* 27(40):10714–10721
23. Sawada M, Suzumura A, Hosoya H, Marunouchi T, Nagatsu T (1999) Interleukin-10 inhibits both production of cytokines and expression of cytokine receptors in microglia. *J Neurochem* 72(4):1466–1471
24. Gao HM, Liu B, Zhang WQ, Hong JS (2003) Critical role of microglial NADPH oxidase-derived free radicals in the in vitro MPTP model of Parkinson's disease. *FASEB J* 17(11):1954–
25. Butovsky O, Talpalar AE, Ben-Yaakov K, Schwartz M (2005) Activation of microglia by aggregated beta-amyloid or lipopolysaccharide impairs MHC-II expression and renders them cytotoxic whereas IFN-gamma and IL-4 render them protective. *Mol Cell Neurosci* 29(3):381–393
26. Zhou XL, Spittau B, Kriegelstein K (2012) TGF beta signalling plays an important role in IL4-induced alternative activation of microglia. *J Neuroinflamm* 9:210–223
27. Ledebor A, Breve JJ, Poole S, Tilders FJ, Van Dam AM (2000) Interleukin-10, interleukin-4, and transforming growth factor-beta differentially regulate lipopolysaccharide-induced production of pro-inflammatory cytokines and nitric oxide in co-cultures of rat astroglial and microglial cells. *Glia* 30(2):134–142
28. Zhao WH, Xie WJ, Xiao Q, Beers DR, Appel SH (2006) Protective effects of an anti-inflammatory cytokine, interleukin-4, on motoneuron toxicity induced by activated microglia. *J Neurochem* 99(4):1176–1187
29. Park KW, Lee DY, Joe EH, Kim SU, Jin BK (2005) Neuroprotective role of microglia expressing interleukin-4. *J Neurosci Res* 81(3):397–402
30. Boche D, Cunningham C, Docagne F, Scott H, Perry VH (2006) TGFbeta1 regulates the inflammatory response during chronic neurodegeneration. *Neurobiol Dis* 22(3):638–650
31. Boche D, Cunningham C, Gaudie J, Perry VH (2003) Transforming growth factor-beta 1-mediated neuroprotection against excitotoxic injury in vivo. *J Cereb Blood Flow Metab* 23(10):1174–1182
32. Bogdan C, Vodovotz Y, Nathan C (1991) Macrophage deactivation by interleukin 10. *J Exp Med* 174(6):1549–1555
33. Buechler C, Ritter M, Orso E, Langmann T, Klucken J, Schmitz G (2000) Regulation of scavenger receptor CD163 expression in human monocytes and macrophages by pro- and anti-inflammatory stimuli. *J Leukoc Biol* 67(1):97–103
34. Fabriek BO, Dijkstra CD, van den Berg TK (2005) The macrophage scavenger receptor CD163. *Immunobiology* 210(2–4):153–160
35. Suh HS, Zhao ML, Derico L, Choi N, Lee SC (2013) Insulin-like growth factor 1 and 2 (IGF1, IGF2) expression in human microglia: differential regulation by inflammatory mediators. *J Neuroinflamm* 10:37
36. Liu Y, Hao W, Letiembre M, Walter S, Kulanga M, Neumann H et al (2006) Suppression of microglial inflammatory activity by myelin phagocytosis: role of p47-PHOX-mediated generation of reactive oxygen species. *J Neurosci* 26(50):12904–12913
37. Gordon S (2003) Alternative activation of macrophages. *Nat Rev Immunol* 3(1):23–35
38. Morris SM (2004) Recent advances in arginine metabolism. *Curr Opin Clin Nutr* 7(1):45–51
39. Bronte V, Zanovello P (2005) Regulation of immune responses by L-arginine metabolism. *Nat Rev Immunol* 5(8):641–654
40. Busch SA, Silver J (2007) The role of extracellular matrix in CNS regeneration. *Curr Opin Neurobiol* 17(1):120–127
41. Wu GY, Bazer FW, Burghardt RC, Johnson GA, Kim SW, Knabe DA et al (2011) Proline and hydroxyproline metabolism: implications for animal and human nutrition. *Amino Acids* 40(4):1053–1063
42. Jenkins CL, Bretscher LE, Guzei IA, Raines RT (2003) Effect of 3-hydroxyproline residues on collagen stability. *J Am Chem Soc* 125(21):6422–6427
43. Thomas T, Thomas TJ (2001) Polyamines in cell growth and cell death: molecular mechanisms and therapeutic applications. *Cell Mol Life Sci* 58(2):244–258
44. Williams K (1997) Interactions of polyamines with ion channels. *Biochem J* 325:289–297
45. Tang Y, Li T, Li J, Yang J, Liu H, Zhang XJ et al (2014) Jmjd3 is essential for the epigenetic modulation of microglia phenotypes in the immune pathogenesis of Parkinson's disease. *Cell Death Differ* 21(3):369–380
46. Colton CA, Mott RT, Sharpe H, Xu Q, Van Nostrand WE, Vitek MP (2006) Expression profiles for macrophage alternative activation genes in AD and in mouse models of AD. *J Neuroinflamm* 3:27
47. Nair MG, Du YR, Perrigoue JG, Zaph C, Taylor JJ, Goldschmidt M et al (2009) Alternatively activated macrophage-derived RELM-alpha is a negative regulator of type 2 inflammation in the lung (vol 206, pg 397, 2009). *J Exp Med* 206(5):1201
48. Holcomb IN, Kabakoff RC, Chan B, Baker TW, Gurney A, Henzel W et al (2000) FIZZ1, a novel cysteine-rich secreted protein associated with pulmonary inflammation, defines a new gene family. *EMBO J* 19(15):4046–4055
49. Munitz A, Seidu L, Cole ET, Ahrens R, Hogan SP, Rothenberg ME (2009) Resistin-like molecule a decreases glucose tolerance during intestinal inflammation. *J Immunol* 182(4):2357–2363
50. Munitz A, Waddell A, Seidu L, Ahrens R, Hogan SP, Rothenberg ME (2008) Resistin-like molecule alpha enhances myeloid cell activation and promotes colitis. *J Allergy Clin Immunol* 122(6):1200–1207
51. Chang NCA, Hung SI, Hwa KY, Kato I, Chen JE, Liu CH et al (2001) A macrophage protein, Ym1, transiently expressed during inflammation is a novel mammalian lectin. *J Biol Chem* 276(20):17497–17506
52. Hung SL, Chang AC, Kato I, Chang NCA (2002) Transient expression of Ym1, a heparin-binding lectin, during developmental hematopoiesis and inflammation. *J Leukocyte Biol* 72(1):72–82
53. Recklies AD, White C, Ling H (2002) The chitinase 3-like protein human cartilage glycoprotein 39 (HC-gp39) stimulates proliferation of human connective-tissue cells and activates both extracellular signal-regulated kinase- and protein kinase beta-mediated signalling pathways. *Biochem J* 365:119–126
54. Raes G, De Baetselier P, Noel W, Beschin A, Brombacher F, Hassanzadeh G (2002) Differential expression of FIZZ1 and Ym1 in alternatively versus classically activated macrophages. *J Leukocyte Biol* 71(4):597–602
55. Welch JS, Escoubet-Lozach L, Sykes DB, Liddiard K, Greaves DR, Glass CK (2002) T(H)2 cytokines and allergic challenge induce YM1 expression in macrophages by a STAT6-dependent mechanism. *J Biol Chem* 277(45):42821–42829
56. Lee E, Yook J, Haa K, Chang HW (2005) Induction of Ym1/2 in mouse bone marrow-derived mast cells by IL-4 and identification of Ym1/2 in connective tissue type-like mast cells derived from bone marrow cells cultured with IL-4 and stem cell factor. *Immunol Cell Biol* 83(5):468–474
57. Choi SH, Aid S, Kim HW, Jackson SH, Bosetti F (2012) Inhibition of NADPH oxidase promotes alternative and anti-inflammatory

- microglial activation during neuroinflammation. *J Neurochem* 120(2):292–301
58. Taylor PR, Gordon S, Martinez-Pomares L (2005) The mannose receptor: linking homeostasis and immunity through sugar recognition. *Trends Immunol* 26(2):104–110
 59. Stahl PD, Ezekowitz RAB (1998) The mannose receptor is a pattern recognition receptor involved in host defense. *Curr Opin Immunol* 10(1):50–55
 60. Lee SJ, Evers S, Roeder D, Parlow AF, Risteli J, Risteli L et al (2002) Mannose receptor-mediated regulation of serum glycoprotein homeostasis. *Science* 295(5561):1898–1901
 61. Chieppa M, Bianchi G, Doni A, Del Prete A, Sironi M, Laskarin G et al (2003) Cross-linking of the mannose receptor on monocyte-derived dendritic cells activates an anti-inflammatory immunosuppressive program. *J Immunol* 171(9):4552–4560
 62. Kerrigan AM, Brown GD (2009) C-type lectins and phagocytosis. *Immunobiology* 214(7):562–575
 63. Kigerl KA, Gensel JC, Ankeny DP, Alexander JK, Donnelly DJ, Popovich PG (2009) Identification of two distinct macrophage subsets with divergent effects causing either neurotoxicity or regeneration in the injured mouse spinal cord. *J Neurosci* 29(43):13435–13444
 64. Madsen DH, Leonard D, Masedunskas A, Moyer A, Jurgensen HJ, Peters DE et al (2013) M2-like macrophages are responsible for collagen degradation through a mannose receptor-mediated pathway. *J Cell Biol* 202(6):951–966
 65. Dauer W, Przedborski S (2003) Parkinson's disease: mechanisms and models. *Neuron* 39(6):889–909
 66. Braak H, Del Tredici K, Rub U, de Vos RA, Jansen Steur EN, Braak E (2003) Staging of brain pathology related to sporadic Parkinson's disease. *Neurobiol Aging* 24(2):197–211
 67. Lashuel HA, Overk CR, Oueslati A, Masliah E (2013) The many faces of alpha-synuclein: from structure and toxicity to therapeutic target. *Nat Rev Neurosci* 14(1):38–48
 68. Polymeropoulos MH, Lavedan C, Leroy E, Ide SE, Dehejia A, Dutra A et al (1997) Mutation in the alpha-synuclein gene identified in families with Parkinson's disease. *Science* 276(5321):2045–2047
 69. Klein C, Westenberger A (2012) Genetics of Parkinson's disease. *CSH Perspect Med* 2(1)
 70. Sanchez-Guajardo V, Barnum CJ, Tansey MG, Romero-Ramos M (2013) Neuroimmunological processes in Parkinson's disease and their relation to alpha-synuclein: microglia as the referee between neuronal processes and peripheral immunity. *ASN Neurol* 5(2):113–139
 71. Zhang W, Wang T, Pei Z, Miller DS, Wu X, Block ML et al (2005) Aggregated alpha-synuclein activates microglia: a process leading to disease progression in Parkinson's disease. *FASEB J* 19(6):533–542
 72. Lee EJ, Woo MS, Moon PG, Baek MC, Choi IY, Kim WK et al (2010) Alpha-synuclein activates microglia by inducing the expressions of matrix metalloproteinases and the subsequent activation of protease-activated receptor-1. *J Immunol* 185(1):615–623
 73. Zhang W, Dallas S, Zhang D, Guo JP, Pang H, Wilson B et al (2007) Microglial PHOX and Mac-1 are essential to the enhanced dopaminergic neurodegeneration elicited by A30P and A53T mutant alpha-synuclein. *Glia* 55(11):1178–1188
 74. Reynolds AD, Kadiu I, Garg SK, Glanzer JG, Nordgren T, Ciborowski P et al (2008) Nitrated alpha-synuclein and microglial neuroregulatory activities. *J Neuroimmune Pharmacol* 3(2):59–74
 75. Gao HM, Kotzbauer PT, Uryu K, Leight S, Trojanowski JQ, Lee VM (2008) Neuroinflammation and oxidation/nitration of alpha-synuclein linked to dopaminergic neurodegeneration. *J Neurosci* 28(30):7687–7698
 76. Rojanathammanee L, Murphy EJ, Combs CK (2011) Expression of mutant alpha-synuclein modulates microglial phenotype in vitro. *J Neuroinflamm* 8:44
 77. Austin SA, Floden AM, Murphy EJ, Combs CK (2006) Alpha-synuclein expression modulates microglial activation phenotype. *J Neurosci* 26(41):10558–10563
 78. Porras G, Li Q, Bezard E (2012) Modeling Parkinson's disease in primates: the MPTP model. *CSH Perspect Med* 2(3)
 79. Wu DC, Teismann P, Tieu K, Vila M, Jackson-Lewis V, Ischiropoulos H et al (2003) NADPH oxidase mediates oxidative stress in the 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine model of Parkinson's disease. *Proc Natl Acad Sci U S A* 100(10):6145–6150
 80. Zhang W, Wang T, Qin L, Gao HM, Wilson B, Ali SF et al (2004) Neuroprotective effect of dextromethorphan in the MPTP Parkinson's disease model: role of NADPH oxidase. *FASEB J* 18(3):589–591
 81. Liberatore GT, Jackson-Lewis V, Vukosavic S, Mandir AS, Vila M, McAuliffe WG et al (1999) Inducible nitric oxide synthase stimulates dopaminergic neurodegeneration in the MPTP model of Parkinson disease. *Nat Med* 5(12):1403–1409
 82. Gao HM, Jiang J, Wilson B, Zhang W, Hong JS, Liu B (2002) Microglial activation-mediated delayed and progressive degeneration of rat nigral dopaminergic neurons: relevance to Parkinson's disease. *J Neurochem* 81(6):1285–1297
 83. Theodore S, Cao S, McLean PJ, Standaert DG (2008) Targeted overexpression of human alpha-synuclein triggers microglial activation and an adaptive immune response in a mouse model of Parkinson disease. *J Neuropathol Exp Neurol* 67(12):1149–1158
 84. Zhang H, Li Y, Yu J, Guo M, Meng J, Liu C et al (2013) Rho kinase inhibitor fasudil regulates microglia polarization and function. *Neuroimmunomodulat* 20(6):313–322
 85. Mattson MP (2004) Pathways towards and away from Alzheimer's disease. *Nature* 430(7000):631–639
 86. Citron M, Oltersdorf T, Haass C, Mconlogue L, Hung AY, Seubert P et al (1992) Mutation of the beta-amyloid precursor protein in familial Alzheimer's-disease increases beta-protein production. *Nature* 360(6405):672–674
 87. Meyer-Luehmann M, Spires-Jones TL, Prada C, Garcia-Alloza M, de Calignon A, Rozkalne A et al (2008) Rapid appearance and local toxicity of amyloid-beta plaques in a mouse model of Alzheimer's disease. *Nature* 451(7179):720–724
 88. Maezawa I, Zimin PI, Wulff H, Jin LW (2011) Amyloid-beta protein oligomer at low nanomolar concentrations activates microglia and induces microglial neurotoxicity. *J Biol Chem* 286(5):3693–3706
 89. Walker DG, Link J, Lue LF, Dalsing-Hernandez JE, Boyes BE (2006) Gene expression changes by amyloid beta peptide-stimulated human postmortem brain microglia identify activation of multiple inflammatory processes. *J Leukoc Biol* 79(3):596–610
 90. Takata K, Kitamura Y, Saeki M, Terada M, Kagitani S, Kitamura R et al (2010) Galantamine-induced amyloid- β clearance mediated via stimulation of microglial nicotinic acetylcholine receptors. *J Biol Chem* 285(51):40180–40191
 91. Mawuenyega KG, Sigurdson W, Ovod V, Munsell L, Kasten T, Morris JC et al (2010) Decreased clearance of CNS beta-amyloid in Alzheimer's disease. *Science* 330(6012):1774
 92. Jimenez S, Baglietto-Vargas D, Caballero C, Moreno-Gonzalez I, Torres M, Sanchez-Varo R et al (2008) Inflammatory response in the hippocampus of PS1M146L/APP751SL mouse model of Alzheimer's disease: age-dependent switch in the microglial phenotype from alternative to classic. *J Neurosci* 28(45):11650–11661
 93. Koenigsknecht-Talboo J, Landreth GE (2005) Microglial phagocytosis induced by fibrillar beta-amyloid and IgGs are differentially regulated by proinflammatory cytokines. *J Neurosci* 25(36):8240–8249
 94. Michelucci A, Heurtaux T, Grandbarbe L, Morga E, Heuschling P (2009) Characterization of the microglial phenotype under specific pro-inflammatory and anti-inflammatory conditions: effects of oligomeric and fibrillar amyloid-beta. *J Neuroimmunol* 210(1–2):3–12

95. Butovsky O, Koronyo-Hamaoui M, Kunis G, Ophir E, Landa G, Cohen H et al (2006) Glatiramer acetate fights against Alzheimer's disease by inducing dendritic-like microglia expressing insulin-like growth factor 1. *Proc Natl Acad Sci U S A* 103(31):11784–11789
96. Lyons A, Griffin RJ, Costelloe CE, Clarke RM, Lynch MA (2007) IL-4 attenuates the neuroinflammation induced by amyloid-beta in vivo and in vitro. *J Neurochem* 101(3):771–781
97. Iribarren P, Chen K, Hu J, Zhang X, Gong W, Wang JM (2005) IL-4 inhibits the expression of mouse formyl peptide receptor 2, a receptor for amyloid beta1-42, in TNF-alpha-activated microglia. *J Immunol* 175(9):6100–6106
98. Tichauer JE, von Bernhardi R (2012) Transforming growth factor-beta stimulates beta amyloid uptake by microglia through Smad3-dependent mechanisms. *J Neurosci Res* 90(10):1970–1980
99. Nathan C, Calingasan N, Nezezon J, Ding A, Lucia MS, La Perle K et al (2005) Protection from Alzheimer's-like disease in the mouse by genetic ablation of inducible nitric oxide synthase. *J Exp Med* 202(9):1163–1169
100. Herber DL, Mercer M, Roth LM, Symmonds K, Maloney J, Wilson N et al (2007) Microglial activation is required for Abeta clearance after intracranial injection of lipopolysaccharide in APP transgenic mice. *J Neuroimmune Pharmacol* 2(2):222–231
101. Herber DL, Roth LM, Wilson N, Wilson N, Mason JE, Morgan D et al (2004) Time-dependent reduction in Abeta levels after intracranial LPS administration in APP transgenic mice. *Exp Neurol* 190(1):245–253
102. DiCarlo G, Wilcock D, Henderson D, Gordon M, Morgan D (2001) Intrahippocampal LPS injections reduce Abeta load in APP + PS1 transgenic mice. *Neurobiol Aging* 22(6):1007–1012
103. Wilcock DM, Lewis MR, Van Nostrand WE, Davis J, Previti ML, Gharkholonareh N et al (2008) Progression of amyloid pathology to Alzheimer's disease pathology in an amyloid precursor protein transgenic mouse model by removal of nitric oxide synthase 2. *J Neurosci* 28(7):1537–1545
104. Chakrabarty P, Ceballos-Diaz C, Beccard A, Janus C, Dickson D, Golde TE et al (2010) IFN-gamma promotes complement expression and attenuates amyloid plaque deposition in amyloid beta precursor protein transgenic mice. *J Immunol* 184(9):5333–5343
105. Chakrabarty P, Jansen-West K, Beccard A, Ceballos-Diaz C, Levites Y, Verbeeck C et al (2010) Massive gliosis induced by interleukin-6 suppresses Abeta deposition in vivo: evidence against inflammation as a driving force for amyloid deposition. *FASEB J* 24(2):548–559
106. Chakrabarty P, Herring A, Ceballos-Diaz C, Das P, Golde TE (2011) Hippocampal expression of murine TNFalpha results in attenuation of amyloid deposition in vivo. *Mol Neurodegener* 6:16
107. Kawahara K, Suenobu M, Yoshida A, Koga K, Hyodo A, Ohtsuka H et al (2012) Intracerebral microinjection of interleukin-4/interleukin-13 reduces beta-amyloid accumulation in the ipsilateral side and improves cognitive deficits in young amyloid precursor protein 23 mice. *Neuroscience* 207:243–260
108. Kiyota T, Okuyama S, Swan RJ, Jacobsen MT, Gendelman HE, Ikezu T (2010) CNS expression of anti-inflammatory cytokine interleukin-4 attenuates Alzheimer's disease-like pathogenesis in APP + PS1 bigenic mice. *FASEB J* 24(8):3093–3102
109. Kiyota T, Ingraham KL, Swan RJ, Jacobsen MT, Andrews SJ, Ikezu T (2012) AAV serotype 2/1-mediated gene delivery of anti-inflammatory interleukin-10 enhances neurogenesis and cognitive function in APP + PS1 mice. *Gene Ther* 19(7):724–733
110. Chakrabarty P, Tianbai L, Herring A, Ceballos-Diaz C, Das P, Golde TE (2012) Hippocampal expression of murine IL-4 results in exacerbation of amyloid deposition. *Mol Neurodegener* 7:36
111. Colangelo V, Schurr J, Ball MJ, Pelaez RP, Bazan NG, Lukiw WJ (2002) Gene expression profiling of 12633 genes in Alzheimer hippocampal CA1: transcription and neurotrophic factor down-regulation and up-regulation of apoptotic and pro-inflammatory signaling. *J Neurosci Res* 70(3):462–473
112. Blalock EM, Geddes JW, Chen KC, Porter NM, Markesbery WR, Landfield PW (2004) Incipient Alzheimer's disease: microarray correlation analyses reveal major transcriptional and tumor suppressor responses. *Proc Natl Acad Sci U S A* 101(7):2173–2178
113. Serrano-Pozo A, Mielke ML, Gomez-Isla T, Betensky RA, Growdon JH, Frosch MP et al (2011) Reactive glia not only associates with plaques but also parallels tau tangles in Alzheimer's disease. *Am J Pathol* 179(3):1373–1384
114. Ikeda M, Shoji M, Kawarai T, Kawarabayashi T, Matsubara E, Murakami T et al (2005) Accumulation of filamentous tau in the cerebral cortex of human tau R406W transgenic mice. *Am J Pathol* 166(2):521–531
115. Zilka N, Stozicka Z, Kovac A, Pilipcinec E, Bugos O, Novak M (2009) Human misfolded truncated tau protein promotes activation of microglia and leukocyte infiltration in the transgenic rat model of tauopathy. *J Neuroimmunol* 209(1–2):16–25
116. Kovac A, Zilka N, Kazmerova Z, Cente M, Zilkova M, Novak M (2011) Misfolded truncated protein tau induces innate immune response via MAPK pathway. *J Immunol* 187(5):2732–2739
117. Sasaki A, Kawarabayashi T, Murakami T, Matsubara E, Ikeda M, Hagiwara H et al (2008) Microglial activation in brain lesions with tau deposits: comparison of human tauopathies and tau transgenic mice TgTauP301L. *Brain Res* 1214:159–168
118. Morales I, Jimenez JM, Mancilla M, Maccioni RB (2013) Tau oligomers and fibrils induce activation of microglial cells. *J Alzheimers Dis* 37(4):849–856
119. Zilka N, Kazmerova Z, Jadhav S, Neradil P, Madari A, Obetkova D et al (2012) Who fans the flames of Alzheimer's disease brains? Misfolded tau on the crossroad of neurodegenerative and inflammatory pathways. *J Neuroinflamm* 9:47
120. Bhaskar K, Konerth M, Kokiko-Cochran ON, Cardona A, Ransohoff RM, Lamb BT (2010) Regulation of tau pathology by the microglial fractalkine receptor. *Neuron* 68(1):19–31
121. Wes PD, Easton A, Corradi J, Barten DM, Devidze N, DeCarr LB et al (2014) Tau overexpression impacts a neuroinflammation gene expression network perturbed in Alzheimer's disease. *PLoS One* 9(8):e106050
122. Yoshiyama Y, Higuchi M, Zhang B, Huang SM, Iwata N, Saido TC et al (2007) Synapse loss and microglial activation precede tangles in a P301S tauopathy mouse model. *Neuron* 53(3):337–351
123. Lewis J, Dickson DW, Lin WL, Chisholm L, Corral A, Jones G et al (2001) Enhanced neurofibrillary degeneration in transgenic mice expressing mutant tau and APP. *Science* 293(5534):1487–1491
124. Asai H, Ikezu S, Woodbury ME, Yonemoto GM, Cui L, Ikezu T (2014) Accelerated neurodegeneration and neuroinflammation in transgenic mice expressing P301L tau mutant and tau-tubulin kinase 1. *Am J Pathol* 184(3):808–818
125. Ikezu S, Ikezu T (2014) Tau-tubulin kinase. *Front Mol Neurosci* 7:33
126. Boillee S, Vande Velde C, Cleveland DW (2006) ALS: a disease of motor neurons and their nonneuronal neighbors. *Neuron* 52(1):39–59
127. Appel SH, Beers DR, Henkel JS (2010) T cell-microglial dialogue in Parkinson's disease and amyotrophic lateral sclerosis: are we listening? *Trends Immunol* 31(1):7–17
128. Swarup V, Phaneuf D, Dupre N, Petri S, Strong M, Kriz J et al (2011) Deregulation of TDP-43 in amyotrophic lateral sclerosis triggers nuclear factor kappaB-mediated pathogenic pathways. *J Exp Med* 208(12):2429–2447
129. Huang C, Tong J, Bi F, Zhou H, Xia XG (2012) Mutant TDP-43 in motor neurons promotes the onset and progression of ALS in rats. *J Clin Invest* 122(1):107–118

130. Henkel JS, Beers DR, Zhao WH, Appel SH (2009) Microglia in ALS: the good, the bad, and the resting. *J Neuroimmune Pharm* 4(4):389–398
131. Philips T, Robberecht W (2011) Neuroinflammation in amyotrophic lateral sclerosis: role of glial activation in motor neuron disease. *Lancet Neurol* 10(3):253–263
132. Xiao Q, Zhao W, Beers DR, Yen AA, Xie W, Henkel JS et al (2007) Mutant SOD1(G93A) microglia are more neurotoxic relative to wild-type microglia. *J Neurochem* 102(6):2008–2019
133. Boillee S, Yamanaka K, Lobsiger CS, Copeland NG, Jenkins NA, Kassiotis G et al (2006) Onset and progression in inherited ALS determined by motor neurons and microglia. *Science* 312(5778):1389–1392
134. Beers DR, Henkel JS, Xiao Q, Zhao W, Wang J, Yen AA et al (2006) Wild-type microglia extend survival in PU.1 knockout mice with familial amyotrophic lateral sclerosis. *Proc Natl Acad Sci U S A* 103(43):16021–16026
135. Beers DR, Henkel JS, Zhao W, Wang J, Appel SH (2008) CD4+ T cells support glial neuroprotection, slow disease progression, and modify glial morphology in an animal model of inherited ALS. *Proc Natl Acad Sci U S A* 105(40):15558–15563
136. Hensley K, Fedynyshyn J, Ferrell S, Floyd RA, Gordon B, Grammas P et al (2003) Message and protein-level elevation of tumor necrosis factor alpha (TNF alpha) and TNF alpha-modulating cytokines in spinal cords of the G93A-SOD1 mouse model for amyotrophic lateral sclerosis. *Neurobiol Dis* 14(1):74–80
137. Liao B, Zhao W, Beers DR, Henkel JS, Appel SH (2012) Transformation from a neuroprotective to a neurotoxic microglial phenotype in a mouse model of ALS. *Exp Neurol* 237(1):147–152
138. Kawamura MF, Yamasaki R, Kawamura N, Tateishi T, Nagara Y, Matsushita T et al (2012) Impaired recruitment of neuroprotective microglia and T cells during acute neuronal injury coincides with increased neuronal vulnerability in an amyotrophic lateral sclerosis model. *Exp Neurol* 234(2):437–445
139. Marden JJ, Harraz MM, Williams AJ, Nelson K, Luo M, Paulson H et al (2007) Redox modifier genes in amyotrophic lateral sclerosis in mice. *J Clin Invest* 117(10):2913–2919
140. Frakes AE, Ferraiuolo L, Haidet-Phillips AM, Schmelzer L, Braun L, Miranda CJ et al (2014) Microglia induce motor neuron death via the classical NF-kappaB pathway in amyotrophic lateral sclerosis. *Neuron* 81(5):1009–1023
141. Kobayashi K, Imagama S, Ohgomori T, Hirano K, Uchimura K, Sakamoto K et al (2013) Minocycline selectively inhibits M1 polarization of microglia. *Cell Death Dis* 4:e525
142. Lewis KE, Rasmussen AL, Bennett W, King A, West AK, Chung RS et al (2014) Microglia and motor neurons during disease progression in the SOD1G93A mouse model of amyotrophic lateral sclerosis: changes in arginase1 and inducible nitric oxide synthase. *J Neuroinflamm* 11:55
143. Majerova P, Zilkova M, Kazmerova Z, Kovac A, Paholikova K, Kovacech B et al (2014) Microglia display modest phagocytic capacity for extracellular tau oligomers. *J Neuroinflamm* 11(1):161
144. Streit WJ, Braak H, Xue QS, Bechmann I (2009) Dystrophic (senescent) rather than activated microglial cells are associated with tau pathology and likely precede neurodegeneration in Alzheimer's disease. *Acta Neuropathol* 118(4):475–485
145. Luo XG, Chen SD (2012) The changing phenotype of microglia from homeostasis to disease. *Transl Neurodegener* 1(1):9
146. Lucin KM, Wyss-Coray T (2009) Immune activation in brain aging and neurodegeneration: too much or too little? *Neuron* 64(1):110–122
147. Lee CK, Weindruch R, Prolla TA (2000) Gene-expression profile of the ageing brain in mice. *Nat Genet* 25(3):294–297
148. Sheng JG, Mrak RE, Griffin WS (1998) Enlarged and phagocytic, but not primed, interleukin-1 alpha-immunoreactive microglia increase with age in normal human brain. *Acta Neuropathol* 95(3):229–234
149. Ye SM, Johnson RW (1999) Increased interleukin-6 expression by microglia from brain of aged mice. *J Neuroimmunol* 93(1–2):139–148
150. Ye SM, Johnson RW (2001) An age-related decline in interleukin-10 may contribute to the increased expression of interleukin-6 in brain of aged mice. *Neuroimmunomodulat* 9(4):183–192
151. Sugama S, Yang L, Cho BP, DeGiorgio LA, Lorenzl S, Albers DS et al (2003) Age-related microglial activation in 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP)-induced dopaminergic neurodegeneration in C57BL/6 mice. *Brain Res* 964(2):288–294
152. Lee DC, Ruiz CR, Lebson L, Selenica MLB, Rizer J, Hunt JB et al (2013) Aging enhances classical activation but mitigates alternative activation in the central nervous system. *Neurobiol Aging* 34(6):1610–1620
153. Bachstetter AD, Norris CM, Sompol P, Wilcock DM, Goulding D, Neltner JH et al (2012) Early stage drug treatment that normalizes proinflammatory cytokine production attenuates synaptic dysfunction in a mouse model that exhibits age-dependent progression of Alzheimer's disease-related pathology. *J Neurosci* 32(30):10201–10210
154. Hesse M, Modolell M, La Flamme AC, Schito M, Fuentes JM, Cheever AW et al (2001) Differential regulation of nitric oxide synthase-2 and arginase-1 by type 1/type 2 cytokines in vivo: granulomatous pathology is shaped by the pattern of L-arginine metabolism. *J Immunol* 167(11):6533–6544
155. Horowitz S, Binion DG, Nelson VM, Kanaa Y, Javadi P, Lazarova Z et al (2007) Increased arginase activity and endothelial dysfunction in human inflammatory bowel disease. *Am J Physiol Gastrointest Liver Physiol* 292(5):G1323–G1336
156. Ding H, Demple B (2000) Direct nitric oxide signal transduction via nitrosylation of iron-sulfur centers in the SoxR transcription activator. *Proc Natl Acad Sci U S A* 97(10):5146–5150