

Widespread adoption of Bt cotton and insecticide decrease promotes biocontrol services

Yanhui Lu¹, Kongming Wu¹, Yuying Jiang², Yuyuan Guo¹ & Nicolas Desneux³

Over the past 16 years, vast plantings of transgenic crops producing insecticidal proteins from the bacterium *Bacillus thuringiensis* (Bt) have helped to control several major insect pests^{1–5} and reduce the need for insecticide sprays^{1,5,6}. Because broad-spectrum insecticides kill arthropod natural enemies that provide biological control of pests, the decrease in use of insecticide sprays associated with Bt crops could enhance biocontrol services^{7–12}. However, this hypothesis has not been tested in terms of long-term landscape-level impacts¹⁰. On the basis of data from 1990 to 2010 at 36 sites in six provinces of northern China, we show here a marked increase in abundance of three types of generalist arthropod predators (ladybirds, lacewings and spiders) and a decreased abundance of aphid pests associated with widespread adoption of Bt cotton and reduced insecticide sprays in this crop. We also found evidence that the predators might provide additional biocontrol services spilling over from Bt cotton fields onto neighbouring crops (maize, peanut and soybean). Our work extends results from general studies evaluating ecological effects of Bt crops^{1–4,6,12,13} by demonstrating that such crops can promote biocontrol services in agricultural landscapes.

Biological control is a valuable ecosystem service^{14,15}, but increasingly intensive farming strongly influences the populations of natural enemies and the biocontrol services they provide^{16–18}. However, landscape biodiversity management and restricted use of pesticides may enhance biocontrol services in agro-ecosystems and could thus favour the development of sustainable farming^{7–9}. Genetically engineered crops that express δ -endotoxins (Cry proteins) from *Bacillus thuringiensis* (Bt) have been increasingly implemented by farmers in many countries since 1996, and more than 6.6×10^7 ha of Bt crops were planted worldwide in 2011 (ref. 19). Bt crops have successfully controlled several major insect pests^{1,2,4,5} and led to a drastic decrease in insecticide use on these crops^{1,5,6}. Because insecticide applications have been gradually reduced in Bt crops, their widespread adoption may benefit natural enemies and may therefore potentially enhance associated ecosystem services such as the control of arthropod pests^{10–12}. This last point has not yet been documented, especially with regard to the long-term landscape-level impacts¹⁰.

From the 1970s, insecticides were applied extensively to control cotton bollworm (CBW), *Helicoverpa armigera*, the most serious insect pest on conventional cotton in China. However, control became almost impossible in the early 1990s because the pest became resistant to most insecticides, and unprecedented outbreaks in 1992 led to a wide overuse of insecticides. Consequently, in 1993, the Chinese government requested systematic insecticide applications in wheat crops for the control of the first-generation CBW; that is, before the following generations colonized cotton crops²⁰. Although insecticide use decreased in cotton, this measure was not sustainable because insecticide applications were increased on wheat crops, resulting in both higher costs and environmental pollution. Bt cotton was therefore approved in 1997 for commercial use to control CBW, and it became the Chinese government's key measure against this cotton pest. It was

rapidly planted on a large scale, rising to 2.4×10^6 ha by 2011 (more than 95% of the cotton crop in northern China). It managed CBW effectively, which led to decreased insecticide use on this pest^{3,21}.

The widespread adoption of Bt cotton may have favoured an increase in generalist natural enemy populations and promoted their associated biocontrol services. We therefore performed two assessments: first, whether implementing Bt cotton on a large scale induced an increase in populations of three groups of key generalist predators in China (ladybirds, lacewings and spiders) in both Bt cotton and three common neighbouring crops, namely maize, peanut and soybean; and second, whether this trend resulted in increased biocontrol services in agricultural landscapes in China. Aphids were selected as a pest model because they are common prey for generalist predators. During 1990–2011, research was conducted in six major cotton-growing provinces (Henan, Hebei, Shandong, Shanxi, Anhui and Jiangsu) in northern China, where about 2.6×10^6 ha of cotton and 3.3×10^7 ha of other crops (notably maize, peanut and soybean) are cultivated annually by more than ten million small-scale farmers.




Figure 1 | Population densities of predators and aphids on cotton with different management regimes at Langfang experimental station (2001–2011). **a**, Predators. **b**, Aphids. The blue and red lines indicate Bt cotton and non-Bt cotton without insecticide sprays, respectively; the green line represents non-Bt cotton with CBW insecticide sprays (chemical control).

¹State Key Laboratory for Biology of Plant Diseases and Insect Pests, Institute of Plant Protection, Chinese Academy of Agricultural Sciences, Beijing, 100193 China. ²National Agro-Technical Extension and Service Center, Beijing, 100026 China. ³French National Institute for Agricultural Research (INRA), UMR1355-ISA, 400 route des chappes, 06903 Sophia-Antipolis, France.

Predators and cotton aphids were sampled from 2001 to 2011 in Bt and non-Bt cotton plots at Langfang experimental station in Hebei province. No significant differences were found for predator ($P = 0.341$) and aphid ($P = 0.555$) abundances between Bt cotton and non-Bt cotton with similar management methods; that is, without application of insecticide (Fig. 1a, b and Supplementary Table 1a, b). However, predator abundance was significantly lower and aphid abundance was significantly higher in plots treated with insecticides for CBW management in comparison with insecticide-free plots ($P < 0.001$) (Fig. 1a, b and Supplementary Table 1a, b), although it varied over years (significant interactions between insecticide application and year). Bt cotton does not itself affect predator and aphid population levels^{10,22}, and generalist predators are clearly susceptible to broad-spectrum insecticides (such as synthetic pyrethroids) used against CBW. Thereafter, insecticide-induced aphid resurgence usually occurs with widespread applications of insecticides.

Predator abundance and insecticide use in cotton were monitored in 36 locations throughout northern China during 1990–2010 (Fig. 2a and Supplementary Table 2). Predator population levels gradually increased over that period, and relatively high population levels were always observed after Bt cotton was implemented in 1997 (Fig. 2b). In 14 selected locations, all three major groups of predators (ladybirds, lacewings and spiders) showed an increasing trend similar to that of the whole predator complex (Fig. 2b). Insecticide use patterns also changed greatly with Bt cotton implementation. After the introduction of Bt cotton, the number of insecticide sprays against CBW (and other

insect pests in general), mainly pyrethroid and organophosphate insecticides (Supplementary Table 3), which have multiple negative effects on natural enemies¹⁷, was lower than during the pre-Bt cotton period, namely 1990–1996 (Fig. 2c). Moreover, predator population level and number of insecticide sprays were positively and negatively related to Bt cotton planting proportions, respectively ($P < 0.001$; Supplementary Fig. 1a, b), and indicated the effect of its large-scale adoption on the predator population trend. Regression analyses showed that fewer insecticide sprays against CBW and all insect pests were correlated to a great extent with an increase in predator populations in northern China ($P < 0.001$) (Fig. 2d, e). The results were consistent in the six provinces, and insecticide use against CBW was a driving factor for predator population level in the cotton agroecosystem (all $P < 0.05$; Supplementary Table 4).

Cotton aphid abundance was surveyed in 24 locations from 1990 to 2010 (Supplementary Table 2) to assess the biocontrol services provided by generalist predators. Linear regression analyses showed that increasing generalist predator populations were correlated with decreasing aphid abundance in northern China in general ($P < 0.001$; Fig. 3a) and in all provinces except Shanxi (Supplementary Fig. 2a–e). During the three main periods studied—that is, without Bt cotton, with less than 90% and more than 90% of Bt cotton planting in the landscapes—aphid populations decreased significantly ($P < 0.001$; Fig. 3b). In addition, aphid population was negatively related to the proportion of Bt cotton planted ($P = 0.003$; Supplementary Fig. 3). Exclusion cage trials in 2010 and 2011 at Langfang and Xinxiang experimental stations (in




Figure 2 | Relationships between predator population density and number of insecticide sprays on cotton in northern China (1990–2010). a, Survey locations, indicated by red dots.

b, Predator population density on cotton in commercial fields in 36 locations (each point represents one-year data; the red arrow indicates the beginning of Bt cotton use). Inset: population abundance of ladybirds (blue), spiders (red) and lacewings (green), collected from 14 locations.

c, Number of insecticide sprays for CBW (grey points) and all insect pests (black points) on cotton; each point represents one-year data. d, Linear relationship between total number of insecticide applications, determined by pooling all treatments against all the insect pests on cotton (x), and the predator abundance (y) in cotton

($y = -1.69x + 30.63$, $F_{1,19} = 71.19$, $R^2 = 0.79$, $P < 0.0001$). e, Linear relationship between number of insecticide applications for CBW only (x) and predator abundance (y)

($y = -1.11x + 16.03$, $F_{1,19} = 137.32$, $R^2 = 0.88$, $P < 0.0001$). The data in d and e are replotted from b and c. All error bars show s.e.m.




Figure 3 | Population abundance of cotton aphid in northern China (1990–2010) and relationship with predator abundance on cotton. **a**, Regression analysis between abundance of aphids (y) (\log_e -transformed) and predator abundance (x) ($y = e^{-0.15x + 8.39}$, $F_{1,19} = 69.67$, $R^2 = 0.79$, $P < 0.0001$). **b**, Aphid population density on cotton in commercial fields in 24 locations (each point represents one-year data, and the red arrow indicates the beginning of Bt cotton use). Red lines show the mean population density of aphids in cotton fields during three main periods, namely before Bt cotton planting (1990–1996), when Bt cotton planting was less than 90% of cotton surfaces planted (1997–2003) and when it was more than 90% (2004–2010). Red lines bearing different letters are significantly different at the $P < 0.05$ level in least-significant-difference post-hoc tests (one-way analysis of variance on \log_e -transformed data: $F_{2,18} = 27.57$, $P < 0.0001$). All error bars show s.e.m.

Hebei and Henan provinces, respectively) further demonstrated the significant effects of predators on aphid population growth in cotton fields (Supplementary Fig. 4a, b). As the cotton aphid populations declined, an invasive whitefly in cotton, *Bemisia tabaci*²⁰, probably served as an alternative prey for the increasing predator populations.




Figure 4 | Relationships between predator abundance on cotton and in three other crops, and between predator and aphid abundances in maize. Data on soybean (2001–2011), peanut (2001–2005 and 2008–2011) and maize (2001–2003 and 2008–2011) were collected at Langfang experimental station. **a**, Linear relationship between predator abundance on cotton (x) and on soybean (y) ($y = 0.10x + 3.38$, $F_{1,9} = 8.11$, $R^2 = 0.47$, $P = 0.0191$). **b**, Linear relationship between predator abundance on cotton (x) and on peanut (y) ($y = 0.09x + 2.66$, $F_{1,7} = 4.38$, $R^2 = 0.38$, $P = 0.0747$). **c**, Linear relationship between predator abundance on cotton (x) and on maize (y) ($y = 0.23x + 14.96$, $F_{1,5} = 2.00$, $R^2 = 0.29$, $P = 0.2164$). **d**, Relationship between predator abundance (x) and abundance of aphids in maize (y ; \log_e -transformed data) ($y = e^{-0.07x + 8.31}$, $F_{1,5} = 5.80$, $R^2 = 0.54$, $P = 0.0610$). All error bars show s.e.m.

All these results indicate that the widespread adoption of Bt cotton ultimately promotes biocontrol services in the agroecosystem because decreased insecticide use leads to an increase in predator populations. Broadly speaking, measures that preserve predators in cotton fields greatly help to control aphid populations; for example, when insecticide applications in wheat were requested by the Chinese government (1993–1996) to prevent CBW outbreaks in cotton (see above), it led to a decreasing trend in aphid abundance (Fig. 3b).

Predator abundance was also monitored from 2001 to 2011 in three neighbouring crops: maize, peanut and soybean at Langfang experimental station. There was a positive relationship between predator abundance in cotton and soybean ($P = 0.019$; Fig. 4a), as well as between cotton and peanut (marginally significant, $P = 0.075$; Fig. 4b). We observed a similar trend in maize but it was not significant ($P = 0.216$; Fig. 4c). The increased predator abundance in maize was linked to a decrease in aphid pest abundance in that particular crop (marginally significant, $P = 0.061$; Fig. 4d).

Biocontrol services are important components in agro-ecosystems and could lead to the development of sustainable agriculture^{7,15,23}. In conventional agricultural practices, insecticides are frequently used to control targeted pests, but they can lead to outbreaks of secondary pests by suppressing their natural enemies²⁴. This so-called insecticide-induced resurgence was first reported for cotton aphid in the 1970s and was regarded as a key factor leading to population outbreaks of this pest in China²⁵. Our work demonstrates the importance of natural enemies in the long-term suppression of the cotton aphid. The widespread adoption of Bt cotton, as a sustainable measure to reduce insecticide use, has indirectly promoted generalist predator abundance in Bt cotton fields but also to a smaller extent in three common adjacent crops in northern China. Bt crops therefore might enhance biocontrol services in agricultural landscapes through an increased abundance of generalist natural enemies. This study provides key information on long-term landscape-level ecological effects of Bt crops as well as useful insights, for example into the management of pest resurgence problems reported for many pests worldwide²⁶.

Generalist predators usually have great dispersal ability and can rely on various food sources. Hence, not only can they synchronously attack different insect pests in one field, but they can also colonize different habitats in different seasons^{27,28}. Furthermore, some habitat management measures, such as inter-planting different crops or wild plants, have been adopted to provide resources such as food supply or shelter for natural enemies, thus increasing conservation biological control in adjacent fields^{7,9,27,28}. We have demonstrated that decreasing insecticide application, through widespread Bt cotton plantings, sustained generalist predators and helped to suppress aphid populations in this

crop. Large-scale insecticide reduction is the key driver in such processes (for example see the period 1993–1996, during which insecticide decrease favoured an increase in predator populations and a decline of aphid populations). Higher generalist predator population levels in Bt cotton lead to lower insect pest levels in the crop, and these predators might provide additional biocontrol services spilling over from cotton fields onto neighbouring crops, although further work should be performed to document this last point. Broadly speaking, the deployment of Bt crops may favour biocontrol services and enhance economic benefits not only in Bt crop fields but also in the whole agricultural landscape. Field studies indicated that Bt crops protected natural enemies in comparison with non-Bt crops, which rely on conventional insecticides^{22,29}. Our present study, demonstrating that biocontrol services are potentially provided by Bt crops throughout the agricultural landscape, may offer new options in developing conservation biological control measures at the landscape level.

Critical concerns about the ecological risk assessment of transgenic crops still remain, especially on a large scale²⁹. The present study confirms no negative effects of one Bt crop, Bt cotton, on generalist predators in agricultural landscapes in China. More particularly, we have demonstrated a marked increase in generalist predator population levels and associated biocontrol services linked to decreased insecticide use owing to the widespread adoption of the Bt crop. Our work provides a comprehensive, long-term and large-scale assessment of the possible ecological and agricultural effects of transgenic crops.

METHODS SUMMARY

The study was based on large-scale surveys of predator and cotton aphid populations in cotton fields of northern China from 1990 to 2010 and on experiments and surveys that were performed at Langfang experimental station of the Chinese Academy of Agricultural Science (CAAS) during the period 2001–2011. The surveys and experiments focused on three major generalist predator groups (ladybirds, lacewings and spiders) and on aphid pests in cotton and in three common cotton-neighbouring crops, namely maize, peanut and soybean.

At the CAAS, we first assessed how cultural practices could affect predator and aphid populations in the long term in cotton fields; cotton plots were established every year and the abundance of predators and cotton aphids was surveyed in three different plot types: Bt cotton, non-Bt cotton and non-Bt cotton with insecticide. Second, we determined the impact of predators on aphid population in cotton by means of exclusion cage trials. Third, we evaluated the impact of implementing Bt cotton on predator and aphid populations in the neighbouring crops. Field plots were established in cotton, maize, peanut and soybean, and population dynamics of predators and aphids were monitored.

Large-scale surveys were conducted in six provinces in northern China (36 locations, 10–20 fields per location) to evaluate the impact of insecticide applications on the abundance of predators and aphids in cotton fields. We tested, first, the relationship between predator abundance and insecticide use during the period 1990–2010 (that is, including the period before and during the widespread adoption of Bt cotton by farmers), and second, how cotton aphid density was related to predator abundance during the same period.

Full Methods and any associated references are available in the online version of the paper at www.nature.com/nature.

Received 3 January; accepted 23 April 2012.

Published online 13 June 2012.

- Shelton, A. M., Zhao, J. Z. & Roush, R. T. Economic, ecological, food safety, and social consequences of the deployment of Bt transgenic plants. *Annu. Rev. Entomol.* **47**, 845–881 (2002).
- Carriere, Y. *et al.* Long-term regional suppression of pink bollworm by *Bacillus thuringiensis* cotton. *Proc. Natl Acad. Sci. USA* **100**, 1519–1523 (2006).
- Wu, K. M., Lu, Y. H., Feng, H. Q., Jiang, Y. Y. & Zhao, J. Z. Suppression of cotton bollworm in multiple crops in China in areas with Bt toxin-containing cotton. *Science* **321**, 1676–1678 (2008).
- Hutchison, W. D. *et al.* Areawide suppression of European corn borer with Bt maize reaps savings to non-Bt maize growers. *Science* **330**, 222–225 (2010).

- Tabashnik, B. E. *et al.* Suppressing resistance to Bt cotton with sterile insect releases. *Nature Biotechnol.* **28**, 1304–1307 (2010).
- Cattaneo, M. G. *et al.* Farm-scale evaluation of the impacts of transgenic cotton on biodiversity, pesticide use, and yield. *Proc. Natl Acad. Sci. USA* **103**, 7571–7576 (2006).
- Landis, D. A., Wratten, S. D. & Gurr, G. M. Habitat management to conserve natural enemies of arthropod pests in agriculture. *Annu. Rev. Entomol.* **45**, 175–201 (2000).
- Crowder, D. W., Northfield, T. D., Strand, M. R. & Snyder, W. E. Organic agriculture promotes evenness and natural pest control. *Nature* **466**, 109–112 (2010).
- Winqvist, C. *et al.* Mixed effects of organic farming and landscape complexity on farmland biodiversity and biological control potential across Europe. *J. Appl. Ecol.* **48**, 570–579 (2011).
- Romeis, J. *et al.* Transgenic crops expressing *Bacillus thuringiensis* toxins and biological control. *Nature Biotechnol.* **24**, 63–71 (2006).
- Bale, J. S., van Lenteren, J. C. & Bigler, F. Biological control and sustainable food production. *Phil. Trans. R. Soc. B* **363**, 761–776 (2008).
- Gatehouse, A. M. R., Ferry, N., Edwards, M. G. & Bell, H. A. Insect-resistant biotech crops and their impacts on beneficial arthropods. *Phil. Trans. R. Soc. B* **366**, 1438–1452 (2011).
- Wolfenbarger, L. L., Naranjo, S. E., Lundgren, J. G., Bitzer, R. J. & Watrud, L. S. Bt crop effects on functional guilds of non-target arthropods: a meta-analysis. *PLoS ONE* **3**, e2118 10.1371/journal.pone.0002118 (2008).
- Bianchi, F. J. J. A., Booij, C. J. H. & Tscharntke, T. Sustainable pest regulation in agricultural landscapes: a review on landscape composition, biodiversity, and natural pest control. *Proc. R. Soc. Lond. B* **273**, 1715–1727 (2006).
- Losey, J. E. & Vaughan, M. The economic value of ecological services provided by insects. *Bioscience* **56**, 311–323 (2006).
- Chapin, F. S. *et al.* Consequences of changing biodiversity. *Nature* **405**, 234–242 (2000).
- Desneux, N., Decourtye, A. & Delpuech, J. M. The sublethal effects of pesticides on beneficial arthropods. *Annu. Rev. Entomol.* **52**, 81–106 (2007).
- Landis, D. A., Gardiner, M. M., van der Werf, W. & Swinton, S. M. Increasing corn for biofuel production reduces biocontrol services in agricultural landscapes. *Proc. Natl Acad. Sci. USA* **105**, 20552–20557 (2008).
- James, C. *Global Status of Commercialized Biotech/GM crops: 2011* (ISAAA brief no. 43, International Service for the Acquisition of Agri-Biotech Applications, 2011).
- Wu, K. M. & Guo, Y. Y. The evolution of cotton pest management practices in China. *Annu. Rev. Entomol.* **50**, 31–52 (2005).
- Lu, Y. H. *et al.* Mirid bug outbreaks in multiple crops correlated with wide-scale adoption of Bt cotton in China. *Science* **328**, 1151–1154 (2010).
- Naranjo, S. E. Long-term assessment of the effects of transgenic Bt cotton on the abundance of nontarget arthropod natural enemies. *Environ. Entomol.* **34**, 1193–1210 (2005).
- Ragsdale, D. W., Landis, D. A., Brodeur, J., Heimpel, G. E. & Desneux, N. Ecology and management of the soybean aphid in North America. *Annu. Rev. Entomol.* **56**, 375–399 (2011).
- Hardin, M. R. *et al.* Arthropod pest resurgence: an overview of potential mechanisms. *Crop Prot.* **14**, 3–18 (1995).
- Wu, K. W. & Liu, Q. X. Study on the resurgence caused by insecticides for cotton aphid, *Aphis gossypii*. *Acta Ecol. Sin.* **12**, 341–347 (1992).
- Rao, C. N., Shivankar, V. J. & Singh, S. in *Encyclopedia of Pest Management* Vol 2 (ed. Pimentel, D.) 597–601 (CRC Press, 2007).
- Symondson, W. O. C., Sunderland, K. D. & Greenstone, H. M. Can generalist predators be effective biocontrol agents? *Annu. Rev. Entomol.* **47**, 561–594 (2002).
- Tscharntke, T. *et al.* Conservation biological control and enemy diversity on a landscape scale. *Biol. Control* **43**, 294–309 (2007).
- Romeis, J., Shelton, A. M. & Kennedy, G. G. *Integration of Insect-resistant Genetically Modified Crops within IPM Programs* (Springer, 2008).

Supplementary Information is linked to the online version of the paper at www.nature.com/nature.

Acknowledgements We thank the agricultural technicians of the 36 surveyed locations for providing data sets from cotton fields; W. Li and L. Wang for providing the predator exclusion cage data sets; Y. Zhang and C. Li for assistance in making the survey map; H. Yuan for help in preparing the list of insecticides in cotton; and R. Senoussi for advice on data analyses. This work was funded by the Key Project for Breeding Genetically Modified Organisms (grant no. 2011ZX08012-004) and the International Science and Technology Cooperation Project (grant no. 2010DFA32200).

Author Contributions K.W., Y.L. and Y.G. designed and performed the experiments. Y.J. performed the surveys. Y.L., K.W. and N.D. analysed the data and shared in the scoping and writing responsibilities.

Author Information Reprints and permissions information is available at www.nature.com/reprints. The authors declare no competing financial interests. Readers are welcome to comment on the online version of this article at www.nature.com/nature. Correspondence and requests for materials should be addressed to K.W. (kmwu@ippcaas.cn).

METHODS

Aphid pests and predator complex considered in the study. In northern China, several aphid species are reported as pests on cotton, maize, peanut and soybean. *Aphis gossypii* Glover (cotton aphid) is the main aphid pest in cotton fields in northern China where there are two key biotypes (called seedling and summer aphids, respectively). Only the summer aphid, which colonizes fields from early July to late August, is considered a major cotton pest^{20,30,31}. The seedling aphid is controlled by insecticide treatments applied on the seeds; these compounds do not last long enough within plants to provide control of the summer aphid²⁰. *Rhopalosiphum maidis* Fitch, *R. padi* Linnaeus and *Sitobion avenae* Fabricius are the three dominant aphid pest species on maize³¹, and *A. glycines* Matsumura and *A. craccivora* Koch are the main aphid species on soybean and peanut, respectively³¹. In our study, the above aphid species were considered in assessing the biocontrol services provided by generalist predators.

In northern China, there are three dominant groups of generalist predators in cotton field (more than 90% of all the predators³⁰): ladybirds, lacewings and spiders. In our study we therefore focused on a predator complex composed of ladybirds (*Propylea japonica* Thunberg, *Harmonia axyridis* Pallas, *Coccinella septempunctata* L. and *Adonia variegata* Goetz), lacewings (*Chrysopa septempunctata* Wesmäl, *Chrysoperla sinica* Tjeder and *Chrysopa formosa* Brauer) and spiders (*Erigonidium graminicolum* Sundevall, *Misumenopus tricuspidata* Fabricius and *Pardosa t-insignita* Boes. et Str.); these compose the most common predators in agricultural landscape of that region. These groups of predators are also common in maize, peanut and soybean fields and were thus also considered as a predator complex for these three crops^{32–34}.

Impact of agricultural practices on predator and cotton aphid populations. Survey experiments were conducted from 2001 to 2011 at Langfang experimental station (39.53° N, 116.70° E), Chinese Academy of Agricultural Sciences (CAAS), Hebei province, China. Fifteen cotton plots (400 m² each) were established every year and were managed with agronomic practices that are standard in northern China. A randomized block design with three replicates was used, which included two Bt and two conventional cotton varieties. One Bt cotton variety expressing *Cry1Ac* (NuCOTN33B) and another Bt cotton varieties expressing *Cry1A* (SGK321) were supplied by Monsanto Co. and the Biotechnology Research Institute, CAAS, respectively. Two conventional cotton varieties (Shiyuan321 and Zhong12) were obtained from the Institute of Plant Protection, CAAS. Shiyuan321 was the non-transgenic isoline of SGK321. Every year, the trial consisted of three treatments: Bt cotton and non-Bt cotton plots without insecticide, and non-Bt cotton (one variety, Zhong12) plots with insecticides. β -Cypermethrin (pyrethroid) and phoxim (organophosphate) were used when insecticides were applied. The choice of these two insecticides, and their frequency of application, were both based on management guidelines for CBW (*Helicoverpa armigera*) used throughout the early 1990s in northern China²⁰ (Supplementary Table 3).

The abundances of predators and cotton aphid were surveyed in the three cotton plot types (Bt cotton, non-Bt cotton and non-Bt cotton with insecticide) every 4 or 5 days from mid-June to late August from 2001 to 2011. At each sampling date, 100 plants at five random locations per plot³⁵ were visually inspected and all predators and aphids were recorded. No significant differences ($P > 0.05$) were found between cotton varieties, so we combined NuCOTN33B and SGK321 as Bt cotton, and Shiyuan321 and Zhong12 as non-Bt cotton, for further analysis. A three-way ANOVA was used to analyse the effects of the cotton variety (Bt cotton and non-Bt cotton), insecticide treatments (chemical control and non-chemical control) and sampling year on predator and aphid abundance, and the interactions between year and cotton variety, and between year and insecticide spray; the means were compared by the least-significant-difference (LSD) test at $P = 0.05$.

Survey of predators and cotton aphid in cotton crops in northern China. From 1990 until 2010, commercial cotton fields in 36 locations in six provinces (Henan, Hebei, Shandong, Shanxi, Anhui and Jiangsu) of northern China were surveyed for predators and cotton aphid (Supplementary Table 2). Insect populations were recorded every 3–10 days from early June to late August every year. For each survey, 10–20 cotton fields were sampled per location. Within each field, a total of 50–100 cotton plants at five random locations were visually inspected for predators³⁰. Among the 36 locations, ladybirds, lacewings and spiders were recorded as a predator complex in 22 locations, whereas in 14 sites the three predator types were recorded individually. The 14 locations included Anxin and Xinji from Hebei province; Dezhou, Binzhou and Chengwu from Shandong province; Ruicheng, Yongji and Linfen from Shanxi province; Dongzhi, Wangjiang and Taihu from Anhui province; and Dafeng, Tongzhou and Haimen from Jiangsu province. The cotton aphid populations were surveyed in 24 locations in five provinces (Supplementary Table 2), using the same sampling schedule as for survey of predators³⁰. On each plant, an upper leaf, a middle leaf and a lower leaf

were examined for aphid presence. At the same time, all insecticide applications (for management of CBW and other arthropod pests) were recorded per field per year.

Linear regression analyses were used to assess the relationship between predator abundance and insecticide use on the data set gathered from 1990 to 2010—that is, including the period during which Bt cotton was increasingly adopted in China by farmers. Both simple and forward stepwise regressions were used to relate predator abundance and insecticide use against CBW and all insect pests for each province and the whole of northern China in the 1990–2010 data set. Simple linear models were used to assess the relationship between aphid density (log-transformed) and predator abundance from early July to late August (in the 1990–2010 data set) for each province and for the whole of northern China. Linear regression analyses were used to assess the relationship between predator abundance and aphid abundance (log-transformed) with Bt cotton planting proportions. In this analysis, the mean abundances of predators and aphids during 1990–1996 were included as the data when the Bt cotton planting proportion was 0.


To evaluate the impact of the predators on cotton aphid population further, exclusion cage trials were conducted in 2010 and 2011 at Langfang experimental station and Xixiang experimental station of CAAS (Henan Province, 35.09° N, 113.48° E). This trial included a caged treatment and an open-field treatment as control^{18,36}. The cage was 2 m wide by 2 m wide by 1.5 m high and made from the insect mesh net, which allowed the emigration and immigration of alate aphids and its parasitoids, but blocked the predators³⁷. Ten cotton plants were covered in each cage. This trial began in July, when almost only apterous aphids were in the field³⁸, and was limited to 15 days to prevent the appearance of alate aphids in the cage^{18,36}. At each site, cage treatments with three or four replicates were established when aphid density reached an average of 2 individuals per plant in 2010 and 20 individuals per plant in 2011. We recorded the aphid abundance 15 days after treatment. Meanwhile, predator densities were surveyed three times, on day 0, day 5 and day 10, in ten randomly selected cotton plants in open field during the whole trial. The aphid abundance in caged and open plants was compared by one-way ANOVA followed by a post-hoc LSD test. Before analysis, the data for aphid abundance were log_e-transformed.

Impact of Bt cotton adoption on populations of predators in neighbouring crops. Population dynamics of the predator complex were monitored in cotton, soybean, peanut and maize field plots from 2001 to 2011 (except for maize, which was monitored during 2001–2003 and 2008–2011, and soybean, which was monitored during 2001–2006 and 2008–2011) at Langfang experimental station. Every year, a total of nine field plots (400 m² each) were established for each crop type and they were managed in the same way, applying the same fertilizers and irrigation treatment, free of any pesticide. A randomized block design with three replicates for each crop type was used. One Bt cotton variety, SGK321, was supplied by the Biotechnology Research Institute (CAAS); the maize (var. Shengshi29), soybean (var. Zhonghuang13) and peanut (var. Huayu16) were provided by Langfang experimental station (CAAS), the Institute of Crop Sciences (CAAS) and the Shandong Peanut Research Institute, respectively. The abundance of predators was recorded in the four different crops (Bt cotton, maize, peanut and soybean) every four or five days from mid-June to late August. At each sampling date, 100 plants in five random spots per plot were visually inspected and all predators were recorded. Linear regression analyses were used to assess the relationship between seasonal density of predators on cotton and soybean/peanut (data set covering the 2001–2011 period) and maize (data set covering the 2001–2003 and 2008–2011 periods).

For the maize plots, maize aphids were also recorded because they are well known as the main pests on maize in northern China. Population levels of aphids on soybean and peanut crops at Langfang experimental station were very low during the course of our study and therefore the data could not be considered in the framework of the study. A simple linear model was used to assess the relationship between aphid abundance (log-transformed) and predator abundance on maize.

30. Qu, X. F. *Cotton Pest Forecast in China: the Criterion, Zoning and Method* (China Scien-tech Press, 1992).
31. Institute of Plant Protection, Chinese Academy of Agricultural Sciences. *Crop Diseases and Pests in China* (China Agricultural Press, 1995).
32. Xu, H. F., Mu, S. M., Xu, Y. Y., Mu, J. Y. & Dong, C. X. On the community structure of major insect pests and natural enemies in summer corn field inlaid in cotton area. *Acta Phytophyl. Sin.* **27**, 199–204 (2000).
33. Yang, Q. M., Sun, M., Xu, Y. F., Shi, A. J. & Mu, J. Y. On the community structure of major insect pests and natural enemies in summer bean field. *J. Shandong Agric. Univ.* **35**, 217–220 (2004).
34. Ding, Y. Q. & Cheng, S. L. Preliminary investigation and utilization of natural enemies of aphid in peanut field. *J. Peanut Sci.* **39**, 45–47 (2010).

35. Wu, K. M. & Guo, Y. Y. Influences of *Bacillus thuringiensis* Berliner cotton planting on population dynamics of the cotton aphid, *Aphis gossypii* Glover, in northern China. *Environ. Entomol.* **32**, 312–318 (2003).
36. Gardiner, M. M. *et al.* Landscape diversity enhances biological control of an introduced crop pest in the north-central USA. *Ecol. Appl.* **19**, 143–154 (2009).
37. Miao, J., Wu, K. M., Hopper, K. R. & Li, G. X. Population dynamics of *Aphis glycines* (Homoptera: Aphididae) and impact of natural enemies in northern China. *Environ. Entomol.* **36**, 840–848 (2007).
38. Lu, Y. H., Qi, F. J. & Zhang, Y. J. *Integrated Management of Diseases and Insect Pests in Cotton* (Golden Shield Press, 2010).



Supplementary Figure 1. Relationship between predator abundance, insecticide use

and Bt cotton plantings in Northern China. a) Linear relationship between Bt cotton


proportion (x) and predator abundance (y), $y=5.93x+7.34$, $F_{1,13}=37.09$, $R^2=0.74$, $P<0.0001$;

b), Linear relationship between Bt cotton proportion (x) and insecticide use (y), insecticide

use against all the insect pests, $y=-2.78x+13.47$, $F_{1,13}=45.62$, $R^2=0.78$, $P<0.0001$;


insecticide use against CBW (*H. armigera*), $y=-6.11x+8.54$, $F_{1,13}=333.95$, $R^2=0.96$,

$P<0.0001$.




Supplementary Figure 2. Relationship between predator and cotton aphid

abundance in Northern China from 1990 to 2010. Regression analysis between (y) mean (\pm SEM) abundance of aphids (Log_e -transformed) and (x) mean (\pm SEM) predator abundance: a), Henan province, $y=e^{-0.06x+8.19}$, $F_{1,19}=9.12$, $R^2=0.32$, $P=0.0070$; b), Hebei province, $y=e^{-0.23x+8.17}$, $F_{1,19}=7.00$, $R^2=0.27$, $P=0.0160$; c), Shandong province, $y=e^{-0.14x+7.48}$, $F_{1,19}=8.92$, $R^2=0.32$, $P=0.0076$; d), Shanxi province, $y=e^{-0.16x+8.63}$, $F_{1,19}=2.36$, $R^2=0.11$, $P=0.1410$; e), Anhui province, $y=e^{-0.05x+7.16}$, $F_{1,19}=24.71$, $R^2=0.57$, $P=0.0001$.



Supplementary Figure 3. Relationship between cotton aphid abundance and Bt cotton plantings in Northern China during the period studied. Linear relationship between Bt cotton proportion (x) and aphid abundance (Log_e -transformed) (y), $y=e^{-0.86x+7.37}$, $F_{1,13}=13.06$, $R^2=0.50$, $P=0.0032$.



Supplementary Figure 4. Aphid abundance (mean \pm SEM) on cotton plants in exclusion cage (without predators) and in open-field (presence of predators) treatments at the Langfang and Xinxiang experimental stations (2010-2011). Grey histograms: without predators, white histograms: with predators. The statistical analysis is based on one-way ANOVA per year and per location. a) Langfang experimental station: 2010, $F_{1,6}=32.55$ ($n=4$); 2011, $F_{1,6}=25.91$ ($n=4$); b) Xinxiang experimental station: 2010, $F_{1,6}=17.77$ ($n=4$); 2011, $F_{1,4}=50.92$ ($n=3$).

Supplementary Table 1. Effects of cotton variety (Bt vs. non-Bt), insecticide spray against CBW (*H. armigera*) and sampling year on predator and aphid abundance at the Langfang experimental station (2001-2011). a) Mean (\pm SEM) numbers (n=3) of aphid and predators in the three treatments i.e. Bt, Non-Bt and Non-Bt-CC. Bt and Non-Bt mean Bt cotton and non-Bt cotton without insecticide sprays, and Non-Bt-CC means non-Bt cotton with chemical control for CBW. b) Statistics from the three-way ANOVA used to analyze the effects of cotton variety (Bt vs. non-Bt), insecticide sprays (Non-Bt vs. Non-Bt-CC) for CBW and sampling year on predator and aphid abundance.

a

Year	Seasonal mean density (Individuals per 100 plants)					
	Predators			Aphid		
	Bt	Non-Bt	Non-Bt-CC	Bt	Non-Bt	Non-Bt-CC
2001	19.63 \pm 0.93	18.50 \pm 1.38	9.17 \pm 0.25	152.81 \pm 59.49	90.67 \pm 21.02	1379.08 \pm 426.40
2002	16.04 \pm 0.69	17.06 \pm 0.41	5.53 \pm 0.67	1961.49 \pm 1120.30	821.41 \pm 216.63	36266.00 \pm 7837.57
2003	61.13 \pm 3.48	63.93 \pm 3.05	25.56 \pm 0.31	1675.47 \pm 573.55	812.32 \pm 47.21	30113.91 \pm 5280.27
2004	27.93 \pm 1.56	28.41 \pm 1.64	11.78 \pm 0.17	852.00 \pm 46.61	780.85 \pm 21.05	13178.56 \pm 381.56
2005	16.07 \pm 0.48	16.65 \pm 1.19	6.57 \pm 0.22	1344.12 \pm 90.17	1292.85 \pm 91.69	3789.30 \pm 161.77
2006	24.91 \pm 1.19	24.93 \pm 1.14	9.98 \pm 0.46	1804.89 \pm 299.68	1665.39 \pm 243.00	8919.58 \pm 22.77
2007	23.67 \pm 0.74	22.35 \pm 1.12	11.84 \pm 1.23	443.38 \pm 93.74	386.11 \pm 40.79	8191.22 \pm 610.00
2008	35.13 \pm 1.12	36.44 \pm 0.55	16.07 \pm 1.15	810.39 \pm 95.81	732.76 \pm 40.29	2134.33 \pm 234.92
2009	20.37 \pm 0.81	19.03 \pm 1.38	8.30 \pm 0.86	3214.58 \pm 183.32	3291.07 \pm 232.69	5556.62 \pm 79.68
2010	39.89 \pm 0.31	42.70 \pm 1.97	27.33 \pm 2.71	1550.33 \pm 392.85	1427.44 \pm 368.12	9829.52 \pm 2060.65
2011	14.57 \pm 0.67	15.30 \pm 0.59	5.10 \pm 0.56	3041.88 \pm 191.86	3067.78 \pm 234.49	5893.57 \pm 170.72

b

Group	Source of variation	df	F value	P value
Predators	Cotton variety	1	0.91	0.3413
	Insecticide application	1	476.71	<0.0001
	Year	10	137.52	<0.0001
	Year x Cotton variety	10	0.59	0.8212
	Year x Insecticide application	10	13.12	<0.0001
Aphid	Cotton variety	1	0.35	0.5552
	Insecticide application	1	464.62	<0.0001
	Year	10	51.19	<0.0001
	Year x Cotton variety	10	0.1	0.9998
	Year x Insecticide application	10	56.45	<0.0001

Supplementary Table 2. Locations for wide-scale survey of predator and aphid population and insecticide use in Northern China (1990-2010).

Province	Location	Longitude	Latitude	Survey contents		
				Insecticide use	Predator abundance	Aphid abundance
Hebei	Fengnan	118.10	39.58	√	√	√
	Anxin	115.91	38.91	√	√	√
	Raoyang	115.75	38.25	√	√	√
	Xinji	115.12	37.54	√	√	
	Gucheng	115.97	37.37	√	√	√
	Pingshan	114.25	38.20	√	√	
	Guantao	115.40	36.46	√	√	√
Shandong	Dezhou	116.30	37.45	√	√	√
	Lingxian	116.58	37.35	√	√	√
	Shanghe	117.15	37.32	√	√	√
	Binzhou	118.00	37.30	√	√	√
	Dongying	118.50	37.47	√	√	√
	Guangrao	118.42	37.05	√	√	√
	Linqing	115.72	36.68	√	√	√
	Guanxian	115.45	35.47	√	√	√
	Liaocheng	115.97	36.45	√	√	√
	Chengwu	115.80	34.90	√	√	√
Jiangsu	Dafeng	120.45	33.20	√	√	
	Tongzhou	121.03	32.05	√	√	
	Haimen	121.15	31.90	√	√	
Anhui	Caohu	117.87	31.62	√	√	
	Chizhou	117.40	30.60	√	√	
	Wuwei	117.75	31.30	√	√	√
	Dongzhi	117.00	30.08	√	√	
	Wangjiang	116.70	30.12	√	√	

	Susong	116.13	30.15	√	√	
	Taihu	116.27	30.42	√	√	
	Taihe	115.62	33.17	√	√	
	Xiaoxian	116.93	34.20	√	√	√
Henan	Anyang	114.35	36.10	√	√	√
	Jixian	114.77	34.57	√	√	√
	Dancheng	115.16	33.63	√	√	√
	Zhenping	112.23	33.03	√	√	√
Shanxi	Ruicheng	110.68	34.71	√	√	√
	Yongji	110.42	34.88	√	√	√
	Linfen	111.50	36.10	√	√	√

Supplementary Table 3. Insecticides used on cotton in Northern China during three periods from 1990 to 2010 (China Pesticide Information Network, <http://www.chinapesticide.gov.cn>). The crosses indicate, for each insecticide, when they were (or not) used.

Insecticide chemical family	Insecticide name	Targeted pest: CBW (<i>H. armigera</i>)			Targeted pest: other insects		
		1990-1996	1997-2003	2004-2010	1990-1996	1997-2003	2004-2010
Organophosphate	Acephate	+	+	+	+	+	+
	Chlorpyrifos	+	+	+	+	+	+
	Chlorpyrifos-methyl		+	+			
	Diazinon						+
	Dichlorvos				+	+	+
	Dimethoate				+	+	+
	Ethion				+	+	+
	Fenitrothion	+	+	+	+	+	+
	Fenthion	+	+	+	+	+	+
	Isocarbophos	+	+	+	+	+	+
	Malathion				+	+	+
	Omethoate				+	+	+
	Phorate				+	+	
	Phosfolan				+	+	+
	Phosmet	+	+	+	+	+	+
	Phoxim	+	+	+	+	+	+
	Phoxim-methyl	+	+				
	Profenofos	+	+	+			
	Pyridaphenthion	+	+	+	+	+	+
	Quinalphos	+	+	+	+	+	+
Triazophos	+	+	+	+	+	+	
Trichlorfon				+	+	+	
Pyrethroid	beta-Cypermethrin	+	+	+	+	+	+

	Bifenthrin	+	+	+			
	Brofenvalerate	+	+		+	+	
	Brofluthrin	+	+		+	+	
	Cyfluthrin	+	+	+	+	+	+
	Fenpropathrin	+	+	+			
	Fenvalerate	+	+	+	+	+	+
	Permethrin	+	+	+			
	tau-Fluvalinate	+	+	+	+	+	+
	Tralomethrin	+	+				
	Valerathrin	+	+		+	+	
Carbamate	Aldicarb				+	+	+
	Carbaryl				+	+	+
	Carbofuran				+	+	+
	Carbosulfan				+	+	+
	Indoxacarb		+	+			
	Methomyl	+	+	+	+	+	+
	Thiodicarb		+	+			
Neonicotinoid	Acetamiprid				+	+	+
	Imidacloprid				+	+	+
	Nitenpyram					+	+
	Thiamethoxam					+	+
Others	Avermectin	+	+	+	+	+	+
	Emamectin benzoate		+	+			
	Endosulfan		+	+			

Supplementary Table 4. Relationship between predator abundance and insecticide

use in cotton in Northern China (1990-2010). No. insecticide sprays: total number of insecticide applications in cotton (i.e. pooling all treatments against all the insect pests on cotton). No. CBW insecticide sprays: number of insecticide applications made for management of only *H. armigera*. RC: regression coefficient. R²: coefficient of determination. df=1,19. For stepwise regression, only variables with significance at the *P*<0.05 were included in the analysis.

Province	Simple regression						Stepwise regression					
	No. insecticide sprays			No. CBW insecticide sprays			No. insecticide sprays			No. CBW insecticide sprays		
	RC	R ²	<i>P</i>	RC	R ²	<i>P</i>	RC	R ²	<i>P</i>	RC	R ²	<i>P</i>
Henan	-2.060	0.78	<0.0001	-1.948	0.79	<0.0001				-1.948	0.79	<0.0001
Hebei	-0.385	0.23	0.0290	-0.349	0.44	0.0011				-0.349	0.44	0.0011
Shandong	-0.767	0.56	0.0001	-0.577	0.78	<0.0001				-0.577	0.78	<0.0001
Shanxi	-0.413	0.19	0.0488	-0.382	0.32	0.0081				-0.382	0.32	0.0081
Anhui	-2.205	0.20	0.0406	-2.143	0.44	0.0010				-2.143	0.44	0.0010
Jiangsu	-1.238	0.21	0.0364	-1.944	0.51	0.0003				-1.944	0.51	0.0003
Northern China	-1.685	0.79	<0.0001	-1.109	0.88	<0.0001				-1.109	0.88	<0.0001