

## **Antimicrobial Activity of Piperine purified from *piper nigrum***

ISSN 1817 - 2695

**Zainab T.K. Aldaly**

*Pharmaceutics and Clinical pharmacy Dep. College of Pharmacy,  
Basrah University*

((Received 28/10/2009 , Accepted 18/8/2010))

### **Abstract:**

An alkaloid piperine was extracted from dry seeds of plant *Piper nigrum* with ethanol by using Soxhlet extraction then isolation and purification by re-crystallization, the structure of Piperine was confirmed by the IR spectroscopy. Antimicrobial activity of Piperine was studied against gram+ve, gram-ve bacteria and *Candida albicans* piperine showed potent antimicrobial activity against tested organisms especially *C. albicans* followed by *E. coli* and less than on *P. aeruginosa* the zone of inhibition ranges from 8 - 23 mm and the minimum inhibitory concentrations (MIC) 3.125 - 100mg/ml.

### **Introduction**

Spices and herbs have been used for centuries by many cultures enhance the flavor and aroma of foods. Early cultures also recognized the value of using spices and herbs in preserving foods and for their medicinal value. Scientific experiments since the late 19<sup>th</sup> century have demonstrated the antimicrobial properties of some spices, herbs and their components. (2, 18) Recently, there has been an increasing interest in the discovery of new natural antimicrobials, because of an increasing risk in the rate of infection with antibiotic-resistant microorganisms and also due to the side and the residual effects of antibiotics. The part of different common herbs, such as *Piper nigrum*, commonly known as black pepper (1) Sometimes called Indian long pepper, is a flowering vine in the family piperaceae, cultivated for its fruit, which is usually dried and used as spice and seasoning, it is a close relative of the black pepper plant, and has a similar, though generally hotter taste. The root and fruit of *piper longum* are used in palsy, gout and lambago. The fruits have a bitter, hot, sharp taste, tonic to liver, stomach,

emmenagogue, abortifacient, aphrodisiac and digestive (11).

The fruits and roots are attributed to numerous medicinal uses and may be used for diseases of respiratory tract, cough, bronchitis, asthma, etc; as counter irritant and analgesic when applied for muscular pains and inflammation, it is also prescribed in case of cholera, dyspepsia, flatulence and diarrhoea. (19) Several alkaloidal and non-alkaloidal constituents have been reported from time to time such as piperine and pipericine (5).

Piperine is the major plant alkaloid present in black pepper *Piper nigrum* and long pepper *Piper longum*, is reported to have bioavailability enhancing activity for some drugs (3, 16). Piperine has good anticonvulsant and antimicrobial properties (13), it also acts as an antioxidant and anticancer agent by its numerous macromolecules associated with them (9).

The purpose of the present study was to determine the antimicrobial activity of piperine that isolated from *Piper nigrum* by taking a selected standard antibiotics.

## Materials and Methods :

### Extraction and identification of the piperine .

Fresh peppercorns ( 25 g ) were ground to affine powder , placed in asoxhlet thimble , and extracted with ethanol ( 250 ml ) for at least 90 minutes , At the end of the extraction the ethanol should be pale Yellow to colourless , the resulting solution was cooled , filtered and concentrated . The residue was dissolved in 10 % Alcoholic potassium hydroxide ( 25 ml ) . The solution was decanted if any residue remained , water drop wise was added

( about 30 ml will be required ) to precipitate the piperine. The piperine was collected and dried . Recrystallization from acetone : hexane ( 3:2 ) produced cleaner material . Structural determination is based on the infrared spectra ( IR spectrum ) and melting point (8) .The structure of Piperine was compared with IR standardize .

### Antimicrobial Screening

As test microorganisms , Gram ( + ) bacteria *Staphylococcus aureus* , Gram ( - ) bacteria *Escherichia coli* , *Proteus vulgaris* , *Klebsiella pneumoniae*, *Pseudomonas aeruginosa* and the yeast *Candida albicans* .Clinical isolates of *Proteus vulgaris* , *Klebsiella pneumoniae* were obtained from MSc.Hussein, K. Abdul-sada , while *Candida albicans* was obtained from Dr. Kawther, T. Khalaf , and *Staphylococcus aureus*, *Escherichia coli*, and *Pseudomonas aeruginosa* from MSc. Eiman, A. Saeed . pharmacology Dept. Pharmacy college .Basrah-university.

Antimicrobial activity was studied by the standard disc diffusion method (4) . Muller hinton medium was used for determining antibacterial activity whereas Potato Dextrose Agar Medium was selected for antifungal activity Standard antibiotic disc of streptomycin ( 10 µg / disc ) and erythromycin

( 15 µg / disc ) were also used for comparison in antibacterial testing respectively only .The piperine 100 mg was dissolved 10 ml of methanol (90%) the control experiment was setup with 2ml of ethanol (90%) which served as control.

The antimicrobial activities were determined by measuring the diameter of the inhibitory zones in mm . The diameter of the growth inhibition zones produced by the piperine on bacteria only were compared with diameter of the zone of the inhibition produced by the standard antibiotic disc .

Minimum inhibitory concentration (MIC) was determined by using the serial dilution method described by ( 15 ) , using the following concentration 400 , 200 , 100 , 50 , 25, 12.5 , 6.25 , 3.125 mg/ml . MIC values were determined as the lowest concentration of the constituent which completely inhibited the growth of the tested microorganisms ( 12 ) .

## Results and Discussion

piperine was obtained as purified crystals as long yellow needles with weight of 1.5 gm, fig 1, having melting point 123. IR spectrum showed characteristic bands for C-H ( aromatic and aliphatic ) ( 2800 , 3000 cm ) , N-C = O ( amide ) ( 1680 cm ) , C = C ( 1600 cm ) C-N ( 1250 cm ) C-O ( 1080 cm ) and C-H ( aromatic ) ( 1000 cm ) , the values of IR to isolated piperine in combination with standardized piperine IR fig ( 2,3 ) and these results in agreement with ( 7 ) .

In the present study piperine exhibited maximum effect against *C. albicans* and *E. coli*. As the diameter of inhibitory zones were 23 , 20 mm respectively and found to be most active antibacterial agent against all bacterial

isolates . Table -1 and fig 4 , 11 , the piperine has higher antimicrobial activity against *E. coli* and *P. vulgaris* in compare to erythromycin and streptomycin table - 1, fig. 4 , 5 while less activity in comparison to streptomycin and erythromycin against *S. aureus* and *P.aeruginosa* . Table -1, fig 6 , 7 , 8 , 9 and piperine showed higher antimicrobial activity against *K. pneumoniae* in compare to Erythromycin but this activity was less than Streptomycin fig.10.

The result of the minimum inhibitory concentrations is shown in table- 2 , all the studied bacteria and *C.albicans* were sensitive to piperine . MIC of piperine was between 3.125 - 100 mg / ml . The most

sensitive microorganism was *C. albicans* with MIC 3.125. Piperine activity may be comes from its ability to interfere with DNA and protein synthesis ( 9 ).

Piperine also act as an inhibitor of Nor A efflux pumps which ac as transport proteins involved in the release of toxic substrate from inside the cells to the external enviroment . These proteins are found in both Gram positive and negative bacteria as well as in eukaryotic organism ( 18 ) .

The higher sensitivity of *C. albicans* to piperine comes from that the piperine, inhibited the lipid accumulation in yeast. ( 6 ) ,the inhibitory effects were quantitatively represented by the total lipid accumulation

amount, triacylglycerol accumulation amount ,decrease in lipid amount and decrease in lipid body size with decrease in cell growth. Inhibitory effect of these compounds lead to decrease in triacylglycerol accumulation without any additional accumulation, of its intermediates suggesting that they will suppress the total carbon inflow into the triacylglycerol biosynthesis.(10).

The sensitivity of *E.coli* to piperine may by resulting from the *E. coil* have ahighly permeable membrane in compaire to other gram negetiv bacteria , piperine alteration in the permeability of the cell wall which contain high level of lipid material (14).

**Table ( 1 ) : Antimicrobial activities of Piperine**

| Test organisms                | piperine<br>100 mg / disc | Streptomycin<br>10 µg / disc | Erythromycin<br>15µ g / disc | Control<br>Ethanol ( 90%) |
|-------------------------------|---------------------------|------------------------------|------------------------------|---------------------------|
| <i>Escherichia coli</i>       | 20                        | 13.5                         | 15                           | -                         |
| <i>Staphylococcus aureus</i>  | 12                        | 23                           | 24                           | -                         |
| <i>Klebsilla pneumoniae</i>   | 15                        | 20                           | 10                           | -                         |
| <i>Proteus vulgaris</i>       | 17                        | 10                           | 6                            | -                         |
| <i>Pseudomonas aeruginosa</i> | 8                         | 22                           | 18                           | -                         |
| <i>Candida albicans</i>       | 23                        | No test                      | No test                      | -                         |


( - ) = not sensitive

**Table( 2): MIC values of the piperine against different target organism.**


| Target organism               | MIC ( mg / ml ) |
|-------------------------------|-----------------|
| <i>Escherichia coli</i>       | 6.25            |
| <i>Staphylococcus aureus</i>  | 50              |
| <i>Proteus vulgaris</i>       | 12.5            |
| <i>Pseudomonas aeruginosa</i> | 100             |
| <i>Klebsilla pneumoniae</i>   | 25              |
| <i>Candida albicans</i>       | 3.125           |




**Figure ( 1 ) : Needle crystal of piperine 16 x**




**Figure(2) : Infra-red spectra of purified piperine standard .**




**Figure(3) : Infra-red spectra of purified piperine**



**Figure ( 4 ) : Activity of piperine , Erythromycine and Streptomycin on *E. coli***



**Figure ( 5 ) : Activity of Piperine , Erythromycine and Streptomycin on *P. vulgaris***



**Figure ( 6 ) : Activity of Erythromycine and Streptomycine on *S. aureus***



**Figure ( 7 ) : Activity of piperine on *S. aureus***




Figure ( 8 ) : Activity of Erythromycine on *K. pneumoniae* and *P. aeruginosa*




Figure ( 9 ) : Activity of piperine on *K. pneumoniae* and *P. aeruginosa*




Figure (10) : Activity of Streptomycin on *K.pneumoniae* and *P.aeruginosa*



Figure (11) : Activity of piperine on *C. albicans*

### Conclusions :

Piperine aclass of compounds derived from a natural plant which is have potent activity against Gram+ve , Gram-ve

bacteria and yeast *Candida albicans* and its may be signifies as a source of therapeutic agents .

### References

1- Ali , M .A , Noor , M .A , Sarmina , Y. , Astag , M . K . and Abu Sayeed , M .( 2007 ) . Antimicrobial Screening of different extracts of *piper longum*

Linn . Re search J : Agriculture and Biological Sciences , 13 (6): 852-857.  
2- Al - Turki , A . I . ( 2007 ) . Antibacterial effect of thyme peppermint, Sage , black pepper and garlic hydrosols

- against *Bacillus subtilis* and *Salmonella enteritidis*. J. of food , Agriculture and Enviroment , 5 ( 2 ) : 93.
- 3- Badmaev, V., Majeed, M. And Norkus, E.(1999). Piperine an alkaloid derived from black pepper, increases serum response of beta - carotene during 14 days of oral beta-carotene supplementation. Nutr.Res. 19:381-388.
- 4- Barry , A .L . ( 1980 ) . Procedures for testing antimicrobial agents in agar media . In antibiotic in laboartory medicine , Eds ., V. Lorian.Williams and wilkins Co., Baltemore ,UsA ., PP : 1-23 .
- 5- Bina , S. , Sabria , B . , Tahsin , G . F . and Fatima , N . (1997 ) . AN Amide from fruits of *piper nigumr* phytochemistry 45 (8):1617-1619.
- 6- Daisuke , M. , Satoshi , O. , Taichi , O. ,Wei , J. and Hiroshi ,T. (2008). Molecular Target of Piperine in the Inhibition of Lipid Droplet Accumulation in Macrophage. Biol .Pharm. Bull. 31 : 1063- 1066.
- 7- Das, B. , Juthika, K. , Sitesh, C. , Mohammad , A. and Joydeb , K. (2007). Antitumor and Antibacterial activity of Ethyl acetate extract of *Ludwi -gita* Hyssopifolia Linn and Its Active Principle Piperine pak.J.Pharm.Sci., 20(2):128-131.
- 8- [http // eb . uvic ca / pmarrs / chem.465 /465 e03 Piperine . Pdf](http://eb.uvic.ca/~pmarrs/chem.465/465e03/Piperine.Pdf)
- 9- Inshad , A.K . , Zahid , M .M . , Ashwani , K . , Vijeshwar , V. and Ghulam , N.O . (2006) . Piperine , aphytochemical potentiator of ciprofloxacin aganist *Staphylococcus aureus* . Antimicrobial Agents and chemothera . 50 (2) :810-812.
- 10- Kazuyoshi, K. , Masakazu , Y. and Yasushi , K . (2006) . Inhibition of Lipid Accumulation and Lipid Body formation in Oleaginous Yeast by Effective Components in Spices , Carvacrol , Eugenol , Thymol , and Piperine . J. Agric . Food Chem . , 54 (10) : 3528-3534 .
- 11- Kirtikar , K . R . and Basu , B. D. (1987 ) . Indian Medicinal Plants , International book distributors, Dehradun , India, (3) :2128- 2129.
- 12- Mazzanti,G. , Mascellino,N.T., Pattinelli,L. ,Coluccia,D. , Manganario , M. and Saso, L. ( 2000 ) Antimicrobial investigaton of semipurified fractions of *Ginkgo biloba* Leaves.J.Ethnopharmacol.,71: 83-88
- 13- Nwinyi , O.C. , Chinedu , N . S . , Ajani, O.O . and Ogunniran , K.O. ( 2009 ) Antibacterial effects of extracts of *Ocimum gratissimum* and *piper guineense* on *Escherichia Coli* and *staphylococcus aureus* . African Journal of food sciene Vol 3 (3) , pp. 077- 081.
- 14- Obafemi, C.A , Sula imon , T.A. , Aki , D.A. and Olug bade , T.A. ( 2006 ) Antimicrobial activity of extracts and agermacranolide . type of sesquiterpene lactone from *Tithonia diversifolia*\_leaf extract. Afr. J. Biotechnol.( 12: 1254-1258 .
- 15- Pepeljnjak, S. , Kosalec, Z. , Kalodera and Kustrak, D. (2003) . Natural Antimycotics from Croation plants , in plant-derived Antimycotics Eds.M.Rai and Haworth press , Binghampton. :49-81.
- 16- Selvcndiran , K. , Princerijeya , J. and Sakthisekaran , D. (2009). In vivo effect of Piperine on serum and tissue glycolprotein levels in benzo (a)pyrene induced lung carcinogenesis in swiss albino mice . Pulmonary Pharmacology & Therapentic , 19 (2) : 107- 111.
- 17- Shelef , L. A. , Naglik , O.A. and Bogen , D.W. ( 1980 ). Sensitivity of some common food - borne bacteria to the spices Sage , rosemary and allspice . J . food .Sci. Vol.45 (4) : pp. 1044-1045.
- 18- Webber , M.A. And Piddock , L.J.V. ( 2003 ).The importance of efflux pumps in bacterial antibiotic resistance , J. of Antimicrobial chemotherapy . 51: 9-11 .
- 19- Young - cheol ,y. L. , Sang , G. , Hee , L. , Kwon , M. , Moo-Key , L., Sang - Hyun and Hoi - seon , L. (2002 ) Apiperidine Amide Extracted from *piper longam* . Fruit Shows Activity against *Aedes aegypti* Mosquito Larvae. J . of Agriculture and food Chemistry . ( 50 ) :3765 -3767 .

## الفعالية ضد ميكروبية للبايبرين المنقى من الفلفل الأسود *piper nigrum*

زينب طعمة خلف الدلي

كلية الصيدلة ، قسم الصيدلانيات والصيدلة السريرية ، جامعة البصرة

### المستخلص

أستخلص المركب الألكلويدي piperine من البذور الجافة لنبات *Piper nigrum* بالميثانول باستخدام جهاز السكوليت وتم عزله وتنقيته باعادة البلورة . حدد التركيب الكيميائي للبايبرين بواسطة طيف الـ IR . درست الفعالية ضد ميكروبية للبايبرين تجاه الجراثيم السالبة والموجبة لصبغة كرام والخمائر *Candida albicans* وقد وجد بانه ذو فعالية تثبيطية تجاه كل العزلات خصوصا ضد *C. albicans* تليها جرثومه *E. coli* و اقل فعالية تجاه جرثومه *P. aeruginosa* إذ تراوحت أقطار التثبيط بين 8 - 23 ملليمتر و التركيز المثبط الأدنى بين 100 - 3.125 مليغرام / مل .